



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



## A propos de ce livre

Ceci est une copie numérique d'un ouvrage conservé depuis des générations dans les rayonnages d'une bibliothèque avant d'être numérisé avec précaution par Google dans le cadre d'un projet visant à permettre aux internautes de découvrir l'ensemble du patrimoine littéraire mondial en ligne.

Ce livre étant relativement ancien, il n'est plus protégé par la loi sur les droits d'auteur et appartient à présent au domaine public. L'expression "appartenir au domaine public" signifie que le livre en question n'a jamais été soumis aux droits d'auteur ou que ses droits légaux sont arrivés à expiration. Les conditions requises pour qu'un livre tombe dans le domaine public peuvent varier d'un pays à l'autre. Les livres libres de droit sont autant de liens avec le passé. Ils sont les témoins de la richesse de notre histoire, de notre patrimoine culturel et de la connaissance humaine et sont trop souvent difficilement accessibles au public.

Les notes de bas de page et autres annotations en marge du texte présentes dans le volume original sont reprises dans ce fichier, comme un souvenir du long chemin parcouru par l'ouvrage depuis la maison d'édition en passant par la bibliothèque pour finalement se retrouver entre vos mains.

## Consignes d'utilisation


Google est fier de travailler en partenariat avec des bibliothèques à la numérisation des ouvrages appartenant au domaine public et de les rendre ainsi accessibles à tous. Ces livres sont en effet la propriété de tous et de toutes et nous sommes tout simplement les gardiens de ce patrimoine. Il s'agit toutefois d'un projet coûteux. Par conséquent et en vue de poursuivre la diffusion de ces ressources inépuisables, nous avons pris les dispositions nécessaires afin de prévenir les éventuels abus auxquels pourraient se livrer des sites marchands tiers, notamment en instaurant des contraintes techniques relatives aux requêtes automatisées.

Nous vous demandons également de:

- + *Ne pas utiliser les fichiers à des fins commerciales* Nous avons conçu le programme Google Recherche de Livres à l'usage des particuliers. Nous vous demandons donc d'utiliser uniquement ces fichiers à des fins personnelles. Ils ne sauraient en effet être employés dans un quelconque but commercial.
- + *Ne pas procéder à des requêtes automatisées* N'envoyez aucune requête automatisée quelle qu'elle soit au système Google. Si vous effectuez des recherches concernant les logiciels de traduction, la reconnaissance optique de caractères ou tout autre domaine nécessitant de disposer d'importantes quantités de texte, n'hésitez pas à nous contacter. Nous encourageons pour la réalisation de ce type de travaux l'utilisation des ouvrages et documents appartenant au domaine public et serions heureux de vous être utile.
- + *Ne pas supprimer l'attribution* Le filigrane Google contenu dans chaque fichier est indispensable pour informer les internautes de notre projet et leur permettre d'accéder à davantage de documents par l'intermédiaire du Programme Google Recherche de Livres. Ne le supprimez en aucun cas.
- + *Rester dans la légalité* Quelle que soit l'utilisation que vous comptez faire des fichiers, n'oubliez pas qu'il est de votre responsabilité de veiller à respecter la loi. Si un ouvrage appartient au domaine public américain, n'en déduisez pas pour autant qu'il en va de même dans les autres pays. La durée légale des droits d'auteur d'un livre varie d'un pays à l'autre. Nous ne sommes donc pas en mesure de répertorier les ouvrages dont l'utilisation est autorisée et ceux dont elle ne l'est pas. Ne croyez pas que le simple fait d'afficher un livre sur Google Recherche de Livres signifie que celui-ci peut être utilisé de quelque façon que ce soit dans le monde entier. La condamnation à laquelle vous vous exposeriez en cas de violation des droits d'auteur peut être sévère.


## À propos du service Google Recherche de Livres

En favorisant la recherche et l'accès à un nombre croissant de livres disponibles dans de nombreuses langues, dont le français, Google souhaite contribuer à promouvoir la diversité culturelle grâce à Google Recherche de Livres. En effet, le Programme Google Recherche de Livres permet aux internautes de découvrir le patrimoine littéraire mondial, tout en aidant les auteurs et les éditeurs à élargir leur public. Vous pouvez effectuer des recherches en ligne dans le texte intégral de cet ouvrage à l'adresse <http://books.google.com>






600030954R



E. BIBL. RADCL.


19151

e.  $\frac{7}{1}$









# **BOTANICON GALLICUM.**

**PARS I.**

*On vend dans la même Librairie :*

**LA FLORE FRANÇAISE**, ou Description de toutes les Plantes qui croissent naturellement en France, disposée selon une nouvelle méthode d'analyse, et précédée d'un exposé des principes de la Botanique; par MM. *de Lamarck*, membre de l'Institut, professeur au Muséum d'Histoire naturelle, et *De Candolle*, professeur de Botanique à Genève; troisième édition, 6 gros vol. grand in-8., fig. et cartes, br. 48 fr.

Cet ouvrage contient la description de six mille espèces; il est accompagné d'une belle Carte botanique sur papier grand-aigle, laquelle indique, par des couleurs diverses, les cantons où croissent les plantes particulières aux différentes contrées de la France, et il est orné de 11 grandes planches, sur lesquelles sont représentées plus de 200 figures servant à l'intelligence de l'étude des végétaux.

---

DE L'IMPRIMERIE DE CRAPELET,  
RUE DE VAUGIRARD, N° 9.

AUG. PYRAMI DE CANDOLLE

**BOTANICON GALLICUM**

SEU

**SYNOPSIS**

**PLANTARUM IN FLORA GALLICA  
DESCRIPTARUM.**

*EDITIO SECUNDA.*

---

EX HERBARIIS ET SCHEDIS CANDOLLIANIS PROPRIISQUE DIGESTUM

A

**J. E. DUBY** V. D. M.

---

**PARS PRIMA**

Plantas Vasculares continens.


---

**PARIS.**

V. DESRAY, RUE HAUTEFEUILLE, N° 4.

~~~~~  
1828.





---

## LECTORI.

---

QUUM mihi summus Candollius, pro suâ erga me amicitia et indulgentiâ, curam concesserit secundam suæ Synopsis plantarum in Florâ Gallicâ descriptorum editionem elaborandi, et mihi cum maximâ benevolentia herbarium et ineditas notas communicaverit, suisque optimis monitis adjuverit, omni ope atque operâ enixus sum ut tantis adminiculis me ostenderem dignum et ut quantum in me esset vacua quæ ex temporis lapsu et Scientiæ progressibus in primâ editione extabant, complerem, Gallicæque Botanices studium facilius redderem. Idcirco, paucis mutatis, ex tribus primis Prodromi systematis naturalis Regni Vegetabilis voluminibus (quorum tertium mihi tantum in manuscripto et labore meo jam pro parte perfecto innotuit), omnia quæ de plantis Gallicis dicta sunt excerpti. Quoad reliquas familias quas Magistri manus nondum pertractaverat, enixe conatus sum ne nimia cum prioribus inesset differentia, et quas, opus meum persequendo, in schedis viri summi notas aut observationes reperii, has sedulo adhibui et designavi signo *DC. mss.* aut *DC. herb.*

Non tantum ex herbariis et schedis Candollianis, undè hocce *Botanicon Gallicum* ederem, desumpsi, sed etiam ex propriis schedis et herbariis nec non ex observationibus quas suppeditarunt mea per Galliam itinera et meum cum multis doctis Botanicis commercium. Hâc occasione datâ liceat animo grato profiteri quantum debeam viris clarissimis *Vaucher, Roeper, Choisy, Seringe, Des Moulins, Prost, Le Prévost, Delise, de Brébisson, Soleirol, Chauvin, Grateloup, Desmazières, Lenormand, Solier, Graves, Monnier, Stire, Despréaux, d'Orbigny, E. Mercier*, etc., etc., qui me aut laboribus ineditis, aut observationibus, aut plantis rarissimis, pro singulari benevolentia adjuvare non dedignati sunt.

Ut pateat quænam fiducia tribuenda sit designatæ specierum stationi, sedulò signum *v. v.* plantis addidi quas ego et celeberrimus Floræ Gallicæ auctor vivas in loco indicato vidimus. Quas solus in loco natali vidit Candollius signo *v. v. DC.*, quas solus ipse signo *v. v. D.*, quas vero nec ille nec ego vivas in provinciis in quibus collectæ fuerunt sed tamen ex his provinciis siccas vidimus, signo *v. s.*, quas denique in provinciâ quâdam viventes testantur auctores, sed quarum specimina in Galliâ collecta non vidi, signo *v. s. ex Hispaniâ, v. s. ex Germaniâ* etc., *v. c.* (vidi cultam), *v. s. c.* (vidi siccam cultam), indicavi. Quum Botanici cujusdam nomen curvæ lineolæ complectuntur, tum specimina ex hoc Botanico proficiscentia vidi; quum vero hoc nomen particula *ex* prægreditur, tum specimina ex loco indicato orta non vidi, ita ut si quis error irrepserit, non mihi sed Botanico laudato tribuendus sit.

Hocce primum volumén tres plantarum Vascularium classes continens, quam citissimè; Deo volente, insequetur volumen secundum, jam prelo subjectum et quod quartam classem seu Cellulares, supplementa varia, clavim analyticam et indicem amborum voluminum continebit.

Die primâ februarii 1828.

J. DUBY v. d. M.

# INDEX.

|                                               |        |
|-----------------------------------------------|--------|
| I. Plantæ VASCULARES seu COTYLEDONEÆ.....     | Pag. 1 |
| Classis prima. DICOTYLEDONEÆ seu EXOGENÆ..... | ibid.  |
| Subclassis I. TRILAMIFLORE.....               | ibid.  |
| Ordo I. Ranunculaceæ.....                     | 2      |
| Trib. I. <i>Clematidea</i> .....              | ibid.  |
| II. <i>Anemoneæ</i> .....                     | 3      |
| III. <i>Ranunculeæ</i> .....                  | 7      |
| IV. <i>Helleboreæ</i> .....                   | 13     |
| V. <i>Pæoniaceæ</i> .....                     | 18     |
| Ordo II. Berberiden.....                      | 19     |
| III. <i>Nymphaeaceæ</i> .....                 | 20     |
| IV. <i>Papaveraceæ</i> .....                  | 21     |
| V. <i>Fumariaceæ</i> .....                    | 24     |
| VI. <i>Cruciferaæ</i> .....                   | 26     |
| Subordo I. <i>Pleurorrhizææ</i> .....         | ibid.  |
| Trib. I. <i>Arabideæ</i> .....                | ibid.  |
| II. <i>Alysiaceæ</i> .....                    | 33     |
| III. <i>Thlaspidææ</i> .....                  | 38     |
| IV. <i>Cakiliaceæ</i> .....                   | 42     |
| Subordo II. <i>Notorhizææ</i> .....           | ibid.  |
| Trib. V. <i>Sicymbreææ</i> .....              | ibid.  |
| VI. <i>Camelinceæ</i> .....                   | 46     |
| VII. <i>Lepidineææ</i> .....                  | 47     |
| VIII. <i>Isatideææ</i> .....                  | 49     |
| Subordo III. <i>Orthoplocææ</i> .....         | 50     |
| Trib. IX. <i>Brassicææ</i> .....              | ibid.  |
| X. <i>Zillææ</i> .....                        | 53     |
| XI. <i>Raphanææ</i> .....                     | 54     |
| Subordo IV. <i>Spirolobææ</i> .....           | 55     |
| Trib. XII. <i>Buniadææ</i> .....              | ibid.  |
| Subordo V. <i>Diplecobolææ</i> .....          | ibid.  |
| Trib. XIII. <i>Subulariææ</i> .....           | ibid.  |
| Ordo VII. <i>Capparidenææ</i> .....           | 56     |
| VIII. <i>Cistineææ</i> .....                  | 57     |
| IX. <i>Violariææ</i> .....                    | 63     |
| X. <i>Rosodaceææ</i> .....                    | 66     |
| XI. <i>Droseraceææ</i> .....                  | 68     |
| XII. <i>Polygalæææ</i> .....                  | 69     |
| XIII. <i>Frankeniaceææ</i> .....              | 70     |
| XIV. <i>Cariophyllæææ</i> .....               | 71     |
| Trib. I. <i>Sileneææ</i> .....                | ibid.  |
| II. <i>Alinceææ</i> .....                     | 80     |
| Ordo XV. <i>Lineæææ</i> .....                 | 88     |
| XVI. <i>Malvaceæææ</i> .....                  | 90     |
| XVII. <i>Tiliaceæææ</i> .....                 | 94     |
| XVIII. <i>Aurantiaçæææ</i> .....              | ibid.  |
| XIX. <i>Hypericineæææ</i> .....               | 96     |
| XX. <i>Acerineæææ</i> .....                   | 98     |
| XXI. <i>Hippocastaneæææ</i> .....             | 99     |
| XXII. <i>Meliaceæææ</i> .....                 | 100    |
| XXIII. <i>Ampelidæææ</i> .....                | ibid.  |
| XXIV. <i>Geraniaceæææ</i> .....               | 101    |
| XXV. <i>Tropæoleæææ</i> .....                 | 105    |
| XXVI. <i>Balsamineæææ</i> .....               | ibid.  |
| XXVII. <i>Oxalidææææ</i> .....                | 106    |

|                                 |          |
|---------------------------------|----------|
| Ordo XXVIII. Zygophylleæ.....   | Pag. 107 |
| XXIX. Rutaceæ.....              | 108      |
| XXIX bis. Coriariæ.....         | xij      |
| Subclassis II. CALYCULOSES..... | 109      |
| Ordo XXX. Celastrineæ.....      | ibid.    |
| Trib. I. Staphylaceæ.....       | ibid.    |
| II. Evonymæ.....                | ibid.    |
| III. Aquifoliaceæ.....          | 110      |
| Ordo XXXI. Rhamneæ.....         | ibid.    |
| XXXII. Terebinthaceæ.....       | 112      |
| Trib. I. Anacardiæ.....         | ibid.    |
| II. Sumachinæ.....              | 113      |
| III. Pteleaceæ.....             | ibid.    |
| Ordo XXXIII. Leguminosæ.....    | 114      |
| Subordo I. Papilionaceæ.....    | ibid.    |
| Trib. I. Sophoreæ.....          | 115      |
| II. Lotææ.....                  | ibid.    |
| III. Hedyarææ.....              | 144      |
| IV. Ficiææ.....                 | 149      |
| V. Phaseolæ.....                | 159      |
| Subordo II. Cæsalpinieæ.....    | 161      |
| Trib. VI. Cassiææ.....          | ibid.    |
| Ordo XXXIV. Rosaceæ.....        | 162      |
| Trib. I. Amygdaleæ.....         | ibid.    |
| II. Spiræaceæ.....              | 164      |
| III. Dryadeæ.....               | 165      |
| IV. Sanguisorbeæ.....           | 173      |
| V. Roseæ.....                   | 175      |
| VI. Pomaceæ.....                | 179      |
| Ordo XXXV. Granatææ.....        | 183      |
| XXXVI. Myrtaceæ.....            | ibid.    |
| Trib. I. Myrteæ.....            | 184      |
| II. Philadelphææ.....           | ibid.    |
| Ordo XXXVII. Cucurbitaceæ.....  | 185      |
| XXXVIII. Onagrarææ.....         | 187      |
| Trib. I. Onagrææ.....           | ibid.    |
| II. Jussieææ.....               | 188      |
| III. Hydrocaryes.....           | 189      |
| Ordo XXXIX. Haloragææ.....      | 190      |
| Trib. I. Cercodianeæ.....       | ibid.    |
| II. Callitricheææ.....          | 191      |
| III. Hippurideææ.....           | ibid.    |
| Ordo XL. Ceratophylleæ.....     | 192      |
| XLI. Lythrarææ.....             | ibid.    |
| XLII. Tamariscinææ.....         | 194      |
| XLIII. Portulacææ.....          | 195      |
| XLIV. Paronychiææ.....          | 196      |
| Trib. I. Teiophieææ.....        | ibid.    |
| II. Illecebreææ.....            | 197      |
| III. Polycarpeææ.....           | 199      |
| IV. Scleranthæææ.....           | ibid.    |
| V. Minuartieææ.....             | ibid.    |
| Ordo XLV. Crassulacææ.....      | 200      |
| XLVI. Nopaleææ.....             | 205      |
| XLVII. Ficoidææ.....            | ibid.    |
| XLVIII. Grossulariææ.....       | 206      |
| XLIX. Saxifragææ.....           | 207      |
| L. Umbelliferææ.....            | 213      |
| Trib. I. Thapsieææ.....         | ibid.    |



## ix

|                                            |              |
|--------------------------------------------|--------------|
| Trib. II. <i>Daucineæ</i> .....            | Pag. 215     |
| III. <i>Carolinææ</i> .....                | 216          |
| IV. <i>Coriandreeæ</i> .....               | 217          |
| V. <i>Silerineæ</i> .....                  | 218          |
| VI. <i>Tordylinesæ</i> .....               | <i>ibid.</i> |
| VII. <i>Pucedaneæ</i> .....                | 219          |
| VIII. <i>Angelicææ</i> .....               | 223          |
| IX. <i>Seselineæ</i> .....                 | 225          |
| X. <i>Scandiceneæ</i> .....                | 238          |
| XI. <i>Smyrneeæ</i> .....                  | 240          |
| XII. <i>Saniculeæ</i> .....                | 242          |
| XIII. <i>Hydrocotylinesæ</i> .....         | 243          |
| Ordo LI. <i>Caprifoliaceæ</i> .....        | 244          |
| Trib. I. <i>Hederaceæ</i> .....            | <i>ibid.</i> |
| II. <i>Sambucineæ</i> .....                | <i>ibid.</i> |
| III. <i>Caprifoliææ</i> .....              | 245          |
| Ordo LII. <i>Loranthææ</i> .....           | 246          |
| LIII. <i>Rubiaceæ</i> .....                | 247          |
| LIV. <i>Valerianeæ</i> .....               | 252          |
| LV. <i>Dipsaceæ</i> .....                  | 255          |
| LVI. <i>Compositæ</i> .....                | 258          |
| Trib. I. <i>Corymbifereæ</i> .....         | 259          |
| II. <i>Cynarocephalææ</i> .....            | 280          |
| III. <i>Cichoraceæ</i> .....               | 294          |
| Ordo LVII. <i>Lobeliaceæ</i> .....         | 310          |
| LVIII. <i>Campanulaceæ</i> .....           | 311          |
| LIX. <i>Vaccinieæ</i> .....                | 315          |
| LX. <i>Ericineæ</i> .....                  | 316          |
| Trib. I. <i>Ericaceæ</i> .....             | <i>ibid.</i> |
| II. <i>Rhodoraceæ</i> .....                | 318          |
| Ordo LXI. <i>Monotropesæ</i> .....         | 319          |
| Subclassis III. <i>COROLLIFLOREÆ</i> ..... | 320          |
| Ordo LXII. <i>Ebenaceæ</i> .....           | <i>ibid.</i> |
| Trib. I. <i>Diospyrææ</i> .....            | <i>ibid.</i> |
| II. <i>Styraceæ</i> .....                  | <i>ibid.</i> |
| Ordo LXIII. <i>Jasminææ</i> .....          | 321          |
| Trib. I. <i>Jasminææ</i> .....             | <i>ibid.</i> |
| II. <i>Lilaceæ</i> .....                   | 322          |
| Ordo LXIV. <i>Apocynææ</i> .....           | 323          |
| Trib. I. <i>Asclepiadeææ</i> .....         | <i>ibid.</i> |
| II. <i>Vincææ</i> .....                    | 324          |
| Ordo LXV. <i>Gentianeææ</i> .....          | 325          |
| LXVI. <i>Polemonideææ</i> .....            | 329          |
| LXVII. <i>Convolvulaceææ</i> .....         | <i>ibid.</i> |
| LXVIII. <i>Borragineææ</i> .....           | 331          |
| LXIX. <i>Solanæææ</i> .....                | 337          |
| LXX. <i>Antirrhineææ</i> .....             | 342          |
| LXXI. <i>Orobanchæææ</i> .....             | 348          |
| LXXII. <i>Rhinanthaceææ</i> .....          | 351          |
| Trib. I. <i>Pedicularineææ</i> .....       | <i>ibid.</i> |
| II. <i>Veroniceææ</i> .....                | 355          |
| Ordo LXXIII. <i>Labiataææ</i> .....        | 359          |
| LXXIV. <i>Verbenaceææ</i> .....            | 377          |
| LXXV. <i>Acanthaceææ</i> .....             | 378          |
| LXXVI. <i>Lentibularieææ</i> .....         | <i>ibid.</i> |
| LXXVII. <i>Primulaceææ</i> .....           | 379          |
| LXXVIII. <i>Globularieææ</i> .....         | 386          |
| Subclassis IV. <i>MONOCHAMYDEÆ</i> .....   | 387          |
| Ordo LXXIX. <i>Plumbagineææ</i> .....      | <i>ibid.</i> |

|                                                                   |          |
|-------------------------------------------------------------------|----------|
| Ordo LXXX. Plantaginæ.....                                        | Pag. 390 |
| LXXXI. Nyctaginæ.....                                             | 393      |
| LXXXII. Amaranthaceæ.....                                         | ibid.    |
| LXXXIII. Chenopodæ.....                                           | 394      |
| Trib. I. Atriplicæ.....                                           | ibid.    |
| II. Phytolacæ.....                                                | 399      |
| Ordo LXXXIV. Polygonæ.....                                        | 400      |
| LXXXIV bis. Thymelææ.....                                         | 405      |
| LXXXV. Laurinææ.....                                              | 407      |
| LXXXVI. Santalacææ.....                                           | 408      |
| LXXXVII. Elmagneæ.....                                            | 409      |
| LXXXVIII. Cytinææ.....                                            | 410      |
| LXXXIX. Aristolochiææ.....                                        | ibid.    |
| XC. Euphorbiacææ.....                                             | 411      |
| XCI. Urticææ.....                                                 | 417      |
| Trib. I. Urticææ.....                                             | ibid.    |
| II. Artocarpææ.....                                               | 419      |
| Ordo XCII. Juglandææ.....                                         | 420      |
| XCIII. Amentacææ.....                                             | ibid.    |
| Trib. I. Collidææ.....                                            | 421      |
| II. Botulinææ.....                                                | ibid.    |
| III. Salicinææ.....                                               | 423      |
| IV. Quercinææ.....                                                | 428      |
| V. Platanææ.....                                                  | 430      |
| VI. Myricææ.....                                                  | ibid.    |
| Ordo XCIV. Coniferææ.....                                         | 431      |
| Trib. I. Taxinææ.....                                             | ibid.    |
| II. Cupressinææ.....                                              | 432      |
| III. Abietinææ.....                                               | 433      |
| Classis secunda. MONOCOTYLEDONEAE sen ENDOGENAE PHANEROGAMAE..... | 435      |
| Ordo XCV. Hydrocharidææ.....                                      | ibid.    |
| XCVI. Alismacææ.....                                              | 437      |
| Trib. I. Butomææ.....                                             | ibid.    |
| II. Alismoidææ.....                                               | ibid.    |
| III. Juncaginææ.....                                              | 438      |
| Ordo XCVII. Potamææ.....                                          | 439      |
| XCVIII. Orchidææ.....                                             | 442      |
| CIX. Iridææ.....                                                  | 451      |
| C. Amaryllidææ.....                                               | 454      |
| CI. Asparagææ.....                                                | 458      |
| CII. Liliacææ.....                                                | 461      |
| Trib. I. Tulipacææ.....                                           | ibid.    |
| II. Asphodelææ.....                                               | 463      |
| III. Homero-callidææ.....                                         | 471      |
| Ordo CIII. Bromeliacææ.....                                       | 472      |
| CIV. Colchicacææ.....                                             | 473      |
| CV. Juncææ.....                                                   | 474      |
| CVI. Palmææ.....                                                  | 480      |
| CVII. Aroidææ.....                                                | ibid.    |
| Trib. I. Aroidææ.....                                             | ibid.    |
| II. Orontiacææ.....                                               | 481      |
| Ordo CVIII. Typhacææ.....                                         | 482      |
| CIX. Cyperacææ.....                                               | 483      |
| Ordo CX. Graminææ.....                                            | 499      |
| CXI ? Lemnæacææ.....                                              | 532      |
| Classis tertia. MONOCOTYLEDONEAE sen ENDOGENAE CRYPTOGRAMAE.....  | 533      |
| Ordo CXII. Characææ.....                                          | ibid.    |
| CXIII. Equisetacææ.....                                           | 534      |

# INDEX.

xj

|                                    |              |
|------------------------------------|--------------|
| Ordo CXIV. Filices.....            | Pag. 536     |
| Trib. I. <i>Ophioglossae</i> ..... | <i>ibid.</i> |
| II. <i>Osmundaceae</i> .....       | <i>ibid.</i> |
| III. <i>Polypodiaceae</i> .....    | 537          |
| Ordo CXV. Marattiaceae.....        | 542          |
| Trib. I. <i>Salviniae</i> .....    | <i>ibid.</i> |
| II. <i>Marattiae</i> .....         | <i>ibid.</i> |
| Ordo CXVI. Lycopodiaceae.....      | 543          |
| Trib. I. <i>Isoetes</i> .....      | <i>ibid.</i> |
| II. <i>Lycopodium</i> .....        | <i>ibid.</i> |

FINIS INDICIS.

*Ordo in loco suo post Rutaceas omissus.*

**Ordo XXIX bis. CORIARIEÆ DC. prod. 1. p. 739.**

Flores hermaphroditi aut sexûs unius abortu monoici dioïcive. Calyx (aut Perigonium) 1-sepalus campanulatus 10-fidus, lobis 5 exterioribus majoribus ovatis, interioribus 5 alternis minoribus callosis. Petala 0. Stamina 10 è toro exserta, 5 inter calycis lobos externos et ovarii angulos, 5 inter calycis lobos internos et ovarii sulcos. Antheræ oblongæ 2-loculares. Ovarium toro crassiusculo insidens 5-loculare 5-angulare. Stylus 0. Stigmata 5 ex apice longa subulata. Carpella 5 ad maturitatem discreta approximata indehiscencia 1-sperma glandulosis lobis amplius cincta. Semen in carpello pendulum. Albumen 0. Embryo rectus, radicula superâ, cotyledonibus 2 carnosus. — Frutices ramis subtetragonis oppositis. Folia opposita simplicia integra. Florum racemi terminales. — Ordini locus in serie naturali dubius.

I. CORIARIA *Niss. act. par. 1711. t. 12.*

Character isdem ac ordinis.

1. C. MYRTIFOLIA (L. sp. 1467.) foliis ovato-lanceolatis acutis triplinerviis subpetiolatis glabris, racemis erectiusculis. ♂ ad sepes in provinciis Mediterraneis. — Fl. fr. 4. p. 920. Lam. ill. t. 822. Duham. arb. 1. t. 73. (v. v.)

---

# BOTANICON

## GALLICUM.

---

### I. PLANTÆ VASCULARES seu COTYLEDONEÆ.

CAR. Plantæ contextu celluloso vasisque lymphaticis constantes; stomatibus donatæ, foliis veris instructæ. Flores sæpius distincti. Embryo intrâ spermodermium inclusus, cotyledonibus instructus.

#### Classis Prima. DICOTYLEDONEÆ seu EXOGENÆ.

CAR. TRUNCUS constans corporibus duobus, ligneo nempe et corticali, juxtâ leges inversas crescentibus. Corpus ligneum exhibens in centro medullam è quâ productiones medullares radiantes trans strata lignea annua, alia (lignum) interiora seniora duriora, alia (alburnum) exteriora juniora molliora albida. Cortex constans complexu celluloso, sub epidermide sito, et stratis corticalibus aliis exterioribus senioribus, aliis interioribus junioribus nomine Libri insignitis. Folia nervis ramosis anastomosantibus. Flores distincti, symmetrici, sæpius numero quinario dispositi. Embryo cotyledonibus, sæpius 2 oppositis, rariùs plurimis verticillatis, constans.

#### Subclassis I. THALAMIFLORÆ.

Calyx poly-sepalus. Petala plurima distincta, staminaque thalamo inserta nec calyci accreta.



Ord. I. RANUNCULACEÆ. *Juss.* 231. *DC. syst.* 1. p. 127.

Perigonium duplex, utrumque liberum hypogynum. Sepala 3-6. Petala sepalorum numero æqualia dupla triplave, libera, æstivatione imbricatâ, rariùs abortû nulla, nunc ex filamentis dilatatis facta et tunc plana, nunc ex antheris transformatis orta et tunc cucullata. Stamina hypogyna, libera, numero indefinita. Antheræ adnatæ. Pistilla  $\infty$  toro inserta, abortu aut coalitione rariùs solitaria. Carpella alia pseudosperma, alia baccata, alia capsularia aut follicularia, 1- $\infty$ -sperma. Semina nunc solitaria erecta aut pendula, nunc plurima secùs carpelli margines seriatiim disposita. Albumen corneum. Embryo minimus, intrâ albuminis foveolam locatus. — Herbæ aut suffrutices. Folia basi in vaginam semi-amplexicaulem sæpiùs dilatata simplicia, sæpè variè secta. Pili o aut simplices.

Trib. I. CLEMATIDÆ. *DC. syst.* 1. p. 131.

Calycis æstivatio valvata aut induplicata. Antheræ lineares extrorsæ. Carpella 1-sperma indehiscencia stylo accreto caudata. — Folia opposita; radices fibrosæ.

I. CLEMATIS. *DC. syst.* 1. p. 131.

Involucrum o aut calyciforme sub flore; sepala 4-8 colorata; petala o aut sepalis breviora; cariopsides  $\infty$  in caud am sæpiùs barbato-plumosam productæ. — Radices perennes.

Sect. I. FLAMMULA. *DC. l. c.*

Invol. o; pet. o; carpellorum caudæ longæ barbato-plumosæ.

1. C. ERRECTA (All. ped. n. 1078.) caule erecto, foliis pinnatisectis, segmentis petiolulatis ovato-lanceolatis integerrimis.  $\mathcal{U}$  in collibus Gallie Mediterraneæ. — Fl. fr. 4. p. 873. C. recta. L. sp. 767. — Lob. ic. 627. f. 2. (v. v.)
2. C. FLAMMULA (L. sp. 766.) caule scandente, foliis pinnatisectis, segmentis glabris integris trilobisve orbiculatis ovalibus oblongis sublinearibusve acutiusculis. in  $\mathfrak{f}$  sterilibus et sepibus Gallie Mediterraneæ. — Fl. fr. 4. p. 873. — Lob. ic. 627. f. 1.
  - a. vulgaris (DC.) fol. segmentis ovali-aut oblongo-lanceolatis. (v. v.)
  - $\beta$ . maritima (DC.) fol. segmentis linearibus. In maritimis prope Monspe-  
lium. C. maritima. L. sp. 767? fl. fr. 5. p. 562. (v. v.)
3. C. VITALBA (L. sp. 766.) caule scandente, foliis pinnatisectis, segmentis ovato-lanceolatis dentato-incisis acuminatis basi truncato-cordatis, pedunculis foliis subbrevioribus.  $\mathfrak{f}$  in sepibus. — Fl. fr. 4. p. 872. — Lob. ic. 626. f. 1. (v. v.)
  - $\beta$ . integrata (DC.) segmentis paucissimis dentatis integerrimisve. (v. v.)
4. C. INTEGRIFOLIA (L. sp. 767.) pedunculis unifloris, floribus nutantibus, foliis integris ovato-lanceolatis glabris.  $\mathcal{U}$  in Pyrenæis ex Lapeyr. — Fl. fr. 5. p. 532. — Lob. ic. 628. f. 1. (v. v. c.)

Sect. II. CHEIROPSIS. *DC. syst.* 1. p. 162.

Involucrum calyciforme è duabus bracteis coalitis sub flore ad apicem pedunculi situm; petala o; caudæ barbatae.

## RANUNCULACEÆ.

3

5. *C. cernua* (L. sp. 766.) foliis ovato-subcoarctatis dentatis fasciculatis, pedunculis 1-floris involucretis, flore minore pallido intra involucrem pedicellato.  $\zeta \cup$  ex Corsicâ acceptam misit Solier. — Bot. mag. t. 1070. *C. cirrhosa*  $\beta$  pedicellata DC. prod. 1. p. 9. (v. s.)
6. *C. BALEARICA* (Rich. in Journ. phys. 1779, fevr. 127 cum ic. non Pers.) pedunculis unifloris sub flore involucretis, foliis ternatum sectis, segmentis petiolatis 3-lobis dentato-incisis.  $\zeta \cup$  in Corsicâ reperiit cl. Soleirol. — *C. calycina* Schrank. mon. t. 15. Involucreum campanulatum subhilobum obtusum florem arcte cingens. (v. s.)

### Sect. III. ATRAGENE. DC. prod. 1. p. 9.

Involucreum 0; sepalâ 4; petalâ 00; caudæ barbatae.

7. *C. ALPINA* (Mill. diet. n. 9.) pedunculis unifloris folio longioribus, foliis biternatum sectis, segmentis ovato-lanceolatis acuminatis serratis, petalis subspathulatis obtusis.  $\zeta \cup$  in dametis asperis subsylvaticis Alpium rarior, et forsan Pyrenæorum. — Fl. fr. 4. p. 874. *Atragene Alpina* L. sp. 764. — J. Bauh. 2. p. 129. f. 2. Fl. cærulei rarò albi. (v. v.)

### Trib. II. ANEMONEÆ. DC. syst. 1. p. 168.

Calycis corollæque æstivatio imbricativa. Carpella 1-sperma indehiscentia in caudas mucronesve sæpiùs desinentia. — Folia radicalia aut alterna.

### II. THALICTRUM. L. gen. 697. DC. syn. 1. p. 168.

Involucreum sub flore 0; petalâ 0; carpella sicca in caudas non desinentia, nunc stipitata, nunc longitudinaliter striato-sulcata. — Herbæ  $\mathcal{L}$ .

#### Sect. I. TRIPTERIUM. DC. l. c.

Fructus triquetri, angulis alatis, substipitatus.

1. *T. AQUILEGIFOLIUM* (L. sp. 770.) stipellis ovatis binis ad basin petioli ramificationum, paniculâ corymbosa, fructibus triquetris.  $\mathcal{L}$  in subsylvaticis montosis. — Fl. fr. 4. p. 878. — Moris. s. 9. t. 20. f. 4. n. 16. (v. v.)

#### Sect. II. EUTHALICTRUM. DC. l. c. p. 172.

Fructus ovato-oblongus striatus sessilis.

2. *T. ALPINUM* (L. sp. 767.) caule simplicissimo subnudo digitali, racemo simplici terminali, floribus natis, foliorum segmentis glabris.  $\mathcal{L}$  in pascuis editioribus Pyrenæorum. — Fl. fr. 4. p. 874. — Moris, s. 9. t. 20. fig. ult. n. 14. (v. v.)
3. *T. FORTIDUM* (L. sp. 768?) caule simplici basi nudo medio folioso apice paniculato, foliis pubescenti-viscidis, segmentis dentibus obtusis.  $\mathcal{L}$  in Alpiis Galloprovinciæ vallibus. — Fl. fr. 4. p. 875. Lam. ill. t. 497. f. 3. — Moris. s. 9. t. 20. f. 10. n. 13. (v. s.)
4. *T. PUBESCENS* (Sahl. in fl. fr. 5. p. 633.) caule simplici foliis sparsis obsoito apice paniculato, foliis pubescenti-viscidulis.  $\mathcal{L}$  in lapidoso calcareis Occitaniz, Galloprov. et Delphin. inferioris. — Gon. hort. 263. Vill. 4. p. 714. (v. v. DC.)
5. *T. MINUS* (L. sp. 767.) caule tereti polline glauco, floribus laxè paniculatis cernuis 4-sepalis, foliorum segmentis glabris subrotundis apice dentatis subtus glaucis, carpellis utrinquè acutis.  $\mathcal{L}$  in pratis montosis subsylvaticis. — Fl. fr. 4. p. 875. Jacq. aust. t. 419. — Tabern. n. t. 55. f. 2. (v. v.)
6. *T. SAXATILE* (DC. Fl. fr. 5. p. 633.) caule tereti polline glauco destituto, floribus paniculatis erectis, fol. segmentis glabris subrotundis apice dentatis

## DICOTYL. SEU EXOGENÆ.

- subtus glaucis, carpellis utrinque acutis.  $\mathcal{U}$  in collibus sylvaticis Alsatiæ, Alpium, Pyrenæorum. — T. minus Poll. n. 522. (v. v.)
7. *T. MAJUS* (Murr. syst. 513.) caule tereti polline glauco destituto, floribus laxè paniculatis cernuis, foliorum segmentis glabris subtus glaucis trifidis, lobulis ovatis submucronatis, carpellis basi obliquè rotundatis.  $\mathcal{U}$  in collibus siecis Delphin. Cebennarum, Pyren. central. ex cl. Benth. — Fl. fr. 4. p. 876. Jacq. aust. t. 420. (v. v.)
  8. *T. GALIODES* (Nestl. in fl. fr. 4. p. 876, 5. p. 633.) caule tereti subsulcato stricto, radice repente, paniculâ strictâ, floribus nutantibus, foliorum segmentis linearibus angustissimis integerrimis margine revolutis, ultimo subsessili.  $\mathcal{U}$  in pratis sylvaticis arenosis Alsatiæ. — C. Bauh. prod. 146. ic. — Moris. s. 9. t. 20 f. 7. n. 8. (v. v. DC)
  9. *T. ANGUSTIFOLIUM* (Jacq. vind. 4. t. 43.) caule stricto tereti subsulcato, radice fibrosâ, paniculâ multiplici erectâ, floribus erectiusculis, foliorum segmentis lineari-lanceolatis integerrimis, ultimo subincurvato remoto.  $\mathcal{U}$  in pratis et sylvarum locis cædis in prov. australibus. — Fl. fr. 4. p. 876. — Pluck. t. 65. f. 3. (v. v.)
  10. *T. LUCIDUM* (L. sp. 770.) caule ramoso tereti subsulcato, radice fibrosâ, paniculâ multiplici erectâ, floribus erectiusculis, foliorum segmentis lineari-lanceolatis integris, ultimo lobato.  $\mathcal{U}$  circa Parisios. — Pluck. t. 65. f. 5. (v. s.)
  11. *T. NIGRICANS* (Jacq. aust. 5. t. 421.) caule stricto sulcato ramoso, radice fibrosâ, paniculâ multiplici erectâ subcorymbosâ, foliorum radicalium segmentis cuneiformibus trifidis caulinarum oblongo-linearibus.  $\mathcal{U}$  in arenosis ad ostia Rhodani, in Galloprov. australi, ad Pyrenæorum radices. — Fl. fr. 5. p. 634. Jacq. aust. 5. t. 421. Sequentis forsitan varietas. (v. v. DC.)
  12. *T. FLAVUM* (L. sp. 770.) caule erecto ramoso sulcato, radice fibrosâ, paniculâ multiplici erectâ subcorymbosâ, foliorum omnium segmentis cuneiformibus trifidis acutis.  $\mathcal{U}$  in pratis humidis. — Fl. fr. 4. n. 877. — Tab. ic. t. 55. f. 1. (v. s.)
  13. *T. SIMPLEX* (L. mant. 78.) caule erecto simplici angulato, radice repente, paniculâ erectâ racemosâ pauciflorâ, foliorum segmentis linearibus aut inferioribus oblongis acutis trifidis.  $\mathcal{U}$  in pratis subpaludosis Occitanie, Galloprov. Delphin. — Fl. fr. 4. p. 877. Fl. dan. t. 244. *T. angustifolium*. Vill. 3. p. 712. (v. v. D.)
  14. *T. GLAUCUM* (Desf. cat. ed. 2. p. 126.) caule erecto tereti striato polline glauco, paniculâ multiplici erectâ confertâ, foliorum segmentis subcordato-ovatis obtusè trifidis subtus glaucis.  $\mathcal{U}$  in Pyrenæis orientalibus ex Lapeyr. — Fl. fr. 4. p. 878. — Moris. s. 9. t. 12. f. 5. n. 1. *T. speciosum*. Poir. (v. v.)
  15. *T. TUBEROSUM* (L. sp. 768.) radice grumosâ, floribus laxè corymbosis solitariisve, involucri nullo, bracteis sessilibus, sepalis petaloideis stamina superantibus.  $\mathcal{U}$  in pascuis sterilibus et pratis collium,  $\alpha$  caule unifloro supra Carcassone,  $\beta$  caule multifloro, in Pyrenæis orientalibus. — Fl. fr. 4. p. 875. Mill. ic. 265. f. 2. — Moris. s. 4. t. 28. f. 4. n. 13. (v. v. DC.)

## III. ANEMONE. DC. syst. 1. p. 188.

Involucrum trifolium à flore distans foliolis incisis; calyx petaloideus, 5-15-sepalus; petala 0; carpella  $\infty$ . Herbæ  $\mathcal{U}$ , folia radicalia petiolata multi-partita aut lobata.

## Sect. I. PULSATILLA. Bauh. pin. 177. DC. l. c.

Cariopsides in caudas longas barbata productæ; involucri folia sessilia in lobos lineares palmati-partita.

1. *A. VERNALIS* (Lin. sp. 759.) foliis pinnatisectis, segmentis cuneato-lanceolatis trifidis glabriusculis, flore erecto, involucri villosissimo, sepalis 6 rectis elliptico-oblongis.  $\mathcal{U}$  in scopulis et pratis ad nivem æternam deliquescentem in Alpiis, Pyrenæis, Arvernâ. — Fl. fr. 4. p. 879. — Hall. helv. t. 21. (v. v.)

2. *A. HALLERI* (All. ped. n. 1922. t. 8o f. 2.) foliis pinnatisectis villosissimis, segmentis tripartitis, lobis lobatis, lobulis lanceolato-linearibus acuminatis, flore erecto, sepalis 6 ovali-lanceolatis.  $\frac{1}{2}$  ad nivem æternam deliquescentem in summis Alpibus Delph. — Fl. fr. 4. p. 879. (v. v.)
3. *A. PULSATILLA* (Lin. sp. 759.) foliis pinnatisectis, segmentis multipartitis, lobis linearibus, flore subnultante, sepalis 6 patentibus.  $\frac{1}{2}$  in campis sylvestribus collibusque apricis Gallie ferè totius. — Fl. fr. 4. p. 880. Bull. herb. 149.  
*ß. rubra* Lam. dict. 1. p. 163. In Arverniâ. (v. v.)
4. *A. PRATENSIS* (Lin. sp. 760.) foliis pinnatisectis, segmentis multipartitis, lobis linearibus, flore pendulo, sepalis 6 erectis apice reflexis.  $\frac{1}{2}$  in campis apricis pratensibus. — Fl. fr. 4. p. 880. — Dod. pempt. 433. f. 2. (v. v.)

Sect. II. *PRÆONANTHUS*. DC. l. c.

Cariopsides in caudas longas barbata productæ. — Involucri foliola petiolata trisecta.

5. *A. ALPINA* (Lin. sp. 760.) foliis petiolo ramoso biternatim sectis, segmentis pinnatim inciso-serratis, involucralibus conformibus, sepalis 6 patentibus.  $\frac{1}{2}$  in pascuis lapideis montanis. — Fl. fr. 4. p. 881. Crantz Austr. 2. t. 3. f. 2 (v. v.)  
*a. major* (DC.) floribus magnis albis extûs subpurpurascens, sepalis ellipticis, foliis glabris aut parè villosis ante florem expansis. — Dalech. Lugd. 850. f. 3. (v. v.)  
*ß. micrantha* (DC.) floribus parvulis albis extûs subpurpurascens, sepalis ellipticis, foliis glabriusculis ante florem expansis. — Dalech. 849. f. 2. (v. v.)  
*γ. flavescens* (DC.) floribus magnis flavis extûs pallidioribus, sepalis ellipticis, foliis parè villosis ante florem expansis. — Jacq. misc. 2. p. 47. f. 4. (v. v.)  
*δ. nivalis* (DC.) floribus mediocribus albis subtus subpurpurasc., sepalis ellipticis, foliis villosissimis cum flore aut post fl. expansis, scapo nudo. (v. v.)  
*ε. intermedia* (DC.) floribus magnis flavis extûs subpallidioribus, sepalis ellipticis, foliis villosissimis ante fl. aut cum fl. nascentibus. (v. v.)  
*ζ. sulphurea* (DC.) floribus mediocribus flavis extûs subpal., sepalis ovatis, foliis villosissimis ante fl. aut cum fl. expansis. — Cam. ep. 393. ic. — A. sulphurea. Lin. mant. 78. (v. v.)

Sect. III. *ANEMONANTHEA*. DC. l. c. p. 196.

Cariopsides ecaudatæ ovoidæ, pedicelli solitarii aut bini, omnes aphylli uniflori. — Involucri foliola sessilia aut petiolata.

6. *A. CORONARIA* (Lin. sp. 760.) foliis ternatim scotis, segmentis multifidis, lobulis linearibus mucronatis, involucralibus sessilibus multifidis, sepalis 6 ovalibus approximatis.  $\frac{1}{2}$  in pascuis subhumidis Gallie Mediterraneæ fl. primo vere. — Fl. fr. 4. p. 882. — Lob. ic. 177. (v. v. DC.)
7. *A. PAVONINA* (Lam. dict. 1. p. 166.) foliis tripartitis, lobis cuneatis inciso-dentatis, involucralibus sessilibus integris subincisive, sepalis 10-12 lanceolatis acutissimis.  $\frac{1}{2}$  in Galloprovincia, circa Olbia et Niceam. fl. vere. — Fl. fr. 5. p. 634. — Lob. ic. 278. (v. s.)  
*ß? fulgens* (DC. prod. 1. p. 18.) foliis tripartitis, lobis cuneatis inciso-dentatis, involucralibus sessilibus oblongis integris subincisive, sepalis oblanceolatis apice latioribus basi attenuatis.  $\frac{1}{2}$  in vineis Vasconie circa Dax. A. hortensis Thor.-chl. Land. 238. A. pavonina Lois. not. p. 87. Flores ampliores quam var. *a.* intensè rubri. (v. s.)
8. *A. STELLATA* (Lam. dict. 1. p. 166.) foliis tripartitis, lobis cuneatis inciso-dentatis, involucralibus sessilibus oblongis integris subincisive, sepalis 10-12 oblongis obtusiusculis.  $\frac{1}{2}$  in sepibus et dumetis Galloprov. austr. — A.

- hortensis*. Lin. sp. 761. Fl. fr. 4. p. 882. 5. p. 635. — Lob. ic. 279. fig. 1. (v. v.)
9. *A. PALMATA* (Lin. sp. 758.) foliis cordatis suborbiculatis obtusè 3-5 lobis dentatis, involucribus sessilibus trifidis, sepalis 10-12 oblongis obtusis.  $\mathcal{U}$  in Galloprov. prope Olbiam. fl. primo vere. — Fl. fr. 5. p. 635. — Lob. ic. 279. f. 2. (v. s.)
10. *A. APENNINA* (Lin. sp. 762.) foliis petiolo ramoso triternatim sectis, segmentis lanceolatis inciso-dentatis acutis, involucralibus petiolatis trisectis inciso-dentatis, sepalis 12-14 oblongis obtusis.  $\mathcal{U}$  in umbrosis submontania Galloprov? Corsicæ. Fl. cærul. aut albi. — Mentz. pug. t. 8. f. 2. Engl. Bot. 1. 106a. J. Bauh. hist. 3. p. 405. f. 2. (v. s.)
11. *A. BALDENSI* (Lin. mant. 78.) foliis petiolo ramoso biternatim sectis, segmentis multipartitis, lobis linearibus, involucralibus breviter petiolatis multifidis, sepalis 8-10 oblongo-subovalibus.  $\mathcal{U}$  in asperis editis Alpium. — Fl. fr. 4. p. 882. Vill. 4. t. 49. *A. fragifera* Jacq. iq. rar. 1. 103. — Sepala alba extus rubentia. (v. s.)
12. *A. NEMOROSA* (Lin. sp. 762.) foliis ternatim sectis, segmentis trifidis inciso-dentatis lanceolatis acutis, involucralibus petiolatis conformibus, sepalis 6 ellipticis.  $\mathcal{U}$  in nemoribus dumetis et collibus umbrosis. fl. albi, rarius rosei cæruleive. — Fl. fr. 4 p. 884. — Clus. hist. 247. fig. 1 et 2. (v. v.)
13. *A. TRIFOLIA* (Lin. sp. 762.) foliis omnibus petiolatis ternatim sectis, segmentis ovato-lanceolatis acutis dentatis, sepalis 5 ellipticis obtusis.  $\mathcal{U}$  in nemorosis submontanis Andegaviæ, Parisiorum circa Chantilly ex Thuil. — Fl. fr. 4. p. 884. — Lob. ic. 281. f. 1. fl. albi. (v. s.)
14. *A. RANUNCULOIDES* (Lin. sp. 762.) foliis radicalibus 3-5-sectis, segmentis subtrifidis inciso-dentatis, involucralibus subpetiolatis tripartitis inciso-dentatis, sepalis 5-6 ellipticis.  $\mathcal{U}$  in nemoribus et pratis montanis. — Fl. fr. 4. p. 885. — Fuchs. hist. 162. ic. fl. lutei. (v. v.)
15. *A. SYLVESTRIS* (Lin. sp. 761.) foliis ternatim quinatimve sectis, segmentis apice inciso-dentatis, involucralibus petiolatis conformibus, pedicello solitario, sepalis 6 ellipticis, fructibus hirsutissimis.  $\mathcal{U}$  in nemoribus et sepibus Alsatiæ, Lotharingiæ, Arvernæ, agri Laudunensis. — Fl. fr. 4. p. 883. — Lob. ic. 280. f. 2. fl. albi. (v. v.)

#### SECT. IV. OMALOCARPUS. DC. l. c. p. 212.

Cariopsides compresso-planæ ovali-orbiculatæ glabræ caudatæ; pedicelli cœ umbellati aphylli 1-flori.

16. *A. NARCISSEFLORA* (Lin. sp. 763.) foliis radicalibus subvillosis palmatim 3-5 partitis, lobis inciso-dentatis, lobulis linearibus acutis, involucralibus apice 3-5-fidis, floribus umbellatis.  $\mathcal{U}$  in pascuis montanis præsertim calcareis. — Fl. fr. 4. p. 885 et 5. p. 635. — J. Bauh. hist. 861. f. 1 et 2. (v. v.)
- fl. monantha* (DC.) floribus solitariis aut binis. — *A. dubia* Bell. app. t. 7. (v. s.)

#### IV. HEPATICA. Dill. giess. p. 108. t. 5. DC. fl. fr. 4. p. 885.

Involucrum calyciforme trifoliatum 1-florum flori approximatum; sepala petaloidea 6-9 duplici triplicive serie disposita; carpella cœcaudata. — Herbar  $\mathcal{U}$ , scapis plurimis radicalibus 1-floris.

1. *H. TRILoba* (DC. fl. l. c.) foliis cordatis trilobis, lobis integerrimis.  $\mathcal{U}$  in sepibus, dumetis, sylvis submontanis. — *Anemone hepatica* L. sp. 758. — Cam. epit. 585. ic. fl. cærulei rarius albi et rosei. (v. v.)

#### V. ADONIS. Dill. giess. 109. t. 4. DC. syst. 1. p. 220.

Calyx 5-sepalus adpressus; petala 6-15 ungue nudo; stamina cœ ad



basin gonophori inserta; cariopsides oo 1-sperma secus gonophorum  
spicatæ ovatæ stylo brevi acuminatæ. — Herbae caule folioso, foliis in  
lobos lineares multipartitis.

(1) Sect. I. ADONIA. DC. *syst.* 1. p. 321.

Cariopsides stylo recto uncinato acuminatæ, petala 5-10. Radices annuæ.

1. *A. AUTUMNALIS* (L. sp. 771.) calyce glabro, petalis 6-8 concavo-conviventibus calyce vix majoribus, carpellis subreticulatis in capitulum ovatum aggregatis stylo brevissimo coronatis, caule ramoso. (I) inter segetes spontanea, in hortis culta. — Eng. bot. t. 308. Weinm. phyt. t. 27. f. b. c. Fl. sanguinei. (v. v.)
2. *A. FLAVA* (Vill. cat. str. 247.) calyce glabro basi soluto, petalis planis oblongis calyce duplò longioribus, carpellis sublaevibus in capitulum oblongum aggregatis, caule subsimplici. (I) inter segetes et in vineis. — Weinm. phyt. t. 28. f. a. b. Fl. flavi. (v. v. DC.)
3. *A. MICRANTHA* (DC. *syst.* 1. p. 222.) calyce glabro basi non soluto, petalis planis oblongis calyce paulò longioribus, carpellis subreticulatis in capitulum ovatum aggregatis, caule subramoso. (I) inter segetes Gallie australis. — Fl. parvi flavi seu flammei. (v. v. DC.)
4. *A. CITRINA* (Hoffm. germ. p. 1. p. 251.) calyce basi hispido, petalis planis oblongis calyce longioribus, carpellis in capitulum ovato-oblongum dispositis, caule subsimplici, flore parvo inter folia subcaesili. (I) inter segetes. — Fl. citrini. An sequentis varietas? (v. v. DC.)
5. *A. ÆSTIVALIS* (L. sp. 772.) calyce basi hispido, petalis planis oblongis obtusis calyce duplò longioribus, carpellis reticulatis in spicam elongatam laxam dispositis conoideo-prismaticis stylo recto concolori rostratis, caule subsimplici. (I) inter segetes Gallie australis. — *A. miniata* Jacq. auct. t. 354. (v. v.)
6. *A. DENTATA* (Dehile ag. 17. descr. t. 53. f. 1.) calyce basi hispido, petalis planis ovali-oblongis calyce paulò longioribus, carpellis basi tuberculato-dentatis reticulatis in spicam elongatam continuam digestis. (I) in Galloprov. superiori inter Digne et Colmar. — Fl. flammei. (v. v. DC.)

Sect. II. CONSILIGO. DC. *syst.* 1. p. 224.

Cariopsides stylo uncinato recurvo acuminatæ; petala 8-15. Radices 2l.

7. *A. VERNALIS* (L. sp. 773.) foliis radicalibus infimæ abortivis ad squamas subvaginantes reductis, mediis et summis sessilibus, carpellis velutinis, petalis 10-12 oblongis subdenticulatis. 2l in collibus et vallibus Alpium, Cebennarum, montium Ligeris (cl. Prost.) — Fl. fr. 4. p. 387. — Moris. s. 6. t. g. f. 1 et 2. Fl. flavi. (v. v.)
8. *A. PYRÆNÆICA* (DC. fl. fr. 5. p. 635.) foliis radicalibus longè petiolatis, petiolo trifido, segmentis multipartitis, summis sessilibus, carpellis glabris, petalis 8-10 oblongo-cuneatis integris. 2l in valle Eynes Pyrenæorum orientaliū. — *A. Apennina* L. sp. 772. excl. syn. — Del. ic. sel. 1. t. 21. Fl. flavi. (v. s.)

Trib. III. RANUNCULÆ. DC. *syst.* p. 228.

Æstivatio calycis et corollæ imbricata. Petala bilabiata aut intus squamulâ ad basin aucta. Carpella 1-sperma sicca indehiscencia. Semen erectum. — Folia radicalia aut erecta.

VI. MYOSURUS. Dill. *giess.* p. 106. t. 4.

Calyx 5-sepalus, sepalis basi solutis seu ultrâ insertionem deorsum

(1) Stirpes sub hoc titulo collectæ nempe sunt forsân varietates notabiles unius speciei, *A. Annuæ*, Mill. dict. n. 1. Fl. fr. 4. p. 387.

longè productis; petala 5 ungue filiformi tubuloso; stamina 5-20; caripsides plurimæ triquetræ confertissimæ, receptaculo longissimo spicatum dispositæ, stylo recto subacuminatæ.

2. *M. minimus* (L. sp. 407.) scapis folio sublongioribus, calycis appendicibus subfoliaceis. ① in apricis pinguibus hyeme inundatis. — Fl. fr. 4. p. 906. — Dod. pempt. 112. f. 1. (v. v.)

VII. CERATOCEPHALUS. *Mærch. meth. p. 218. DC. syst. 1. p. 230.*

Calyx 5-sepalus subpersistens, sepalis basi non productis; petala unguiculata; stamina 5-15; carpella ∞ secus rachin persistentem longè spicata, basi utrinquè gibba, apice producta in cornu compressum semine 4-gono sextuplò longius; embryo rectus.

1. *C. falcatus* (Pers. ench. 1. p. 341.) pericarpiorum cornubus falcatis subascendentibus. ① inter segetes provinc. Mediterr. — Fl. fr. 4 p. 900. — Bocc. Sic. t. 14. f. 2. 3. (v. v.)

VIII. RANUNCULUS. *C. Bauh. pin. 180. DC. prod. 1. p. 46. Ser. mel. helv. 2. p. 45.*

Calyx 5-sepalus, sepalis basi non solutis deciduis; petala intus basi squamulâ foveolari nectariferâ instructa; stam. ∞; caripsides ovatæ subcompressæ, in mucronem aut cornu semine vix longius desinentes, in capitulum globosum cylindraceumve dispositæ.

Sect. 1. BATRACHIUM. *DC. syst. p. 1. 233.*

Pericarpia transversè rugoso-striata; petala alba ungue flavo foveâ nectariferâ notato.

1. *R. menziesii* (L. sp. 781.) caule repente, foliis subreniformibus sub-3-5 lobis, lobis latis integris obtusissimis, petalis oblongis calyce vix longioribus, staminibus 5-12, carpellis glabris transversè rugoso-striatis, stylo brevi crassiusculo subtruncato vix deflexo. ② ad scaturigines et fontes in aquis vadosis et locis inundatis provinc. septentr. et occident. — Fl. fr. 4. p. 894, Drev. et Hayn. Eur. t. 106. (v. v. DC.)
2. *R. tripartitus* (DC. ic. rar. 1. t. 49 et Fl. fr. 5. p. 637.) caule natante, foliis submersis capillaceo-multifidis, emersis tripartitis, lobis cuneatis apice dentatis, carpellis glabris transversè rugoso-striatis, stylo brevi crassiusculo subtruncato vix deflexo. ② in fossis, scaturiginibus inundatisque Gallie occident. — Herba sequentis cum flore prioris. (v. v. DC.)
3. *R. aquatilis* (L. sp. 781.) caule natante, foliis submersis capillaceo-multifidis, emersis tripartitis, lobis cuneiformibus apice dentatis, petalis obovatis calyce majoribus, carpellis setis rigidis hispidis glabrisve transversè rugoso-striatis, stylo brevi crassiusculo subtruncato vix deflexo. ② in aquis aquosisve. — Fl. fr. 4. p. 894. Non diversam a 2 præcedentibus contendit cl. Seringe mel. helv. t. 2. p. 49.
  - a. *heterophyllus* (DC.) foliis emersis tripartitis. — Tabern. ic. 54. f. 2. Variat foliis emersis peltatis aut non peltatis. (v. v.)
  - β. *capillaceus* (DC.) foliis petiolatis, omnibus immersis in lacinias filiformes tenues dissectis. — Barr. ic. 57. t. 566. (v. v.)
  - γ. *cæspitosus* (DC.) foliis petiolatis, omnibus emersis circumscriptione suborbiculatis in lacinias divergentes rigidulas dissectis, petiolis basi latè vaginato-auriculatis. — Schk. handb. 2. t. 152. (v. v.)
  - δ. *stagnatilis* (DC.) foliis sessilibus, omnibus immersis capillaceo-multifidis circinnatis, lacinii abbreviatis, vaginis obsolete auriculatis. (v. v.)
  - ε. *peucedanifolius* (DC.) foliis petiolatis, omnibus immersis in lacinias longas parallelas dissectis. — Oed. dan. t. 376. (v. v.)

Sect. II. RANUNCULASTRUM. DC. prod. 1. p. 27.

Carpella lævia compressa in spicam digesta; radices grumosæ.

4. *R. BULLATUS* (L. sp. 744.) foliis omnibus radicalibus petiolatis ovatis dentatis plus minusve bullatis, scapis nudis unifloris, carpellis tenuiter reticulatis, stylo brevi apice tumidiusculo subuncinato.  $\frac{1}{2}$  in arvis incultis Cornicæ. — Fl. fr. 4. p. 903. — Clas. hist. 1. p. 238. f. 1. et 2. Flores latei, odorati. (v. s.)
5. *R. CUNEOFOLIOS* (L. sp. 780.) foliis radicalibus petiolatis subvillosis trisectis, segmentis semel aut bis tripartitis acutis, primordialibus ovatis dentatis trilobisve, caule erecto 1-2-floro, calyce patente subreflexo, carpellis punctatis, stylo rostrato compresso.  $\frac{1}{2}$  in collibus Galliæ occidentalis mediz australisque. — Fl. fr. 4. p. 900. — Barr. ic. t. 581. (v. v.)
6. *R. MOSSULIACUS* (L. sp. 778.) foliis lanuginosis, radicalibus trilobis, lobis cuneatis trifido-dentatis, superioribus tripartitis, lobis integris linearibus, caule erecto paucifloro, calyce reflexo, capitulo ovato, carpellis punctatis, stylo rostrato compresso acuto recto.  $\frac{1}{2}$  in pascuis apricis saxatilibusve regionis mediterraneæ. — Fl. fr. 4. p. 899 et 5. p. 688. ic. rar. t. 50. — R. Illyricus Beal. cyst. syst. vern. ord. 1. t. 13. f. 1. Variat foliis angustis, cuneatis, rotundisve. (v. v.)
7. *R. AMATICUS* (L. sp. 777.) foliis ternatim biternatimve sectis, segmentis dentatis aut inciso-trifidis, caule erecto simplici aut inferne ramoso, calyce patente demum reflexo, capitulo cylindrico, carpellis minutissime punctatis, stylo elongato teretiusculo acuto patenti-deflexo.  $\frac{1}{2}$  ex Oriente indigena, in hortis culta. — Fl. fr. 4. p. 892. — Weinm. phyt. 849-852. Variat caule ramoso, foliis 3-sectis multifidiisve, lobis acutis obtusisve, florum colore. (v. v. c.)

Sect. III. THORA. DC. l. c. p. 28.

Carpella lævia subglobosa; radices grumosæ.

8. *R. THORA* (L. sp. 775.) foliis glaberrimis, radicalibus petiolatis caulinoque scassili reniformibus crenatis, floralibus incisiss, caule 2-3-floro glabro, carpellis longitudinaliter nervosis, stylo filiformi apice subcircinnato.  $\frac{1}{2}$  in scopulis ad nictum in summis Alpibus et Pyrenæis. — Fl. fr. 4. p. 903. — Clas. hist. 1. p. 239. f. 3. (v. v.)

Sect. IV. HECATONIA. DC. l. c.

Carpella lævia, ovato-subrotunda in capitulum subrotundum digesta; radices fibrosæ.

\* *Floribus albis, foliis dissectis.*

9. *R. RUTEPOLIUS* (L. sp. 777.) foliis pinnatim sectis, lobis 3-lobato-incisiss multifidis, caule sub 1-floro, calyce glabro, petalis 8-10, ungue discolore.  $\frac{1}{2}$  in scopulis ad nictum in summis Alpibus et Pyrenæis. — Fl. fr. 4. p. 893. All. ped. t. 67. f. 1. (v. s.)
10. *R. GLACIALIS* (L. sp. 777.) foliis radicalibus petiolatis palmatim tripartitis 3-sectisve, lobulis subobtusis crassiusculis, caule subunifloro, calyce hirsutissimo, carpellis compressis marginatis, stylo compresso.  $\frac{1}{2}$  inter scopulos prope nives in Alpibus et Pyrenæis. — Fl. fr. 4. p. 892. — J. Bauh. 3. p. 862. f. 3. (v. v.)
11. *R. SEQUIERI* (Vill. dauph. 4. t. 49.) foliis tripartitis, partitionibus acutis trifidis aut integris, caule subunifloro, calyce glabro, petalis 5 integris.  $\frac{1}{2}$  inter scopulos prope nives in Alpibus. — Fl. fr. 4. p. 893. R. Columnæ. All. ped. t. 67. f. 3. 4. (v. v.)
12. *R. ALPESTRIS* (L. sp. 778.) foliis orbiculatis trilobis, lobis apice lobulato-crenatis obtusiusculis, caule subunifloro, calyce glabro, petalis 5 obcordatis trilobatisve, carpellis obovatis tenuiter venosis, stylo rostrato compresso apice uncinato.  $\frac{1}{2}$  inter scopulos et in pascuis editioribus Alpium, Pyren.,

Jurassi, Vogesorum. — Fl. fr. 4. p. 892. Jacq. aust. t. 110. Seg. ver. t. 12. f. 1. (v. v.)

13. *R. ACONITIFOLIUS* (L. mant. 79.) foliis palmatim 3-7-partitis, partitionibus inciso-dentatis, supremis sessilibus in lobos lineari-lanceolatis fissis, caule ramoso multifloro, calycibus adpressis glabris, carpellis venosis, stylo conico apice circinnato.  $\mathcal{U}$  in montanis. — Fl. fr. 4. p. 890.

*a. humilis* (DC.) foliis radicalibus 3-5 partitis, bracteis lanceolatis serratis, caule gracili glabro. — In pratis udibus excelsis. — Lob. ic. 668. f. 2. (v. v.)

*$\beta$ . crassicaulis* (DC.) foliis radicalibus 3-partitis, bracteis lanceolatis serratis, caule crassiusculo apice adpressè villosulo. — Secus rivulos et in udibus Pyrenæorum. — R. heterophyllus, Lapeyr. abr. 316 non Sm. (v. v.)

*$\gamma$ . intermedius* (DC.) elatior, foliis radicalibus tripartitis, bracteis linearibus integris. — In sylvaticis Cebennæ. (v. s.)

*$\delta$ . platanifolius* (DC.) elatior, foliis radicalibus 5-7-lobis acuminatis, bracteis linearibus integris. — In dumetis et sylvaticis montium demissiorum. — R. platanifolius L. mant. 79. — Lob. ic. 668. f. 1. et 667. f. 2. (v. v.)

*$\epsilon$ . dealbatus* (DC.) foliis radic. 7-lobis dentatis subtus candicantibus, supremis trilobis. — In rupibus calcareis Pyren. — R. dealbatus Lapeyr.  $\dagger$ .

14. *R. LACINIOSUS* (Bell. act. taur. 5. t. 8.) foliis cuneiformibus apice irregulariter acutè incisis, caule glabro ramoso multifloro, calycibus adpressis.  $\mathcal{U}$  rarissimus in pascuis fertilibus Alpium Delphinatûs ex Vill. — Valdè affinem *R. aconitifolio* affirmat cl. Seringe mel. bot. 2. p. 54, hybridum inter *R. aconitifolium* et *Pyrenæum* cl. DC. syst. 1. 242. Fl. fr. 4. p. 891. (v. v. DC. in Ped.)

\*\* *Floribus albis, foliis integris.*

15. *R. PYRÆNEUS* (L. mant. 248.) foliis linearibus lanceolatisve integerrimis, scapo pedunculatisve apice tomentosis, capitulis ovatis, stylis subpatentibus apice uncinatis.  $\mathcal{U}$  in pascuis Alpium et Pyrenæorum. — Fl. fr. 4. p. 889. Deless. ic. vol. 1. t. 27. f. B. (v. v.)

*$\beta$ . buplevirifolius* (DC.) caule unifloro, foliis lanceolatis. *R. Pyrenæus* Jacq. misc. 1. t. 18. f. 1. *R. buplevirifolius* Lapeyr. abr. 314. (v. v.)

*$\gamma$ . plantagineus* (DC.) caule multifloro, foliis lato-lanceolatis. — *R. plantagineus* All. ped. t. 76. f. 1. (v. v.)

16. *R. ANGUSTIFOLIUS* (DC. fl. fr. 5. p. 636.) foliis linearibus sublancoelatis glaucescentibus non amplexicaulibus, scapo pedunculisque glabris, capitulis globosis, stylis conico-subteretibus acutis rectis ascendentibus.  $\mathcal{U}$  in turfosis montanis Pyr. orient. circa Mont-Louis. — Deless. ic. sel. 1. t. 27. f. A. Variat caule 1- $\infty$ -floro. (v. s.)

17. *R. AMPLEXICAULIS* (L. sp. 774.) foliis ovali-lanceolatis acuminatis amplexicaulibus, scapo pedunculatisve glabris.  $\mathcal{U}$  in locis excelsis udibus scopulosis Alpium et Pyren. — Fl. fr. 5. p. 636. Deless. ic. sel. 1. t. 27. f. C. Moris. s. 4. t. 30. f. 2. n. 37. Variat caule 1 aut  $\infty$ -floro. (v. v.)

18. *R. PARNASSIIFOLIUS* (L. sp. 774.) foliis radicalibus petiolatis subcordatis ovato-subrotundis, caulinis sessilibus ovato-lanceolatis, pedunculis hirsutis, capitulis ovato-globosis, carpellis obovato-globosis, stylis teretibus circinnatis.  $\mathcal{U}$  in Alpium et Pyrenæorum scopulis nivis æternæ contiguus. — Fl. fr. 4. p. 890. Jacq. coll. 1. t. 9. f. 3. (v. v.)

\*\*\* *Floribus luteis, foliis indivisis.*

19. *R. GRAMINEUS* (L. sp. 773.) foliis lanceolatis linearibusve integerrimis, caule erecto glaberrimo ad collum fibrilloso, radice fasciculatâ, squamulâ petalorum tubulosâ, capitulis ovatis, carpellis lævibus, stylo conico crassiusculo apice vix flexo.  $\mathcal{U}$  in pratis inundatis campisque subargillosis Galliæ mediæ et australis. — J. Bauh. 3. p. 866. f. 2 et 3. (v. v.)

20. *R. LINGUA* (L. sp. 773.) foliis lanceolatis subserratis sessilibus semi-amplexicaulibus, caule erecto glabro, capitulo globoso, carpellis lævibus;

stylo lato rostrato ascendente. *℥* in palæstris inter arundines, ad fontes et fossas. — Fl. fr. 4. p. 904. — J. Bauh. 3. p. 865. ic. (v. v.)

21. *R. FLAMMULA* (L. sp. 772.) foliis glabris linearibus lanceolatis subovatisve subintegris, inferioribus petiolatis, caule declinato pleno basi subradicante, pedunculis oppositifoliis, capitulo ovato-globoso, carpellis lævibus, stylo conico deflexo brevissimo. *℥* in humidis frequens. — Fl. fr. 4. p. 905. fl. dan. t. 575. *R. ovatus* Pers. — Dod. pempt. t. 43a. f. 2. (v. v.)

*β. reptans* (Sm.) caule repente ad quemque nodum radicales agente, foliis linearibus integerrimis. — In glareosis udis rarius. — *R. reptans*. L. Fl. fr. p. 905. fl. dan. t. 108. (v. v.)

\*\*\*\* *Floribus flavis, foliis dissectis.*

22. *R. AURICOMUS* (L. sp. 775.) foliis glabris, radicalibus petiolatis cordatis plerisque tripartitis lobatisve, caulinis in lobos lineares integros subdentatoeve partitiis, calyce pubescente petalis brevioris, capitulis subglobosis, stylo teretiusculo recurvo brevi. *℥* in pascuis humidis et umbrosis. — Fl. fr. 4. p. 899. Variat lobis foliorum superiorum integris (Fuchs. hist. 156. ic.) aut serratis (Weinm. 857. A.) et inferioribus indivisis. (All. ped. 82. f. 2.) (v. v.)

23. *R. SCLELERATUS* (L. sp. 776.) foliis glabris, radicalibus petiolatis tripartitis, lobis trilobatis obtusè subincisis, summis tripartitis, lobis oblongo-linearibus integris, floralibus oblongis, calyce glabro, carpellis minutissimis in capitulum ovato-oblongum dispositis, stylo vix conspicuo. (i) in fossis et paludibus. — Fl. fr. 4. p. 897. Fuchs. 159. ic. (v. v.)

24. *R. MONTANUS* (Wild. sp. 2. p. 131.) foliis radicalibus glabris tripartitis orbiculatis, laciniis trifidis obtusis, caulinis sessilibus in lobos lineares integerrimos 3-5-partitis, caule 1-floro apice adpressè pubescente, calyce glabrisculo. *℥* in pratis editioribus Alpium et Pyren. — Fl. fr. 4. p. 895. *R. nivalis* Jacq. anat. t. 325 et 326 non L. (v. v.)

*β. tenuifolius* (DC.) foliorum laciniis acutis. — In Jurasso et Galloprov. (v. s.)

25. *R. VILLARSH* (DC. fl. fr. 4. p. 896.) foliis pubescentibus, radicalibus suborbiculatis tripartitis laciniis trifidis dentatis acutis, caulinis sessilibus in lobos lineares integros 3-5-partitis, caule 1-floro apice pubescente, calyce patentim villosulo. *℥* in pratis asperisque Alpium Delph. et Galloprov. — *R. Lapponicus* Vill. 4. p. 743. (v. v. DC.)

26. *R. GOUANI* (Wild. sp. 2. p. 132.) foliis radicalibus orbiculatis 5-lobatis, lobis inciso-dentatis, caulinis sessilibus palmato-partitis lobis dentatis, caule pubescente, calyce subvillosulo. *℥* in pratis Pyren. et rarius Alpium. — Fl. fr. 4. p. 896. et 5. p. 627. *R. Pyrenæus* Gou. ill. t. 17. f. 1 et 2 non L. (v. v. DC.)

27. *R. ACERIS* (L. sp. 779.) foliis pubescentibus subglabrisve palmato-partitis, lobis inciso-dentatis acutis, summis linearibus, caule erecto multifloro subpubescente; pedunculis teretibus, calyce subvillosulo, capitulis globosis, carpellis punctis numerosissimis notatis, stylo brevi rostrato recto vel apice vix curvato. *℥* in pratis pascuis et sylvestribus. — Fl. fr. 4. p. 899. Fuchs. 879. ic. (v. v.)

*β. sylvaticus* (DC.) petiolis foliisque subtus velutino-villosis. — *R. lanuginosus*. Fl. fr. 4. p. 899. non L. (v. s.)

*γ. multifidus* (DC.) foliorum lobis inciso-multifidis. (v. s.)

28. *R. POLYANTHEROS* (L. sp. 779.) foliis 3-5-palmato-partitis inciso-multifidis, lobis sublinearibus, caule erecto multifloro petiolisque patentim pilosis, pedunculis subacutis, calyce hirsuto, stylo brevissimo rectiusculo. *℥* in pratis, dumetis et sylvis caducis Gallie borealis. — Fl. fr. 5. p. 638. *R. napellifolia* var. Crantz. aust. 2. t. 4. f. 1. sup. (v. s.)

29. *R. TUNBERGII* (Lapeyr. abr. 320.) foliis radicalibus amplis trilobis lobis distantibus, lateralibus bilobis, intermedio cuneiformi trilobo, lobulis inci-

- ais, caule subscescente pubescente, pedunculis divaricatis teretibus, calyce adpresso.  $\mathcal{L}$  in Pyrenæorum pratis. — Affinis. R. lanuginoso, an satis a R. nemoroso distincta. DC. *sys.* 1. p. 281. R. polyanthemo referendam speciem censet cl. Walker-Arnolt qui hanc in herb. Lapeyr. vidit. (v. s. DC.)
30. R. *nemorosus* (DC. *sys.* 1. p. 284.) foliis radicalibus ultrà medium 3-5-fidis lobis cuneiformibus trifidis, lobulis apice dentatis, caule erecto petiolisque patentim pilosis, pedunculis sulcatis, carpellis stylo incurvato uncinatis.  $\mathcal{L}$  in sylvis.
- a. multiflorus* (DC.) caule, multifloro, foliis superioribus in lobos lineares integros partitis. — In montanis ad radices Jurassi et Cebennarum. (v. v.)
- β. pauciflorus* (DC.) caule 1-3-floro, foliis superioribus sublobatis inciso-dentatis, summis minimis. — In subalpinis Cebenn. Pyren., circa Aginnum (St. Am.), Bardigalam (Desmoul.). — R. villosus St. Am. bouq. t. 5. (v. s.)
31. R. *lanuginosus* (L. sp. 779.) foliis trifidis holosericeis, lobis latis dentatis subincisis, caule erecto multifloro petiolisque reflexo-pilosis, pedunculis teretibus, carpellis stylo uncinatis.  $\mathcal{L}$  in sylvis montanis. — Fl. dan. t. 397. — J. Bauh. *hist.* 3. p. 417. f. 3. (v. v.)
32. R. *repens* (L. sp. 779.) foliis pinnatim trisectis, segmentis cuneatis trilobatis inciso-dentatis, caule flagella prostrato-repentina e collo agente, florifero suberecto, calyce adpresso, capitulo subrotundo, carpellis punctis minimis notatis, stylo tenui deflexo.  $\mathcal{L}$  in pratis fossis plateis oleraceisque. — Fl. fr. 4. p. 898. — Dod. *pempt.* 425. ic. (v. v.)
- β. erectus* (DC.) caulibus floridis erectis, flagellis repentibus nullis. (v. v.)
33. R. *palustris* (Sm. in Rees. *cyclop.* n. 52.) foliis subhirsutis petiolatis ultrà medium trifidis, sinubus obtusis, lobis ovatis incisis dentatis, caule erecto ramoso subnundo, capitulis globoso-depressis, carpellis punctis numerosis notatis stylo brevi recto acuminatis.  $\mathcal{L}$  in Corsicâ. — Caulis dichotomi. DC. *sys.* 1. p. 294. R. Corsicus. Fl. fr. 5. p. 637. (v. s.)
34. R. *bulbosus* (L. sp. 778.) foliis radicalibus petiolatis trisectis subpinnatisectis, segmentis trifidis inciso-dentatis, medio petiolulato, caule erecto ad collum bulboso, calyce reflexo, capitulis globoso-depressis, carpellis punctatis stylo brevi apice uncinato acuminatis.  $\mathcal{L}$  in pratis pascuisque frequena. — Fl. fr. 4. p. 901. — Fuchs. *hist.* 160. ic. (v. v.)

Sect. v. ECHINELLA. DC. *prod.* 1. p. 41.

Carpella tuberculis aculeisve scabra.

\* *Folius dissectis lobatisve.*

35. R. *philoxotis* (Retz. obs. 6. p. 31.) foliis trilobatis trisectisve, lobis obtusis inciso-dentatis, medio petiolato, calyce reflexo, caule basi non bulboso, capitulis globosis, carpellis ad margines tuberculorum minimorum seriem unicam gerentibus, stylo brevi lato apice inflexo. ① in pratis, aquilegiis, dametis. — Fl. fr. 4. p. 901 et 5. p. 639. — J. Bauh. 3. p. 417. f. 2. (v. v.)
- β. intermedius* (DC.) foliis subglabris. — In humidioribus. R. intermedius Poir. (v. v.)
- γ. parvulus* (DC.) caule nudo subunifloro. — In siccis, lapidosis, apricis, aridis. — R. parvulus L. mant. 79. — Barr. ic. t. 791. (v. v.)
36. R. *cornigerus* (Viv. cors. 8. et app. 2.) sericeo-hirsutus, foliis radicalibus omnibus cordatis subrotundo-ovatis dentato-crenatis integris trilobisve, caulibus incisis simplicibusque, caule subunifloro pedunculisque radicalibus nudis unifloris, calycibus reflexis, carpellis obsoletè tuberculatis, stylo brevissimo patenti-deflexo. — In monte Coscione Corsicæ. — R. alpinus, etc. Bocconius. t. 124. (v. s.)
37. R. *arvensis* (L. sp. 780.) foliis glabris, primordialibus apice dentatis, radi-

calibus tripartitis, caulinis multifidis, lobis linearibus, caule erecto multifloro, carpellis utrinque echinatis stylo longo recto ascendente acuminatis. ① in arvis. — Fl. fr. 4. p. 902. — Fuchs. 157. ic. (v. v.)

38. *R. muricatus* (L. sp. 780.) foliis glabris petiolatis suborbiculatis trilobis grossè dentatis, caule erectiusculo aut diffuso, pedunculis oppositifoliis, calyce patente, carpellis utrinque punctatis et tuberculoso-aculeatis stylo lato compresso recto acuminatis. ① in depressis humidis prov. Mediterr. — Fl. fr. 4. p. 902. — Clus. 233. f. 2. (v. v.)

39. *R. parviflorus* (L. sp. 780.) foliis villosis orbiculatis trilobis grossè dentatis incisive, caulibus molliter pilosis decumbentibus, pedunculis oppositifoliis, calyce demum reflexo petala æquante, carpellis punctatis tuberculosisque stylo compresso brevi apice uncinato rostratis. ① in humidis lapidosis provinc. occident. et australium. — Fl. fr. 4. p. 902. *R. parvulus* Lapeyr. abr. non L. *R. Xatardi* Lapeyr. supp. 77. — Moris. s. 4. 28. f. 12. n. 21. (v. v.)

40. *R. subapetalus* (Vict. Aug. in ann. soc. lin. 1. p. 193.) caule erecto substriato, florum pedunculis oppositifoliis, petalis 2-3-calyce minoribus interdum nullis, foliis superioribus 3-5 fidis, inferioribus subrotundis lobatis omnibus hirsutis, seminibus elevato-punctatis stylo subhamato coronatis. ① in sepibus ad ostium flum. Ain. — An satis à precedenti distinctus?

41. *R. raiolus* (Desf. atl. 1. t. 113.) foliis subvillosis glabrativæ, primordiis subrotundis crenatis, cæteris tripartitis, lobis divergentibus basi cuneatis apice dentatis, caule erecto, pedunculis oppositifoliis, calyce demum subreflexo petalis brevioræ, carpellis tuberculatis et minute punctatis, stylo brevi patente vix uncinato. ① in agris et humidis Corsicæ, Telonis, Perpiuiani. — Fl. fr. 5. p. 639. *R. parviflorus* Lapeyr. abr. 321. non L. ex cl. Walker-Arnolt. (v. v. DC.)

\* \* *Foliis integris.*

42. *R. ornithoglossifolius* (Vill. dauph. 4. t. 49.) foliis inferioribus petiolatis cordatis obtusis, superioribus oblongis sessilibus, caule erecto fistuloso, pedunculis oppositifoliis, carpellis tuberculato-spinosis, stylo latissimo crasso obtuso brevi uncinatis. ① in udis pratis et fossis provinciarum australium, agri Bordigalensis et Andegavensis. — Fl. fr. 5. p. 639. (v. v.)

43. *R. moufflorus* (L. sp. 773.) foliis radicalibus petiolatis ovali-oblongis, floribus sessilibus oppositifoliis, carpellis granulato-tuberculosiis stylo vix rostellatis. ① in humidis pluvii subinundatis sylvæ Fontainebleau et Andegaviæ ex Bast. — Fl. fr. 4. p. 903. — Vaill. act. par. 1719. t. 4. f. 4. (v. v. DC.)

IX. *FICARIA*. Dill. nov. gen. p. 108. t. 5. DC. syst. 1. p. 304.

Calyx 3-phyllus caducus; petala 9 intus basi foveolata; stamina ovariaque oo; carpella lævia compressa obtusa.

1. *F. ranunculoides* (Moench. meth. 315.) radice grumosa, caule folioso, foliis cordatis. ② in pratis sepibus dumetisque humidiusculis. — Fl. fr. 4. p. 886. *Ranunculus Ficaria* L. Bull. t. 43. — Fuchs. 867. ic. (v. v.)

Trib. IV. *HELLEBOREÆ*. DC. prod. 1. p. 42.

Æstivatio calycis et corollæ imbricata. Petala nunc o, nunc irregularia bilabiata nectarifera. Calyx petaloideus. Carpella capsularia intus dehiscentia oo sperma.

X. *CALTHA*. Pers. ench. 2. 107. DC. prod. 1. p. 44.

Calyx coloratus 5-sepalus, sepalis suborbiculatis petaloideis; petala o; stamina oo; capsulæ 5-10 compressæ patentés 1-loculares oo spermæ.

1. *C. PALUSTRIS* (L. sp. 784.) caule erecto, foliis cordatis suborbiculatis rotundè crenatis, auriculis rotundatis.  $\mathcal{U}$  in humidiusculis, secùs sepes fossas et rivulos, primo vere. — Fl. fr. 4. p. 918. — Dod. pempt. 598. f. 1. 2. (v. v.)

XI. *TROLLIUS*. L. gen. n. 700.

Calyx coloratus 5-15-sepalus, sepalis deciduis petaloïdeis; petala 5-20 parva basi tubulosa unilabiata; stamina ovariaque  $\infty$ ; capsulæ oo sessiles subcylindraceæ polyspermæ.

1. *T. EUROPAEUS* (L. sp. 782.) sepalis 15 in globum ferè conniventibus, petalis 5-10 longitudine staminum.  $\mathcal{U}$  in pratis montanis subfrigidis. — Fl. fr. 4. p. 906. Lam. ill. t. 499. — Lob. ic. 675. (v. v.)

XII. *ERANTHIS*. Salisb. trans. lin. 8. p. 303. DC. prod. 1. p. 46.

Involucrum multipartitum subflore. Flos sessilis 5-8 sepalus, sepalis coloratis petaloïdeis oblongis deciduis; petala 5-8 tubulata brevissima; capsulæ pedicellatæ; semina globosa simplici serie disposita.

1. *E. HYEMALIS* (Salisb. l. c.) sepalis 6-8 oblongis.  $\mathcal{U}$  in umbrosis humidiusculis ad radices montium, primo vere. — Helleborus hyemalis L. sp. 783. Fl. fr. 4. p. 929. — Lob. ic. 676. (v. v.)

XIII. *HELLEBORUS*. Adans. fam. 458. DC. prod. 1. p. 46.

Calyx persistens 5-sepalus, sepalis subrotundis obtusis magnis sæpè viridibus; petala 8-10 brevissima tubulata inferne angustiora nectarifera; stam. 30-60; ovaria 3-10; stigmata orbiculata; capsulæ coriaceæ, semina elliptica duplici serie disposita.

\* *Scapo aphyllò 1-2 floro.*

1. *H. NIGRA* (L. sp. 783.) foliis radicalibus pedatisectis glaberrimis, scapo aphyllò 1-2-floro bracteato.  $\mathcal{U}$  in montosis sylvaticis subalpinis Galloprovinciæ, hyeme florens. — Fl. fr. 4. p. 908. — Lob. ic. 681. (v. v.)

\*\* *Caule paucifloro subdichotomo folia floralia subsessilia fusa aut incisa gerente.*

2. *H. VIRIDIS* (L. sp. 784.) foliis radicalibus glaberrimis pedatisectis, caulinis subsessilibus palmati-partitis, pedunculis sæpè bifidis, sepalis subrotundo-ovatis viridibus.  $\mathcal{U}$  in sylvis et dumetis, aprili et maio in planitiibus florens, jun. jul. in montibus excelsis. — Fl. fr. 4. p. 908. Jacq. aust. t. 106. — Fuchs. hist. 274. ic. Garid. six. t. 47. (v. v.)

\*\*\* *Caule folioso multifloro, pedunculis bracteatis.*

3. *H. FETIDUS* (L. sp. 784.) caule multifloro folioso, foliis pedatisectis glaberrimis, segmentis oblongo-linearibus.  $\mathcal{U}$  in pascuis glareosis præsertim calcareis, dumetis et secùs vias planitierum, autumno et hyeme florens. — Fl. fr. 4. p. 907. — Dod. pempt. 386. ic. (v. v.)
4. *H. LIVIDUS* (Ait. kew. 2. p. 272.) caule multifloro folioso, foliis trisectis glaberrimis subtùs glaucis, segmentis ovato-lanceolatis.  $\mathcal{U}$  in Corsicâ hyeme et primo vere florens. — Fl. fr. 4. p. 907. *H. argutifolius* Viv. cors. 8. — Moris. s. 12. t. 4. f. 5. n. 7. (v. v.)

XIV. *ISOPYRUM*. L. gen. 701.

Involucrum o. Calyx 5-sepalus deciduus regularis petaloïdeus; petala 5 æqualia tubulata bilabiata labio exteriorè bifido; stam. 15-20; ovaria 2-20; capsulæ sessiles 1-loculares oblongæ compressæ membranaceæ.

1. *I. THALICTROIDES* (L. sp. 783.) capsulis 1-3, sepalis obtusis, radice repente



fasciculatâ aut grumosâ, petiolis basi in aurículas membranaceas dilatatis. 2/ in montanis umbrosis Pyren., Arvernâ, Delph. — Fl. fr. p. 929. — Moris. s. i. t. 1. f. ult. n. 5. (v. v.)

XV. GARIDELLA. *Tourn. inst. 655. t. 43. L. gen. n. 571.*

Calyx 5-sepalus deciduus vix petaloideus; petala 5 bilabiata 2-fida; stam. 10-40; ovaria 3 connexa; styli brevissimi; capsulæ 2-3 polyspermæ connatæ in unicam 2-3-locularem vix 2-3-cornutam.

1. G. NIGELLASTRUM (L. sp. 753.) petalis sessilibus patentibus, staminibus 10-12. ① in apricis inter vineas et oliveta Galloprov. — Fl. fr. 4. p. 911. — Garid. aix. t. 39. (v. v. DC.)

XVI. NIGELLA. *Tourn. inst. 248. t. 134. L. gen. n. 685.*

Calyx coloratus deciduus petaloideus patens 5-sepalus; petala 5-10 bilabiata; stam. ∞; ovaria 5-10 plus minusve basibus concreta, in stylis simplices longos desinentia; capsulæ totidem plus minusve concretæ polyspermæ, stylis elongatis rostratæ.

1. N. ARVENsis (L. sp. 753.) antheris apiculatis, stylis 5-7 extus circumnathim revolutis, capsulis lævibus in fructum obconicum basi angustatum ultra medium connexis, caule glabro, ramis subdivergentibus. ① in arvis inter segetes. — Fl. fr. 4. p. 910 et 5. p. 640. Bull. t. 126. (v. v.)
2. N. SATIVA (Lin. sp. 753.) antheris muticis, capsulis muricatis in fructum ovatum ad apicem usque connexis, caule erecto pilosiusculo, floribus nudis. ① in arvis Monapellii. — Fl. fr. 5. p. 640. — Fuchs. 503. ic. fl. albid. (v. v. c.) *β. citrina*, seminibus luteis, floribus plena. — Clus. 2. p. 207. ic.
3. N. DAMASCENA (Lin. sp. 753.) antheris muticis, capsulis 5 lævibus bilocularibus in unicam ovato-globosam ad apicem usque connexis, floribus involucro folioso cinctis, sepalis patentibus. ① in arvis inter segetes Gallie australis. — Fl. fr. 4. p. 910. — Fuchs. hist. 504. ic. Var. sepalis albis cæruleisve. (v. v.)

XVII. AQUILEGIA. *Tourn. inst. t. 242. Lin. gen. n. 275.*

Calyx deciduus colorato-petaloideus 5-sepalus, petala 5 deorsum producta in calcaria cava apice callosa inter sepala exserta; ovaria 5; capsulæ totidem erectæ stylis acuminatæ.

1. A. VULGARIS (Lin. sp. 752.) calcaribus incurvis, capsulis villosis, caule folioso multifloro, foliis glabrinisculis, stylis stamina non superantibus. 2/ in pratis sepius umbrosis. — Fl. fr. 4. p. 911. — Cam. ep. 405. ic. (v. v.)
2. A. VISCOsa (Gou. ill. t. 19.) calcaribus incurvis, capsulis villosis caule pauci aut unifloro subnudo foliisque pubescenti-viscosis, stylis stamina non superantibus. 2/ in asperis montanis apricis Gallie australis. — Fl. fr. 4. p. 912. 5. p. 640. Lach. act. Helv. 8. t. 5. (v. v.)
3. A. ALPINA (Lin. sp. 752.) calcaribus rectis apice subincurvis limbo petalorum dimidiò brevioribus, caule 2-3-floro folioso, foliorum segmentis in lobos lineares profundè multifidis. 2/ in locis umbrosis et humidis Alpium. Deless. ic. sel. 1. t. 48. A. Alpina var. α. Fl. fr. 4. p. 912. All. ped. t. 66. (v. v. in Helv.)
4. A. PYRÆNICA (DC. fl. fr. 5. p. 640.) calcaribus rectis limbo vix brevioribus, caule subnudo subunifloro, foliorum segmentis in lobos lineares profundè multifidis. 2/ in pratis elatis Pyren. — A. Alpina var. β. fl. fr. 4. p. 912. Priori dimidiò minor. (v. v.)

XVIII. DELPHINIUM. *Tourn. inst. 426. t. 241. L. gen. n. 681.*

Calyx deciduus petaloideus irregularis, sepalo nempè superiore in

calcar deorsum producto ; petala 4 , 2 superiora basi in appendices intrā calcar contentas producta.

Sect. I. CONSOLIDA. DC. syst. I. p. 341.

Ovaria 1; petala 4 in unum coalita; calcar internum 1-petalum. — Species annuæ.

1. D. AJACIS (Lin. sp. 748.) caule erecto subglabro subsimplici, ramis vix divergentibus longè floriferis, floribus densè racemosis, pedicellis bractearum longitudine, capsulis pubescentibus. (1) in hortis cultum, in Corsicâ reperit Requien, in Santoniâ Desmoulins. — Fl. fr. 4. p. 913. Weinm. phys. t. 409. 411. Lob. ic. 740. t. 1. (v. v.)
2. D. CONSOLIDA (Lin. sp. 748.) caule erecto subglabro divaricatum ramoso, floribus paucis laxè racemosis, pedicellis bractearum longioribus, capsulis glabris. (1) in arvis et segetibus. — Fl. fr. 4. p. 913. — Lob. ic. 739. f. 2. (v. v.)
3. D. PUBESCENS (DC. fl. fr. 5. p. 641.) caule recto pubescente apice divaricatum subramoso, floribus laxè racemosis, pedicellis bractearum longioribus, foliis capsulisque pubescentibus. (1) in agris et segetibus circa Narbonam, Monspelium, Nemausim, Avenionem, Massiliam (Solier). (v. s.)

Sect. II. DELPHINELLUM. DC. l. c. p. 346.

Ovaria 3; petala libera glabra; calcar internum dipetalum. Species annuæ.

4. D. CARDIOPETALUM (DC. syst. I. p. 347.) caule erecto subramoso, foliis glabris membranaceis trisectis, segmentis multifidis, lobis linearibus, rameis bracteolae inferioribus multifidis, racemis confertis, petalis stipitatis, limbo orbiculato basi cordato. (1) in Pyren. vallibus. — D. peregrinum Lam. dict. 2. p. 264. Fl. fr. 4. p. 914. excl. var.  $\beta$  et All. syn. 5. p. 641. non L. D. Garumnæ Lapeyr. supp. (v. v.)
5. D. JUNCUM (DC. fl. fr. 5. p. 641.) caule erecto ramosissimo, foliis glabris rigidis, inferioribus multifidis, rameis bracteisque linearibus integris, racemis laxis, petalis stipitatis limbo elliptico. (1) in Galloprov? Nicæ. — D. peregrinum All. ped. 425. f. 3. (v. s. ex Barbariâ.)

Sect. III. DELPHINASTRUM. DC. syst. I. p. 351.

Ovaria 3-5; petala libera, inferiora in disco barhata bifida; calcar elongatum internum dipetalum. — Species perennes.

6. D. INTERMEDIUM (Ait. kew. 2. p. 243.) petiolis basi non dilatatis, foliis cordatis 5-7 fidis, superioribus 3-lobis, omnium lobis inciso-serratis, pedicellis bracteolis calycibus ovarisque glabris.  $\gamma$  in vallibus Pyren. (v. v. DC.)
  - a. pilosissimum (DC.) caule pilosissimo, foliis villosis, nervis supernè pilosis racemo ramoso. — Gm. sib. 4. t. 80.
  - $\beta$ . leptostachyum (DC.) caule petiolisque glabris, foliis pubescentibus, racemo simplici brevi. D. intermedium Lapeyr. ?
  - $\gamma$ . ranunculifolium (DC.) caule glabro, petiolis pilosisculis, foliis tenuibus pubescentibus, racemo simplici laxo.
  - $\delta$ . laxum (DC.) caule foliisque glabris, lobis elongatis acutius incisis, racemo laxo racemoso.
7. D. MONTANUM (DC. fl. fr. 5. p. 641.) petiolis basi non dilatatis, foliis pubescentibus 5-lobatis, lobis basi cuneatis trifidis incisis, racemo simplici, bracteis calycibus capsulisque pubescentibus, calcare inflexo.  $\gamma$  in vallibus excelsis Pyren. et Alp. Galloprov. — Clus. hist. 2. p. 94. f. 2. (v. v.)

Sect. IV. STAPHYSAGRIA. C. et J. Bauh. DC. prod. I. p. 56.

Ovaria 3-5, petala libera; calcar breve internum dipetalum; capsulae ventricosae, semina pauca magna. — Species biennes.

8. D. REQUIENII (DC. fl. fr. 5. p. 642.) calcare calycis ferè longitudine, bracteolis medio pedicelli insertis, petiolis pilosis. (2) in insulis Stæzchadum, et Corsicâ (cl. Soleirol.). Deless. ic. sel. 1. t. 63. (v. s.)

9. *D. PICTUM* (Willd. enum. 574.) calcare calyce vix brevior, bracteolis ad basin pedicelli insertis, petiolis pubescentibus, pedicellis flore vix longioribus. ② in Gallopr. maritimā. — *D. Staphysagria* J. Baub. 3. p. 641. f. 2. (v. s.)

10. *D. STAPHYSAGRIA* (Lin. sp. 750.) calcare brevissimo, bracteolis ad basin pedicelli insertis, petiolis pilosis, pedicellis flore duplò longioribus. ② in ruderatis Occitanis et Gallopr. — Fl. fr. 4. p. 913. — J. Baub. 3. p. 642. f. 1. (v. v.)

XIX. *ACONITUM*. *Tourn. inst. t. 239 et 240. L. gen. 682. DC. syst. 1. p. 364.*

Calyx petaloideus irregularis deciduus vel marcescens, sepalo nempe superiore concavo cassideiformi; petala 2 superiora (cuculli, nectaria) longè unguiculata apice in saccum expansa sub galeâ recondita.

Sect. I. *ANTHORA*. *DC. l. c.*

Sepala persistentia, galea conica vel semicircularis, ovaria 5; — folia in lobos lineares palmatisecta.

1. *A. ANTHORA* (L. sp. 751.) floribus paniculatis, cucullorum sacco ferè nullo, calcare crasso spirali, staminum alis evanescentibus, foliis multifidis, laciniis linearibus acutis. ½ in Jurasso et Pyrenæis. — Fl. fr. 4. p. 917. — Moris. s. 12. t. 2. n. 7. (v. v.)

ß. *atrovirens* (DC. syst.) foliis caulibusque glabratiss. In Pyrenæis.

γ. *cucullum* (Ser. in DC. prod. 1. p. 57.) paniculâ floribusque puberulis, galeâ conicâ suprâ rostrum incumbente. — In Jurasso. *A. anthoroidenum* DC. syst. 1. p. 366.

Sect. II. *LYCOCTONUM*. *DC. syst. 1. p. 367.*

Sepala decidua; galea conico-cylindracea, ovaria 3; — lobi foliorum cuneiformes pinnati rarò bipinnatisecti.

2. *A. LYCOCTONUM* (L. sp. 750.) caule piloso, foliis petiolatis ciliatis ultrâ medium 3-5-lobatis, lobis cuneatis 3-fidis, floribus spicatis paniculatisve, galeâ conico-cylindraceâ ovarisque glabriusculis. ½ in sylvis dumetis pratensque montosis. — Fl. fr. 4. p. 916. — Moris. s. 12. t. 2. n. 1. (v. v.)

3. *A. PYRÆNICUM* (Lam. dict. 1. p. 33.) densè velutino-pubescent, caule ramissimo, foliis petiolatis ultrâ medium 5-9-lobatis, lobis cuneatis trifidis, floribus paniculatis, galeâ conico-cylindraceâ angustatâ ovarisque velutinis. ½ in Pyrenæis. — DC. fl. fr. 4. p. 916. syst. 1. p. 368. *A. Lycoctonum* µ. Ser. in DC. prod. 1. p. 57. — Camer. epit. 831. ic. (v. v. DC.)

Sect. III. *CAMMARUM*. *DC. l. c. p. 374.*

Sepala decidua, galea conica compressa ovaria 3-5; — lobi foliorum pinatisecti.

4. *A. PANICULATUM* (Lam. dict. 1. p. 33.) paniculâ divaricatâ, ramis puberulis, galeâ conico-semicirculari, rostro inclinato, sacco cucullorum subconicoideo, calcare crasso brevi spirali, alis staminum angustis evanescentibus, folliculis elongatis. ② in asperis dumetis abruptis Alpium. — Fl. fr. 4. p. 918. 5. p. 643. — Ser. mus. helv. 1. t. 15. f. 20 et 21. (v. v.)

Sect. IV. *NAPELLUS*. *DC. syst. 1. p. 371.*

Sepala decidua, galea semicircularis rarò navicularis; ovaria 3-7. — Foliorum lobi bipinnatisecti.

5. *A. NAPELLUS* (L. sp. 751.) floribus densè spicatis vel laxè paniculatis, sacco cucullorum subconico, calcare brevi crasso inclinato, alis staminum cuspidatis vel evanescentibus, foliorum lobis cuneatis pinnatisectis, ovaris 3, rarò 5, glabris vel pilosis. ② in montosis. — Fl. fr. 4. p. 917. Lob. ic. 679.

f. 1. Inflorescentiâ, galæ formâ, floris colore, foliis, staturâ admodum variabilis. (v. v.)

Trib. V. *PÆONIACEÆ*. DC. prod. 1. p. 64. — Ranunculaceæ spuria. DC. syst. 1. p. 281.

Antheræ introrsæ. — An ordo proprius? ex DC. prod: l. c.

XX. *ACTÆA*. L. sp. ed. 2. p. 722. DC. syst. 1. p. 381.

Calyx deciduus 4-sepalus; petala 4; carpella ∞-sperma.

1. *A. SPICATA* (L. sp. 722.) monogyna, baccis subglobosis, petalis staminum longitudine, racemo ovato, foliis bi-triternatimve sectis segmentis ovato-lanceolatis serratis incisive. ½ in umbrosis asperis montanis. — Fl. fr. 4. p. 920. Bull. t. 83. — Lob. ic. 682. f. 1. (v. v.)

XXI. *PÆONIA*. L. gen. n. 678.

Calyx 5-sepalus foliaceus inæqualis; petala 5-10; stam. ∞; discus carnosus ovaria cingens; carpella 2-5 grossa stigmatibus bilamellatis crassis instructa, in folliculos capsulares conversa; semina subglobosa nitida. — Folia caulina biternatim secta; flores ampli.

1. *P. CORALLINA* (Retz. obs. 3. p. 34.) herbacea, carpellis tomentosis, foliorum segmentis ovatis integris glabris. ½ circa Aureliam, Blois et propè Alais. — Fl. fr. 5. p. 643. — P. mas. Dod. pempt. 194. f. 1. (v. v.)
2. *P. OFFICINALIS* (Retz. obs. 3. p. 35.) herbacea, carpellis tomentosis rectiusculis, foliorum segmentis inæqualiter laciniatis glabris, laciniis ovato-lanceolatis. ½ in pratis nemorosis Delphinatûs, Galloprov? Pyrenæorum ex Lapeyr. — Fl. fr. 4. p. 919. — P. femina Dod. 195. f. v. Lob. ic. 682. (v. v.)
3. *P. RUSSÆ* (Biv. man. sic. 4. p. 12.) herbacea, carpellis pilosis recurvatis, foliorum segmentis ellipticis integris subtus vix pubentibus. ½ in Corsicæ sylvâ Perticato dietâ reperiit cl. Soleirol. — DC. prod. 1. p. 66. Petala chermesina. (v. s. fructigeram.)
4. *P. PARADOXA* (And. trans. Linn. 12. p. 288.) herbacea, carpellis tomentosis rectis, foliorum segmentis multipartitis subundulatis subtus glaucis pilosis. ½ in Galliæ australis montibus apricis rarè. — DC. prod. 1. p. 66. P. promiscua Lob. ic. 683. f. 2. P. peregrina DC. fl. fr. 5. p. 643. (v. v. DC.)

Ordo II. BERBERIDEÆ. *Vent. tabl. 3. p. 83. DC. fl. fr. ed. 3. v. 4. p. 627. syst. 2. p. 1. prod. 1. p. 105.*

Sepala 3-4 aut sæpius 6 decidua, oblonga aut ovalia, sæpè subcolorata, duplici serie alternatim disposita, extùs squamis sæpius stipata. Petala tot quot sepala et iis opposita, aut rarius numero dupla, hypogyna, sæpius basi intùs glandulâ squamulâve aucta. Stam. tot quot petala iis opposita, filam. brevibus, antheris oblongis adnatis bilocularibus, loculis à basi ad apicem valvulâ subelastica dehiscens. Ovarium (abortivum) solitarium ovatum subobliquum 1-locul., stylo sublaterali brevissimo et stigmate suborbiculato coronatum. Fructus baccatus capsularisve. Semina imæ placentæ laterali adfixa 1-3, ovata aut globosa. Albumen carnosum sæpè subcorneum. Embryo rectus axilis, radiculâ ad apicem plus minùsve inkrassatâ, cotyledonibus planis. — Frutices aut Herbæ perennes; folia aut foliola aristato-serrata.

I. BERBERIS. *L. gen. 442.*

Sepala 6 squamis 3 stipata; pet. 6 intùs biglandulosa; bacca 2-3-sperma. — Frutices foliis primariis abortivis in spinam sæpius mutatis, secundariis in axillis fasciculatis. Flores flavi.

1. *B. vulgaris* (Lin. sp. 472.) spinis tripartitis, foliis fasciculatis subobovatis ciliato-serratis, racemis multifloris pendulis, petalis integris.  $\mathfrak{f}$  in dumetis sepibus incultis et apricis præsertim calcareis Gallie totius. — Fl. dan. t. 904. Fl. fr. 4. p. 627. Fructus ruber et in var. cultis luteus violaceus purpureus niger aut albus. (v. v.)

II. EPIMEDIUM. *L. gen. n. 148.*

Sepala 4 extùs basi bibracteolata; pet. 4 intùs appendice discolore aucta; capsula siliculæformis oo-sperma. — Herbæ perennes, foliis multisectis, floribus atropurpureis.

1. *E. alpinum* (Lin. sp. 171.) foliis radicalibus nullis, caulino binternatim secto, segmentis cordato-lanceolatis acuminatis aristato-serratis.  $\mathcal{U}$  in umbrosis et dumetis montanis Alpium, Vagesorum ubi forsân à Lindernio introductum. — Fl. fr. 4. p. 628. Lam. ill. t. 83. (v. v. in Helv.)

Ordo III. NYMPHÆACEÆ. DC. *prop. med. ed.* 2. p. 119.  
*syst.* 2. p. 39. *prod.* p. 113.

Torus floris expansus in urceolum ovaria includentem et stigmatibus coronatum. Sepala 4-6 toro inserta persistentia colorata. Petala pluriserialia, toro inserta, oblonga, plana, serie quâque totidem petalis constante quot sunt sepala, alternâ cum iis alternâ. Stam.  $\infty$  multiserialia toro cum petalis inserta, filamentis planiusculis, antheris adnatis introrsis linearibus bilocularibus rimâ duplici longitudinaliter dehiscentibus. Carpella 8-24 intrâ torum inclusa 1-styla membranacea evalvia polysperma, stigmatibus extrâ torum urceolatum peltatim radiatis basi connatis apice liberis. Semina  $\infty$  parietibus carpellorum adfixa inversa ovato-globosa, arillo folliculæformi cincta. Embryo ad basin albuminis farinacei situs, parvus, turbinato-globosus, inclusus in sacculo proprio membranaceo undè monocotyledoneus videtur, sed sacculo fissocotyledones 2 foliaceas exhibens. — Herbæ aquaticæ, truncotereti horizontali repente, foliis longè petiolatis limbo natantibus, floribus longè pedunculatis radicalibus solitariis.

I. NYMPHÆA. DC. *syst.* 2. p. 49.

Sepala ad basin tori; petala staminaque cum toro carpella tegente longè adnata et bacca ideò quasi semi-infera cicatrisata. — Flores albi.

1. *N. ALBA* (Lin. sp. 729.) foliis cordatis integerrimis, stigmate 16-radiato, radiis adscendentibus.  $\mathcal{L}$  in fossis lacubus fluminibusque lentè fluentibus Galliæ totius. — Fl. dan. t. 602. Fl. fr. 4. p. 630. (v. v.)

*β. minor* (Beal. cyst. vern. ord. VII. t. 3. f. 2.) flore minore. — In Alsatia circa Argentinam. (v. s.)

II. NUPHAR. DC. *syst.* 2. p. 59.

Sepala petala staminaque ad basin tori inserta et ideò bacca quasi supera lævis. — Flores lutei.

1. *N. LUTEA* (Sm. prod. fl. gr. 1. p. 361.) calyce 5-sepalo, stigmate integro 16-20-radiato profundè umbilicato, foliis ovali-cordatis, lobis approximatis, petiolis acutangulis triquetris.  $\mathcal{L}$  in fossis et fluminibus lentè fluentibus Galliæ ferè totius. — *Nymphaea lutea* Lin. sp. 729. engl. bot. t. 159. Fl. fr. 4. p. 630. (v. v.)
2. *N. FUMILA* (Sm. engl. bot. 2292.) calyce 5-sepalo, stigmate lobato 10-radiato, foliis oblongo-cordatis punctatis subpubescentibus, lobis approximatis, petiolis basi semiteretibus apice subtriquetris.  $\mathcal{L}$  in lacubus montanis Vogesorum. — DC. *syst.* 2. p. 60. Dimidiò minor quam *N. lutea*. (v. s. ex Germania.)

Ordo IV. PAPAVERACEÆ. *DC. syst. 2. p. 67. prod. 1. p. 117.*  
— *Papaveracearum. gen. Juss.*

Calyx 2-sepalus foliaceus, sepalis concavis florem juniorem sæpius includentibus. Pet. regularia eglandulosa, antè evolutionem irregulariter plicata, sæpius 4 unicâ serie disposita et inter se cruciata. Stam. inter se libera, rariùs 4 serie unicâ petalis opposita, sæpius seriebus multiplicatis 8, 16 20... etc. Filam. filiformia; antheræ biloculares basi insertæ sulco duplici dehiscentes. Ovarium liberum constans carpellis nunc paucis (2-3) nunc plurimis (10-12) intrâ thalami productionem membranaceam sæpè amplexis. Stylus 0 aut brevis. Stigmata radiatim patentia. Capsula ovata aut elongata siliquæformis, carpellis margine seminifero inter se connexis, et placentas ideò utrinquè seminiferas nunc bifiliformes nunc septiformes conficientibus. Semina  $\infty$  horizontalia subglobosa exarillata. Albumen carnosum-oleosum. Embryo minimus ad basin albuminis rectus. — Herbæ succo lacteo albo aut croceo fætæ. Radices fibrosæ. Folia alterna penninervia dentata aut pinnatilobata. Pedunculi elongati 1-flori.

I. PAPAVER. *Tourn. inst. t. 119. L. gen. n. 448. DC. fl. fr. 4. p. 631.*

Sepala 2 decidua; pet. 4; stam.  $\infty$ ; stylus 0; stigmata 4-20 radiantia sessilia super discum ovarium coronantem; capsula obovata 1-loc. septis incompletis, carpellis 4-20 in thalami productione membranaceâ inclusis constans, sub stigmatum coronâ valvulis brevibus dehiscent. — Pedunculi antè florescentiam apice inflexi.

### §. 1. Capsulis hispidis.

1. *P. PARNASSIUM* (Willd. enum. 563) capsulâ hispidâ obovatâ, sepalis setosis, pedunculis radicalibus, foliis pilosis pinnatilobatis, lobis incisive dentatisve subobtusis.  $\mathcal{L}$  in apricis calcareis montium regionem olearum circumdantium nempe in Pyrenæis orient. Cebennis, Alpibus aust.—*DC. syst. 2. p. 71.*  
*a. luteum* (DC.) floribus flavis defloratis subaurantiacis. — *Argemone Pyrenaica* Lin. *P. aurantiacum* Lois. not. 84. *DC. fl. fr. 5. p. 585* *P. alpinum* et *P. suaveolens* Lapeyr. (v. v.)  
*ß. puniceum* (DC.) floribus puniceis. In Pyrenæis loco dicto Port de Plan ex Lapeyr.
2. *P. ALPINUM* (Lin. sp. 724.) capsulâ hispidâ obovato-oblongâ, sepalis pilosisculis, pedunculis radicalibus, foliis glabriusculis bipinnatisectis, lobulis tenuibus subacutis.  $\mathcal{L}$  in apricis inter lapides et saxa Alpium editiorum. — *Jacq. austr. t. 83. Fl. fr. p. 93a. Flores albi.* (v. v.)
3. *P. HYADUM* (Lin. sp. 725.) capsulâ hispidâ obovato-globosâ torosâ, sepalis pilosis, caule folioso multifloro, foliis pinnatisectis, lobis multifidis linearibus.  $\textcircled{1}$  in agris versuris et oleraceis Gall. ferè totius. — *Lob. ic. t. 276. f. 1. Fl. fr. 4. p. 631. Fl. rubri.* (v. v.)
4. *P. ARGEMON* (Lin. sp. 725.) capsulâ hispidâ elongato-clavatâ, sepalis glabriusculis, caule folioso multifloro, foliis bipinnatisectis, lobis linearibus.

① in campis arenosis, agris, oleraceis Gall. totius. — Lob. ic. t. 276. f. 2. Fl. fr. 4. p. 631. Fl. parvi pallide coccinei angue nigro. (v. v.)

§. 2. *Capsulis glabris.*

5. *P. DURIUM* (Lin. sp. 726.) capsulâ glabrâ oblongo-clavatâ, sepalis pilosis, caule multifloro setis patentibus hispido, setis pedunculorum adpressis, foliis pinnatipartitis, lobis inciso-dentatis. ① in arvis sabulosis oleraceisque. Gall. tot. — Schkuhr. handb. 2. t. 140. Fl. fr. 4. p. 633. Flores coccinei aut rarius albi. (v. v.)
6. *P. RUMEX* (Lin. sp. 726.) capsulâ glabrâ obovatâ, sepalis pilosis, caule multifloro setis patentibus scabro, foliis pinnatipartitis, lobis elongatis inciso-dentatis acutis. ① in arvis frequens inter segetes. — Blackw. herb. t. 2. et 560. Fl. fr. 4. p. 632. Flores coccinei variantes albi aut carnei aut marginati. (v. v.)
7. *P. ROUSIER* (Vig. diss. 39. t. 1. f. 1.) capsulâ glabrâ subrotundâ, sepalis pilosis, caule multifloro piloso, foliis bipinnatipartitis villosis, lobis linearibus pilo terminatis. ① in arenosis ad Frontignan propè Monapellum legit Roubien, in Corsicâ Ph. Thomas. — Fl. fr. 5. p. 585. (v. s.)
8. *P. STRICTURUM* (DC. fl. fr. 5. p. 586) capsulâ glabrâ obovatâ, sepalis subsetosis, caule glabro paucifloro, foliis amplexicaulibus glaucis inciso-repandis, dentibus setâ terminatis. ① in insulis Stœschadum obs. Requier. et in Corsicâ Ph. Thomas. — Deless. ic. sel. 2. t. 7. An stirps sylvestris *P. somniferi nigri*? (v. s.)
9. *P. SOMNIFERUM* (Lin. sp. 726.) capsulâ obovatâ aut globosâ calycibusque glabris, caule glabro glauco, foliis amplexicaulibus inciso-repandis, dentibus subobtusis. ① ex Oriente ortum nunc freq. cultum.
  - a. *nigrum* (DC.) capsulis globosis sub stigmate foraminibus apertis, semin. nigris. — Cultum præsertim in Alsatiâ ad confic. oleum *huile d'aillette* dictum. Lam. ill. t. 451. (v. v. c.)
  - β. *album* (DC.) capsulis ovato-globosis sub stigmate foraminibus nullis seu oblitteratis, seminibus albis. — Cultum ad opium extrahendum. An species propria? Wood. med. bot. t. 185. (v. v. c.)

II. *MECONOPSIS*. Vig. diss. f. 3. DC. fl. fr. 5. p. 586.

Sep. 2 decidua; pet. 4; stam. ∞; stylus brevis; stigm. 4-6 radiantia convexa libera; capsula obovata 1-loc. valvis 4-6 apice dehiscens, placentis tenuibus vix intus in membranulas angustas productis. —

1. *M. CAMBRICA* (Vig. diss. 48. f. 3.) capsulis levibus, foliis plurimis petiolatis. ② in pratis umbrosis dumetis et sepibus Pyrenæorum et montium Arvernæ. — *Papaver cambricum* Lin. sp. 727. Fl. fr. 4. p. 633. Succus flavescens. Flores flavi. (v. v.)

III. *ROEMERIA*. DC. Syst. 2. p. 92. prod. 1. p. 122.

Pet. 4; stam. ∞; capsula elongata 3-4-valvis, valvis ab apice ad basin dehiscentibus; sem. reniformia scrobiculata cristâ glandulosâ destitutâ. — Herbæ ①.

1. *R. HYBRIDA* (DC. l. c.) siliquis 3-4-valvibus erectis apice rigide setosis. ① in agris cultis et vineis regionis Mediterraneæ. — *Chelidonium hybridum* Lin. sp. 724. Fl. fr. 4. p. 636. — Clus. hist. 2. p. 92. f. 2. Succus flavus. Flores violacei. (v. v. DC.)

IV. *GLAUCIUM*. Tourn. inst. t. 130. DC. syst. 2. p. 94. prod. 122.

Pet. 4; stam. ∞; capsula elongata bivalvis dissepimento cellulari biloculari; semina ovato-reniformia. — Herbæ ②.

1. *G. FLAVUM* (Crantz. austr. 2. p. 141.) caule glabro, foliis caulinis repandis,



siliquâ tuberculato-scabriusculâ. ② in glareosis et arenosis secus littora marium et fluminum. — *Chelidonium Glaucium* Lin. sp. 724. Fl. fr. 4. p. 635. — Cam. epit. 805. ic. Flores flavi. Succus croceus. (v. v.)

2. *G. corniculatum* (Curt. lond. 6. t. 3a.) caule piloso, foliis caulinis pinnatifidis incis, siliquâ setosâ. ① in agris et segetibus regionis Mediterraneæ rariis. — *Chelidonium corniculatum* Lin. sp. 724. Fl. fr. 4. p. 635. Flores phœnicei. (v. v. DC.)

V. *CHELIDONIUM*. *Cartn. fruct. 2. t. 115. f. 5. DC. syst. 2. p. 98.*

Sep. 2 glabra; pet. 4; stam. ∞; capsula elongata bivalvis 1-locul. valvis à basi ad apicem dehiscentibus; semina cristâ glandulosâ instructa. — Herbar. 2.

1. *C. MAJUS* (Lin. sp. 723. var. a.) pedunculis umbellatis, foliis pinnatisectis, segmentis subrotundis dentato-lobatis, petalis ellipticis integris. 2 frequens in umbrosis etruderatis. — Fl. fr. 4. p. 634. Lam. ill. t. 450. f. 1. Flores flavi. Succus croceus actis. (v. v.)

VI. *HYPECOUM*. *Tourn. inst. t. 115. L. gen. 171.*

Pet. 4, interiora sæpius 3-loba; stam. 4; capsula siliquæformis bivalvis transversè nodosa seu articulata; semina inter articulationes solitaria.

1. *H. procumbens* (Lin. sp. 181.) siliquis articulatis compressis arcuatis, petalis 3-lobis, externis dorso glabris. ① in arvis et arenosis secus mare Mediterraneum. — Lam. ill. t. 88. Fl. fr. 4. p. 640. Flores flavi. (v. v.)
2. *H. pendulum* (Lin. sp. 181.) siliquis nodosis teretibus pendulis, petalis glabris, 2 exterioribus ovato-oblongis, 2 interioribus 3-partitis. ① in arvis Galloprovinciæ, Monspelii, Andegavim. — Lob. ic. 743. f. 2. — Fl. fr. 4. p. 641. fl. flavi. (v. s.)

Ordo V. FUMARIACEÆ. DC. *syst.* 2. p. 105. *prod.* 1. p. 125.

Calyx 2-sepalus membranaceus deciduus parvus. Pet. 4 inter se cruciata, nunc libera, nunc basi coalita, nunc 1 liberum et 3 coalita, 2 exteriora sepalis alterna nunc ambo nunc unicum in calcar gibbositateve producta, 2 interiora oblongo-linearia apice callosa. Stam. Filam. 6, in phalanges 2 oppositas ante petala exteriora sitas disposita. Antheræ 6 parvæ, laterales cujusque phalangis 1-loculares, mediæ biloculares; et ideo antherarum loculi 8, et in quâque phalange antheræ 2 strictiori sensu habendæ. Ovarium 1 liberum. Stylus filiformis. Stigma bilamellatum petalis interioribus parallelum. Fructus siccus nunc siquosus bivalvis polyspermus dehiscens, valvulis oppositis, placentis 2 ad suturas nerviformibus persistentibus, nunc bivalvis 2-spermus valvulis coalitis indehiscens, nunc evalvis? indehiscens, abortu? 1-spermus. Semina placentis lateralibus adfixa horizontalia ovato-globosa nitida arillata aut carunculata. Albumen carnosum. Embryo basilaris. Cotyled. oblongæ. — Herbæ succo aqueo. Flores racemosi.

I. CORYDALIS. DC. *fl. fr.* 4. p. 636. *syst.* 2. p. 113.

Pet. 4; unicum basi calcaratum; siliqua bivalvis compressa polysperma. — Herbæ perennes aut annuæ.

SECT. I. CAPNITES. DC. *prod.* 1. p. 126.

*Radice tuberosâ; caule simplici, folia alterna pauca gerente.*

1. C. TUBEROSA (DC. *fl. fr.* 4. p. 637.) caule simplici exsquamato, foliis 2 biternatim sectis, segmentis cuneatis inciso-multifidis, bracteis ovatis integris, radice tuberosâ cavâ.  $\mathcal{L}$  in sepibus et dumetis umbrosis submontanis. — *Fumaria bulbosa*  $\alpha$  Lin. sp. 933. — Lob. ic. 759. f. 1. Fl. purpurei aut albi. (v. v.)
2. C. FABACKA (Pers. ench. 2. p. 269.) caule subsimplici erecto sub infimo folio squamigero, foliis 3-4 petiolatis biternatim sectis, segmentis oblongis obtusiusculis, bracteis ovatis acutis pedicello longioribus, radice tuberoso-bulbosâ solidâ.  $\mathcal{L}$  in sylva Compiegne ex Lois. — Fl. dan. t. 1394. DC. *syst.* 2. p. 118. *fl. fr.* 5. p. 586. *Fumaria bulbosa*  $\beta$  Lin. *fl. succ.* n. 83 $\alpha$  (v. s. ex Vallesia.)
3. C. BULBOSA (DC. *fl. fr.* 4. p. 637.) caule simplici erecto sub infimo folio squamigero, foliis 3-4 petiolatis biternatim sectis, segmentis cuneatis oblongisve bracteisque apice incisiss, radice tuberoso-bulbosâ solidâ.  $\mathcal{L}$  in sepibus, umbrosis, humidiusculis submontanis. — Lob. ic. 759. f. 2. *Fumaria bulbosa*  $\gamma$  Lin. sp. 983. Fl. purpurascens. (v. v.)

SECT. II. CAPNOIDES. DC. *prod.* 1. p. 128.

*Radice fibrosâ; caule ramoso folia alterna plurima gerente.*

4. C. CAPNOIDES (Pers. ench. 2. p. 270.) caule ramoso diffuso, foliis bipinnatim sectis, segmentis obovato-cuneatis trifidis, bracteis lineari-subulatis pedicello triplo brevioribus, siliquis linearibus vix pedicello longioribus.  $\mathcal{L}$  in fissuris rupium circa Narbonam et in muris Argentinæ et Basilæ. — DC. *syst.* 2. p. 125. — Lob. ic. 758. f. 2. — *Fumaria lutea* Lin. mant. 258. *Corydalis lutea* Fl. fr. 4. p. 638. — Flores flavi in var. Europæâ, albi in Mauritanicâ nondum apud nos repertâ. (v. s.)

5. *C. CLAVICULATA* (DC. fl. fr. 4. p. 638.) caule ramoso diffuso scandente, foliis bipinnatim sectis, petiolis cirrhosis, segmentis ovalibus integris, bracteis oblongis pedicello-longioribus. ① in sepibus saxosisque prov. australium et occidentalium.—Lob. ic. 758. f. 1. *Fumaria claviculata* Lin. sp. 985. Flores pallide flavi. (v. v.)

II. *SARCOCAPNOS*. DC. syst. 2. p. 129. prod. 1 p. 129.

Pet. 4, unicum basi calcaratum; capsula bivalvis indehiscens 2-sperma, valvulis 3-nerviis planiusculis.

1. *S. ENNEAPHYLLA* (DC. l. c.) foliis petiolo ramoso triternatim sectis, segmentis ovato-obiculatis. ② in fissuris rupium agri Ruscinonensis.—*Fumaria enneaphylla* Lin. sp. 984. Lam. ill. t. 597. f. 4. DC. Fl. fr. 5. p. 587. Flores albidio ochroleuci apice purpurei. (v. s.)

III. *FUMARIA*. Toura. inst. p. 237. DC. syst. 2. p. 131.

Pet. 4, unicum basi gibbum aut breviter calcaratum; cariopsis indehiscens 1-sperma stylo post anthesin deciduo mutica. — Herba annua.

Sect. I. *PLATYCAPNOS*. DC. l. c.

Silicula seu cariopsides compresso-planæ.

1. *F. SPICATA* (Lin. sp. 985.) siliculis compressis ovalibus lævibus, racemis spicatis, pedicellis bractea multò brevioribus, caulibus erectis. ① in agris oleraceisque regionis Mediterraneæ. — Clus. hist. 2. p. 208. f. 1. Fl. fr. 4. p. 640. Flor. purpurei. (v. v.)

Sect. II. *SPHEROCAPNOS*. DC. l. c.

Silicula seu cariopsides globosæ.

2. *F. CAPREOLATA* (Lin. sp. 985.) siliculis globosis, pedicellis fructiferis recurvatis bractea longioribus, racemis oblongis, caulibus subscandentibus, foliis bipinnatisectis, petiolis subcirrhosis. ① in sepibus, dumetis et arvis asperis prov. australium.—DC. fl. fr. 4. p. 639. ic. rar. 1. t. 34. Fl. carnei apice atropurpurei. (v. v.)
3. *F. MEDIA* (Lois. not. p. 101.) siliculis globosis depressiusculis, pedicellis fructiferis erectis bractea duplò longioribus, racemis laxiusculis, caulibus erectis, foliis suprà decompositis, petiolis subcirrhosis. ① in agris oleraceisque Galliae australis et circa Parisios, Andegavum, Lauzertam, etc.—Vaill. bot. t. 10. f. 4. — Fl. fr. 5. p. 587. Flores albidii apice atropurp. (v. v.)
4. *F. OFFICINALIS* (Lin. sp. 984.) siliculis globoso-retusis, pedicellis fructiferis erectis bractea duplò longioribus, racemis laxiusculis, caule erecto, foliis suprà decompositis, lobis linearibus. ① ubique in cultis, arvis, hortis. — Fuchs. hist. 338. ic. Fl. fr. 4. p. 539. Flor. purpurascens. (v. v.)
5. *F. PARVIFLORA* (Lam. dict. 2. p. 567.) siliculis globosis submucronulatis, pedicellis fructiferis erectis bractea longioribus, racemis laxis, caule diffuso, foliis suprà decompositis, lobis linearibus canaliculatis. ① in cultis oleraceisque sæpè cum priore mixta. — Flores pallidi. Sm. engl. bot. t. 590. Fl. sc. 4. p. 639. (v. v.)
6. *F. VAILLANTII* (Lois. not. 102.) siliculis globosis vix mucronulatis, pedicellis fructiferis erectis bractea longioribus, racemis brevibus, caule erectiusculo, foliis suprà decompositis, lobis linearibus planis. ① in agris arenosis circa Parisios, Andegavum, Monspeliam, etc.—Vaill. bot. t. 10. f. 6. Fl. fr. 5. p. 587. Herba glaucescens. Flores purpurascens. (v. v.)
7. *F. DENSIFLORA* (DC. cat. h. Monsp. 113.) siliculis globosis, pedicellis fructiferis erectis bractea longioribus, racemis densis, calycibus dentatis, caule erectiusculo, foliis suprà decompositis, lobis linearibus crassiusculis. ① in agris lapideis et maris circa Telonem et Monspelium. — Fl. fr. 3. p. 588. syst. 2. p. 137. Var. floribus purpureis et albidis. (v. v.)

Ordo VI. CRUCIFERÆ. *Juss. gen.* 257. *DC. fl. fr.* 4. p. 641.  
*syst.* 2. p. 139. *prod.* 1. p. 131.

Calyx 4-sepalus, sepalis cruciatis, 2 placentariis (superiore nempe et infer.) oppositis planis angustioribus, 2 lateralibus seu valvaribus latioribus basi concavis gibbis aut rariùs calcaratis, glandulis è toro ortis sepalorum basi imminentibus. Petala 4 hypogyna, sepalis alterna, sæpiùs inter se æqualia, decidua. Stam. 6 è toro orta tetradynama, 2 ante sepala lateraliter sita breviora, 4 longiora antè sepala placentaria geminata interdum inter se coalita. Antheræ biloculares introrsæ. Torus glandulas callosas inter petala et genitalia gerens. Ovarium 1 biloculare, elongatum et tunc stylus subnullus, aut abbreviatus et tunc stylus elongatus. Stigmata 2. Fructus elongatus (tunc *siliqua* dictus) aut abbreviatus (et tunc *silicula*), rariùs 1-locularis evalvis, sæpiùs bilocularis bivalvis, loculis septo longitudinaliter divisus, valvulis planis concavis aut carinatis. Semina in quoque loculo ad utramque suturam adfixa, rarissimè abortu solitaria, exalbuminosa, oleosa. Embryo (1) formâ varius semper plus minùs curvato-plicatus. — Herbæ aut suffrutices. Folia ferè omnium alterna. Racemi oppositifolii aut terminales, sæpiùs ebracteati.

Subordo I. PLEURORHIZÆ (0 =). *DC. syst.* 2. p. 146.

Cotyledones planæ accumbentes. Radicula lateralis nempe rimæ cotyledonum incumbens. Semina compressa interdum marginata.

Trib. I. ARABIDEÆ seu Pleurorhizæ siliquosæ. *DC. l. c.*

Siliqua elongata bivalvis dehiscens, septo lineari, stylo abbreviato, valvulis planis concavis aut subcarinatis. Semina in loculo quoque plurima. Cotyledones planæ septo parallèlæ accumbentes. (0 =)

I. MATHIOLA. *Brown. Kew. ed. 2. vol. 4. p. 119. DC. syst.* 2. p. 162, non L.

Calyx erectus basi bisaccatus; petala unguiculata; stam. libera edentula; siliqua teres vel compressa elongata, stigmatè connivente bilobo, lobis dorso incrassatis vel cornigeris; semina 1-serialia.

Sect. I. PACHYNOTUM. *DC. l. c.*

Petalorum laminæ obovatæ planæ purè purpureæ aut albæ; stigmatum dorsa incrassata aut gibba.

1. *M. INCANA* (Brown. h. Kew. ed. 2. v. 4. p. 119.) caule basi suffruticoso erecto ramoso, foliis lanceolatis integerrimis incanis, siliquis subcylindricis eglandulosis. 5 2/3 in rupibus maritimis Nicæ et insularum Stæchadam reperi,

(1) Embryonis structura faciliùs videtur in seminibus nondum absolutè maturis.

- in hortis frequenter culta. — *Cheiranthus incanus* Lin. sp. 924. DC. fl. fr. 4. p. 656. — Weinn. phyt. t. 643. f. a-e. et 644. f. a-b. Variat flore purp. coccineo albo et varieg., simplici et pleno. (v. v. sp. D.)
2. *M. ANNUA* (DC. syst. 2. p. 165.) caule herbaceo erecto ramoso, foliis lanceolatis obtusis incanis, siliquis subcylindricis eglandulosis. ① in maritimis Europæ australis ex Lin., in hortis freq. culta. — *Cheiranthus annuus* Lin. sp. 925. Fl. fr. 4. p. 656. — J. Bauh. hist. 2. p. 875. f. 1. Var. fl. purp. et albo. (v. v. c.)
3. *M. SINUATA* (Brown. h. Kew. 4. p. 120) caule erectiusculo herbaceo ramoso, foliis oblongis tomentosis, inferioribus sinuatis, siliquis compressis velutinis et glanduloso-muricatis. ② in arenosis maritimis secus Mediterraneum et secus Oceanum in prov. occid. — *Cheiranthus sinuatus* Lin. sp. 926. Fl. fr. 4. p. 657. — Lob. ic. t. 330. f. 2. Flores roseo-purpurei vespere odorati. (v. v.)

Sect. II. LUPERIA. DC. syst. 2. p. 169.

Stigmatis dorsa crassa non cornigera; petala oblonga sordide à flavo purpurea.

4. *M. TRISTIS* (Brown. h. Kew. 4. p. 120.) caule basi suffruticoso ramoso erecto, foliis tomentosis linearibus integris dentatisve, floribus subæsilibus, siliquis teretiusculis. ② in apricis lapideis Galloprovinciæ et rariis in Occidentia. — *Cheiranthus tristis* Lin. sp. 925. Fl. fr. 4. p. 625. excl. var. Sabauda. — Barr. ic. t. 803. Flores vespere odoratissimi. (v. v. DC.)

Sect. III. ACINOTUM. DC. syst. 2. p. 175.

Petala obovata; siliqua apice tricuspidata stigmatum dorsis nempe in cornua exsertis.

5. *M. TRICUSPIDATA* (Brown. l. c.) caule suberecto ramoso, foliis sinuato-pinnatifidis, siliquæ cuspidibus 3 acutis subæqualibus. ① in arenis maritimis secus mare Mediterraneum, nec unquam Candollina vidit secus Oceanum ubi tamen à cl. Laterrade indicatur. — *Cheiranthus tricuspidatus* Lin. sp. 926. Fl. f. 4. p. 655. Habitus et flores *M. sinuata*. (v. v.)

II. CHEIRANTHUS. Brown. kew. ed. 2. v. 4. p. 118. DC. syst. 2. p. 176.

Calyx basi bisaccatus clausus; siliqua teres aut compressa elongata; stigma bilobum aut capitatum, lobis dorso non gibbis; semina 1-serialia ovata compressa, in nostris immarginata.

1. *C. CHEIRI* (Lin. sp. 924.) foliis lanceolatis integerrimis, pube bipartitâ adpressâ aut nullâ, siliquis linearibus, stigmatibus lobis recurvis. ② interdum in muris tectis et lapideis præsertim prov. austral., in hortis frequenter culta. — Fl. fr. 4. p. 637. — Lob. ic. 330. f. 1 et 331. f. 2. Flores flavo-ferruginei, in hort. facile pleni, odorati. (v. v.)

III. NASTURTIUM. DC. syst. 2. p. 187.

Calyx patens æqualis; petala integra interdum 0; stigma subbilobum; siliqua teretiuscula aut abbreviata, valvis concavis enervibus; semina parva irregul. biserialia immarginata.

Sect. I. CARDAMINUM. DC. l. c.

Petala alba calyce majora; siliquæ teretiusculæ; glandulæ 4 ad basim staminum.

1. *N. OFFICINALE* (Brown. h. Kew. 4. p. 110.) foliis pinnatisectis, segmentis ovatis subrotundo-cordatis repandis. ② in rivulis fontibus acuturiginibus Galliæ totius. Culta ad usus pharmaceuticos et culinares in Normannia. — *Sisymbrium nasturtium*. Lin. sp. 916. Fl. fr. 4. p. 661. — Lob. ic. t. 209. f. 1. (v. v.)

## Sect. II. BRACHYLOBOS. DC. l. c.

Petala flava; siliquæ teretiusculæ aut ellipsoideæ; glandulæ tori parvæ.

2. *N. SYLVESTRÆ* (Brown. l. c.) foliis pinnatisectis, segmentis lanceolatis serratis incisive.  $\mathcal{L}$  in pascuis humidis, ruderatis, glareosis secus fluviorum ripas. — *Sisymbrium sylvestre*. Lin. sp. 916. Fl. fr. 4. p. 662. Engl. bot. t. 2324. Radix repens. Petala aurea calyce majora. (v. v.)
3. *N. PALUSTRE* (DC. syst. 2. p. 191.) foliis pinnati-lobatis, auriculis amplexicaulibus ciliatis, lobis confluentibus dentatis glabris, radice fusiformi, petalis calycem æquantibus. ① in inundatis ubique. — *Sisymbrium palustre*. DC. fl. fr. 4. p. 662. Schk. handb. t. 187. — J. Bauh. hist. 2. p. 866. f. 1. (v. v.)
4. *N. PYRÆNAÏCUM* (Brown. l. c.) foliis radicalibus petiolatis obovatis lyratisve, caulinis amplexicaulibus pinnatipartitis, lobis linearibus integris, siliculis ovalibus stylo apiculatis.  $\mathcal{L}$  in pascuis siccis collibus montosisque a Pyrenæis ad Alsatiam, ab Armoraciâ australi ad Galloprov. — *Sisymbrium Pyrenaicum*. Lin. sp. 916. Fl. fr. 4. p. 663. Myagram Pyren. Berger. phyt. t. 143. (v. v. DC.)
5. *N. AMPHIBIUM* (Brown. l. c.) foliis oblongo-lanceolatis pinnatifidis serratisve, radice fibrosâ, petalis calyce majoribus, siliculis ellipsoideis.  $\mathcal{L}$  in aquosis et inundatis Galliæ totius. — *Sisymbrium amphibium*. Lin. sp. 917. Fl. fr. 4. p. 663. Myagram aquaticum Berger. phyt. t. 145. (v. v.)
  - a. *indivisum* (DC.) foliis omnibus integrisculis serratisve. — C. Bauh! prod. 38. f. 1.
  - β. *varifolium* (DC.) foliis aliis serratis, aliis pectinato-pinnatifidis, aliis capillaceo-multifidis. — Lob. ic. 319.

## IV. BARBAREA. Brown. in h. Kew. ed. 2. v. 4. p. 109. DC. syst. 2. p. 205.

Calyx erectus basi subæqualis; petala unguiculata limbo integro; siliqua tetragono-anceps, valvis concavo-carinatis; semina 1-serialia. — Herbarum  $\mathcal{L}$  glaberrimæ. Flores flavi.

1. *B. VULGARIS* (Brown. h. Kew. 4. p. 109.) foliis inferioribus lyratis lobo terminali subrotundo, superioribus obovatis dentatis.  $\mathcal{L}$  in ruderatis humidinaculis, sepius et palustribus Gall. tot. — *Erysimum barbarea*. Lin. sp. 922. var. α et γ. Fl. fr. 4. p. 560. — Lob. ic. t. 207. f. 2. (v. v.)
2. *B. PRÆCOX* (Brown. l. c.) foliis inferioribus lyratis, lobo terminali ovato, superioribus pinnatipartitis, lobis lineari-oblongis integerrimis.  $\mathcal{L}$  in herbis uliginosis Pyrenæorum, Cebennarum, Alpitum austral. et in pratis subsalsis maritimis Armoraciæ. — *Erysimum præcox* Smith! engl. bot. t. 1129. Fl. fr. 4. p. 661. Edulis. (v. v. DC.)

## V. TURRITIS. Dill. nov. gen. giess. p. 120. t. 6. DC. syst. 2. p. 211.

Calyx laxus; pet. unguiculata, limbo oblongo integro; siliqua elongata anceps stricta, valvis planis nervosis; semina in quoque loculo biserialia numerosissima. — Flores albid.

1. *T. GLABRA* (Lin. sp. 930.) foliis radicalibus dentatis pilosis, caulinis amplexicaulibus integerrimis glabris, siliculis strictissimis pedicello sextuplo longioribus. ② in pascuis, siccis, apricis et sylvulis arenosis Gall. tot. — Clus. hist. 2. p. 126. f. 5. *Arabis perfoliata* Fl. fr. 4. p. 675. (v. v.)

## VI. ARABIS. L. gen. n. 818.

Calyx erectus; pet. unguiculata limbo patente integro; siliqua linearis stigmatibus sessilibus coronata, valvulis planiusculis nervosis vel venosis, septo lineari; semina 1-serialia. — Flores albi rarius rosei.

## Sect. I. ALOMATIUM. DC. syst. 2. p. 214.

Semina immarginata, seu parvâ alâ membranaceâ cincta.

§. 1. *Foliis caulinis cordato-amplexicaulibus, petalorum limbis obovatis ab ungue distinctis patentibus.*

1. A. VARNA (Brown. h. Kew. 4. p. 105. non Desf.) foliis caulinis cordatis amplexicaulibus dentatis pube tripartita scabris, pedicellis calyce brevioribus, stigmate subemarginato. ① in arvis agris et asperis prov. Mediterranearum. — *Hesperis verna* Lin. sp. 928. Fl. fr. p. 653. — Barr. ic. t. 875 et 876. Flores parvi purpurei (v. v.)
2. A. ALPINA (Lin. sp. 928) foliis pluridentatis pube ramosa villosis lanceolatis acutis, radicalibus subpetiolatis, caulinis cordato-amplexicaulibus, pedicellis calyce glabriusculo longioribus. ② in rupibus saxosisque apricis montium calcareorum. — Clus. hist. 1. p. 125. f. 1 et 2. Fl. fr. 4. p. 674. Flores albi uti sequentes. (v. v.)

§. 2. *Foliis caulinis cordato-amplexicaulibus, petalis oblongo aut cuneato-linearibus erectis.*

3. A. AURICULATA (Lam. dict. 1. p. 219.) foliis subdentatis pube ramosa scabris, inferioribus ovalibus in petiolum subattenuatis, caulinis obtuse cordato-auriculatis, pedicellis calyce vix longioribus, siliquis maturis à caule divergentibus strictis. ① in collibus, muris, glareosis Delphinatus, Occitanis, Galloprovinciis. — DC. fl. fr. 4. p. 673. syst. 2. p. 220. (v. v.)  
  - a. A. aspera All! ped. auct. 18. t. 2. f. 2.
  - β. A. recta Villd. dauph. 3. p. 319. t. 37.
4. A. SAXATILIS (All. ped. 2. p. 973.) foliis subdentatis pube ramosa scabris, inferioribus in petiolum subattenuatis caulinis acutiusculè cordato-auriculatis, pedicellis demum calyce quadruplò longioribus, siliquis maturis patulis subflexuosis. ① in saxosis rupestribus asperis montium Delphinatus, Galloprovinciis et Pyren. valle Eynes ex cl. Benth. — Fl. fr. 4. p. 647. Ar. nova Vill. dauph. 3. t. 37. (v. v.)
5. A. SAGITTATA (DC. syst. 2. p. 221.) foliis subdentatis pube sæpius ramosa scabris, radicalibus ovatis aut oblongatis in petiolum attenuatis, caulinis lanceolatis sagittato-cordatis, pedicellis calycis longitudine, siliquis erectis strictis. ② in asperis muris, et campis præsertim prov. austral.  
  - a. *Gerardiana* (DC.) foliis scabris, caulinis basi in aurículas productis. — A. sagittata Fl. fr. 5. p. 592. — Ger. gallopr. 367. n. 2. (v. v.)
  - β. *subglabrata* (DC.) foliis fere glabris, caulinis auriculato-sagittatis, racemo composito. — In muris ubi Mont de Marsan legit. L. Dufour. (v. s.)
6. A. HIRSAUTA (DC. Fl. fr. 4. p. 675. 5. p. 592.) foliis dentatis pube sæpius ramosa scabris, radicalibus obovato-oblongis in petiolum attenuatis, caulinis ovato-lanceolatis, pedicellis calycis longitudine, siliquis erectis. ② in campis et asperis Galliæ ferè totius, præsertim australis. — *Turritis hirsuta* Lin. sp. 930. Jacq. ic. var. 1. t. 126. Boiss. fl. eur. t. 456. f. 1. (v. v.)

§. 3. *Foliis caulinis sessilibus, petalis oblongo aut cuneato-linearibus erectis.*

7. A. MURALIS (Bertol! deo. ital. 2. p. 137.) foliis pube ramosa hirsutis, radicalibus spatulatis obtuse dentatis, caulinis ovatis acutè dentatis, racemo stricto, pedicellis calyce demum duplò longioribus. ② in rupibus montanis Cevennarum et Alpium demissiorum, ad Vallemclausam. — Fl. fr. 5. p. 592. (v. v.)
8. A. STRICTA (Huds. angl. 592.) foliis pube sparsa bifida ciliato-scabris, radicalibus obovatis dentatis, caulinis oblongis subintegris, racemo stricto, pedicellis calyce vix longioribus, siliquis laxiusculis. ② in rupibus calcareis apricis Alpium demissiorum et Pyrenæorum. — Fl. fr. 4. p. 677. var.  
  - a. *Turritis Rayi* Vill. dauph. 4. p. 326. t. 38. (v. v.)
9. A. CILIATA (Brown. h. Kew. 4. p. 107.) foliis subdentatis glabris ciliatis, radicalibus subsessilibus ovali-oblongis, caulinis oblongis, racemo stricto,

- pedicellis calycis longitudine. ② in rupibus apricis Alpium et Pyrenæorum ex Benth. — *A. stricta*  $\beta$  Fl. fr. 4. p. 667. priori valdè affinis. Dubia civis. (v. s. sp. ex Helv.)
10. *A. INCANA* (Roth. cat. bot. 1. p. 79.) foliis omnibus sessilibus subdentatis pilis ramosis incanis, radicalibus obovato-oblongis, caulinis oblongis, racemo erecto, pedicellis demum calyce quadruplò longioribus, siliquis strictis. ② in campestribus sticis circa Nevers legit cl. Reynier. — DC. syst. 2. p. 225. (v. s.)
11. *A. ? THALIANA* (Lin. sp. 929.) foliis pilosis subdentatis, radicalibus petiolatis ovato-oblongis, caule ramoso, siliquis laxis adscendentibus, pedicellis calyce multò longioribus. ① frequens in muris, tectis, arenosis, aridis. — Fl. fr. 4. p. 618. *Sisymbrium Thalianum* Gay in ann. sc. natur. avril. 1826. — Thal. herc. 84. t. 7. D. (v. v.)
12. *A. SERPYLLIFOLIA* (Vill. dauph. 3. p. 18. t. 37.) foliis subintegris pilis ramosis ciliato-subscabris, radicalibus caulisque ovalibus basi attenuatis, racemo laxiusculo, pedicellis calycis longitudine. ② in rupestribus montosis Alpium Delphinatù et Galloprovinciæ. — Fl. fr. 4. p. 678. (v. v. DC.)
- §. 4. *Foliis caulinis sessilibus aut petiolatis, petalorum limbo obovato seu cuneato ab ungue distincto.*
13. *A. PETRAEA* (DC. syst. 2. p. 229.) foliis glabris aut pilis simplicibus bifurcatis ciliatis vel scabris, radicalibus incis, caulinis oblongo-linearibus integris, caule sæpius ramoso, siliquis erecto-patulis, stigmatibus capitato.  $\mathcal{U}$  in rupibus et saxosis montium Arvernæ et Pyrenæorum. — Fl. fr. 4. p. 679. *A. runcinata* Lapeyr. ex cl. Benth. (v. s.)
14. *A. ARENOSA* (Scop. carn. ed. 2. n. 837. t. 40.) foliis pube bifurcà villosis, radicalibus lyrato-pinnatifidis, caulinis inciso-dentatis, caule ramoso pilis simplicibus hispidis, pedicellis siliquisque patulis. ① in arenosis vineisque Alsatie et prope Dolam (Cordienne). — *Sisymbrium arenosum* Lin. sp. 919. Fl. fr. 4. p. 668. — C. Bauh! prod. 40. f. 1. Flores rosei. (v. s.)
15. *A. CEBENNENSIS* (DC. syst. 2. p. 234.) foliis omnibus petiolatis angulato-ovatis acuminatis grossè dentatis pube minimà subvelutinis, pedicellis siliquisque patulis. ② in locis asperis subumbrosis montium Cebennorum et Aubrac. — Deless! ic. sel. 2. t. 26. *Hesperis inodora* var. Lin! sp. 927. (excl. syn.) Gou. fl. monsp. 167. Fl. fr. 5. p. 589. Flores pallidè violacei. (v. v. DC.)

## Sect. II. LOMASPORA. DC. syst. 2. p. 234.

Semina marginata seu alà membranacè late cincta.

16. *A. TURBITA* (Lin. sp. 930.) foliis amplexicaulibus acuminatis subdentatis pubescentibus, pedicellis calycis longitudine, siliquis secundis decurvis. ② in sepiibus et dumetis montosis Gall. ferè tot. — Clus. hist. 2. p. 126. f. 2. — Fl. fr. 4. p. 675. Flores albedo-subochrolei. (v. v.)
17. *A. PUMILA* (Jacq. fl. austr. t. 281.) foliis scabris subintegris, radicalibus obovatis, caulinis sessilibus ovatis oblongisve, racemo nutante, pedicellis calyce duplò longioribus, siliquis erectis pedicello triplò longioribus.  $\mathcal{U}$  in asperis lapideis rupestribusve Alpium editiorum Delphin. et Gallopr. — *A. scabra* Fl. fr. 4. p. 677. Flores albi. Folia nunc ferè glabra, nunc pilis tripartitis subsimplicibusve scabra. (v. v. DC.)
18. *A. BELLIDIFOLIA* (Jacq. fl. austr. t. 280.) foliis glabris subintegris, radicalibus obovatis, caulinis sessilibus ovatis, racemo erecto, pedicellis calyce triplò longioribus, siliquis pedicello quadruplò longioribus.  $\mathcal{U}$  in graminosis lapideis Alpium et Pyrenæorum oriental. — Lin. mant. 94. Fl. fr. 4. p. 676. — Clus. hist. 2. p. 129. f. 2. — Fl. albi. (v. v.)
19. *A. CÆRULÆA* (Jacq. coll. 2. p. 56.) foliis glabris subintegris, radicalibus oblongo-obovatis, caulinis paucis oblongis, racemo nutante, pedicellis calycis



longitudine, siliquis erectis.  $\mathcal{Y}$  in asperis et pascuis editioribus Alpium Galloprov. superioris et Delphinatūs. — Fl. fr. 4. p. 678. — Turritis cœrulea All. ped. n. 981. t. 40. f. 2. Tota glabra. Flores ex albo subcœrulei. (v. v. DC.)

VII. CARDAMINE. DC. *syst.* 2. p. 245. — *Cardamines* sp. L.

Calyx basi æqualis; pet. unguiculata, limbo integro; siliquæ sessiles lineares compressæ, valvis planis enerviis dissepimento incrassato subangustioribus sæpius elasticè dehiscentibus, stylo brevi aut subnullo; semina ovata immarginata 1-serialia, funiculis tenuibus. — Flores albi, rarius sublacini.

§. 1. *Foliis ferè omnibus indivisis.*

1. *C. ASARIFOLIA* (Lin. sp. 913.) foliis glabris petiolatis cordato-orbiculatis subsinuato-dentatis, caule erecto, siliquis erectis pedicello duplò longioribus.  $\mathcal{Y}$  in rivulis humidis montanis Alpium Galloprov. suprà vallem Barcinonensem ex Barr. ic. t. 1163. — Fl. fr. 4. p. 681. Radix repens. Flores albi. (v. v. sp. ex Pedemontii montibus Galloprov. conterminis.)
2. *C. BELLIDIFOLIA* (Lin. sp. 913.) foliis glabris crassiusculis, radicalibus petiolatis ovatis integris, caulinis paucis integris aut subbilobis, siliquis erectis, stigmate subsessili.  $\mathcal{Y}$  in fissuris rupium humidis Alpium edit. et Pyrenæorum. — All. ped. n. 949. t. 18. f. 3. *C. alpina* Willd. sp. 3. p. 481. Fl. fr. 4. p. 680. Variat in montibus Lapponiæ petiolis longioribus, in nostris limbo ferè brevioribus, et limbo integro aut trilobo. (v. v.)

§. 2. *Foliis plerisque trilobis.*

3. *C. BRASCHIFOLIA* (Lin. sp. 913.) foliis glabris membranaceis petiolatis, radicalibus indivisis, caulinis in auriculam acutam basi utrinque productis, inferioribus trisectis, superioribus pinnatim 5-lobis, siliquis erectis stylo brevissimo apiculatis.  $\mathcal{Y}$  in umbrosis humidis Alpium, Cebennarum, Pyrenæorum et montium Arvernæ. — Jacq. fl. aust. suppl. t. 31. Fl. fr. 4. p. 680. An fortè prioris var? (v. v.)
4. *C. TRIFOLIA* (Lin. sp. 913.) foliis glabriusculis trisectis, segmentis sessilibus rhombæo-subrotundis dentatis, scapo nudo, ramis infimis radiciformibus repentibus.  $\mathcal{Y}$  in umbrosis sylvaticis montanis Arvernæ ex Delarb. et alibi in Galliâ ex Lam. sed vereor ne var. foliis trilobis *C. amaræ* pro eâ descriptam fuerit. — Fl. fr. 4. p. 682. — Lob. ic. t. 211. f. 1. (v. v. sp. ex Germaniâ et Ital.)

§. 3. *Foliis plerisque pinnatipartitis.*

5. *C. AMARA* (Lin. sp. 915.) foliis pinnatisectis, radicalium segmentis subrotundis, caulinorum dentato-angulatis, stylo filiformi acutiusculo, caule basi radicante.  $\mathcal{Y}$  in aquosis umbrosis et ad rivulos Gall. ferè totius præ. in submontanis. — Fl. fr. 4. p. 683. Engl. bot. t. 1000. Variat foliis trisectis (an *C. trifolia* Delarb?) et subvillosis (*C. umbrosa* Lejeun. spa. 2. p. 63.) (v. v.)
6. *C. PRATENSI* (Lin. sp. 913.) foliis pinnatisectis, segmentis radicalium subrotundis, caulinorum linearibus lanceolatisve integris, stylo brevissimo viz siliquâ tenuiore, stigmate capitato.  $\mathcal{Y}$  in pratis humidiusculis Galliæ totius frequens. — Fl. fr. 4. p. 684. — Lob. ic. t. 210. f. 1. et fors. f. 2? Flores lilacini rarius albi. (v. v.)
7. *C. HIRSA* (Lin. sp. 915.) foliis pinnatisectis, radicalium segmentis subrotundis petiolulatis, superiorum oblongis subsessilibus, petalis oblongis, siliquis erectis. ① in sylvaticis umbrosis asperis sepibus Galliæ totius. — Fl. fr. 4. p. 684. — Barr. ic. t. 455. Variat foliis subhirsutis et glabriusculis. Hujus cl. Saint-Hilaire observ. monstrum calycibus 3 flores tetrandros gerentibus. (v. v.)
8. *C. SYLVATICA* (Link. in Hoffm. blatt. 1. p. 50.) foliis pinnatisectis, seg-

- mentia irregulariter sinuato-dentatis mucronatis, petalis oblongis, siliquis patentibus latitudine stylum æquantibus glabris, rachi hirsuta. ① in sylvaticis Galliæ ferè totius sed cum priore, forsàn non immeritò, sæpè confusa. — *C. impatiens* fl. dan. t. 735. (v. s.)
9. *C. UMBROSA* (Andrz. in DC. syst. 2. p. 260.) foliis pinnatisectis, segmentis integris mucronatis, petalis oblongis, siliquis subpatulis stylum crassum latitudine duplò superantibus rachique glaberrimis. ① in Cebennorum montium sylvaticis et forsàn ubique cum 2 prioribus confusa. (v. v.)
10. *C. PARVIFLORA* (Lin. sp. 919.) foliis pinnatipartitis, lobis sessilibus oblongis linearibusve integris, infimis a caule remotis, petalis oblongo-linearibus, pedicellis subpatulis, siliquis erectis. ① in herbosis humidis Galloprov., Occitanis, et prov. occident. — Fl. fr. 4. p. 685. — Gmel. fl. sib. 3. t. 64. *C. hirsutæ* affinis sed gracilior glabra, etc. (v. v.)
11. *C. IMPATIENS* (Lin. sp. 914.) foliis pinnatisectis, segmentis ovali-oblongis subdentatis, infimis cauli approximatis acutis stipulæformibus, ① in umbrosis sylvaticis montosis Gall. ferè totius. — J. Bauh. hist. 2. p. 886. f. 1. Fl. fr. 4. p. 685. Variat petalis nullis aut parvis oblongis, foliis glabris aut subciliatis. (v. v.)
12. *C. LATIFOLIA* (Vahl. symb. 2. p. 77.) foliis pinnatisectis glabris, segmentis 3-7 suborbiculatis dentato-angulatis, siliquis erectis pedicello paulò longioribus stylo acuminatis. ② secus rivulos et in humidis umbrosis Pyrenæorum orient. et central. — Herm. parad. p. 203. t. 69. — Fl. fr. 4. p. 683. Flores purpurascetes magn. *C. pratensis*. (v. v. DC.)
13. *C. THALICTROIDES* (All! ped. n. 951. t. 57. f. 1.) foliis petiolatis glabris, superioribus pinnatisectis, segmentis 3-5 petiolulatis ovatis trilobatis, caule debili, siliquis patulis. ② in sylvaticis asperis subhumidis Alpium Delphinatús. — Fl. fr. 4. p. 681. *C. Plumieri* Vill. dauph. 3. t. 38. Flores albi cum ungue subflavescente, nec flavi ut incautè dixit All. (v. v. DC.)

## VIII. PTERONEURUM. DC. syst. 2. p. 269.

Calyx basi æqualis; pet. unguiculata integra; siliqua sessilis lanceolaris, valvis planis septo angustioribus enerviis à basi elasticè dissilientibus, placentis extús alatis nervo acuto secus stylum ideò ancipitem producto; funiculi umbilicales alato-dilatati.

1. *P. GRÆCUM* (DC. l. c.) foliorum segmentis petiolulatis suborbiculatis dentato-lobatis subæqualibus. ① in umbrosis humidis Corsicæ teste Boccone Sic. p. 84. t. 44. f. N. et t. 45. f. 2. mus. p. 171. t. 116. — *Cardamine græca* Lin! sp. 915. Fl. fr. 4. p. 683. Fl. albi magn. *Cardamines amara*. (v. s. sp. ex Italiâ.)

## IX. DENTARIA. Tourn. inst. t. 111. L. gen. n. 811.

Calyx basi æqualis erectus; pet. unguiculata limbo obovato aut orbicordato; siliqua sessilis lanceolata apice attenuata compressa, valvis planis enerviis septo angustioribus elasticè sæpè revolutis, stylo filiformi, stigmate subsimplici; funiculi umbilicales alato-dilatati; semina ovata immarginata 1-serialia. — Trunci radicales carnosì horizontales dentati. Folia petiolata secta.

1. *D. DIGITATA* (Lam. dict. 2. p. 268.) foliis caulinis plurimis alternis petiolatis palmatis 5-sectis, segmentis oblongo-lanceolatis acuminatis grossè serratis. ② in sylvaticis umbrosis montanis Gall. ferè totius. — Clus. hist. 2. p. 122. f. 1. 2. Fl. fr. 4. p. 686. — *D. pentaphyllos* β et γ Lin! sp. 912. Flor. albi aut subpurpurascetes. (v. v.)
2. *D. PINNATA* (Lam. ill. t. 562 f. 1.) foliis caulinis alternis pinnatisectis, segmentis oblongis acuminatis serrato-dentatis. ② in umbrosis sylvaticis

montanis Gall. ferè tot. — Cus. hist. 2. p. 123. ic. Fl. fr. 4. p. 687. D. pentaphyllos & Lin. sp. 912. flores prioris. (v. v.)

3. *D. bulbifera* (Lin. sp. 912.) foliis caulinis alternis pinnatisectis, summis indivisis, plerisque in axillis bulbiferis ♀ ad radices montium, in locis humentibus umbrosis in Arvernâ, Alsatiâ, Delphinatâ, circa Parisios, Abbativillam, etc. — Fl. fr. 4. p. 687. Cus. hist. 2. p. 121. f. 1. (v. s.).

Trib. II. *ALYSSINÆ* seu *Pleurorhizæ* latiseptæ. DC. syst. 2. p. 280.

Silicula bi- (rariùs uni-) locularis, bivalvis, ovata aut oblonga, compressa aut turgida, valvis planis aut concavis non carinatis, septo ovato aut oblongo in majore fructus diametro. Semina ovata aut compressa. Cotyledones planæ accumbentes. (o =)

X. *LUNARIA*. L. gen. n. 809. DC. syst. 2. p. 280.

Calyx clausus basi bisaccatus; petala unguiculata, limbo obovato; stam. libera edentula; silicula pedicellata elliptica aut oblonga, placentis nerviformibus marginata, plana, bilocul., septo membranaceo persistente, valvis planis enerviis, funiculis longis septo adnatis, stylo filiformi; semina distantia marginata.

1. *L. rediviva* (Lin. sp. 911) siliquis lanceolatis utrinquè attenuatis. ♀ in sylvis montanis Alpium, Juraasi, Vogesorum, Pyrenæorum, etc. — Fl. fr. 4. p. 688. — Lob. ic. t. 278. f. 1. Flores purpurascens odor. (v. v.)

β. *alpina* (DC.) foliis superioribus minus cordatis, siliculis angustioribus. — *L. alpina* Berg. phyt. 3. t. 115. In Pyrenæis et Vogesis. An spec. prop? (v. v.)

2. *L. annua* (Moench. meth. 126.) siliculis ellipticis utrinquè obtusis. ② in sylvis montanis Alsatiæ, et Galloprovinciæ ex Gerard. In hortis sæpè culta. — Lob. ic. t. 322. f. 2. *L. annua* Lin. sp. 911. Fl. fr. 4. p. 688. Flores inodori violaceo-lilacini aut albi. (v. v. c.)

XI. *FARSETIA*. Desv. journ. bot. 3. p. 173. DC. syst. 2. p. 286.

Calyx erectus basi bisaccatus; pet. unguiculata, limbo integriusculo; stam. minora dente aucta; silicula stylo brevi apiculata, elliptica, sessilis, compresso-plana, septo membranaceo elliptico, valvulis planis; semina horizontalia compressa suborbiculata marginata.

1. *F. clypeata* (Brown. h. Kew. 4. p. 96.) caulibus herbaceis erectis, foliis oblongis repandis, siliculis pube brevi velutinis, stigmatibus capitatis. ② in collibus saxosis montanisque propè *S.-Amand* in ditone dictâ *Cher* ex cl. Gay, circa Monspelium ex Gouan, et Nicæam ex All., sed mihi dubia civis. — Lob. ic. t. 323. f. 1. *Alyssum clypeatum* Lin. sp. 909. Fl. fr. 4. p. 696. Flores flavi. (v. s. sp. ex Oriente.)

XII. *BERTEROA*. DC. syst. 2. p. 286.

Calyx erectus basi æqualis; pet. unguiculata, limbo bilobo; stam. libera, 2 minora basi intus dentata; silicula bilocularis elliptica, valvis convexo, planis membranaceis, septo elliptico, stylo persistente; semina ovalia, plana submarginata.

1. *B. incana* (DC. l. c.) siliculis pubescentibus subventricosis. ② in rudatis arenosis apricis inultis in Alsatiâ et Galloprovinciâ. — Lob. ic. 216. f. 2.

*Alyssum incanum* Lin. sp. 978. Fl. fr. 4. p. 694. Herba erecta incana, fol. oblongo-lin., flor. albis. (v. v. DC.)

XIII. VESICARIA. Lam. ill. t. 559. DC. syst. 2. p. 295.

Calyx clausus; pet. unguiculata limbo integriusculo; stam. libera; silicula globosa inflata immarginata dehiscens stylo apiculata, valvis membranaceis convexissimis, septo membranaceo; semina 4-6 in quoque loculo.

1. *V. UTRICULATA* (Lam. ill. t. 559. f. 1.) calyce bisaccato, foliis oblongis integerrimis glabris, inferioribus ciliatis subspathulatis. ♂ in rupibus calcareis Delphinatus et in Burgundiâ propè Semur ex Dur. — Fl. fr. 4. p. 696. — Barr. ic. t. 883. *Alyssum utriculatum* Lin. mant. 265. Flor. flavi. Stam. edentula. (v. v. sp. in Vallesii. D.)

XIV. ALYSSUM. DC. syst. 2. p. 301. — *Alyssi* sp. L.

Calyx æqualis; pet. unguiculata limbo integriusculo; stam. filamenta omnia aut nonnulla dentata; silicula orbiculata ovalisve stylo tenui apiculata, compressa, valvis planis aut centro convexis, loculis 1-2-spermis, septo tenuissimo; semina ovalia compressa.

Sect. I. ADYSETON. DC. l. c.

Flores flavi; stamina dentata; semina marginata.

1. *A. CORNICUM* (Duby Mss.) caulibus basi suffruticosis glabris ramosissimis, ramis sterilibus foliisque inferioribus spathulato-ovatis utrinquè argenteis, siliculis orbiculatis compressis glaberrimis lucidis, seminibus subimmarginatis. ♂ in Corsicâ circa Bastia. (v. s.)
2. *A. ALPESÆ* (Lin. mant. 92.) caulibus basi suffruticosis diffusis canescentibus, foliis obovatis incanis, racemis simplicibus, siliculis obovato-oblongis canescentibus, valvis planis. ♂ ♀ in scopulis apricis montanis Alpium Gallo provincie, Delphinatus, in Lotharingiâ ex Will. et circa Senlis ex Thuil. — Ger. Galloprov. 352. t. 13. f. 2. Fl. fr. 4. p. 693. (v. v.)
3. *A. MONTANUM* (Lin. sp. 907.) caulibus subherbaceis diffusis pubescentibus, foliis subincanis, infer. obovatis, super. oblongis, racemis simplicibus, siliculis orbiculatis subemarginatis canescentibus. ♀ in collibus apricis montibusque demissioribus calcareis Gall. ferè tot. — Fl. fr. 4. p. 694. — Lob. ic. t. 220. f. 1. (v. v.)  
*β. arenarium*, foliis rotundioribus. — *A. arenarium* Lois. Fl. gall. 401. not. 96. (v. v.)
4. *A. DIFFUSUM* (Tenor. app. h. neap. 1815. p. 58.) caulibus subherbaceis diffusis canescentibus, foliis incanis, infimis subobovatis, cæteris lineari-lanceolatis utrinquè attenuatis, racemis simplicibus, siliculis ovatis subemarginatis stylo subæqualibus. ♀ in summis Pyrenæis orient. monte dicto Cambres d'Aze. — *A. montanum* var. Benth. cat. 59. (v. v. DC.)
5. *A. CAMPESTRÆ* (Lin. sp. 909.) caulibus herbaceis diffusis, foliis lanceolatis sublinearibusve hirsutis, siliculis orbiculatis tuberculato-hirsutis longè racemosis stylo sextuplò longioribus. (1) in campis arenosis apricis Gallie ferè totius, præsertim australis. — Fl. fr. 4. p. 694. — Barr. ic. t. 912. f. 2. (v. v.)
6. *A. CALYXIMUM* (Lin. sp. 908.) caulibus herbaceis diffusis, foliis lineari-lanceolatis canescentibus, calycibus persistentibus, siliculis orbiculatis subemarginatis pubescentibus stylo quadruplò longioribus. (1) in campis siccis apricis arenosis lapidosis Gall. ferè tot.; præsertim australis. — Fl. fr. 4. p. 694. — Cam. epit. 558. f. 1. (v. v.)

Sect. II. LOBULARIA. DC. syst. 2. p. 318.

Flores albi; semina in loculis sæpius solitaria; stam. edentula.

7. *A. MARITIMUM* (Lam. dict. 1. p. 98.) caulibus basi suffrutescentibus procumbentibus, foliis lineari-lanceolatis acutis subincanis, siliculis ovalibus glabris stylo brevi apiculatis. ♂ ♀ in arenosis maritimis et apricis ruderalis secus mare Mediterraneum et secus Rhodanum usque ad Aransonem. — Fl. fr. 4. p. 592. — Barr. ic. t. 844. t. 907. f. 1. et 908. f. 1. — *A. minimum*, *A. halimifolium* et *Clypeola maritima* Lin. (v. v.)
8. *A. SPINOSUM* (L. sp. 907.) caule fruticoso, ramis pedunculisque senescentibus spinosis, foliis oblongo-linearibus argenteis, siliculis orbiculatis glabris stylo brevi apiculatis. ♂ in rupestribus calcareis apricis regionis olearum (et circa Epernon et Senlis. ex Mér.?). — Fl. fr. 4. p. 692. — Barr. ic. t. 808. (v. v.)
9. *A. MACROCARPUM* (DC. syst. 2. p. 321.) caule fruticoso ramoso subspinescente, foliis oblongis obtusis argenteis, siliculis obovato-orbiculatis subemarginatis glabris stylo apiculatis 8-ovulatis. ♂ in rupibus calcareis montium regionem olearum circumdantium, nempe in Cebennis propè Roquelaure et Mende, in Pyrenæis orient. propè Villefranche. — Deless. ic. sel. 2. t. 41. (v. v. DC.)
10. *A. PYRÆNAÏCUM* (Lapeyr. abr. 371. suppl. 91.) caulibus fruticosis ramosis inermibus, foliis obovatis basi attenuatis cano-tomentosis, siliculis ellipticis villosulis stylo longè acuminatis. ♂ in rupibus Pyrenæorum orientalium ad montem Conat legit Xatard. — DC. syst. 2. p. 322. *A. Halimifolium* Fr. Fl. fr. 5. p. 594. (v. s.)

XV. CLYPEOLA. *Garten. fruct. 2. f. 141. f. 6. DC. syst. 2. p. 326.*

Calyx basi æqualis; pet. integra; stam. filamenta dentata; silicula orbicularis plana subemarginata indehiscens 1-loc.-1-sperma; stigma sessile; semen lateraliter ope funiculi horizontalis adfixum, centrale, compressum.

1. *C. JONTHLASPE* (Lin. sp. 910.) siliculis margine ciliatis, caulibus diffusis aut adscendentibus. ♀ in sabulosis apricis, muris, sterilibus calcareis aut gypsaceis prov. oleiferarum, et in Arvernâ et Delph. infer. — Fl. fr. 4. p. 630. — Lob. ic. t. 215. f. 1. Flores flavi minimi. (v. v.)

XVI. PETROCALLIS. *Brown. DC. syst. 2. p. 330.*

Cal. æqualis; pet. integra; stam. edentula; stylus brevissimus; silicula ovalis, septo elliptico membranaceo, valvis planiusculis medio nervigeris, loculis 2-spermis; funiculi umbilicales septo adnati? sem. immarginata.

1. *P. PYRÆNAÏCA* (Brown. h. Kew. 4. p. 93.) ♀ in rupibus apricis inter saxa et lapides Pyrenæorum, Alpium Galloprov. et Delphin. — *Draba Pyrenaica* Lin. sp. 896. Fl. fr. 4. p. 698. All. ped. t. 8. f. 1. Flores rosei. Folia cuneata apice 3-5-fida. (v. v.)

XVII. DRABA. *DC. syst. 2. p. 331. — Drabæ sp. L.*

Cal. basi æqualis suberectus; pet. integra; stam. edentula; silicula ovalis aut oblonga, integra valvis planiusculis, septo tenui valvarum latitudine; sem. oo biseriata immarginata.

Sect. 1. AIZOPSIS. *DC. l. c.*

Stylus filiformis; pet. apice retusa; silicula apice attenuata. — Folia rigida nervo medio carinata, pilis simplicibus ciliata; flores flavi. Herbæ ♀.

1. *D. AIZOPSIS* (Lin. mant. 91.) scapis nudis glabris, foliis rigidis lanceolato-linearibus carinatis ciliatis, staminibus petala subæquantibus, stylo latitudinem siliculæ æquante. ♀ in apricis rupestribus glareosis Alpium, Jurassi, montium Arvernæ, Pyrenæorum. — Fl. fr. 4. p. 697. — Lob. ic. 381. f. 1. (v. v.)

- β. minor* (DC.) foliis brevissimis, siliculâ constanter glabrâ. — *D. ciliaris*. Fl. fr. 4. p. 697. (v. v.)  
*γ. diffusa* (DC.) caulibus elongatis diffusis. — Ger. Gallopr. 344. t. 13. f. 1. — *D. ciliaris* Lin. mant. 91. (v. s.)

## Sect. II. LEUCODRABA. DC. l. c.

Stylus brevissimus; pet. obtusa aut subemarginata; silicula ovali-oblonga. — Folia mollinescula subradicalia; pili ramosi. Flores albi; scapi subnudi. Herbar. 2.

2. *D. NIVALIS* (Willd. sp. 3. p. 427.) scapis nudis monophyllisve glabris, foliis oblongo-linearibus ciliatis subpilosus, siliculis elliptico-oblongis pedicellisque glaberrimis. 2. in rupestribus editioribus Alpium Delphinatus, ad montem Galibier, etc. — DC. syst. 2. p. 344. Fl. fr. p. 699. excl. var. Pyren. — Fl. dan. t. 142. (v. v. DC.)  
 3. *D. TOMENTOSA* (Wahlenb. helv. 123. n. 672. t. 3.) scapo 1-2-phylo velutino, foliis ovali-oblongis pube stellatâ brevi tomentosis, pedicellis pubescentibus, siliculis ovalibus ciliatis. 2. in rupibus Pyrenæorum centralium ad vallis d'Aure montem Penne Blanque legit cl. Marchand. — DC. syst. 2. p. 346. (v. v.)  
 4. *D. STELLATA* (Jacq. obs. n. 54. t. 4. f. 3.) scapis 1-phyllis pubescentibus, foliis ovali-oblongis pube stellatâ brevi tomentosis, pedicellis puberulis, siliculis oblongis. 3. in fissuris rupium et scopulis Pyrenæorum et Alpium. — Fl. fr. 4. p. 699. Deless. ic. sel. 2. t. 46. f. B. D. hirta Vill. dauph. 3. p. 282. (v. v.)  
*β. hebecarpa* (DC.) siliculis undiquè velutino-pubescentibus. DC. syst. 2. p. 346. Hab. cum var. *α*. (v. v.)  
 5. *D. LEVIGATA* (DC. syst. 2. p. 346.) scapis nudis 1-phyllisve pubescentibus, foliis ovatis pube stellatâ brevi tomentosis, siliculis elongato-linearibus pedicellisque glaberrimis. 2. in rupibus Pyrenæorum centralium editiorum ad Port d'Oo et montem Maladetta. — Deless. ic. sel. 2. t. 46. f. A. (v. v.)

## Sect. III. HOLARGES. DC. syst. 2. p. 348.

Siliculæ elongatæ stylo distincto brevi apiculatæ. — Herbar. ① aut ②; caules foliosi sæpius ramosi; flores albi.

6. *D. CONTORTA* (Ehr. beitr. 7. p. 155.) caule folioso ramoso pube stellatâ velutino, foliis ovatis dentatis, siliculis oblongis glabris subcontortis. ① ② in rupibus calcareis montanis Pyrenæorum. — DC. syst. 2. p. 348. *D. incana* var. *β*. Lin. sp. 897. Fl. fr. 4. p. 700. Engl. bot. t. 388. (v. v.)

## Sect. IV. DRABELLA. DC. syst. 2. p. 351.

Siliculæ elliptico-oblongæ, stigmate sessili. — Herbar. ①; caules foliosi ramosi; flores minimi.

7. *D. NEMORALIS* (Ehr. beitr. 7. p. 154. non Delarb.) caule ramoso folioso pubescente, foliis ovatis dentatis pubescentibus, siliculis elliptico-oblongis 32-36-spermis pube minimâ velutinis. ① in Pyrenæis ad montem Canigon legit Thomas. — *D. nemorosa* Houtt. pfl. 4. t. 60. f. 1. non All. Petala flava. (v. s.)  
 8. *D. MURALIS* (Lin. sp. ed. 1. p. 643.) caule ramoso folioso pubescente, foliis ovatis dentatis subcordato-amplexicaulibus subhirsutis, siliculis elliptico-oblongis glabris 12-16-spermis. ① in umbrosis, muris, rupestribus prov., præsertim australium, circa Monapellum, etc. — Fl. fr. 4. p. 699. — J. Bauh. hist. 2. p. 939. f. 1. (v. v.)

## XVIII. EROPHILA. DC. syst. 2. p. 356. — Drabæ sp. L.

Cal. æqualis laxiusculus; petala bipartita; stam. edentula; silicula ovalis aut oblonga, septo membranaceo, valvis planiusculis, stigmate sessili; semina parva immarginata in quoque loculo ∞ biseriata.

1. *E. VULGARIS* (DC. l. c.) siliculis ellipticis pedicellis brevioribus, scapis 5-15-floris.

① in maris, aggeribus, pascuis, aridis, apricis primo vere vulgaris. — *Draba verna* Lin. sp. 896. Fl. fr. 4. p. 698. — Lob. ic. t. 469. f. 1. flor. albi. (v. v.)

XIX. COCHLEARIA. *Tourn. inst. t. 101. L. gen. 803.*

Cal. patens basi æqualis, sepalis concavis; pet. limbo obovato obtuso; stam. edentula; silicula globosa ovata aut oblonga, septo tenui, valvulis ventricosus crassiusculis, loculis rarius di-sæpius polyspermis; stylus brevissimis; semina immarginata. — Flores albi.

SECT. I. KERNERA. *DC. syst. 2. p. 359.*

Silicula globosa, valvis rigidioribus, stylo brevi, stigmatē bilobo.

1. *C. SAXATILIS* (Lam. dict. 2. p. 165.) siliculis globosis lævibus, foliis radicalibus oblongis dentatis pilosis, caulinis lineari-oblongis. ½ in rupestribus et asperis præsertim calcareis montanis prov. præsertim australium. — *Myagrum saxatile* Lin. sp. 894. Fl. fr. 4. p. 718. — Cam. epit. 338. ic. (v. v.)

*A. incisa* (DC.) foliis lyratis pinnatifidisque. — In Jurasso, etc. (v. s.)

2. *C. AURICULATA* (Lam. dict. 2. p. 165.) siliculis globosis lævibus, foliis radicalibus oblongis dentatis pilosis, caulinis cordato-sagittatis. ½ in rupestribus calcareis Pyrenæorum, Cebennarum, Arvernæ. — *Myagrum montanum* Berg. phyt. 3. t. 140. *Myagrum auriculatum* Fl. fr. 5. p. 597. (v. v. DC.)

SECT. II. ARMORACIA. *Rupp. jen. p. 87. DC. l. c.*

Silicula ellipsoidea, stylo brevi filiformi, stigmatē capitato ferè disciformi.

3. *C. ARMORACIA* (Lin. sp. 904.) siliculis ellipsoideis, foliis radicalibus oblongis crenatis, caulinis elongato-lanceolatis dentatis incisive, radice carnosā maximā. ½ in humidis Delphinatū ex Vill., Lotharingiæ ex Willem., Tolosæ ex Tournon, agri Syrtici propè la Teste ex Laterrade, forsā Armoraciæ ex nomine, mihi nunquā obvia nisi in hortis ubi frequenter culta ob radicem culinariam. — Fl. fr. 4. p. 701. — Fuchs. hist. 660. ic. (v. v. c.)

SECT. III. COCHLEAR. *DC. l. c. p. 362.*

Silicula globosa ovata aut oblonga, apice non emarginata, stigmatē subsessili coronata.

4. *C. ELASTIFOLIA* (Lin. sp. 904.) siliculis subglobosis, foliis caulinis cordato-sagittatis amplexicaulibus acuminatis integerrimis. ① ② in agris et cultis Delphinatū propè Sisteron ex Vill., Corsicæ. — Fl. fr. 4. p. 702. — Lob. ic. t. 321. f. 2. (v. s.)
5. *C. ANGLICA* (Lin. sp. 903.) siliculis ovato-subglobosis venoso-reticulatis, pedicello dimidiō brevioribus, foliis radicalibus petiolatis integris, caulinis oblongis. ½ in littoribus maritimis Armoraciæ obs. cl. Bonnemaïson, Bardigalæ Laterr., ad ostium sequanæ cl. Le Prévost. — DC. syst. 2. p. 364. — Lob. ic. t. 294. f. 1. (v. s.)
6. *C. OFFICINALIS* (Lin. sp. 903.) siliculis ovato-globosis pedicello dimidiō brevioribus, foliis radicalibus petiolatis cordatis, caulinis ovatis dentato-angulatis. ½ in littoribus maritimis cœnosis petrosive Belgii gallici circa Dnnkerque, Neustriæ, etc. — Fl. fr. 4. p. 700. — Lob. ic. t. 293. f. 2. (v. v.)
7. *C. PYRÆNICA* (DC. syst. 2. p. 365.) siliculis obovatis subglobosis pedicelli longitudine, foliis radicalibus cordato-reniformibus integris, caulinis paucis ovatis dentatis ½ secus rivulos in Pyrenæis centralibus editioribus ad montem Néouvielle, — Deless. ic. sel. 2. t. 48. *C. officinalis* Lapeyr. abr. 368. (v. v. DC.)
8. *C. DANICA* (Lin. sp. 903.) siliculis ellipsoideis pedicelli longitudine, foliis omnibus petiolatis subdeltoideis. ½ in maritimis cœnosis petrosive secus Oceanum hinc inde a Pictaviā ad Belgium. — Fl. fr. 4. p. 701. — Lob. ic. t. 615. f. 2. (v. v. DC.)

Trib. III. *THLAPSIDEÆ* seu *Pleurorrhizæ* angustiseptæ.  
*DC. prod.* 1. p. 175.

Silicula bilocularis bivalvis, septo angustissimo lineari, valvis carinatis aut navicularibus. Semina ovalia compressa sæpè marginata.

XX. *THLASPI*. *Dill. giss.* p. 123. t. 6. *DC. syst.* 2. p. 373.

Calyx basi æqualis; pet. æqualia integra; stam. edentula; silicula depressa apice emarginata, septo ovali-oblongo, stylo filiformi aut brevissimo, valvis navicularibus dorso membranâ foliaceâ alatis; semina in quoque loculo oo. rarius 2, ovata, immarginata. — Flores albi.

Sect. I. *NOMISMA*. *DC. l. c.*

Silicula plana suborbiculata, sinn angust apice emarginata, stylo subnullo, valvis compressissimis dorso per totam longitudinem alatis; semina in quoque loculo 6-8, striis curvis periphærieis insculpta.

1. *T. ARVENSE* (Lin. sp. 901.) foliis oblongis dentatis, caulibus erectis, siliculis obovato-orbiculatis pedicello brevioribus.  $\frac{1}{2}$  in arvis, agris, cultis et arenosis Galliæ totius. — Fl. fr. 4. p. 709. Lam. ill. t. 557. f. 1. (v. v.)

Sect. II. *PTEROTROPIS*. *DC. l. c. p.* 377.

Silicula subobovata emarginata aut truncata; valvis dorso in alam expansis; semina non striata.

2. *T. ALLIACEUM* (Lin. sp. 901.) foliis oblongis obtusis subdentatis, infer. petiolatis, super. amplexiculis sagittato-amplexicaulibus, siliculis subobovatis ventricosis, stigmate subsessili. (1) in agris et arvis agri Lugdunensis ex Latour, Burgundiæ ex Dur., Lotharingiæ ex Buch., Armoraciæ ex Aubri. — Fl. fr. 4. p. 710. — J. Bauh. hist. 2. p. 923. f. 3. (v. s.)
3. *T. FRASOLIATUM* (Lin. sp. 902.) foliis subdentatis, radicalibus petiolatis, caulibus cordato-amplexicaulibus, caule ramoso, petalis calyci æqualibus, siliculis obcordatis 8-spermis, stigmate subsessili. (2) in agris et arvis apricis prov. præsertim calcareorum. — Fl. fr. 4. p. 710. — Barr. ic. t. 915. (v. v.)
4. *T. MONTANUM* (Lin. sp. 902.) foliis subcaernosis integris, radicalibus obovatis petiolatis, caulibus oblongis sagittato-amplexicaulibus, petalis calyce majoribus, siliculis obcordatis 4-spermis, stylo filiformi.  $\frac{1}{2}$  in montanis apricis Pyrenæorum, Cebennarum, Jurassi, Alpium. — Fl. fr. 4. p. 711. — J. Bauh. hist. 2. p. 928. ic. (v. v.)
5. *T. ALPESTRÆ* (Lin. sp. 903.) foliis subintegris, radicalibus ovatis petiolatis, caulibus oblongis amplexicaulibus, petalis calyci subæqualibus, siliculis obcordatis 8-12-spermis, stylo filiformi.  $\frac{1}{2}$  in agris et pratis montanis Pyrenæorum, Cebennarum, Alpium. — Fl. fr. 4. p. 711. *T. pumilum*. 111. Clus. hist. 2. p. 131. ic. (v. v.)  
 $\beta$ . *arenarium*, scapo subnudo, pedicellis inferioribus elongatis. In sylvis sabulosis agri Syrtici reperiit Des Moulins. (v. s.)
6. *T. HETEROPHYLLUM* (DC. fl. fr. 4. p. 712.) foliis radicalibus petiolatis patulis integris dentatis et lyratis, caulibus erectis cordato-sagittatis, petalis calyci subæqualibus.  $\frac{1}{2}$  in Pyrenæis forsan Hispanicis legit. cl. Clemente. *Lepidium heterophyllum* Benth? cat. 95? qui plantæ suæ tribuit siliculas ellipticas alatas vix emarginatas glabras et stylum exsertum filiformem. (v. s. olim in herb. Claron DC.)

XXI. *HUTCHINSIA*. *Brown. Kew. v.* 4. p. 82. *DC. syst.* 2. p. 384.

Calyx erectus æqualis; pet. æqualia integra; stam. edentula; silicula oblonga aut elliptica, apice acuta aut truncata, depressa, utrinquè m-



## CRUCIFERÆ.

39

tegra; valvis navicularibus apteris, septo oblongo, loculis 2-4-rarissimè 6-spermis; semina è quaque placenta alterna pendula.

### Sect. I. IBERIDELLA. DC. l. c.

Stylus filiformis. — Folia integra aut subdentata.

1. *H. ROTUNDIFOLIA* (Brown. h. Kew. 4. p. 82.) foliis subcarnosis integerrimis, infimis petiolatis obovatis, caulinis ovato-oblongis subamplexicaulibus, staminibus petalis et stylo siliculâ dimidiò brevioribus. ½ inter lapides et in rupium fissuris prope nives in Alpibus Galloprov. et Delph. — Iberis rotundifolia. Lin. sp. 905. *Lepidium rotundifolium*. All. ped. t. 55. f. 2. Fl. fr. 4. p. 706. Flores rosei. (v. v.)
2. *H. BREVISETA* (DC. syst. 2. p. 387.) foliis subcarnosis petiolatis obovatis subdentatis, siliculâ obovato-cuneatâ truncato-emarginatâ, stylo brevissimo. ½ in Corsicâ montibus Bastilica reperiunt cl. Soleirol et Ph. Thomas. — *H. pygmæa* Viv. Cors. App. 3. Fl. parvi albi. (v. s.)

### Sect. II. NASTURTIODEA. DC. l. c.

Stylus brevissimus aut nullus. — Folia pinnato-lobata. Flores albi.

3. *HP ALPINA* (Brown. l. c.) foliis pinnatipartitis glabris, petalis calyce deciduo duplò longioribus, siliculis utrinquè acutis, stylo brevissimo exserto. ½ in rupibus humidiusculis et scopulis Alpium et Pyrenæorum editiorum. — *Lepidium Alpinum* Lin. amen. 4. p. 321. Fl. fr. 4. p. 705. — Clus. hist. 2. p. 128. f. 1. Verum *Lepidium* esse affirmat cl. Gay in ann. sc. nat. avr. 1826. (v. v.)
4. *H. PETRAEA* (Brown. l. c.) foliis pinnatipartitis glabris, petalis calyce vix longioribus, siliculis utrinquè obtusis 4-spermis, stigmate sessili. ① in petrosis, lapidosis, arenosis submontanis apricis Gall. ferè totius. — *Lepidium petraeum* Lin. sp. 899. Fl. fr. 4. p. 706. Boiss. fl. eur. t. 440. f. 1. (v. v.)
5. *HP PROCUMBENS* (Desv. journ. 3. p. 168.) foliis sinuato-pinnatifidis aut subintegris, petalis vix calyce majoribus, siliculis utrinquè obtusis 10-12-spermis, stigmate sessili. ① in ruderatis maritimis secùs vias et semitas ad oras salinarum in prov. præsertim australibus. — *Lepidium procumbens* Lin. sp. 898. Fl. fr. 4. p. 706. — Magn. bot. Monsp. 184. ic. Verum *Lepidium* contendit cl. Gay l. c. (v. v.)

β. foliis subintegris. ① Masailie. (v. s.)

XXII. *TEESDALIA*. Brown. Kew. ed. 2. v. 4. p. 83. DC. syst. 2. p. 391. — *Guepinia*. Bast.

Calyx deciduus sepelis basi tamen subcoarctatis; pet. integra aequalia aut inæqualia; stam. intus basi squamulâ aucta; silicula depressa ovalis apice emarginata, valvis navicularibus in carinâ subalatis, septo oblongo angusto, stylo nullo; semina in quoque loculo 2 orbicularia. — Flores albi minimi. Herbæ ①, foliis radicalibus rosulatis pinnatilobatis.

1. *T. IBERICA* (DC. syst. 2. p. 392.) petalis inæqualibus, exterioribus majoribus. ① in arenosis, sylvaticis, nudis et sterilibus Gallie ferè totius. — Iberis nudicaulis Lin. sp. 907. *Guepinia iberica* Fl. fr. 5. p. 596. — Lob. ic. t. 221. f. 2. (v. v.)
2. *T. LEPIDIUM* (DC. l. c.) petalis aequalibus. ① in arenosis sterilibus apricis subsylvaticis Corsicæ, Occitanie, Delphinatus australis et verosim. Galloprovincie. — *Lepidium nudicaule* Lin. sp. 898. *Guepinia lepidium* Fl. fr. 5. p. 596. — Magn. bot. Monsp. 186. ic. (v. v.)

### XXIII. IBERIS. L. gen. n. 804.

Calyx basi æqualis; pet. 4 inæqualia, 2 exteriora majora; stam. eden-

tula; silicula valvis carinatis navicularibus depressissima, basi ovata, apice valvis in lobulos productis-emarginata, stylo persistente filiformi, septo angustissimo interdum bipartibili, loculis 1-spermis; semina ovata pendula. — Flores albi aut subpurpurascens.

Sect. 1. IBERIDIUM. DC. *prod.* 1. p. 179.

Radicula descendens; semen immarginatum; septum simplex.

\* *Pedicellis fructiferis corymbosis.*

1. *I. SPATHULATA* (Berg. phyt. ic.) herbacea glabra, foliis spathulatis integris subcarnosis, siliculis corymbosis sinu angusto acuto emarginatis. ① in Pyrenæorum lapidosis calcareis schistosive aquâ nivali irrigatis. — Fl. fr. 4. p. 716. Ib. carnosa Willd. sp. 3. p. 455. Flores purpurascens. (v. v. DC.)
2. *I. NANA* (All. auct. 15. t. 2. f. 1.) herbacea glabra, foliis subrotundo-spathulatis integris subcarnosis, siliculis corymbosis emarginatis sinu latiusculo obtuso. ① ② in rupibus apricis Alpium austral. edit., in montibus Aaroux, Ventoux, et Alpib. Sedenensibus Galloprovinciæ. — Fl. fr. 4. p. 717. Ib. aurescens Vill. Flores purpurascens. (v. v. DC.)
3. *I. CILIATA* (All. ped. auct. 15. non Willd.) herbacea glabriuscula, foliis linearibus integris basi ciliatis, siliculis corymbosis emarginatis lobulis obtusis stylo subæqualibus. ② in rupestribus apricis Galloprovinciæ circa Brignoles, et propè Niceam. — Fl. fr. 5. p. 597. Flores albi aut subpurpurei. (v. v.)
4. *I. LIMPOLIA* (Lin. sp. 905.) herbacea glabra, foliis linearibus integerrimis, radicalibus subdentatis, siliculis corymbosis bidentatis. ① ② in apricis cultis Galloprovinciæ, Nicææ, Occitanicæ infer. et Pyren. orient. ex cl. Benth. — Fl. fr. 4. p. 716. — Garid. aix. 459. t. 105. Flores purpurei conferti. (v. v. D.)
5. *I. UMBELLATA* (Lin. sp. 906.) herbacea glabra, foliis lanceolatis acuminatis, inferioribus serratis, superioribus integerrimis, siliculis umbellatis acutissime bifidis. ① Nicææ spontanea dicitur; in hortis culta. — Fl. fr. 4. p. 716. — Lob. ic. t. 216. f. 1. Flores sæpius purpurascens (v. v. sp. in Etruriâ DC.)

\* \* *Pedicellis fructiferis racemosis, caulibus herbæcis.*

6. *I. PINNATA* (Gouan. hort. Monsp. 319.) herbacea glabra, foliis pinnatifidis, racemis corymbosis post anthesin parum elongatis. ① inter segetes prov. australium. — Fl. fr. 4. p. 715. — Lob. ic. t. 217. f. 2. Flores albi. (v. v.)
- β. *crenata* (Lam. dict. 3. p. 213.) foliis inciso-crenatis. ① cum var. α. Ib. pandureformis Pourr. (v. v.)
7. *I. INTERMEDIA* (Guers. bull. philom. n. 82. t. 1. h.) herbacea, foliis lanceolatis glabris obtusis integris aut radicalibus subdentatis, floribus mox racemosis, siliculis ovatis truncatis seu latissime emarginatis. ② in rupibus calcareis secus Sequanam inter Rothomagum et Duclair. — Fl. fr. 4. p. 715. Flor. albi. (v. v.)
8. *I. AMARA* (Lin. sp. 906.) herbacea, foliis lanceolatis acutis subdentatis, floribus corymbosis mox racemosis, siliculis orbiculatis angustè emarginatis. ① in arvis agris et segetibus Gall. ferè totius. — Fl. fr. 4. p. 714. — Riv. tetr. it. t. 112. Fl. alb. (v. v.)

\* \* \* *Pedicellis fructiferis racemosis, caulibus frutescentibus.*

9. *I. GARRICIANA* (All. Ped. n. 920. t. 40. f. 3. et 54. f. 2.) frutescens, foliis oblongis basi attenuatis obtusis integerrimis glabris, floribus corymbosis. ③ in saxosis apricis Pyrenæorum orient. et centralium. — Fl. fr. 4. p. 714. Flor. albi. (v. v.)
10. *I. SAKATILIS* (Lin. amœn. 4. p. 321.) frutescens, foliis linearibus integerrimis subcarnosis subacutis glabris ciliatisve, floribus corymbosis. ③ in saxosis apricis collinis Pyrenæorum orient., Occitanicæ, Galloprovinciæ. — Fl. fr. 4. p. 714. — Lob. iq. t. 217. f. 1. (v. v. DC.)

*β. corifolia* (DC.) foliis glabris obtusis. *β* in monte Ventoso. Sims. bot. mag. t. 164a. (v. s. DC.)

Sect. II. *IBERIDASTRUM*. DC. l. c.

Radicula horizontalis; semen submarginatum; septum ferè duplex.

11. *I. SEMPERFLORENS* (Lin. sp. 904.) frutescens, foliis cuneatis spathulatisve obtusis integerrimis glabris, floribus corymbosis, siliculis apice truncato-submarginatis lobulis obsoletis. *β* ex Sicilia orta, in hortis sæpè culta. — Fl. fr. 4. p. 713. — Riv. tetr. irr. l. 224. f. 2. (v. v. c.)

XXIV. *BISCUTELLA*. L. gen. n. 808.

Cal. sepala nunc basi æqualia nunc lateraliter ad basim gibboso-calcarata; pet. æqualia integra; stam. edentula; silicula bilocularis, stylo longè persistente superata, loculis compressissimis planis orbiculatis scutiformibus lateraliter axi adnatis, à basi ad apicem ab axi separabilibus, vix aut non marginatis, 1-spermis; semina compressa; embryo inversus nempe radícula descendens. — Flores flavi.

Sect. I. *JONDRABA*. DC. syst. 2. p. 407.

Calyces basi bicalcarati; glandulæ tori valdè prominentes.

1. *B. APRICULATA* (Lin. sp. 911.) calycibus obtusè bicalcaratis, siliculis glabris in disco punctis elevatis scabris, lobulis apice in stylum coeuntibus. ① in agris cultis Galloprovinciæ ex Ger., Delphinatus ex Vill. sed forsàn cum seq. confusa. — DC. diss. t. 1. f. 2. Fl. fr. 4. p. 689. Schkuhr. handb. 2. t. 82. (v. v. c.)
2. *B. HISPIDA* (DC. diss. n. 3. t. 1. f. 1.) calycibus acutè bicalcaratis, siliculis glabris in disco punctis elevatis scabris, lobulis apice in stylum non coeuntibus, caule hispido. ① in apricis montosis Galloprovinciæ. — Barr. ic. t. 230 et 1219. (v. s.)
3. *B. CICHORIIFOLIA* (Lois. add. p. 167.) calycibus acutiusculè bicalcaratis, siliculis glabris in disco punctis elevatis scabris, lobulis in stylum non coeuntibus, caule villosa. ② in saxosis apricis ad radices Pyrenæorum centralium propè Bagnères de Luchon detexit Berger et in Pyren. orient. Thomas. — DC. diss. n. 4. t. 2. Folia infer. runcinata. (v. s.)

Sect. II. *THLASPIDIUM*. DC. l. c.

Calyces basi æquales; glandulæ tori vix prominentes.

4. *B. CHLATA* (DC. diss. n. 9. syst. 2. p. 410.) siliculis lævibus margine ciliatis, caule erecto elongato folioso, foliis sessilibus oblongis remotè dentatis. ① in Corsicâ reperit Ph. Thomas. — *B. coronopifolia*. DC. ic. rar. 439 non L. Fl. fr. 4. p. 692. (v. s.)
5. *B. LÆVIGATA* (Lin. mant. 225.) siliculis glabris lævibus, foliis subradicalibus piloso-scabris oblongis in petiolum attenuatis subdentatis integrisve, caulibus linearibus paucis integerrimis. ② in asperis apricis Alpium, Jura, Pyrenæorum, etc. — Fl. fr. 4. p. 689. ic. rar. t. 38. (v. v.)
6. *B. CORONOPHOLIA* (All. ped. n. 907.) siliculis glabris lævibus, foliis piloso-scabris plerumquè radicalibus pinnatifidis, lobis utrinquè 2-3 remotis. ② in apricis sterilibus collium et montium dentissimum Occidentis, Galloprov., Delphinatus australis. — DC. diss. n. 2. t. 18. non Fl. fr. (v. v. DC.)
7. *B. AMBIGUA* (DC. diss. n. 23. t. 11. f. 1.) siliculis glabris lævibus, foliis piloso-scabris, radicalibus sinuato-dentatis basi attenuatis, caulibus paucissimis basi subcoarctatis semi-amplexicaulibus. ② in sterilibus apricis collibusque Galloprovinciæ et Occitanie. — DC. syst. 2. p. 415. Var. foliorum sinibus dentibus revolutis aut planis. (v. v.)

8. *B. SAXATILIS* (DC. fl. fr. 4. p. 690. diss. t. 10.) siliculis glabris in disco punctis elevatis scabris, foliis piloso-scabris plerumque radicalibus oblongis. 2/ in sterilibus apricis montosis provinc. ab Alpibus ad Pyrenæos. — *B. picridifolia* Lapeyr. ex Arnolt. *B. apula* Lapeyr. non L. ex cl. Benth.

α. Foliis integris. — *B. longifolia* Vill. Dauph. 3. 365. (v. v.)

β. Foliis inciso-dentatis aut dentato-pinnatifidis. — *B. mollis* Loia. not. 168. (v. s.)

γ. Foliis subdentatis, caule humillimo. — *B. coronopifolia* Lapeyr. non L. *B. intermedia* Gon. ill. 42. (v. v.)

Trib. IV. *CAKILINEÆ* seu *Pleurorrhizæ* *Lomentaceæ*. DC.  
*syst.* 2. p. 427.

Siliqua aut silicula transversè in articulos 1-2-loculares 1-2-spermos secedens. Semina immarginata. Cotyledones planæ accumbentes, septo (dùm adsit) parallelæ.

XXV. *CAKILE*. *Tourn. inst. t.* 483. DC. *syst.* p. 185.

Calyx basi bigibbosus; pet. limbo obovato; stam. edentula; silicula biarticulata compressa, articulo superiore ensiformi aut ovato; semen in loculis solitarium, superioris erectum, inferioris pendulum. — *Herbæ carnosæ floribus albis aut subpurpurascensibus.*

1. *C. MARITIMA* (Scop. carn. n. 844.) siliculæ articulo superiore ensiformi. ① frequens in arenis maritimis secus Oceanum et mare Mediterraneum. — Fl. fr. 4. p. 718. *Bunias cakile*. Lin. sp. 936.

α. *vulgaris* (DC.) foliis pinnatifidis, lobis angustis distantibus. — Lob. ic. t. 223. f. 1. (v. v.)

β. *sinuatifolia* (DC.) foliis dentato-repandis seu sinuatis. — *C. Ægyptiaca* Wild. sp. 3. p. 417. (v. s. ex Ital.)

#### Subordo II. *NOTORHIZÆ* (o || .)

Cotyledones planæ incumbentes. Radicula dorsalis nempe medio dorso cotyledonum incumbens. Semina ovata non marginata.

Trib. V. *SISYMBREÆ* seu *Notorhizæ* *siliquosæ*. DC. *syst.* 2.  
p. 438.

Siliqua bilocularis bivalvis dehiscens valvis concavis carinatisve. Cotyledones planæ incumbentes septo contrariæ.

XXVI. *MALCOMIA*. *Brown. Kew. II. v.* 4. p. 121. DC. *syst.* 2. p. 438.

Calyx clausus; pet. limbo obovato aut leviter emarginato; stam. edentula; siliqua teres stigmate acutissimo simplici terminata. — Flores albi aut purpurascens.

1. *M. AFRICANA* (Brown. h. Kew. 4. p. 121.) caule ramoso diffuso, foliis lanceolatis subdentatis, pube 2-4-partita, pedicellis calyce persistente brevioribus, siliquis scabris. ① in agris vineis et arenosis prov. oleiferarum. — Bocc. Sic. t. 42. f. 1. *Hesperis Africana* Lin. sp. 928. Fl. fr. 4. p. 562. (v. v. DC.)

2. *M. MARITIMA* (Brown. l. c.) caule erecto ramoso, foliis ellipticis obtusis integris basi attenuatis, pube adpressâ 2-4-partita, pedicellis calyce subbrevioribus, siliquis pubescentibus apice longè acuminatis. ① in litoribus arenosis propè

Narbonam, Massiliam (Solier) et ex Thore in ericetis Bardigalensis. — Barr. ic. t. 1127. *Cheiranthus maritimus* Lin. sp. 924. *Hesperis maritima* Fl. fr. 4. p. 654. (v. v.)

3. *M. PARVIFLORA* (DC. syst. 2. p. 442.) caule erecto humili ramoso, foliis oblongis obtusis subintegris, pube tomentosa stellata, pedicellis demum calycis longitudine, siliquis pubescentibus. ① in arenosis maritimis Corsicæ, agri Ruscinonensis, Occitania, Galloprov. — *Hesperis parviflora* Fl. fr. 4. p. 654. 5. p. 390. ic. rar. 1. t. 35. (v. v.)
4. *M. LITTORÆA* (Brown. l. c.) caule erecto multiplici, foliis lanceolato-linearibus subintegris tomento brevi canis, pedicellis calycis longitudine, siliquis incanis. ② in arenosis maritimis à Galloprovincia ad Nannetes frequens. — Lob. ic. t. 331. f. *Cheiranthus littoreus* Lin. sp. 925. Fl. fr. 4. p. 656. (v. v.)

#### XXVII. HESPERIS. *L. gen. n. 817. DC. syst. 2. p. 447.*

Calyx clausus bisaccatus; pet. subintegra; Stam. edentula; siliqua teretiuscula aut tetragona, stigmatibus 2 erectis conniventibus; semina oblonga subtriquetra.

1. *H. LACINATA* (All. ped. n. 985. t. 82. f. 1.) pedicellis calyce brevioribus, petalis obovato-oblongis, foliis obovatis inciso-dentatis, caule hispido. ② in rupibus apricis Galloprovinciæ, Delphinatûs australis, et Pyren. orient. propè Font de Comps ex cl. Benth. — Fl. fr. 4. p. 652. Flores nunc flavescentes nunc purpurascens interdum in iisdem plantis. (v. v. DC.)
2. *H. MATRONALIS* (Lam. dict. 3. p. 321. ill. t. 564. f. 1.) pedicellis calycis longitudine, petalis obovatis, siliquis erectis torosis glabris margine non incrassatis, foliis ovato-lanceolatis dentatis, caule erecto subsimplici. ④ in dumetis et sepibus submontanis. — Fl. fr. 4. p. 652.
  - a. *hortensis* (DC.) caule firmo recto, floribus sæpius odoratis, petalis frequentius emarginato-mucronatis, foliis basi non cordatis. — Var. fl. albis purpureis aut variis, simplicibus plenisque. (v. v. c.)
  - β. *sylvestris* (DC.) habitu laxo, foliis saltem inferioribus basi plus minus cordatis et grossius dentatis, floribus semper purpurascens variis odoratis, petalis obtusis. — *H. inodora* Lin. sp. 917. Jacq. fl. austr. t. 347. Ad radices Pyrenæorum, etc. (v. v. sp.)

#### XXVIII. SISYMBRIUM. *All. ped. 1. p. 274. DC. syst. 2. p. 458.*

Calyx basi æqualis; petala integra; Stam. edentula; siliqua sessilis teretiuscula, valvis concavis, stylo vix ullo, septo membranaceo; stigmata 2 distincta aut in capitulum connata; semina ovata aut oblonga 1-serialia.

##### Sect. I. VELARUM. *DC. l. c.*

Siliquæ subuliformes basi latiores apice in stylum brevissimum desinentes, pedicello brevissimo post anthesin incrassato secus axiû adpressæ. — Flor. flavi.

1. *S. OFFICINALE* (Scop. carn. n. 824) foliis runcinatis pilosis, caule piloso, siliquis subulatis rachi adpressis. ① ubique in incultis ruderalis secus vias et muros. — Fl. fr. 4. p. 672. — Lob. ic. t. 206. f. 1. — *Erysimum officinale* Lin. sp. 922. (v. v.)

##### Sect. II. NORTA. *DC. l. c.*

Siliquæ teretes; calix patens; semina oblonga. — Flores flavi; racemi cbractati.

2. *S. STRICTISSIMUM* (Lin. sp. 922.) foliis lanceolatis petiolatis dentatis pubescentibus. ④ in apricis montosis asperis circa Parisios ex Merat, in vallibus

*Alpium Delphinatūs* ex Vill. — Fl. fr. 4. p. 972. — Cam. epit. 342. ic. (v. v. c.)

Sect. III. IRIO. DC. l. c.

Siliquæ teretes; semina ovata subtriquetra. — Flores flavi; pedicelli ebracteati.

3. *S. OBTUSANGULUM* (Scheich. in Fl. fr. 4. p. 671.) foliis pinnatipartitis, lobis ovali-oblongis obtusis sinuato-dentatis, sinibus rotundatis, caule basi retrorsum subhispido. ② in arenosis, cultis, glareosis et ruratis ferè ubique. — *S. jacobæefolium* Berg. phyt. ic. Non diversum esse a *Brassicâ Erucastro* vult cl. Gay. (v. v.)
4. *S. ACUTANGULUM* (DC. fl. fr. 4. p. 670.) caule foliisque glabris, radicalibus runcinatis, caulinis pinnatifidis lobis sinibusque acutis, calyce patentissimo, siliquis scabris. ② in asperis apricis et ruderatis vallium Pyrenæorum et Alpium, in collibus prov. austr. — *Sinapis Pyrenæica*. Lin. sp. 934. All. ped. t. 55. f. 1. (v. v.)
5. *S. TARAXACIFOLIUM* (DC. fl. fr. 4. p. 670. ic. rar. t. 37.) foliis runcinato-pinnatifidis, lobis sinibusque acutis, siliquis declinatis glabris, calyce erectiusculo. ② in asperis montium Sedenensium Galloprovinciæ, et edit. Pyren. orientalium ex cl. Benth. — Caulis glaber infernè pilo spatuloso gerens. (v. s.)
6. *S. AUSTRIACUM* (Jacq. austr. 3. t. 262.) caule siliquis foliisque glabris, fol. radicalibus runcinatis, caulinis incisim pinnatifidiæ, lobis sinibusque acutis, calyce patente. ② in asperis apricis et ruderatis collium et vallium Delphinatūs, Galloprovinciæ, Occitaniæ et Pyrenæorum orient. — *S. erysimifolium*. Pourr. Fl. fr. 4. p. 671. (v. v. DC.)
7. *S. IRIO* (Lin. sp. 921.) caule foliisque glabris runcinato-pinnatipartitis, lobis dentatis, terminali elongato, calyce siliquisque patenti-erectis. ① in ruderatis, cultis, secus vias et muros ferè ubique. — Jacq. fl. austr. t. 322. Fl. fr. 4. p. 669. Flor. parvi. (v. v.)
8. *S. COLUMNÆ* (Jacq. fl. austr. t. 323.) caule villosissimo subincano, foliis runcinatis pubescentibus, lobis dentatis, integrisve acutis, siliquis magnis erectiusculis, calyce laxo. ① in ruderatis, cultis, viis et hortis Argenterati, Montpellier, Narbonæ, agri Buscinonensis. — Fl. fr. 5. p. 590. (v. v.)  
*β. villosissimum* (DC. syst. t. 2. p. 469.) siliquis pubescentibus, caule foliisque villosissimis, foliorum lobis dentatis. — Parisiis forsan ex hort. Bot. migratum. S. Læselii. Thuil. fl. par. 335. Fl. fr. 4. p. 670. (v. v.)
9. *S. PANNOXICUM* (Jacq. ic. rar. t. 1. t. 123.) foliis inferioribus runcinatis hispidis, lobis dentatis, superioribus pinnatisectis glabris, lobis linearibus integerrimis, siliquis patentibus. ① in Alsatiâ ad pagum Mutzig propè Argentoratū legit Nestler. — Fl. fr. 5. p. 591. (v. s.)
10. *S. ASPERUM* (Lin. sp. 920.) foliis glabris pinnatipartitis, lobis oblongis obtusis subdentatis, pedicellis brevissimis, siliquis muricato-scabris stylo brevi acuminatis. ① in arenosis glareosisque humidis subpaludosis Galloprov., Occitaniæ, Delphinatūs austr., Pictaviæ, Arvernæ, Burgundiæ. — J. Bauh. hist. 2. p. 858. f. 3. — Fl. fr. 4. p. 668. Flores parvi. (v. v. DC.)
11. *S. SOPHIA*. (Lin. sp. 922.) foliis bipinnatisectis, lobis oblongo-linearibus incisim, pedicellis calyce quadruplò longioribus, petalis calyce minoribus. ① in maceris, muris, tectis, ruderatis Gall. ferè totius. — Fl. fr. 4. p. 669. — Fuchs. hist. t. 2. (v. v.)
12. *S. TANAGTIFOLIUM* (Lin. sp. 916.) foliis pinnatisectis, segmentis lanceolatis inciso-serratis, extimis confluentibus, petalis calyce majoribus, siliquis pedicello brevioribus. ② in asperis apricis, saxosis frigidis Alpium et centralium Pyrenæorum. — Fl. fr. 4. p. 664. — Zan. hist. 86. t. 33. (v. v. sp. in Alp.)

Sect. IV. KIBERA. DC. syst. 2. p. 477.

Calyx clausus; glandulæ tori nullæ aut minimæ; stylus brevis crassus

retasus. — Pedicelli ex bracteis axillis solitarii aut plurimi. Flores parvi, flavi aut albi.

13. *S. SUPINUM* (Lin. sp. 907.) pedicellis axillaribus brevissimis solitariis, siliquis erectis puberulis, foliis sinuato-pinnatifidis, caule retrorsum pubescente. ① in arenosis humidis et ad agrorum margines secus ripas Sequanae propè Parisios, Rhodani ad Lugdunum, in Jurasso circà lacum Jovis. — Isuard. act. acad. sc. 1724. t. 18. Fl. fr. 4. p. 668. Cor. alba. (v. v. DC.)
14. *S. POLYCRATIUM* (Lin. sp. 918.) pedicellis subternis axillaribus brevissimis, siliquis erectis glabris, foliis sinuato-runcinatis, lobis acutis dentatis, inferioribus majoribus. ① in muris humidis et rupium fissuris prov. oleiferarum et Arvernæ. — Fl. fr. 4. p. 667. — Lob. ic. t. 206. f. 2. Fl. flavi. (v. v.)

Sect. VI. *ARABIDOPSIS*. DC. l. c.

Siliquæ lineares compressæ stigmate sessili truncatæ. — Flores albi brevissimè pedicellati; pedicelli ebracteati.

15. *S. BURSIFOLIUM* (Lin. amon. 7. p. 322.) foliis lyratis pinnatifidis glabris, caule erecto folioso, pedicellis crassis calyce brevioribus. ① in Pyrenæorum vallibus orientalibus ex Pourret. — Fl. fr. 4. p. 668. — Dill. h. elsh. f. 177. (v. s. sp.?)
16. *S. PINNATIFIDIUM* (DC. fl. fr. 4. p. 667.) foliis radicalibus lyratis, caulinis pinnatipartitis, lobis linearibus integris, terminali majore, pedicellis gracilibus calyce ferè brevioribus. ② in pascuis lapideis rupestribusque montium Arvernæ, Alpium, Pyrenæorum. — S. dentatum All. ped. t. 57. f. 3. *Arabis pinnatifida* Lam. ill. t. 563. f. 3. (v. v.)

XXIX. *ALLIARIA*. Adans. fam. 2. p. 418. DC. syst. 2. p. 488.

Calyx laxus basi æqualis caducus; pet. limbo obovato; stam. edentula; glandulæ 4 inter stam. et pistillum; siliqua subtetragona nempè teres valvis striâ carinali et lineis prominulis angulata, stylo brevissimo; semina subcylindrica. — Flores albi.

1. *A. OFFICINALIS* (Andrz. in DC. syst. 2. p. 489.) foliis cordatis, siliquis prismaticis pedicello multoties longioribus. ② ad sepes, dumeta et fossas Gall. ferè totius. — Fuchs. hist. 105. ic. *Erysimum Alliaria* Lin. sp. 922. *Nesperis Alliaria* Fl. fr. 4. p. 651. (v. v.)

XXX. *ERYSIMUM*. DC. syst. 2. p. 490.

Calyx clausus; pet. integra; stam. edentula; siliqua valvis carinatis tetragona, sessilis; stylus vix ullus; stigmata 2 patula; semina ovata aut oblonga 1-serialia. — Flores flavi aut rarissimè albi.

Sect. I. *ERYSIMASTRUM*. DC. l. c.

Calyx cum petalis deciduus. — Folia nec cordata nec amplexicaulia.

1. *E. STRICTUM* (Gærtn. fl. welt. 2. p. 451.) foliis lanceolatis dentatis pube tripartita pubescentibus, caule stricto, siliquis erectis, stylo brevissimo, stigmate bilobo. ② in ruderatis submontanis et arvis glareosis Delphinatûs. — *E. hieracifolium* Jacq. austr. t. 73. (v. v.)
2. *E. VIRGATUM* (Roth. cat. bot. 75.) foliis oblongo-lanceolatis subdentatis pube tripartita pubescentibus, caule stricto tereti, siliquis erectis, styli longitudine latitudinem siliquæ vix excedente. ② in Jurasso, ad vias Belgii. — DC. syst. 2. p. 496. (v. v.)
3. *E. CHEIRANTHOIDES* (Lin. sp. 923.) foliis lanceolatis subdenticulatis scabriusculis viridibus, siliquis erecto-patulis pedicello duplò longioribus, stigmate parvo subsessili. ① in ruderatis, arvis, ad margines agrorum et viarum, juxtà sepes

et interdum in humidis Gall. ferè totius. — Fl. fr. 4. p. 659. — Lob. ic. t. 225. f. 1. (v. v.)

4. *E. HELVETICUM* (DC. fl. fr. 4. p. 658.) foliis linearibus integris, cauleque erecto vix ramoso, pube adpressâ bipartitâ cinereis, siliquis erectiusculis, stigmate pedicellato emarginato. ② in Pyren. central. ex cl. Benth. — *Cheiranthus Helveticus* Jacq. vind. t. 9. (v. v. in Helv. D.)
5. *E. CANESCENS* (Roth. cat. bot. 1. p. 76.) foliis linearibus aut subdentatis pube bipartitâ subcanescentibus, unguibus petalorum calyce longioribus, laminis obovato-oblongis, siliquis erectis pedicello quintuplò longioribus, stigmate subsessili. ② in collibus et aggeribus siccis et apricis Niceæ et Galloprovinciæ speciatim ad rupem Sanctæ Victoriæ ex DC. syst. 2. p. 501. — *Cheiranthus alpinus* Jacq. austr. t. 75. (v. v.)
6. *E. LANCEOLATUM* (DC. syst. 2. p. 502.) foliis inferioribus lanceolatis dentatis superioribus sublinearibus integris, unguibus calyce longioribus, laminis orbiculato-obovatis, siliquis erectis, stigmate subsessili. ② in apricis rupestribus campestribus et muris in Gall. ferè totâ.  
*a. majus* (DC.) caule simplici aut ramoso pedali rigido. — *Erysimum majale*. Fl. fr. 4. p. 657. *Cheiranthus erysimoides* Lin. sp. 923. Jacq. fl. austr. t. 74. (v. v.)  
*β. minus* (DC.) caule simplici interdum multicauli subascendente. — *Cheiranthus Alpinus* Lam. dict. 2. p. 716. *E. ochroleucum β.* Fl. fr. 4. p. 658. (v. v.)
7. *E. OCHROLEUCUM* (DC. fl. fr. 4. p. 658.) foliis oblongo-lanceolatis subdentatis pube bipartitâ aut nullâ, caule decumbente ramoso, petalis obovatis, siliquis erectis stylo pedicelli longitudine apiculatis. ½ in Jurassi montibus Helvetiæ conterminis. — *Cheiranthus ochroleucus* Hall. fil. ex Schl. DC. syst. 2. p. 181. — Hall. Helv. n. 449. t. 14. (v. v.)

#### Sect. II. CORINGIA. DC. syst. 2. p. 507.

Petala erectiuscula alba; glandula inter pistillum et stam. lateral. — Folia caulina cordata amplexicaulia lævia.

8. *E? ALPINUM* (Baumg. trans. 2. p. 263.) foliis membranaceis glabris, caulinis cordato-sagittatis amplexicaulibus oblongis, radicalibus petiolatis ovatis. ½ in sylvaticis lapidosis montanis Alpium Delphinatus, Cebennarum et Alsatiæ. — *Brassica alpina* Lin. mant. 95. Vill. dauph. 3. p. 330. t. 36. Fl. fr. 4. p. 647. *Arabis brassicæformis* Wallr. Sched. crit. 359. Gay. in ann. sc. nat. 1826. (v. v.)
9. *E. PERFOLIATUM* (Crantz. austr. 27. DC. syst. 2. p. 508.) foliis radicalibus obovatis, caulinis cordato-amplexicaulibus, omnibus obtusis glabris glaucis, siliquæ tetragonæ lateribus enerviis. ½ in agris saxis et clivis Gall. ferè tot. — Lob. ic. 396. f. 2. *Brassica orientalis* Lin. sp. 931. *Brassica perfoliata* Fl. fr. 4. p. 696. (v. v.)

#### Trib. VI. CAMELINEÆ seu Notorhizæ latiseptæ. DC. syst. 2. p. 513.

Silicula 2-locul. aut abortu 1-locul. bivalvis, valvis plus minùs concavis sæpiùs dehiscens, septo elliptico in majore fructus diametro. Semina plura ovata immarginata. Cotyled. planæ, incumbentes.

#### XXXI. CAMELINA. Crantz. fl. aust. 1. p. 17. DC. syst. 2. p. 514.

Calyx basi æqualis; pet. integra; stam. edentula; silicula obovata aut globosa, stylo filiformi persistente superata, valvis ventricosis, loculis polyspermis; semina oblonga immarginata. — Flores flavi.

1. *C. SATIVA* (Crantz. austr. 10.) siliculis cuneato-pyriformibus 4-rostratis stylo longiusculo terminatis, foliis subintegerrimis lanceolatis. ① inter



segetes et Lina hinc inde spontanea. — *Myagrum sativum* Lin. sp. 894. Cav. ic. t. 66. Fl. fr. 4. p. 117. *Alyssum utriculatum* Lapeyr. (v. v.)

α? *pilosa* (DC.) foliis integris pilosis. — *C. sylvestris* Wallr. sched. 347. Semper sylvestris. (v. v.)

β. *glabrata* (DC.) foliis integris subglabris. — Lind. als. t. 1. Culta ad oleum è seminibus extrahendum. (v. v. c.)

2. *C. DENTATA* (Pers. ench. 2. p. 191.) siliculis subgloboso-pyriformibus 4-costatis stylo longiusculo superatis, foliis repando-dentatis. ① in arvis Alsatiæ arenosis. — J. Bauh. hist. 2. p. 893. ic. — DC. syst. 2. p. 516. (v. s. DC.)

XXXII. NESLIA. Desv. journ. 3. p. 162. DC. syst. 2. p. 516.

Cal. æqualis patens; pet. integra; stam. edentula; silicula coriacea indehiscens subgloboso-compressa, septo tenui in majore diametro interdum evanido, valvis concavis indistinctis; semina in quoque loculo solitaria, pendula, subglobosa. — Flores flavi.

1. *N. PANICULATA* (Desv. journ. bot. 3. p. 162.) ① in arvis et arenosis Gall. ferè totius. — Bunnias paniculata Fl. fr. 4. p. 721. *Myagrum paniculatum* Lin. sp. 894. — C. Bauh. prod. 52. ic. Folia amplexicaulia sagittata oblongo-lanceolata integra. (v. v. DC.)

Trib. VII. *LEPIDINEÆ* seu *Notorhizæ angustiseptæ*.  
DC. syst. 2. p. 521.

Silicula septo angustissimo, valvis carinatis aut valdè concavis. Semina in loculis solitaria aut pauca ovata immarginata. Ccyledones planæ incumbentes.

XXXIII. SENEBIERA. DC. syst. 2. p. 521.

Calyx æqualis patens; pet. integra; stam. edentula, interdum abortu 4 aut 2; silicula didyma, subcompressa, evalvis, aptera, bilocularis, indehiscens, stigmata sessili, valvis subglobosis rugosis aut subcristatis, loculis 1-spermis; semina pendula, subgloboso-triquetra. — Flores albi minimi.

Sect. I. *NASTURTIOLUM*. DC. l. c.

Silicula didyma apice emarginata, septo nempè valvulis globosis brevior.

1. *S. PINNATIFIDA* (DC. mem. soc. h. n. Par. 144. t. 9. Fl. fr. 4. p. 703.) foliis pinnatilobatis, lobis oblongis dentatis aut subincis, siliculis compressis didymis reticulatis. ① ② in ruderalis præsertim maritimis in Armoraciâ circa Quimper et Nannetes, in Vasconiâ, Mospelii. — *Lepidium didymum* Lin. mant. 92. Caules plurimi diffuso-procumbentes. (v. v.)

Sect. II. *CARARA*. DC. l. c.

Silicula apice non emarginata, septo nempè valvis paulò longiore, valvulis compressis dorso cristato-rugosis.

2. *S. CORONOPUS* (Poir. dict. 7. p. 76.) foliis pinnatilobatis, lobis integris dentatis pinnatifidiæ, siliculis acutiusculis compressis, valvis dorso cristato-rugosis. ① ② in ruderalis apricis secus vias in calcareis et arenosis ferè ubique. — *Cochlearia Coronopus* Lin. sp. 904. *Coronopus vulgaris* Fl. fr. 4. p. 704. 5. p. 595. — Lob. ic. 238. f. 1. (v. v.)

XXXIV. CAPSELLA. DC. syst. 2. p. 383.

Calyx æqualis; pet. integra; stam. edentula; stigma sessile; silicula depresso-complanata, triangularis, apice truncata, septo ferè lineari, loculis 8-10-spermis, valvis carinatis compressis dorso non alatis. — Flores albi.

1. *C. bursa-pastoris* (DC. l. c.) ① in agris, oleraceis, ruderalisque ubique frequens. — *Thlaspi bursa-pastoris* L. sp. 903. Fl. dan. t. 729. Fl. fr. 4. p. 709. Ples herb. fasc. 5. ic. — Planta valde varians quoad foliorum formam. Petala in stamina mutata vidit cl. Jacquin. (v. v.)

XXXV. *LEPIDIUM*. Brown. Kew. ed. 2. v. 4. p. 85.

Cal. æqualis; pet. integra; stam. edentula; silicula ovalis depressa dehiscens, valvis carinatis, septo membranaceo angustissimo; semina in quoque loculo solitaria, pendula, subtriquetra aut compressa. — Flores albi.

Sect. I. *CARDARIA*. DC. syst. 2. p. 528.

Silicula ovato-cordata acuta subdepressa, valvis concavis suburgidis dorso apteris, stylo filiformi experto superata.

1. *L. DRABA* (Lin. sp. ed. 1. p. 645.) siliculis cordatis suburgidis apice integris stylo superatis, foliis amplexicaulibus lanceolatis dentatis. ① in arvis agris oleraceis Gall. ferè totius, præsertim in Occitaniâ frequens. — *Cochlearia draba* Lin. sp. ed. 2. p. 904. Fl. fr. 4. p. 702. — Lob. ic. 224. f. 1. (v. v.)

Sect. II. *CARDAMON*. DC. l. c.

Silicula suborbiculata apice emarginata, valvis carinato-navicularibus dorso subulatis, stylo brevissimo intrâ emarginaturam incluso; cotyledones tripartitæ.

2. *L. SATIVUM* (Lin. sp. 899.) siliculis orbiculatis alatis emarginatis, foliis variè divis incisisque, ramis non spinescentibus. ① ex Oriente ortum nunc in hortis frequens cultum. — *Thlaspi sativum* Fl. fr. 4. p. 708. — Lob. ic. 212. f. 1. (v. v. c.)  
*β. crispum*. C. Bauh. prod. 43. et 44. ic. (v. v. c.)

Sect. III. *LEPIA*. DC. l. c.

Silicula suborbiculata apice emarginata, valvis navicularibus apice alatis, alis stylo sæpius adnatis, stylo brevissimo intrâ emarginaturam incluso; cotyled. integre.

3. *L. CAMPESTRE* (Brown. h. Kew. 4. p. 465.) siliculis ovatis alatis emarginatis lepidoto-punctatis, foliis caulinis sagittatis dentatis. ① ② in viis, agris argillosis apricis arvis Gall. ferè tot. — *Thlaspi campestre* Lin. sp. 902. Fl. fr. 4. p. 712. *Lepidium cristatum* Lapeyr. ex Arnolt. — Fuchs. hist. 306. ic. (v. v.)  
 4. *L. HIRTUM* (Smith. comp. ed. 3. p. 98.) siliculis ovatis alatis emarginatis hirtis, foliis caulinis sagittatis villosis subintegris. ② in arvis, agris apricis, lapidosis prov. oleiferarum. — *Thlaspi hirtum* Lin. sp. 901. Fl. fr. 4. p. 713. — C. Bauh. prod. 47. ic. (v. v. DC.)  
 5. *L. NUMIFUUM* (Req. in ann. sc. nat. t. 5. p. 385.) siliculis ovatis alatis emarginatis glabris, foliis radicalibus lyratis ovatisque, caulinis sagittatis integris, caulibus prostratis pedicellisque pubescentibus. ④ in Corsicæ montibus. (v. s.)

Sect. IV. *DILEPTIUM*. DC. l. c. p. 538.

Silicula subelliptica brevissimè emarginata, valvis carinatis nec dorso nec apice alatis, stylo vix ullo. — Flores minimi interdum apetalis 2-4-andri.

6. *L. RUDERALE* (Lin. sp. 900.) siliculis ovatis emarginatis patulis pedicello brevioribus, foliis glabris, radicalibus pinnatilobatis, rameis linearibus integris. ① in ruderalis et secus vias ferè ubique. — *Thlaspi ruderale* Fl. fr. 4. p. 707. *L. subulatum* et *graminifolium* Lapeyr. ex cl. Benth. — Lob. ic. 214. f. 1. Variat floribus apetalis et 4-petalis, 2-4-andr. (v. v.)

Sect. v. LEPIDIASTRUM. DC. l. c.

Silicula ovata aut elliptica integerrima nec ullo modo emarginata stigmate subcassili apiculata, valvis carinatis apteris.

7. *L. LATIFOLIUM* (Lin. sp. 899.) siliculis ovatis stigmate apiculatis, foliis ovato-lanceolatis indivisis subserratis, infimis longe petiolatis. 2½ in umbrosis succulentis humidis ferè ubique. — Fl. fr. 4. p. 704. — Lob. ic. 318. f. 2. (v. v.)
8. *L. IRRIS* (L. sp. 900.) siliculis ovatis stigmate apiculatis, foliis radicalibus incisis pinnatisectisve, caulinis linearibus integris, caule ramosissimo. 2½ in ruderalis et secus vias ferè ubique. — *L. graminifolium* Lin. sp. 900. Lob. ic. 223. f. 2. Fl. fr. 4. p. 705. (v. v.)

XXXVI. ÆTHIONEMA. Brown. Kew. l. c. p. 80. DC. syst. 2. p. 557.

Calyx basi inæqualis; pet. integra; stam. majora aut inter se connata aut intus dentata; silicula ovalis sæpius emarginata, valvis navicularibus dorso slatis, loculis oligospermis; semina ovato-oblonga. — Flores ex albo rosei.

1. *Æ. SAXATILE* (Brown. h. Kew. 4. p. 80.) siliculis bilocularibus obeordatis, stylo brevissimo, valvis dorso alatis integris, stam. majoribus dentatis, racemis fructiferis laxis. 5 in rupestribus agri Ruscinonensis, Occitanie, Galloprov., Delphin. austr. — *Thlaspi saxatile* Lin. sp. 901. Fl. fr. 4. p. 710. *Thlaspi marginatum* Lapeyr. supp. ex cl. Arnott. — Barr. ic. 845. (v. v.)

Trib. VIII. ISATIDEÆ seu NOTORHIZEÆ NUCA-MENTACEÆ. DC. syst. 2. p. 563.

Silicula valvis indistinctis aut indehiscentibus carinatis, septo evanido 1-locul. 1-sperma. Semina ovato-oblonga immarginata. Cotyled. planæ oblongæ incumbentes.

XXXVII. ISATIS. Lin. gen. 834. DC. syst. 2. p. 564.

Calyx æqualis patens; pet. integra; stam. edentula; stigma sessile; silicula ovali-oblonga 1-locul. depresso-plana, valvis carinatis vix dehiscentibus dorso subalatis; semen pendulum. — Flores flavi. Fol. integra, sup. basi acutè sagittata sessilia.

1. *I. TINCTORIA* (Lin. sp. 936.) siliculis cuneatis basi acuminatis apice subspathulatis obtusissimis glabris latitudine majore triplò longioribus. ② Fl. fr. 4. p. 722.
  - a. *sylvestris*, foliis præc. inferioribus parè villosis. — In ruderalis apricis secus vias ferè ubique. Lob. ic. 352. f. 1. (v. v.)
  - β. *sativa* (DC.) foliis glabris latioribus. Culta in agris pinguibus circà Albi, etc. ad usum tinctorum. (v. v. c.)
  - γ. *hirsuta* (DC.) foliis hirsutis angustioribus. *I. hirsuta* Vill. Dauph. 3. p. 308. non All. — In apricis saxosis. (v. s.)
2. *I. CAMESCHENS* (DC. fl. fr. 5. p. 598.) siliculis elongato-cuneatis pubescentibus infernè angustatis supernè subspathulatis obtusis, latitudine quadruplò longioribus, pedicelli apice obconici longitudinem duplò excedentibus. ② in apricis maritimis Galloprovinciæ circà Telonem. — Caulis et fol. infer. pilis patulis mollibus hirsuta. (v. s.)

XXXVIII. MYAGRUM. Tourn. inst. t. 99. DC. syst. 2. p. 573.

Cal. suberectus; petala oblonga; stam. majora inter se imà basi sub-

connata; stylus brevis; silicula compressa ferè cuneata, apice lacunas 2 vacuas, infernè loculum 1 monospermum gerens; semen pendulum oblongum. — Flores pallide flavi. Fol. integra, super. sagittato-amplexicaulia.

1. *M. PERFOLIATUM* (Lin. sp. 893.) ① in agris et arenosis prov. præsertim australium. — *Cakile perfoliata* Fl. fr. 4. p. 720. — C. Bauh. prod. 51. ic. (v. v.)

Subordo III. ORTHOPLOCEÆ. (o>>) DC. syst. 2. p. 438.

Cotyledones incumbentes, conduplicatæ seu secus nervum medium longitudinaliter plicatæ et radiculam dorsalem in sinu foventes. Semina immarginata sæpiùs globosa.

Trib. IX. *BRASSICÆ* seu *ORTHOPLCEÆ SILIQUOSÆ*. DC. syst. 2. p. 581.

Siliqua valvis longitudinaliter dehiscentibus, septo lineari.

XXXIX. *BRASSICA*. DC. syst. 2. p. 582.

Calyx erectus; pet. obovata; stam. edentula; siliqua teretiuscula stylo parvo brevi obtuso; semina uniseriata subglobosa.

Sect. I. *BRASSICARIA*. DC. syst. 2. p. 582.

Siliqua sessilis, rostro nullo aut conico aspermo superata.

1. *B. OLERACEA* (Lin. sp. 932.) foliis polline glaucis subcarnosis repandis lobatisve, etiam junioribus glaberrimis. ② DC. in ann. ag. franc. vol. 19. diss. 10. t. 2. p. 582.

*a. sylvestris* (DC.) in rupibus maritimis Normanniæ? *B. oleracea* Sm. engl. bot. t. 637. (v. v. in littore Angliæ DC.)

*β. acephala* (DC.) caule tereti elongato, foliis expansis, racemis paniculatis. — Lob. ic. 243. f. 1. Culta in hortis et agris prov. præsertim occid. (v. v. c.)

*γ. bullata* (DC.) caule tereti subelongato, foliis junioribus subcapitatis demùm patulis bullatis crispatisve, racemis paniculatis. — Lob. ic. 244. f. 1. Culta in hortis. (v. v. c.)

*δ. capitata* (L.) caule tereti abbreviato, foliis concavis non bullatis in capitulum antè florescentiam congestis, racemis paniculatis. — Lob. ic. 243. f. 2. Culta in hortis. Variat fol. viridibus et purpureis, capite spherico ovato aut depresso. (v. v. c.)

*ε. caulorapa* (DC.) caule ad originem foliorum tumido subgloboso. — Lob. ic. 246 f. 1. Variat alba et purpurascens, in hortis culta. (v. v. c.)

*ζ. botrytis* (L.) racemorum pedunculis corymbosis confertis antè florescentiam valdè carnosus abbreviatis, floribus sæpè abortivis. — Lob. ic. 245. f. 1. Culta in hortis. (v. v. c.)

2. *B. CAMPESTRIS* (DC. syst. t. 2. p. 588.) foliis polline glaucis subcarnosis, inferioribus novellis subhispidis ciliatisve lyratis dentatis, cæteris cordato-amplexicaulibus acuminatis. ②

*a. oleifera* (DC.) radice fusiformi exili, caule elongato. — *Br. campestris* Lin. sp. 931. Sm. engl. bot. t. 224. Culta in agris sub nom. *Colsat* seu *Colsat*. (v. v. c.)

*β. pabularia* (DC.) radice fusiformi exili, caule abbreviato. *Chou à faucher*. (v. v. c.)

*γ. napobrassica* (DC.) radice tumidâ napiformi. Culta in agris, variat radice alba (*chou navet blanc*), purpurascens (*chou navet rouge*) et flavescens (*Rutabaga*). (v. v. c.)

3. *B. RAPA* (Lin. sp. 931.) foliis radicalibus lyratis polline glauco destitutis

setoso-scapris, caulinis mediis incisiss, summis integerrimis lævibus. (2) DC. syst. 2. p. 590.

- a. *depressa* (DC.) radice sub collo tumida globoso-depressa in radiculam tenuem subtus abruptè desinente. — Lob. ic. t. 197. f. 1. Culta in agris, variat radios albâ, flavescente, nigricante, rubente, etc. (v. v. c.)
- β. *oblonga* (DC.) radice oblongâ sensim apice attenuatâ. — Lob. ic. 197. f. 2. Culta rariss. Radix alba. (v. v. c.)
- γ. *oleifera* (DC.) radice exili. — Lob. ic. 298. f. 1. Culta ad oleum è seminibus in Delphin. sub nom. *navette*. (v. v. c.)
4. B. *NAPUS* (Lin. sp. 951. DC. syst. 2. p. 592.) foliis glabris polline cæsie glaucescentibus, radicalibus lyratis, caulinis pinatifidis crenatisque, summis cordato-lanceolatis amplexicaulibus, siliquis divaricato-patentibus. (2) B. *arvensis* Lapeyr. non L. (cl. Arnott. ex autopsiâ in herb. Lap.)
  - a. *oleifera* (DC.) radice exili. — Lob. ic. 200. f. 2. Culta in agris sub nomine *navette d'hiver*. (v. v. c.)
  - β. *esculenta* (DC.) radice fusiformi sub collo incrassatâ. — Lob. ic. 200. f. 1. Culta in agris et hortis, variat radice albâ, flavâ, nigricante. (v. v. c.)
5. B. *PRÆCOX* (Waldst. et Kit. in DC. syst. 2. p. 593.) foliis glabris polline cæsie glaucescentibus, radicalibus cauliniisque inferioribus lyratis, summis cordato-lanceolatis amplexicaulibus crenatis, siliquis erectis. (1) culta ad oleum in Alsatia sub nom. *navette d'été*. (v. v. c.)
6. B. *NUMILIS* (DC. syst. 2. p. 598.) foliis radicalibus carnosiss pinnatifidis, lobis integris apice saepius piliferis, scapis nudis, siliquis patulis basi attenuatis stylo crassiusculo terminatis. ½ in planitie argilloss non longè à Monapello ponè montem Lupi inter Londres et Laroque aprili flor. (v. v. DC.)
7. B. *REPANDA* (DC. syst. 2. p. 598.) foliis radicalibus carnosiss glabris repando-dentatis, scapis nudis, stylo tenui à siliqua distincto. ½ in apricis montium Genevre et Galibier in Delphinatu, et in Alpibus Barcinonenais Galloprov. — *Sisymbrium monense* Vill. Dauph. 3. t. 39. non Lin. *Sis. repandum* Fl. fr. 4. p. 665. (v. v. DC.)

Sect. II. ERUCASTRUM. DC. syst. 2. p. 598.

*Siliqua sessilis, rostro conico basi 1-2-spermo superata.*

8. B. *RICHARDI* (Vill. Dauph. 3. p. 331. t. 36.) foliis glabris, inferioribus petiolatis oblongis subdentatis, summis lineari-lanceolatis paucis. ½ in asperis lapidosis pratis Alpium Delphinatûs et Galloprovinc. — All. ped. t. 58. f. 1. et 76. f. 2. Fl. fr. 4. p. 649. Petala flava. (v. v. DC.)
9. B. *ERUCASTRUM* (Lin. sp. 932.) foliis runcinatis subglabris, lobis inæqualibus obtusè sinuatis, caule basi hispido. (1) in arenosis hinc indè. — Bull. herb. t. 331. Fl. fr. 4. p. 650. Petala flava venis nigricantibus reticulata. (v. s.)
10. B. *CHEIRANTHOS* (Vill. Dauph. 3. p. 332. t. 36.) foliis petiolatis hispidis pinnatipartitis, lobis sinuato-dentatis ovali-oblongis, caule basi hispido, siliquis rostro quadruplò longioribus. (2) seu ½ Fl. flavi.
  - a. *arvensis*, caule elongatò ramoso erecto. — In arenosis asperis et ad ripas à Pictaviâ ad Delphinatum et in prov. austral. — J. Bèth. hist. 3. p. 858. f. 2. B. *Cheiranthos* Fl. fr. 4. p. 650. (v. v.)
  - β. *montana*, caule humili brevissimo cæspitoso. — In montibus excelsioribus Pyren. et Alpium. — Br. *montana* Fl. fr. 4. p. 651. *Sisymbrium vimineum* Lapeyr. non L. excl. Arnott. (v. v.)
11. B. *CHEIRANTHIFLORA* (DC. syst. 2. p. 601.) foliis radicalibus petiolatis lyrato-pinnatis-lobatis subhispidis, caulinis paucis lobis integris acutis, radice exili, siliquis rostro triplò longioribus. (1) in arenosis circa *tête de Buch*, *Sables d'Olone*, *Tarare* et in Burgundiâ. — *Raphanus cheiranthiflorus* Willd. hort. herol. t. 19. Flor. flavi. (v. v. DC.)

XL. SINAPIS. *Lin. gen. 821. DC. syst. 2. p. 607.*

Calyx basi æqualis patens. Cætera Brassicæ. — Flores flavi.

Sect. I. MELANOSINAPIS. *DC. l. c.*

Siliqua teres aut subtetragona, stylo brevi parvo apiculata.

1. *S. NIGRA* (Lin. sp. 933.) siliquis glabris lævibus subtetragonis rachi adpressis, foliis infimis lyratis, summis lanceolatis integerrimis petiolatis. ① in agris, arvis, aggeribus, ruderatis ubique. — Fl. fr. 4. p. 644. — Lob. ic t. 202. f. 2. (v. v.)

β. *torulosa* (Pers. ench. 2. p. 207.) foliis latis hastato-lobatis, super. ovatis repandis, siliquis torulosis. — Parisiis, etc. (v. s.)

γ. *turgida* (Pers. l. c.) foliis lobatis repandis basi auriculatis, siliquis turgidis venosis divergenti-adpressis, rostro conico striato. — Parisiis, etc. (v. s.)

δ? *villosa* (Mer. fl. par. 265.) foliis infer. ovatis dentatis subsinuatis glabris, superioribus lanceolatis, siliquis villosis. — Parisiis. An eadem ac *S. geniculata* Desf. (v. s.)

Sect. II. CERATOSINAPIS. *DC. syst. 2. p. 611.*

Siliqua rostro conico aspermo superata.

2. *S. ARVENSIS* (Lin. sp. 935.) siliquis glabris multangulis torulosis rostro tenui subancipiti triplo longioribus, caule foliisque pilosis. ① in arvis vineis agris ubique. — Fl. fr. 4. p. 644. — Fuchs. hist. 257. ic. (v. v.)

β. *foliis incis. Lob. ic. 198. f. 2. (v. s.)*

3. *S. ORIENTALIS* (Lin. sp. 933.) siliquis retrorsum pilosiusculis subtetragonis torulosis rostro tenui brevioribus. ① in arvis et agris Andegaviæ, Arduennæ, etc. — Schkuhr. handb. 1. t. 186. Fl. fr. 4. p. 645. (v. s.)

Sect. III. HIRSCHFELDIA. *DC. l. c.*

Siliqua teres, loculo utroque sub-4-spermo, rostro ovato 1-spermo lævi indurata.

4. *S. INCANA* (Lin. sp. 934) siliquis glabris rachi adpressis subtorulosis, caule ramoso infernè scabro, foliis lyratis scabris. ② in arvis, ruderatis, collinis lapidosisque prov. australium. — Fl. fr. 4. p. 646. Jacq. hort. vind. t. 169. *Myagrum hispanicum* Lin. sp. 893. (v. v.)

Sect. IV. LEUCOSINAPIS. *DC. l. c.*

Siliqua rostro ensiformi superata, valvis torulosis. — An *Eruca* sect? an genus?

5. *S. ALBA* (Lin. sp. 933.) siliquis hispidis patentibus rostro ensiformi subangustioribus, foliis lyratis cauleque subglabris. ① inter segetes et in ruderatis prov. austr., culta sub nom. *moutarde blanche* pro condimento. — Fl. fr. 4. p. 645. — Fuchs. hist. 538. ic. Fl. flavi. (v. v.)

XLI. MORICANDIA. *DC. syst. 2. p. 626.*

Calyx clausus basi bisaccatus; petala integra; stam. edentula; siliqua compresso-tetragona elongata, stylo compresso-conico; semina ovata parva biserialia. — Herba glabra glauca. Flores purpurascens.

1. *M. ARVENSIS* (DC. syst. 2. p. 626.) siliquis subtetragonis, foliis caulinis cordato-amplexicanlibus integerrimis. ② in arvis humentibus et glareosis Galloprovinciæ et forsan Alsatiae ex Mapp. — Bocc. sic. t. 25. f. 3-4. — *Brassica arvensis* L. mant. 95. Fl. fr. 4. p. 657. (v. v.)

β. *suffruticosa* (Desf. atl. 2. p. 94.) caulibus basi suffrutescentibus. ② in collibus Niceæ. — Fl. fr. 5. p. 589. (v. v. sp. et c. DC.)

XLII. DIPLTAXIS. DC. *syst.* 2. p. 628.

Calyx laxus basi æqualis; pet. integra; stam. edentula; siliqua compressa linearis, valvis planis nervo medio donatis, stylo conico semina 1-2 gerente aut vacuo; semina ovata parva in quoque loculo biseriaria aut rarius abortu subuniseriaria. — Flores flavi aut albi. Siliquæ in sp. Gall. erectæ, in calyce sessiles aut pedicellatæ.

1. *D. ERUCOIDES* (DC. l. c.) pubescens, siliquis sessilibus erectiusculis, stylo coniformi, foliis sessilibus runcinato-lyratis dentatis. ① in agris vineis et ad vias agri Rusciniensis et Niceæ. — Barr. ic. t. 132. *Sinapis erucoides* Lin. sp. 934. Fl. fr. 4. p. 644. Fl. albi. (v. s.)
2. *D. TENUIFOLIA* (DC. l. c.) glabra, siliquis subpedicellatis erectis, stylo filiformi brevi aspermo, foliis supremis integris, infimis pinnatifidis, lobis linearibus integris pinnatifidisve. ② in ruderatis, glareosis, lapidosis, apricis ferè ubique. — Fuchs. hist. 262. ic. *Sisymbrium tenuifolium* Lin. sp. 917. Bull. herb. t. 335. Fl. fr. 4. p. 665. Flor. flavi. (v. v.)
3. *D. MURALIS* (DC. l. c.) subpilosiuscula, siliquis sessilibus erectis, stylo brevi subfiliformi, foliis radicalibus dentatis lyratisve glabris, caulibus subnudis ascendentibus. ① ② in ruderatis, muris, glareosis, arenosis apricis hinc-indè. — Barr. ic. t. 131. *Sisymbrium murale* Lin. sp. 918. Fl. fr. 4. p. 664. Fl. flavi. (v. v.)
  - β. foliorum lobis profundioribus acutioribus dentatis. *S. erucistrum* Gou. ill. 12. t. 20. (v. s.)
  - γ. *minor*, foliorum lobis profundioribus obtusis. *S. Barrelieri* Thuil. par. 334. non Lin. (v. s. sp.)
  - δ. *pygmæa*. — *S. monense* Thuil. par. 333 non Lin.
4. *D. VIMINEA* (DC. l. c.) glabra, siliquis sessilibus erectis, stylo brevi filiformi, foliis radicalibus lyratis obtusissimis levibus, caulibus nudis subdecumbentibus. ① in vineis, cultis et arenosis Galloprov., Occitanie, Delphinatûs, agri Parisiensis, etc. — Bocc. sic. 19. t. 10. *Sisymbrium vimineum* Lin. sp. 919. Fl. fr. 4. 665. Fl. flavi. (v. v.)
5. *D. SAXATILIS* (DC. l. c.) subpilosiuscula, siliquis erectis sessilibus basi attenuatis, stylo brevi conico, foliis radicalibus pinnatilobatis crassiusculis, lobis integris, caulibus subnudis erectis. ② in asperis rupestribus collinis Galloprovinciæ propè Digne et ad montem Sanctæ Victoriæ. — Ger. gallopr. 360. n. 8. *Sisymbrium monense* Lin. sp. ed. 2. non ed. 1. *Sis. saxatile* Fl. fr. 4. p. 665. Fl. flavi. (v. v. DC.)

XLIII. ERUCA. Tourn. *inst. t. III. DC. syst.* 2. p. 637.

Calyx erectus; pet. integra; stam. edentula; siliqua ovali-oblonga, valvis concavis lævibus, rostro ensiformi aspermo vix valvulis brevior; sem. globosa. — Petala alba aut flava venis fuscis reticulata.

1. *E. SATIVA* (Lam. fl. fr. 2. p. 496.) foliis lyrato-pinnatifidis, lobis dentatis acutis, caule hirsuto, pedicellis calyce deciduo brevioribus. ① ad vias, in scrobibus, oleraceis et segetibus prov. austral. — Lob. ic. 204. f. 1. *Brassica Eruca* Lin. sp. 932. Fl. fr. 4. p. 649. Variat fl. albo et flavo, siliquis glabris aut pilosis, fructibus turgidis glabris (*B. turgida* Pers. ench. 2. p. 207.) etc. (v. v.)

Trib. X. ZILLEÆ seu ORTHOPLOCEÆ NUCAMEN-TACEÆ. DC. *syst.* 2. p. 646.

Silicula indehiscens, subglobosa, valvis indistinctis. Semina globosa in loculis solitaria. Cotyl. conduplicatæ.

XLIV. CALEPINA. Adans. *fam.* 2. p. 423. DC. *syst.* 2. p. 648.

Calyx subpatens basi æqualis; pet. obovata, exteriora paulò majora;

stam. edentula; silicula coriacea 1-locul. 1-sperma; semen pendulum. — Flores albi.

1. *C. CORVINI* (Desv. journ. 3. p. 158.) ① in arvis, vineis, arenosis et ruderalis prov. præsertim australium. — Barr. ic. 894. f. 1. et t. 1152. *Bunias cochlearioides* Fl. fr. 4. p. 721. (v. v.)

Trib. XI. *RAPHANÆ* seu *ORTHOPOCEÆ LOMENTACEÆ*. DC. syst. 2. p. 649.

Silicula seu siliqua transversè in articulos secedens. Semina globosa. Cotyl. conduplicatæ.

XLV. *CRAMBE*. Lin. gen. n. 825. DC. syst. 2. p. 650.

Calyx patens basi subæqualis; pet. æqualia integra; stam. majora apice dente laterali aucta; silicula coriacea biarticulata, articulis indehiscentibus 1-spermis, inferiore abortiente pedicelliformi brevi, sup. globoso. — Flores albi.

1. *C. MARITIMA* (Lin. sp. 937.) filamentis longioribus bifurcis, silicula mutica, foliis subrotundis crassiusculis sinuatis undulatis dentatis glaucis cauleque glaberrimis. ② in arenosis maritimis Oceani et Mediterranei. — Lob. ic. 245. f. 2. — Fl. fr. 4. p. 722. (v. v.)

XLVI. *RAPISTRUM*. Boerh. lugd. 406. DC. syst. 2. p. 430.

Calyx laxus; pet. integra; stam. edentula; silicula biarticulata, articulo superiore ovato rugoso; semina in loculis solitaria, superioris erectum, inferioris pendulum. — Flores flavi.

1. *R. PERENNE* (Berg. phyton. ic.) siliculis glabris, articulo superiore ovato stylo longiore, foliis pinnatifidis, lobis dentato-incisis acutis. ② in agris arenosis Galloprov., Lugduni, Lotharingæ, Alsatiæ? — *Myagrum perenne* Linn. sp. 893. Jacq. austr. t. 414. *Cakile perennis* Fl. fr. 4. p. 719. (v. s.)
2. *R. RUCCOSUM* (Berg. phyt. ic.) siliculis pubescentibus, articulo superiore globoso rugoso stylo brevior, foliis obtusis dentatis, radicalibus sublyratis. ① in agris et arenosis prov. præsertim australium. — *Myagrum rugosum* Lin. sp. 893. *Cakile rugosa* Fl. fr. 4. p. 719. (v. v.)
- β. *stylosum* (DC.) foliis ferè omnibus radicalibus, siliculis villosis, artic. infer. sæpè abortivo. Fl. fr. 5. p. 598. in Alsatiâ, Occitaniâ. (v. s.)
- γ. *venosum* (DC.) siliculis adultis glabris, artic. inferiore rariùs abortivo. — *Myagrum venosum* Pers. ench. 2. p. 183. Tolosæ. (v. v. DC.)
3. *R. ORIENTALE* (DC. syst. 2. p. 433.) siliculis glabris, articulo superiore globoso rugoso stylo sublongiore, inferiore sæpè abortivo, foliis oblongis, caulinis profunde dentato-sinuatis. ① in Corsicâ reperiit Ph. Thomas. (v. s.)
4. *R? HYPOGÆUM* (Duby Mss.) siliculis hispidulis, articulis 2 ovatis, scapis 1-floris foliis pinnatis brevioribus. ①? in littore australi Corsicæ (Viv.) — *Erucaria hypogæa* Viv. Cors. 11. App. 3. t. 1. f. 2. Planta valdè singularis, silicula 1-2-nodosa, locus superior 1-spermus aut aspermus aut nullus, inferior 2-spermus, contractione 2-locularis. Nec minime referri potest *Erucariæ* cum in *Erucariâ* cotyledones replicatæ subspirales et in nostra incumbentes conduplicatæ sint, ut ipse vidi. An genus novum? (v. s. D.)

XLVII. *RAPHANUS*. Lin. gen. n. 1098. DC. syst. 2. p. 662.

Calyx erectus basi subbisaccatus; pet. limbo obovato aut obcordato; stam. edentula; silicula teres stylo conico acuminata, evalvis, coriaceo-suberosa; semina 1-serialia globosa pendula.

Sect. I. *RAPHANIS*. DC. l. c.

Siliqua vix ac nevix substrangulata fungosa bilocularis; semina septulis transversis distincta.



1. *R. SATIVUS* (Lin. sp. 935.) siliquis teretibus torosis acuminatis vix pedicello longioribus. ① ex Oriente ortus, in hortis vulgatissimus. — Lam. ill. t. 566. Fl. fr. 4. p. 641. (v. v. c.)

A. RADICULA radice plus minùs carnosâ albâ rosâ aut rubrâ.

a. *rotunda* (DC.) radice subglobosa. — Lob. ic. t. 201. f. 1. Esculenta.

β. *oblonga* (DC.) radice oblongâ carnosâ. — Lob. ic. 201. f. 2. Esculenta.

γ. *oleifera* (DC.) radice exili elongata vix subcarnea. Culta rarissimè ad oleum seminum. R. Chinensis Mill.

B. *NIERA* radice carnosocompactâ duriusculâ sapore acerrimâ. — Lob. ic. 202. f. 1. Variat radice oblonga et variâs rotunda, nigrâ griseâ et variâs alba.

#### Sect. II. RAPHANISTRUM. DC. l. c.

Siliqua per maturitatem moniliformis, articulis 1-spermis.

2. *R. RAPHANISTRUM* (Lin. sp. 953.) siliquis unilocularibus articulatis striatis 3-8-spermis stylo longioribus, foliis simpliciter lyratis. ① in arvis, agris, cultis et arenosis ubique. — Fl. fr. 4. p. 643. — Lob. ic. 199. f. 1. Variat flore albo flavo aut purpurascens, unicolori aut venis fuscis reticulato. (v. v.)

3. *R. MARITIMUS* (Sm. engl. bot. t. 1643.) siliquis 1-locul. articulatis striatis 2-6-spermis, stylo conico articulo ultimo fere brevior, foliis interruptè lyratis. ② in maritimis Armoraciz inter Brest et Quimper. — Fl. fr. 5. p. 588. Flor. flavi. (v. s.)

Subordo IV. SPIROLOBEÆ. (o || ||) DC. syst. 2. p. 670.

Cotyledones incumbentes lineares circinnatim convolutæ. Semina subglobosa.

#### Trib. XII. BUNIADEÆ seu SPIROLOBEÆ NUCAMENTACEÆ. DC. syst. 2. p. 670.

Silicula nucamentacea indehiscens 2-4-locul., loculis 1-spermis.

##### XLVIII. BUNIAS. Brown. DC. syst. 2. p. 670.

Calyx basi æqualis; pet. unguiculata, limbo obcordato; stam. edentula; silicula nucamentacea. — Flores flavi.

1. *B. BAUCAGO* (Lin. sp. 935.) siliculis tetragonis, angulis cristatis, foliis radicalibus runcinatis. ① in arvis et agris prov. austr. — Fl. fr. 4. p. 720. — C. Bauh. prod. 41. ic. (v. v.)

Subordo V. DIPLECOLOBEÆ. (o || || ||) DC. syst. 2. p. 776.

Cotyledones incumbentes lineares bicrures seu biphlicatæ nempè bis transversè plicatæ. Semina depressa.

#### Trib. XIII. SUBULARIÆ. DC. syst. 2. p. 697.

Silicula ovalis, septo elliptico, valvis convexis, loculis polyspermis, stigmatè sessili.

##### XLIX. SUBULARIA. Lin. gen. n. 799. DC. syst. 2. p. 697.

Cal. suberectus; pet. ovalia; stam. edentula; silicula in quoque loculo 4-sperma. — Flores albi. Folia radicalia subulata.

1. *S. AQUATICA* (Lin. sp. 896.) ① in laenibus et inundatis Vogesorum ex Lois., in piscinâ propè Basileam ex Lach. — *Draba subularia* Lam. ill. t. 556. f. 3. DC. fl. fr. 5. p. 595. (v. s. sp. ex Campinâ Leodensi.)

Ordo VII. CAPPARIDÆ. *Juss. gen. 242. DC. prod. 1. p. 237.*

Sepala 4, nunc ferè libera, æqualia aut inæqualia, nunc basi in tubum coalita, limbo vario. Petala 4 cruciata sæpè unguiculata et inæqualia. Stamina ferè perigyna nempe toro aut imò calyci inserta, rariùs tetradynama, sæpiùs ordine quaternario multiplici disposita, defnita aut indefinita. Torus hemisphæricus aut elongatus sæpè glandulifer. Thecaphorum gracile sæpè basi toro vaginatum. Ovarium stipitatum ovato-oblongum carpellis 2 arctè coalitis compositum. Stylus 0 aut filiformis. Fructus varius, siliquosus baccatusve, intùs 1-locularis, placentis 2 polyspermis in fructibus dehiscentibus intervalvularibus. Semina sæpiùs reniformia exalbuminosa (endoplevrà tamen tumidà). Embryo incurvus. Cotyledones foliaceæ planiusculæ subincumbentes. — Herbæ frutices aut arbores. Stipulæ veræ nullæ. Spinæ stipulares. Folia alterna petiolata. — Ordo vix in Florà gallicà recipiendus.

I. CAPPARIS. *Lin. gen. n. 643.*

Calyx 4-partitus; pet. 4; torus parvus; thecaphorum gracile; stam. ∞; siliqua subbaccata stipitata. — Frutices foliis simplicibus integris.

1. *C. spinosa* (Lin. sp. 720.) spinis stipularibus uncinatis foliis subrotundis retusis, pedicellis solitariis 1-floris. ♂ in fissuris rupium mris et lapidosis culta in Galloprovincià, et rariùs in Occitanià, agro Ruscinnonensi et Turienensi. Blackw. herb. t. 417. Fl. fr. 4. p. 724. (v. v.)

Ordo VIII. CISTINEÆ. *Dunal in DC. prod.* 1. p. 263. — *Cisti DC. fl. fr.* 4. p. 811.

Calycis sepala 5 cum pedicello continua persistentia sæpius inæqualia, 2 exteriora cæteris minora interdum evanida, 3 interiora per æstivationem contorta. Petala 5 hypogyna caduca æqualia per æstivationem contorta sed directione petalis contraria. Stam. numero indefinita pauca aut plurima hypogyna erecta, filam. liberis, antheris ovatis bilocul. birimosi basi insertis. Ovarium liberum. Stylus 1 filiformis. Stigma simplex; capsula constans valvis 3-5 rarius 10 nunc medio nervum longitudinalem placentarium gerentibus et tunc capsula 1-locul., nunc nervo medio in septum plus minusve completum abeunte et tunc capsula completè aut incompletè multilocularis. Semina idèò aut placentæ nunc quasi parietali nunc septo intus adnatæ inserta, plurima, parvula. Albumen farinosum. Embryo spiralis aut curvus intrà albumen. — Suffrutices aut Herbæ. Folia simplicia penninervia integra aut subdentata, primordialia semper opposita, cætera sæpius opposita interdum alterna, basi nunc nuda, nunc stipulis 2 foliaceis instructa. Racemi floribus sæpius 1-lateralibus successivis. Flores petalis fugacissimis diurnis flavis albis aut purpureis.

I. CISTUS. *Tourn. inst.* t. 136. (DC.) *fl. fr.* 4. p. 811.

Capsula 5-aut 10-locularis, valvis 5-10 medio septiferis; seminum embryo filiformis spiralis. — Frutices suffruticesve.

Sect. I. ERYTHROCISTUS. *Dun. in. DC. prod.* 1. p. 264.

Sepala exteriora angustiora sæpè minora, interna basi concava margine scariosa; petala rosea aut purpurea; capsulæ 5-loculares.

1. *C. INCANUS* (Lin. sp. 737.) foliis spathulatis tomentosis rugosis subtrinerviis sessilibus subconnatis, superioribus angustioribus, pedunculis 1-floris.  $\frac{3}{4}$  in collibus incultis apricis et lapidosis Occitanie. — *Fl. fr.* 4. p. 812. — *Clus. hist.* 1. p. 69. f. 1. (v. v.)
2. *C. castrus* (Lin. sp. 738.) foliis sessilibus lineari-lanceolatis undulato-crispis trinerviis rugosis pubescentibus, floribus subsessilibus ternis quaternisve umbellatis.  $\frac{3}{4}$  in collibus apricis et lapidosis prov. oleiferarum. — *Fl. fr.* 4. p. 811. *Cav. ic. t.* 174. — *Clus. hist.* 1. p. 69. f. 2. (v. v.)
3. *C. ALPIDUS* (Lin. sp. 737.) foliis sessilibus oblongo-ellipticis incano-tomentosis subtrinerviis, floribus 3-8 terminalibus subumbellatis, sepalis externis majoribus.  $\frac{3}{4}$  in apricis lapidosis collibus prov. oleiferarum. — *Fl. fr.* 4. p. 812. *Dun. in DC. prod.* 254. — *Clus. hist.* 1. p. 68. *ic.* Petala dilutè purpurea basi maculâ parvâ lateâ. (v. v.)

Sect. II. LEDONIA. *Dun. l. c.*

Sepala 2 exteriora majora valdè acuminata vel nulla; petala alba aut albida; stam. oo pistillo longiora; stigma subsessile capitatum magnum; capsulæ 5-10 locul. — Folia sæpè glutinosa.

\* *Calycibus 5-sepalis.*

4. *C. SALVISOLIS* (Lin. sp. 738.) foliis petiolatis ovatis obtusis rugosis subtilis

- tomentosis, pedunculis longis tomentoso-albicantibus 1-floris supernè articulatis solitariis.  $\S$  in aridis subsylvaticis dumetis Pictaviæ, Vasconia, Delphin., Galloprov., Corsicæ. — Fl. fr. 4. p. 813. var.  $\alpha$ . — Clus. hist. 1. p. 70. ic. (v. v.)
5. *C. CORBARIENSIS* (Pourr. ex Dun. l. c.) foliis petiolatis subcordatis ovatis acuminatis margine fimbriatis utrinquè rugosis leviter glutinosi, pedunculis longis 1-5-floris.  $\S$  in montibus Narbonensibus dictis les Corbières. — *C. hybridus*. Pourr. chl. 36. *C. salvifolius*  $\beta$  Fl. fr. 4. p. 813. Rami autumnales *C. populifolii* ex cl. Benth. (v. v. DC.)
6. *C. MONSPELIENSIS* (Lin. sp. 737) foliis lineari-lanceolatis sessilibus trinerviis utrinquè villosis, pedunculis cymosis subsecundis.  $\S$  in apricis aridis prov. oleiferarum. — Lam. ill. t. 477. f. 4. DC. fl. fr. 4. p. 814. (v. v.)
7. *C. LEDON* (Lam. dict. 2. p. 17.) foliis connatis oblongo-lanceolatis nervosis suprâ glabris nitidis subtus villosis-sericeis, floribus corymboso-cymosis, pedunculis calycibusque villosis-sericeis.  $\S$  in aridis dumetis Occitanie et Galloprov. — Duham. arb. ed. nov. 1. p. 168. t. 66. Fl. fr. 4. p. 814. (v. v.)
8. *C. HIRAUTUS* (Lam. dict. 2. p. 17.) foliis sessilibus oblongis obtusis hirsutis, pedunculis brevibus 1-floris aut cymoso-multifloris, capsulis parvis calyce maximo hirsuto et pyramidalit tectis.  $\S$  in Armoraciâ propè Landerneau. — Fl. fr. 5. p. 621. excl. syn. Ait. ex Dun. DC. prod. 1. p. 265. (v. s.)
9. *C. LONGIFOLIUS* (Lam. dict. 2. p. 16.) foliis breviter petiolatis oblongo-lanceolatis margine pubescentibus et undulatis subtus venosis, pedunculis cymosis bracteolis infimis caducis.  $\S$  in sterilibus ad Donos et Fontfroide circâ Narbonam. — Fl. fr. 4. p. 813. (v. s.)
10. *C. POPULIFOLIUS* (Lin. sp. 736) foliis petiolatis cordatis acuminatis lævibus, floribus cymosis, pedunculis bracteatis, bracteolis oblongis infimis caducis.  $\S$  var.  $\alpha$  pedunculis longè hirsutis in Mauritanîâ. Var.  $\beta$  calycibus glabriusculis ad Fontlaurier in montibus Corbariensis agri Narbonensis. — Fl. fr. 5. p. 620. — Clus. hist. 1. p. 78. f. 1. 2. (v. s.)

\*\* *Calycibus 3-sepalis.*

11. *C. LAURIFOLIUS* (Lin. sp. 736.) foliis petiolatis ovato-lanceolatis trinerviis suprâ glabris subtus tomentosis, petiolis basi dilatatis connatis, capsulis 5-locularibus, stigmate sessili.  $\S$  in collibus aridis subsylvaticis Occitanie, Vasconie, agri Ruscinonensis. — Fl. fr. 4. p. 814. — Clus. hist. 1. p. 78. f. 1. (v. v.)
12. *C. LADANIFERUS* (Lin. sp. 737.) foliis subsessilibus basi connatis lineari-lanceolatis suprâ glabris subtus tomentosis, capsulis 10-locularibus.  $\S$  circâ Forum-Julii (cl. Solier.) (v. s.)

## II. HELIANTHEMUM. Tourn. inst. 128. DC. fl. fr. 4. p. 815.

Calyx 3-sepalus aut 5-sepalus, sepalis 2 exterioribus minoribus; capsula trivalvis valvis medio septi aut seminiferis; embryo uncinato-inflexus. — Herbæ aut suffrutices.

### Sect. 1. HALIMIUM. Dun. in. DC. prod. 1. p. 267.

Calyx 3 rarè 5 sepalus, sepalis 2 externis minutis; stigma capitatum sub 3-lobum. — Suffrutices. Folia exstipulata.

1. *H. UMBELLATUM* (Mill. dict. p. 5.) caule ramoso, ramis junioribus tomentoso-pilosis viscosis, foliis sessilibus lineari-oblongis margine revolutis viscosis subtus tomentosis, pedunculis unifloris racemoso-verticillatis terminalibus umbellatis, calyce trisepalo villosis.  $\S$  in dumetis circâ Fontembleaudi Cænomanum, Aureliam, Burdigalam (Laterr.), Aginnum et in agro Ruscinonensi. — Fl. fr. 5. p. 815. (v. v. DC.)
2. *H. ALYSSOIDES* (Vent. choix. t. 20.) caule ramosissimo diffuso patulo, foliis sessilibus basi attenuatis oblongo-ovatis obtusiusculis breviter hirsutis, pedunculis terminalibus solitariis seu umbellatis 1-2 floris folio longioribus,

calycibus 3-sepalis acuminatis hirsutis. ♂ in Galliâ occidentali. — Fl. fr. 4. p. 818. (v. v.)

β. *rugosum* foliis margine subdenticulatis subseriis utrinque tomentoso-rugosis subtus incano-sericeis, pedunculis folio brevioribus. In Galliâ occidentali circa Burdigalam. — H. alyssoides L. Fl. fr. 5. p. 622. H. rugosum Dun. in DC. prod. 1. p. 268. (v. s.)

3. H. *HALIMPOLIUM* (Willd. enum. 569.) caule fruticoso erecto ramoso, foliis utrinque leproso-candidis subpetiolatis ovato-oblongis, pedunculis longis subpaniculatis, calycibus 5-sepalis. ♂ in Corsicâ. — Fl. fr. 5. p. 621. — Lob. ic. 2. p. 113. f. 1 et 2. (v. s.)

Sect. II. TUBERARIA. Dun. l. c. p. 270.

Calyx 5-sepalus, sepalis externis minoribus vel majoribus; stamina numerosa pistillo multo longiora; stylus subnullus. — ♀ ant. ☉. Folia inferiora opposita exstipulata, superiora interdum alterna sæpius stipulata, stipulis longis linearibus acutis.

4. H. *TUBERARIA* (Mill. dict. n. 10.) caulibus subsimplicibus, foliis radicalibus in petiolum attenuatis ovato-oblongis canescentibus subtus nervosis, canaliculis sessilibus subglabris lanceolatis, calycibus glabris nitidis. ♀ in rupetribus Nicæ, Galloprov., Corsicæ. — Fl. fr. 4. p. 818. — J. Bauh. 2. p. 12. f. 4. et p. 13. f. 1. (v. v.)
5. H. *PLANTAGINEUM* (Pers. ench. 2. p. 77.) caule hirsuto, foliis elliptico-lanceolatis oppositis sessilibus, subtus in nervis hirsutis, supra hirsutis, summis oblongo-linearibus stipulatis subalternis, racemis brevibus ebracteatis, sepalis externis subglabris angusto-linearibus internos villosos-incanos subquantibus, petalis denticulatis. ♂ in Corsicâ. — Dun. l. c. An a seq. satis distincta? (v. s.)
6. H. *GUTTATUM* (Mill. dict. n. 18.) caule subhirsuto, foliis oppositis sessilibus oblongo-linearibus villosis-hirsutis, extremis alternis, racemis laxis ebracteatis, pedicellis filiformibus subnudis, sepalis externis internis dimidio brevioribus. ♂ in Galliâ mediâ, australi et occidentali. — Fl. fr. 4. p. 819. 5. p. 623. Cum hac specie conjungit cl. Benth. sp. præcedentem et 3 subsequentes, forsitan non immerito. (v. v.)
7. H. *XERIOCAULON* (Dun. l. c.) caule ramoso 2-3-chotomo hirsutissimo, foliis oblongo-linearibus angustis hirsutis oppositis, superioribus stipulatis, extremis alternis, racemis simplicibus ebracteatis, pedicellis filiformibus longis pilosis, sepalis externis angustis. ♂ in Galliâ ex Dun. (v. s. ex Hispaniâ.)
8. H. *INCONSPICUUM* (Pers. ench. 2. p. 77.) caule ramoso 2-3-chotomo parce hirsuto gracili, foliis oblongo-linearibus angustis hirsutis oppositis, superioribus stipulatis, extremis alternis, racemis longis filiformibus, pedicellis brevibus secundis, floribus minutis, petalis oblongo-linearibus calyce minoribus altero margine pilosis. ♂ in Corsicâ. — Dun. l. c. (v. s.)
9. H. *PUNCTATUM* (W. enum. 570.) caule ramoso dichotomo puberulo subcinereo, foliis oblongis 3-5-nerviis viridi-cinereis breviter piloso-scabriusculis, pilis stellatis minimis, superioribus alternis acutiusculis stipulatis, racemis longis pubescentibus cinereis paucifloris. ♂ in Galliâ occidentali circa Dax. — Dun. l. c. An sp. distincta? (v. s.)

Sect. III. BRACHYPETALUM. Dun. l. c. p. 271.

Calyx 5-sepalus, sepalis externis minutis, internis 3-nerviis acuminatis, petala sæpe calyce breviora; stamina 10-20; capsula 3-quetra subnitida. — Herbæ ☉. Folia petiolata, stipulata, stipulis oblongo-linearibus.

10. H. *LADIPOLIUM* (W. enum. 571.) caule erecto subglabro, foliis petiolatis oblongo-ellipticis subdenticulatis glabrescentibus tomentosisve oppositis, superioribus alternis floribus oppositis, pedunculis glabrescentibus tomentosisve erectis calyce brevioribus, calycibus acuminatis subpubescentibus. ☉ in siccis

- Nicæ, Galloprov., Occitaniz. — Fl. fr. 4. p. 819. — Lob. ic. 2. p. 118. f. 2. H. Niloticum Pers. Dun. in DC. prod. 1. p. 272. (v. s.)
11. H. *INTERMEDIUM* (Thib. ex Dun. l. c.) caule erecto ramoso, foliis obovato-oblongis petiolatis obtusiusculis subdenticulatis, summis alternis oppositifloris, pedunculis folio longioribus calycibusque villosis-cinereis, calycibus fructiferis oblongo-lanceolatis erectis. ① ad vias circa Celleneuve propè Monspelium ex cl. Benth. cat. 84. — Cav. ic. t. 144. (v. s. ex Hisp.)
12. H. *DENTICULATUM* (Pers. ench. 2. p. 78.) caule ramoso erecto vel patulo, ramis tomentoso-pubescentibus apice subcinereis, foliis breviter petiolatis obovato-oblongis acutiusculis subdenticulatis tomentosis subtus incano-sericeis, stipulis linearibus, pedunculis calycibusque oppositis bracteis, bracteis alternis subincisis. ① in Occitania, et agro Ruscini. ex Benth. Bractez subovatz. (v. s.)
13. H. *SALICIFOLIUM* (Pers. l. c.) caule ramoso, ramis subhirsutis, foliis breviter petiolatis obovato-oblongis acutiusculis denticulatis subtomentosis, stipulis lineari-oblongis, pedunculis calycibusque hirsutis suboppositibracteis, bracteis ovatis acumine integris. ① in sterilibus Nicæ, Galloprov., Lugduni, Pictaviæ. — Fl. fr. 4. p. 820. — Seg. ver. 3. t. 6. f. 3. (v. v.)

Sect. IV. *FUMANA*. Dun. l. c. p. 274.

Calyx ante anthesin apice tortus 5-sepalus, sepalis externis angustis parvulis, internis 3 ovato-acuminatis 4-5-venosis margine scariosis; petala calyce subduplò longiora; stamina pauca; stigma sub-3-lobum; capsula 3-valvis. — Caules 5. Folia linearia.

14. H. *FUMANA* (Mill. dict. n. 6.) caule ramoso tortuoso subdiffuso erectiusculo, foliis exstipulatis alternis margine piloso-scabriusculis subinvolutis, inferioribus brevibus confertis, pedunculis folio longioribus, capsulis apertis nudis. 5 in collibus apricis. — Fl. fr. 4. p. 816. — Desf. atl. 1. t. 205. — Barr. ic. 286 et 446. (v. v.)
15. H. *PROCUMBENS* (Dun. l. c.) caule procumbente ramoso, ramis elongatis junioribus canescentibus, foliis exstipulatis alternis sublaixis margine et subtus pilosis, pilis strigosis, pedunculis folio brevioribus, capsulis apertis semina gerentibus. 5 in Gallia australi. — Barr. ic. 445. An præcedentis var. ut vult cl. Benth. (v. s.)
16. H. *LÆVIPS* (W. enum. 370.) caule ascendente, foliis alternis setaceis glaucis subglabris stipulatis, stipulis filiformibus longis, pedunculis longis racemosis secundis, pedicellis glabris basi bracteatis, calycibus hirsutis. 5 inter saxa aprica in provinciis mediterraneis. — Fl. fr. 4. p. 816. Ger. Gallopr. t. 14. (v. v.)
17. H. *GLUTINOSUM* (DC. fl. fr. 4. p. 821.) caule ascendente ramis villosoglutinosus subcinereis, foliis sublinearibus margine revolutis villosoglutinosus oppositis, summis alternis, stipulis mucronatis erectis, pedunculis calycibusque villosoglutinosus, seminibus tenuissimè excavato-punctatis. 5 in Gallia australi. — Barr. ic. t. 415. (v. v.)
- β. thymifolium* (DC. l. c.) foliis glabriusculis brevioribus. H. thymifolium Pers. Dun. in DC. prod. 1. p. 276. — Barr. ic. t. 444. (v. s.)
- γ. juniperinum* (Benth.) foliis inferioribus glabris. H. juniperinum Lag. ex Dun. l. c. — Barr. ic. t. 443. (v. s.)

Sect. V. *PSEUDOCISTUS*. Dun. l. c.

Calyx 5-sepalus, sepalis externis angustis minutis, internis 4-venis; petala calyce vix duplò longiora; stylus basi contortus; stigma capitatum 8-lobum. — Herbæ 2 aut 3. Folia petiolata opposita rarò stipulata.

18. H. *ORIGANIFOLIUM* (Pers. ench. 2. p. 76.) caule suffruticoso 2-3-chotomo, foliis petiolatis ovatis utrinquè pilosis, racemis brevibus terminalibus, petalis calyce vix majoribus. 5 circa Alais et Monspelium. — Cav. ic. 3. t. 262. f. 1. (v. v. DC.)

19. *H. OBLANDICUM* (DC. fl. fr. 4. p. 817.) caule suffruticoso procumbente ramoso, foliis lanceolato-ellipticis obtusiusculis utrinque viridibus sæpè glabris interdum ciliatis petiolatis, summis sessilibus, racemis simplicibus paucifloris, calycibus subgloboso-ovatis.  $\zeta$  in Alpibus? — *C. celandicus* L. sp. 741. Clus. hist. p. 73. ic. Petala calyce vix longiora. (v. s. ex Succia.)
20. *H. ALPESTRIS* (Dun. in DC. prod. 1. p. 276.) caule suffruticoso procumbente ramoso, ramis piloso-hirsutis, foliis utrinque subviridibus oblongo-ellipticis subglabris vel fasciculatim piloso-hirsutis petiolatis, summis subsessilibus, pedicellis calycibusque piloso-hirsutis, pilis albedo-cinereis.  $\zeta$  in petrosis montosis Jurassi, Galliae australis, occidentalisque. — *C. alpestris* Crantz, Aust. t. 6. f. 1. Petala calyce duplò longiora. (v. v.)
21. *H. FENICILLATUM* (Thib. ex Dunal. l. c.) caule suffruticoso, ramis procumbentibus longis foliisque viridibus utrinque hispidis, inferioribus petiolatis ovatis, superioribus lineari-oblongis subsessilibus, calycibus hispidis, floribus minutis.  $\zeta$  circa Monspelium, et in aridis apricis prov. medit. An sp. satis distincta? (v. s.)
22. *H. ITALICUM* (Pers. ench. 2. p. 76.) caule suffruticoso, ramis simplicibus erectis longis piloso-tomentosis, foliis piloso-hispidis, pilis strigosis adpressis, inferioribus ovatis, superioribus lanceolatis oblongis, calycibus piloso-hispidis canescentibus.  $\zeta$  in siccis Cebennarum et probabiliter omnis regionis mediterraneæ. — Barr. rar. t. 366. (v. s.)
23. *H. CANUM* (Dun. in DC. prod. 1. p. 277.) caule suffruticoso procumbente, ramis ascendentibus piloso-tomentosis incanis, foliis ovatis ellipticisve pilosis subtus tomentoso-canis, pedicellis calycibusque piloso-canescensibus.  $\zeta$  in rupibus apricis submontanis. — *Cistus canus* L. sp. 740. *H. vineale* Pers. Dunal. l. c. *Cistus piloselloides* Lapeyr. Abr. 301. — Clus. hist. 1. p. 74. f. 1. An satis à sequenti distinctum? (v. v.)
24. *H. MARIFOLIUM* (DC. fl. fr. 4. p. 817.) caule suffruticoso procumbente, foliis exstipulatis petiolatis ovato-cordatis vel ovatis acutiusculis suprà viridibus pilosis subtus incanis, racemis solitariis simplicibus paucifloris terminalibus elongatis.  $\zeta$  in montosis australioribus. — Barr. t. 441. (v. v.)

#### Sect. VI. EUHELIANTHEMUM. Dunal. l. c.

Calyx antè anthesin apice subortus, 5-sepalus, sepalis externis multò minoribus, internis 2 sæpius 4-costatis sulcatis; petala calycibus 2-4-plo longiora; stylus basi flexus; stigma simplex; capsula calyce oblecta 1-localaris. —  $\zeta$  foliis oppositis breviter petiolatis, stipulis lineari-lanceolatis.

#### \* *Petalis luteis.*

25. *H. LAVANDULIFOLIUM* (DC. fl. fr. 4. p. 820.) caule erecto ramoso, ramis canescentibus, foliis oblongo-linearibus margine revolutis subtus tomentoso-incanis, junioribus utrinque canescentibus, stipulis bracteisque lineari-acutis ciliatis, floribus confertis, sepalis glaucis ciliatis, internis acutis.  $\zeta$  in Corsicà et circà Massiliam. — Barr. ic. t. 288. (v. v.)
26. *H. STACHADIFOLIUM* (Pers. ench. 2. p. 79.) caule erecto, ramis tomentosis, canescentibus, foliis oblongo-linearibus obtusiusculis utrinque subtomentosis subtus incanis margine revolutis, stipulis subvillosis lineari-lanceolatis, floribus confertis, sepalis villosis externis ciliatis viridibus, internis acuminatis incanis.  $\zeta$  in Corsicà (Salzmann.) — Dunal. in DC. prod. 1. p. 279. (v. s.)
27. *H. TOMENTOSUM* (Dun. in DC. prod. 1. p. 729.) caule suffruticoso ramoso, ramis elongatis subcanescentibus, foliis lanceolato-oblongis margine sæpè revolutis subtus incano-tomentosis suprà glabriusculis, calycibus sulcatis, nervis elevatis pilosis.  $\zeta$  in Gallià ex Dunal. (v. s. in Hisp.)
28. *H. ACUMINATUM* (Pers. ench. 2. p. 79.) ramis erectis, foliis longè petiolatis oblongis margine revolutis utrinque viridibus pilosis subtus tomentosisculis, stipulis subglabris linearibus petiolo longioribus, racemis laxis, calycibus glabris nitidis.  $\zeta$  in agro Niceensi. An cum sequenti conjungendum ut vult cl. Benth? (v. s.)

29. *H. VULGARE* (Gærtn. fruct. 1. p. 371. t. 6.) caule suffruticoso procumbente ramoso, foliis margine vix revolutis subtus incano-cinereis suprà viridibus pilosis subciliatis, inferioribus suborbiculatis, superioribus oblongis, stipulis oblongo-linearibus ciliatis petiolo longioribus pedicellis calycibusque pilosis. ♂ in sylvis siccis frequens. — Fl. fr. 4. p. 821. Fl. dan. t. 101. Cistus hirsutus Lapeyr. Abr. 303. ex autopsiâ cl. Arnott. (v. v.)
30. *H. GRANDIFLORUM* (DC. fl. fr. 4. p. 801.) caule suffruticoso ascendente, foliis superioribus subplanis oblongis pilosiusculis subtus nunc viridibus nunc dilutè cinereis, stipulis ciliatis petiolo sublongioribus, floribus magnis, calycibus subhirsutis. ♂ in Pyrenæis et Galloprovinciæ alpestribus. — Scop. Carn. ed. 2. t. 25. (v. v.)
31. *H. OSCURUM* (Pers. ench. 2. p. 79.) caule suffruticoso ascendente ramosissimo, foliis utrinquè hirsutis subviridibus, superioribus ellipticis, stipulis ciliato-hirsutis petiolo longioribus, racemis longis, calycibus hirsutis. ♂ in sylvis. — Fl. fr. 5. p. 624. var. *α* (v. v.)
- β. nummularium* (DC.) foliis inferioribus orbiculatis, superioribus oblongo-linearibus. — In Galliâ australi. — Fl. fr. 5. p. 624. — J. Bauh. hist. 2. p. 20. f. 3. An hæc 4 species meritò distinguuntur? (v. s.)
32. *H. RIGIDUM* (Pers. l. c.) caule suffruticoso ramoso ramis cinereis, foliis ovatis oblongiæ margine revolutis tomentoso-hirtis subtus canescentibus suprà viridi-cinereis, stipulis angustis petiolo sublongioribus, calycibus hirsutissimis albidis. ♂ in aridis Galliæ australis. — Fl. fr. 4. p. 822. Cav. ic. 2. t. 146. Calyces parvi. (v. v.)
- \*\* *Petalis coloratis.*
33. *H. PILOSUM* (Pers. ench. 2. p. 79.) caule ramoso ramis erectiusculis, foliis linearibus lineari-oblongiæ utrinquè incanis apice setosis, stipulis subulatis, calycibus pilosiusculis nervoso-striatis. ♂ in apricis Galliæ australis. — Fl. fr. 4. p. 823. — All. ped. t. 45. f. 2. (v. v. DC.)
34. *H. AFRICANUM* (DC. fl. fr. 4. p. 824.) caule ramoso ramis patulis, foliis petiolatis oblongo-linearibus margine vix revolutis subtus tomentosis suprà glaucescentibus, stipulis subulatis petiolo longioribus, calycibus brevissimè villosis striatis glauco-cinereis obtusiusculis. ♂ circà Parisios, in Jurasso australi, Occitania, Andegaviâ. (Bast.) (v. s.)
- β. hispidum*, calycibus piloso-hirsutis. — In Galliâ australi. (Dun.) — *H. hispidum*. Dun. in DC. Prod. 1. p. 282. (v. s. ex Liguriâ.)
35. *H. PULVERULENTUM* (DC. fl. fr. 4. p. 823.) caule ramosissimo prostrato, foliis oblongo-linearibus margine revolutis obtusis subtus incanis suprà glaucis, stipulis subulatis petiolo longioribus, calycibus canescentibus minutè tomentoso-pubescentibus. ♂ in collibus sterilibus Galliæ mediæ et australis. — Petala alba. (v. v.)
36. *H. POLYFOLIUM* (DC. fl. fr. 4. p. 823.) ramis procumbentibus glabriusculis, foliis oblongis ovatis obtusiusculis subplanis subtus tomentoso-incanis suprà glabris, stipulis bracteisque linearibus viridibus ciliatis, calycibus striatis glabriusculis subnitidis. ♂ in sylvis Galliæ ex Lam. et Dun. — Dill. elth. t. 145. f. 172. Petala alba. (v. s. ex Angliâ.)
37. *H. ROSEUM* (DC. fl. fr. 4. p. 822.) caule subprocumbente tomentosiusculo, foliis ovato-lanceolatis utrinquè tomentosiusculis subtus pallidè cinereis, stipulis linearibus, pedicellis calycibusque piloso-subhirsutis. ♂ circà Nicæam et in Pyrenæis. — All. ped. t. 45. f. 4. *H. vulgare γ versicolor* Benth. An *H. vulgare* varietas? (Dun.) (v. s.)
38. *H. MAJORANÆ-FOLIUM* (DC. fl. fr. 5. p. 625. *α.*) caule erecto ramosissimo, foliis petiolatis ovato-oblongis acutiusculis margine revolutis subtus incano-tomentosis suprà viridi-glaucescentibus tomentoso-hirtis, stipulis subulatis setosis, calycibus hirsutissimis, pilis albidis. ♂ circà Monspelium, Viudomagum, Arelatem. (v. s.)



Ordo IX. VIOLARIEÆ. DC. fl. fr. 4. p. 801. Juss. ann. mus. 18. p. 476.

Calyx sepalis 5 persistentibus liberis aut basi inter se sub-coalitis sæpius deorsum productis, æstivatione imbricatâ. Petala 5 cum sepalis alterna, hypogyna, thalamo inserta, (in nostris) inæqualia; inferius basi calcaratum cucullatumve. Stam. 5 thalamo inserta cum petalis (in nostris) alterna; antheræ 2-loculares, sæpè liberæ, ovario adpressæ interdum basi plus minusve in discum monadelphum solutæ; filamenta sæpissimè dilatata. Ovarium 1-loculare, placentis 3 parietalibus, stylus 1; capsula 3-valvis polysperma. Albumen carnosum. Embryo rectus.—Herbæ aut suffrutices foliis sæpius alternis, stipulatis.

I. VIOLA. *Toung. inst.* 419. t. 286. *Ging. in DC. Prod.* 1. p. 291.

Calyx sepalis in appendices basi productis; petala inæqualia, inferius deorsum in calcar cavum plus minus productum; stamina approximata aut coarctata (nec coalita), filamentis basi dilatatis.

Sect. I. NOMIMUM. *Ging. l. c.*

Stigma rostratum, foraminulo ad extremitatem rostri sito; stylus ab apice ad basin attenuatus; capsula sæpius 3-gona.

§. 1. *Stigmatibus apice depressis; caulibus subnullis.*

1. V. *FINNATA* (L. sp. 1323.) stigmate 3-angulari marginato, radice divisâ sublignosâ, foliis pinnatis multipartitis partitionibus multilobatis, sepalis ovatis.  $\mathcal{L}$  in *Alpibus Delphinatûs* ex Vill. — Fl. fr. 4. p. 802.—J. Bauh. 3. p. 544. f. 2. (v. s. ex Helv.)
2. V. *PALUSTRIS* (L. sp. 1324.) stigmate marginato, radicibus articulatis squamosis, foliis reniformi-cordatis, stipulis lato-ovatis acuminatis, sepalis ovatis obtusis, capsulâ oblongo-trigonâ.  $\mathcal{L}$  in muscosis humidis montanis — Fl. fr. 4. p. 804. — Moris. s. 5. t. 35. n. 5. (v. v.)

§. 2. *Stigmatibus convexis, deorsum immarginatis.*

3. V. *HIRT*A (L. sp. 1324.) villosa, stigmate uncinato acuto nudo, foliis cordatis, stipulis glandulosi dentatis, sepalis ovatis obtusis margine ciliatis, petalis 2 mediis subbarbatis, calcare subconico nectariis glabris, capsulis turgidis hirtis.  $\mathcal{L}$  in nemorosis et pratis. — Fl. fr. 4. p. 802. — Moris. s. 5. t. 35. n. 4. (v. v.)
4. V. *PYRENEICA* (Ram. in DC. fl. fr. 4. p. 803.) stigmate uncinato nudo, foliis latè subcordatis, petiolis apice dilatatis, sepalis ovatis obtusis.  $\mathcal{L}$  in petrosis *Pyreneorum* (v. s.)
5. V. *ODORATA*. (L. sp. 1324.) stigmate uncinato nudo, foliis rotundato-cordatis, sepalis ovatis obtusis, calcare obtusissimo, capsulâ turgidâ hirtâ, stolonibus flagelliformibus.  $\mathcal{L}$  in nemorosis et sepibus. — Fl. fr. 4. p. 803. Bull. herb. t. 160. (v. v.)
6. V. *MIRABILIS* (L. sp. 1326.) stigmate subreflexo nudo, caulibus rigidis petiolisque villosis, pilis secundis, foliis reniformi-cordatis acuminatis, sepalis 3 maximis oblongis acuminatis, calcare cylindrico obtuso longo, capsulæ valvis peracuminatis.  $\mathcal{L}$  in sylvis et dumetis montium *Delphinatûs* ex Vill. — Fl. fr. 4. p. 805. — Dill. elth. t. 303. f. 390. (v. s. ex Pedem.)

7. *V. CANINA* (L. sp. 1324.) stigmatē papilloso subreflexo, caule adaltiore ascendente ramoso glabro, foliis cordatis, stipulis acuminatis pectinatim tenuiter laceris, sepalis subulatis, pedunculis glabris, capsulæ elongatæ trigonæ valvis acuminatis, seminibus pyriformibus fuscis.  $\mathcal{U}$  in sylvis dumetis sepibusque. — Fl. 4. p. 806. — J. Bauh. hist. 3. p. 544. f. 1. (v. v.)
- $\beta$ . *minor* (DC. fl. fr. 5. p. 617.) caule brevissimo, foliis reniformi-cordatis glabresculis. In Jurasso, Alpibus, Pyrenæis. (v. v.)
8. *V. ARENARIA* (DC. fl. fr. 4. p. 806.) pruinoso-pubescentis, stigmatē papilloso subreflexo, caulibus simplicibus erectiusculis, foliis cordato-subrotundis, stipulis erectis inciso-dentatis, sepalis lanceolato-oblongis, capsulis turbinato-ellipticis obtusis pubescentibus, semin. ovatis castaneis.  $\mathcal{U}$  in montibus Galloprov. superioris. — Pio diss. t. 1. f. 2. (v. v.)
9. *V. PUMILA* (Vill. dauph. 2. p. 266. cat. strasb. t. 5.) glabrescula, stigmatē papilloso subreflexo, caulibus ramosis diffusis procumbentibus, foliis ovato-lanceolatis, petiolis marginatis, stipulis ovato-lanceolatis dentatis incisive, calcare cylindrico apice subrecurvo obtuso sepala vix æquante, valvis subrectis glabris apice truncatis, sem. ovatis nigro-castaneis.  $\mathcal{U}$  in Delphinatū monte Corps, in arenosis Vasconiz, turfosis Jurassi. — Fl. fr. 5. p. 618.
- $\alpha$ . *lancifolia* (Ging.) foliis lanceolatis basi in petiolum gradatim attenuatis. — In arenosis Vasconiz.  $\mathcal{V}$ . *lancifolia* Thore. DC. fl. fr. p. 807. (v. v.)
- $\beta$ . *ericetorum* (Ging.) foliis basi truncatis vel subcordatis in petiolum abrupte attenuatis. — In ericetis turfosis Jurassi, Delph., Vasconiz. (v. v.)
- $\gamma$ . *littoralis* (Ging.) caulibus elongatis, floribus spetalis. — Circā Burdigalam reperiit cl. Desmoulins. (v. s.)
10. *V. MONTANA* (L. sp. 1325.) glabrescula, stigmatē papilloso subreflexo, caulibus simplicibus erectis, foliis imis cordatis, superioribus ovatis acutis, petiolis marginatis, stipulis oblongis dentatis incisive, calcare conico truncato recto sepalis breviori, valvis arcuatis subacutis.  $\mathcal{U}$  in pratis humidis Galliz borealis et mediz. — Fl. fr. 4. p. 807. — Camer. epit. 911. ic. Flores albi demum cœrulescentes. (v. v. DC.)
11. *V. ANDROSCEUS* (L. sp. 1325.) stigmatē uncinato nudo, caulibus fruticulosis ramosis erectis teretiusculis pubescentibus, foliis lanceolatis basi attenuatis, stipulis oblongis adherentibus, calcare cucullato obtuso auriculas vix superante, valvis naviculatis obtusis oligospermis.  $\mathcal{J}$  in saxosis et sabuletis maritimis circā Telonem, Narbonam et in Corsicā. — Fl. fr. 5. p. 618. — Barr. ic. t. 568. (v. s.)

#### Sect. II. DISCHIDIUM. Ging. l. c. p. 300.

Stigma erostratum, apice plus minùs 2-lobum, foraminulo inter lobos sito; stylus ab apice ad basin attenuatus; capsula sæpè 3-gona oligosperma.

12. *V. BIFLORA* (L. sp. 1326.) stigmatē bifido, lobis crassis divergentibus, foliis reniformibus, stipulis ovatis, sepalis linearibus, petalis glabris, calcare brevissimo.  $\mathcal{U}$  in humidis montanis. — Fl. fr. 4. p. 808. — J. Bauh. hist. 3. p. 545. f. 1. — Pluck. t. 233. f. 7. (v. v.)

#### Sect. III. MELANIUM. DC. prod. 1. p. 301.

Stigma urceolatum utrinquē fasciculis pilorum onustum, aperturā magnā infernè labello munitā; stylus ab apice ad basin attenuatus; capsula obsoletè hexagona.

13. *V. NUMMULARIFOLIA* (All. ped. t. 9. f. 4.) glaberrima, caulibus ramosis decumbentibus, foliis subrotundis integris, stipulis lanceolatis setoso-dentatis, sepalis oblongis, calcare breviusculo.  $\mathcal{U}$  in saxosis Alpium Delph., Corsicæ. — Fl. fr. 4. p. 804. — Bocc. mus. 2. p. 163. t. 127. (v. s.)
14. *V. CORNUTA* (L. sp. 1325.) radice fibrosā diffusā, caulibus ascendentibus, foliis cordato-ovatis crenatis ciliatis, stipulis oblique cordatis inciso-dentatis ciliatis, sepalis subulatis, calcare subulato calyce longiore.  $\mathcal{U}$  In Pyrenæis. — Fl. fr. 4. p. 810. bot. mag. t. 791. (v. v.)

15. *V. Cenisia* (All. ped. n. 1641.) radice trunco subfusiformi lignoso, caulibus diffusis procumbentibus indivisis, foliis apathulato-ovatis oblongiave integris, stipulis integris vel palmatisectis segmentis obovatis folio simillimis integris, calycibus pubescentibus, calcare tubuloso acuto calycem æquante.  $\frac{1}{2}$  in Alpibus. — Ging. in DC. prod. 1. p. 301.
- a. *ovatifolia* (Ging.) foliis omnibus ovatis vix pubescentibus. — In saxosis Alpium Galloprovinciæ. *V. Cenisia* All. ped. t. 22. f. 6. Fl. fr. 4. p. 805. (v. s.)
- ß. *diversifolia* (Ging.) foliis imis ovatis, superioribus oblongis tomentoso-pubescentibus. — In Pyrenæis. *V. Cenisia* Lapeyr. abr. (v. v.)
16. *V. CALCARATA* (L. sp. 1325.) radice fibrosâ diffusâ, caulibus simpliciusculis cespitosis, foliis spathulato-subrotundis elongatiave crenatis, stipulis palmatifidis integriave, sepalis oblongis glanduloso-denticulatis, calcare subulato calyce longiore, nectario staminibus vix duplò longiore.  $\frac{1}{2}$  in editioribus montibus. — Fl. fr. 5. p. 810. — Barr. ic. 692. (v. v.)
- ß. *Bertolonii* (Ging.) caulibus elongatis, foliis imis ovatis, extimis elongatis stipulisque palmatifidis glabriusculis. — In montibus Corsicæ. — V. Bertoloni. Pio. diss. t. 3. f. 2. (v. s.)
- γ. *decipiens* (Ging.) caulibus elongatis, foliis stipulisque hispidis, sepalis lanceolatis hispidulis. — In Pyrenæis orient. *V. hispida* Lapeyr. abr. 123. excl. syn. (v. s.)
17. *V. SUBETICA* (W. enum. supp. 12.) radice fibrosâ diffusâ, caulibus simpliciusculis, foliis ovato-oblongis, stipulis palmatifidis lobis integris, sepalis lanceolatis, petalis cuneatis longè unguiculatis discretis, calcare subulato acutiusculo porrecto nectario staminibus vix longiore.  $\frac{1}{2}$  in pratis Vogesorum, montium Arvernæ, Galloprov. et agri Ligericini. — V. Grandiflora. Fl. fr. 5. p. 620. — Hall. helv. t. 17 f. 1. (v. v.)
18. *V. ROTHOMAGENSIS* (Desf. cat. 153.) hispida, radice subfusiformi, caulibus ramosis diffusis, foliis ovalibus, imis subcordatis, stipulis pinnatifidis, calcare tubuloso obtuso sepalis brevior.  $\frac{1}{2}$  in rupibus et aggeribus cretaceis à Rothomago ad Mimatem. — Fl. fr. 4. p. 809. Pio diss. t. 2. Vix ac ne vix differt à *V. tricolore*. (Ging.) (v. s. D.)
19. *V. TRICOLOR* (L. sp. 1826.) radice subfusiformi, caulibus ramosis diffusis, foliis imis ovato-cordatis, stipulis runcinato-pinnatifidis, lobo medio crenato, petalis breviter unguiculatis incumbenibus, calcare crasso obtuso non porrecto, nectariis brevibus. ① ② in arvis. — Fl. fr. 4. p. 808. — Cam. ep. 912. ic.
- a. *hortensis* (DC. prod. 1. p. 303.) petalis intensè velutinis calyce multò majoribus. In hortis culta. (v. v.)
- ß. *degener* (DC.) subramosa, floribus 3-coloribus subvelutinis calyce paulò majoribus, stipulis maximis. In oleraceis. (v. v.)
- γ. *alpestris* (DC.) elongata, foliis remotis, floribus sulfureis purpureo-maculatis calyce majoribus. In pratis Alpium. (v. v.)
- δ. *purpurea* (DC.) foliis ovatis, sepalis breviusculis latis valdè appendiculatis, floribus purpureis calyce majoribus. In Armoricæ inferioris campis. (v. s.)
- ε. *arvensis* (DC.) caulibus ramosis assurgentibus, petalis calyce vix longioribus flavescentibus cæruleis vel purpureo-maculatis. In agris et arenosis frequens. — V. arvensis Fl. fr. 4. p. 808. — Cam. ep. 918. ic. (v. v.)
- ζ. *sabulosa* (DC.) caulibus plurimis diffusis, foliis remotis ovatis elongatiave angusto-lanceolatis corollâ vix brevioribus. — In arenosis maritimis. (v. s.)
- η. *bellioides* (DC.) hispidula, caulibus brevissimis, foliis subrotundis confertis, petalis calyce brevioribus. In arenosis Corsicæ et circâ Mons-pelium. (v. s.)
- θ. *nana* (DC.) glabriuscula, caule brevissimo, cotyledonibus per anthesim persistentibus. In arenosis maritimis Neustriæ. (v. s.)

Ordo X. RESEDACEÆ. DC. *theor.* 214. *ess. prop. ed.* 2. p. 110. — Capparidum. *gen. Juss.*

Sepala 4-6 cum pedicellis continua persistentia. Petala 4-6 sepalis alterna, hypogyna, inæqualia, superiora ungue squamiformi limbo palmati-partito, lateralia 2-3-loba, infima minora integra. Stam. 10-24 hypogyna, filamentis sæpius basi plus minusve inter se coalitis mono-aut polyadelphis, antheris 2-locularibus erectis. Squama nectarifera obtusissima toro sub staminibus inserta. Torus nunc brevissimus nunc stipitiformis. Ovaria 3-6 puncto libera summo toro inserta monostyla, nunc valvatis coalita in ovarium unicum stylis brevibus 3-6 conicis cum suturis ipsis continuis coronatum. Carpella ideò nunc libera folliculiformia intus dehiscentia oligosperma, nunc coalita in capsulam suturis indehiscentibus 3-6-valvem apice hiantem intus unilocularem polyspermam stylis 3-6 coronatam. Placentæ 3-6 in medio carpellorum aut valvularum adnatæ sæpius polyspermæ, interdum (in carpellis liberis) 1-2-spermæ. Semina secus placentas biserialia subpendula testâ crustaceâ instructa. Albumen nullum aut tenue carnosomembranaceum. Embryo arcuatus, cotyledones carnosæ, radícula supera. — Herbæ foliis alternis, floribus breviter pedicellatis. (DC. mss.)

An potius, ut vult cl. Hooker, calyx auctorum pro involucrio, squamæ et petala pro perigonio et appendicibus perigonii habendæ, et inflorescentia ut capitulum ex floribus sterilibus et perfectis compositum describenda. Ordo tunc Euphorbiaceis et Datiscæ affinis.

I. RESEDA. *Neek. elem. n.* 991. DC. mss. — *Reseda*, sp. L.

Petala 4-6 inæqualia, superiora laciniata; stam. 10-24, carpella 3-6 coalita in capsulam 3-6-gonam apice hiantem; 3-6 stylis coronatam, intus 1-locularem; placentæ 3-6 stylis alternæ, in medio valvularum coalitarum adnatæ semina 60 biserialia gerentes. (DC. mss.)

#### Sect. I. RESEDASTRUM.

Sepala 6, stigmata 3, rarissimè 4. — Folia indivisa aut 3-loba.

1. R. PHYTUMA (L. sp. 645) foliis obtusis aut mucronulatis, radicalibus integerrimis spathulatis caulinis semi-trilobis, calycibus maximis, fructibus laxis. ① in arvis sabulosis. — Fl. fr. 4. p. 727. — C. Bauh. prod. 42. f. 1. Antheræ pallide flavæ; flores inodori. (v. v.)
2. R. ODORATA (L. sp. 646.) foliis oblongis obtusis aut mucronulatis undulatis integerrimis trilobisque, calycibus corollam æquantibus, fructibus laxis. ① ex Aegypto et Barbariâ orta, in hortis culta. — Fl. fr. 4. p. 727. Curt. bot. mag. t. 29. Mill. ic. t. 217. antheræ crocæ; flores odori. (v. v. c.)

3. *R. LUTEA* (L. sp. 645.) foliis pinnatifidis undulatis superioribus 3-fidis, fructibus laxis oblongis triangularibus apice truncatis.  $\mathcal{U}$  ad vias et muros, in sabulosis. — Fl. fr. 4. p. 727. eng. bot. t. 327. — Lob. ic. 222. f. 1. (v. v.)

Sect. II. LEUCORESEDA. DC. l. c.

Sepala 5, stigmata 4, rarisimè 3. — Folia pinnatipartita, flores albi.

4. *R. ALBA* (L. sp. 645.) foliis pinnatipartitis, lobis oblongis integris sæpè undatis, floribus 4-gynis, fructibus confertis, capsulis stylis persistentibus apice conniventibus coronatis.  $\textcircled{1}$   $\mathcal{U}$  in sabulosis maritimis Gallis mediterraneis passim. — Fl. fr. 4. p. 726. *R. undata* L. sp. 644. Fl. fr. 4. p. 726. — Barr. rar. t. 587. — J. Bauh. hist. 3. p. 2. p. 467. f. 1. (v. v.)

Sect. III. LUTEOLA. DC. *ms.*

Sepala 4, stigmata 3. — Folia indivisa.

5. *R. LUTEOLA* (L. sp. 643.) caule erecto, foliis lanceolatis undulatis integris, fructibus confertis, capsulis supernè hiantibus eornibus 3 foliaceis erectis 3 alternis carnosis introflexis coronatis.  $\textcircled{2}$  ad vias. — Fl. fr. 4. p. 725. eng. bot. t. 320. — Lob. ic. 353. f. 1. (v. v.)

Sect. IV. GLAUCORESEDA. DC. l. c.

Sepala 5, stigmata 2, capsula subglobosa. — Folia indivisa.

6. *R. GLAUCA* (L. sp. 644.) caulibus diffusis, foliis linearibus acutis angustissimis ad basim utrinquè dente acuto exserto stipulaceo auctis, capsulis subglobosis brevissimè cornatis.  $\mathcal{U}$  in saxosis apricis vallium Pyrenæorum præsertim centralium, et montis Lupi propè Monapellum ex ol. Benth. — Fl. fr. 4. fr. 725. — Pluck. alm. t. 107. f. 2. (v. v. DC.)

II. ASTROCARPUS. Neck. elem. n. 99a. DC. *ms.*

Petala 4-6 inæqualia, superiora laciniata; stam. 12-15; carpella 4-6 toro stipitiformi insidentia stellatim patentia follicularia intus rimâ dehiscentia; semina 1-2 in medio carpellorum propè basim adnata. (DC. *ms.*)

1. *A. SESAMOIDES* (DC. l. c.) caulibus diffusis, foliis caulinis linearibus integerrimis, seminibus lenticularibus in medio capsula insertis.  $\mathcal{U}$  *Reseda sesamoides* DC. fl. fr. 4. p. 725. ic. rar. t. 40.

*a. stellata*, foliis radicalibus obovato-oblongis. — In rupestribus Cebenn., Pyren., montium Arvernæ. — *R. sesamoides* L. sp. 644. Lam. ill. t. 400. f. 2. (v. v.)

*β. purpurascens*, foliis radicalibus oblongo-linearibus. — In sabulosis planities Gallis austro-occident. — *R. purpurascens* L. — Lob. ic. 353. f. 2. (v. v.)

Ordo XI. DROSERACEÆ. *DC. theor.* 214. *prod.* 1. p. 317.

Calyx sepalis 5 persistentibus per æstivationem imbricatis. Petala 5 distincta hypogyna sepalis alterna. Stamina nunc petalis alterna et iis numero æqualia, nunc petalis dupla tripla quadrupla, libera; antheræ 2-loculares. Ovarium 1; styli 3-5 nunc basi coaliti nunc distincti, bifidi ramosive. Capsula 1-3, locularis, 3-5-valvis, valvulis margine plus minus introflexis, ab apice dehiscentibus, intus nunc in nervo medio, nunc ad basin seminiferis. Semina nuda aut arillo tenui folliculari involuta. Albumen cartilagineum carnosumve: Embryo rectus, radicula ad hylum versâ. — Herbæ foliis alternis, petiolis basi ciliis stipularibus sæpius instructis.

I. DROSERA. *L. gen.* 391. *Lam. ill.* t. 220.

Sepala petalaeque 5 inappendiculata; stam. 5; styli 3-5, bipartiti.

1. *D. rotundifolia* (L. sp. 402.) foliis orbiculatis, petiolis pilosis limbo longioribus, scapis rectis folia triplò superantibus, seminibus arillatis. ① in humidis turfosis. — Fl. fr. 4. p. 729. Drev. et Hayn. eur. 3. t. 74. — Lob. ic. 802. f. 3. (v. v.)
2. *D. intermedia* (Drev. et Hayn. pl. eur. 3. p. 43. t. 75. B.) foliis obovatis, petiolis glabris limbo longioribus, scapis ascendentibus folia paulò superantibus, seminibus exarillatis. ①? in turfosis et humidis. — DC. prod. 1. p. 318. *D. longifolia*. L. sp. 403. Fl. fr. 4. p. 729. eng. bot. t. 868. (v. v.)
3. *D. anglica* (Huds. ang. 135.) foliis oblongis obtusis basi attenuatis, petiolis glabris limbo vix longioribus, scapis erectis folia duplò superantibus, seminibus arillatis. ½ in paludosis Galliz borealis. — Fl. fr. 4. p. 729. *D. longifolia* Drev. et Hayn. 3. t. 75. A. — Moris. s. 15. t. 4. n. 1. (v. v.)

II. ALDROVANDA. *Monti. act. bon.* 2. p. 3. p. 404. t. 12. *L. gen.* 390.

Sepala petalaeque 5 inappendiculata; stam. 5; styli 5 filiformes breves; stigmata obtusa; caps. globosa 5-valvis 1-locularis 10-sperma.

1. *A. vesiculosa* (L. sp. 402.) ½ All. ① Sav. — In aquis circa Arelatem et Burdigalam (Dunal). Folia verticillata, petiolo in cilia 6 setacea desinente, limbo complicato ferè vesiculari. — Fl. fr. 4. p. 730. (v. v. DC.)

III? PARNASSIA. *Tourn. inst.* t. 127. *L. gen.* 384.

Sepala 5; petala 5; squamæ 5 (an stamina abortiva?) unguibus petalorum oppositæ, in setas apice glandulosas desinentes; stam. 5; anth. posticæ; stigmata 4 sessilia; capsula 4-valvis 1-locularis valvis medio septiferis; semina arillata.

1. *P. palustris* (L. sp. 391.) appendicibus 9-13-setis, petalis subsessilibus, foliis radicalibus cordatis, caulino amplexicauli. ½ in paludibus submontanis. — Fl. fr. 4. p. 728. fl. dan. t. 784. (v. v.)

Ordo XII. POLYGALÆÆ. *Juss. ann. mus.* 14. p. 386. *DC.*  
*prod.* 1. p. 321.

Calyx sepalis 5 per æstivationem imbricatis, 2 interioribus sæpius petaliformibus, 3 externis minoribus, quorum 2 anteriora interdum coalita et tertium posticum. Petala 3-4 hypogyna, mediante tubo stamineo plus minusve connexa, rarius distincta. Stamina filamenta cum petalis coalita, monadelphia, apice in phalanges 2 oppositas æquales divisa. Antheræ 8, 1-loculares, basi insertæ, apice poris dehiscentes. Ovarium 1 liberum 2-loculare, rarius 1-3-loculare. Stylus 1. Stigma 1. Pericarpium capsulare aut drupaceum 2 aut 1-loculare, valvis medio septigeris. Semina in loculis solitaria pendula; sæpè basi carunculato-arillata. Embryo rectus nunc in albumine carnoso axilis, nunc rarius exalbuminosus endoplevra tunc tumidiore. — Herbæ aut suffrutices; foliis in plerisque alternis super caulem articulatis integris.

I. POLYGALA. *Tourn. inst.* 4. 79. *DC. l. c.*

Calycis sepala persistentia, 2 interiora alæformia; petala 3-5 tubo stamineo connexa, inferiore carinæformi; capsula compressa elliptica obovata aut obcordata; semina pubescentia, hylo carunculata, comâ destituta.

Sect. 1. POLYGALON. *DC. l. c.*

Carina cristata; capsula glabra; bractæ 3 mox deciduæ ad basin pedicellorum.

1. *P. SAXATILIS* (Desf. atl. t. 175.) foliis oblongo-lanceolatis mucronatis, caule basi fruticoso decumbente, racemis extra-axillaribus paucifloris, alis ellipticis. ♂ in rupestribus apricis circa Narbonam, Massiliam. — *Fl. fr.* 3. p. 457. (v. s.)
2. *P. VULGARIS* (L. sp. 986.) foliis lineari-lanceolatis obtusiusculis, caulibus ascendentibus, alis ovatis capsulâ parum longioribus corollâ subbrevioribus, ovario subsessili. ♀ in pratis et sylvaticis. — *Fl. fr.* 3. p. 456.
  - a. *vera* (DC.) caulibus erectiusculis, foliis inferioribus obovatis obtusis, superioribus linearibus acutis. — *Fl. dan.* t. 516. (v. v.)
  - β. *elata* (DC.) caulibus erectis, foliis ex ovali-oblongis. *P. comosa* Schkuhr. handb. t. 194. (v. s.)
  - γ. *angustifolia* (DC.) caulibus erectis, foliis linearibus. (v. v.)
3. *P. AMARA* (L. sp. 987.) foliis radicalibus obovatis obtusissimis, caulinis linearibus, caulibus erectiusculis, alis ellipticis corollam æquantibus, capsulâ suborbiculatâ. ♀ in pratis montanis. — *Fl. fr.* 3. p. 456.
  - a. *Austriaca* (Crantz aust. t. 2. f. 4.) amarissima, floribus minoribus albicantibus, alis post anthesin viridescentibus. (v. v.)
  - β. *alpestris* (DC.) insipida, alis fructus coloratis, radice magis cespitosa. Jacq. aust. t. 412. (v. v.)
  - γ. *cespitosa* (DC.) floribus cæruleis, caulibus cespitoso-procumbentibus, foliis linearibus summis ad flores approximatis. (v. v.)
  - δ. *Alpina* (Poir. dict. 5. p. 488.) floribus cæruleis paucis, caulibus cespitoso-procumbentibus brevissimis, foliis inferioribus obovatis. (v. v.)

4. *P. MONSPELIACA* (Lin. sp. 987.) glabra erecta subsimplex, foliis linearibus acuminatis, alis oblongis utrinque acutis trinerviis capsulâ multò longioribus. ① in sterilibus prov. mediterranearum. — Fl. fr. 3. p. 457. DC. ic. rar. 1. p. 3. t. 9. (v. v.)
5. *P. EXILIS* (DC. cat. Monsp. 133.) erecta glabra ramosissima, foliis linearibus crassiusculis, alis ovalibus obtusis 1-nerviis capsulæ longitudine, racemo laxiusculo. ① in arenosis circa Monspelium, Avenionem, in Gallo-provincia et supra Lugdunum propè Château-Gaillard. — Fl. fr. 5. p. 386. Boiss. fl. eur. 1. t. 474. f. 1. (v. s.)

Sect. II. *CHAMÆBUXUS*. *Dill. gen. t. 9.*

Calycis sepalum inferius concavo-uncullatum intus ad basin glandulam fovens; corollæ carina cristata.

6. *P. CHAMÆBUXUS* (L. sp. 989.) caulibus suffruticulosis ramosis procumbentibus, foliis oblongo-lanceolatis mucronatis, racemis 1-2-floris, carinâ cristatâ. ② 5 in Alpibus, montibus Alsatiae, etc., Pyrenæis orientalibus ex Lapeyr. — Fl. fr. 3. p. 458. Jacq. Austr. t. 233. (v. v.)

Ordo XIII. *FRANKENIACEÆ*. *St.-Hil. mem. plac. cent. p. 39.*

*DC. prod. 1. p. 349.*

Cal. sepala 4-5 erecta basi coalita in tubum sulcatum, persistentia æqualia. Petala tot quot sepala, iis alterna, hypogyna, unguiculata, fauce squamulis subcoronata. Stamina hypogyna, nunc petalis æqualia et iis alterna; nunc 1-2 addita petalis opposita. Filamenta filiformia. Antheræ subrotundæ. Ovarium 1, liberum. Stylus filiformis 2-3-fidus. Capsula calyce persistente stipata, bi aut sæpius tri-imò quadri-valvis, 1-locularis; valvulis ad marginem utrumque placentiferis, polyspermis. Embryo rectus in medio albuminis, radiculâ brevi ad umbilicum spectante. — Herbæ aut suffrutices; foliis oppositis aut verticillatis, exstipulatis, basi productis in membranulam amplexicaulem.

I. *FRANKENIA*. *Lin. gen. n. 445.*

Stylus 3-fidus, lobis oblongis intus stigmatosis; capsula 3-4-valvis polysperma.

1. *F. PULVERULENTA* (Lin. sp. 474.) foliis obovatis retusis glabris subtus pulverulentis in petiolo ciliatis, radice exili. ① in arenosis maritimis à Nicæâ ad Hispaniam. — Fl. fr. 4. p. 766. — Clus. hist. 2. p. 186. f. 2. (v. v.)
2. *F. LEVIS* (Lin. sp. 473.) foliis linearibus margine revolutis glabris basi ciliatis, caulibus prostratis calycibusque glabris. ② ④ in maritimis Oceani à Nannete ad Bayonam et Mediterranei secus Occitaniam. — Fl. fr. 4. p. 765. — Mich. gen. t. 22. f. 1. (v. v.)
3. *F. INTERMEDIA* (DC. prod. 1. p. 349.) foliis linearibus margine revolutis glabris basi ciliatis, caulibus prostratis velutinis, calycibus hispidis. ③ ④ in maritimis Mediterræis. — *F. hirsuta*. Fl. fr. 4. p. 766. — Mich. gen. t. 22. f. 2. (v. v.)



Ordo XIV. CARIOPHYLLÆ. *Juss. gen. 299. Ser. in DC. prod. 1. p. 351.*

Calyx sæpius persistens sepalis 4 aut sæpius 5 cum pedicello continuis nunc liberis, nunc inter se in tubum 4-5-dentatum coalitis, æstivatione imbricatis. Petala totidem hypogyna toro plus minùsve elevato (Anthophoro dicto) inserta, sepalis alterna, unguiculata, ad faucem interdum squamis petaloideis aucta, rarissimè abortu nulla. Stamina petalorum numero dupla toro inserta; petalis alia alterna, alia opposita; filamenta subulata interdum basi submonadelpha; antheræ 2-loculares. Ovarium tori apice insertum simplex 2-5-valve, stylis totidem superatum. Capsula valvis 2-5 basi coalitis apice dehiscentibus, nunc sæpius 1-locularis, nunc 2-5-locularis, valvis proferentibus è medio septa, nunc incompleta, nunc ad axim continua. Placenta centralis. Semina  $\infty$  (rarissimè pauca et definita). Albumen farinosum sæpius centrale. Embryo nunc periphericus plus minùsve incurvus, rarissimè rectus centralis, radiculâ ad hilum versa. — Herbæ aut suffrutices, foliis oppositis integris.

Trib. I. *SILENÆ*: *DC. prod. 1. p. 351. — Cariophyllæ. Lam.*

Sepala coalita in tubum cylindraceum apice 4-5-dentatum.

I. *GYPSOPHILA*. *Linn. gen. n. 768.*

Calyx campanulatus sub 5-lobus, laciniis margine membranaceis; petala 5 non unguiculata; stamina 10; styli 2; capsula 1-locularis.

Sect. I. *STRUTHIUM*. *Ser. in DC. prod. l. c.*

Calyces squamis destituti.

1. *G. FASTIGIATA* (Lin. amæn. 3. p. 23.) floribus corymboso-fastigiatis, caule adscendente, foliis lanceolato-linearibus absolute triquetris lævibus obtusis secundis, staminibus exsertis.  $\frac{1}{2}$  propè Monspelim ex Gouan. — Fl. fr. 4. p. 735. — Gm. sib. 4. t. 61. f. 1. (v. v. c.)
2. *G. REPENS* (Lin. amæn. 3. p. 23. spec. 581.) caulibus subpaniculatis paucis floris, staminibus petalis emarginatis brevioribus, foliis linearibus glabris.  $\frac{1}{2}$  in petrosis ad vias et secus torrentes Alpium, Pyren. et montium Arvernæ. — Fl. fr. 4. p. 736. Ger. Galloprov. t. 15. f. 2. (v. v.)
3. *G. MURALIS* (Lin. amæn. 3. p. 24. spec. 583.) caule dichotomo paniculato ramosissimo, floribus axillaribus solitariis, petalis crenatis, foliis linearibus planis longitudine pedicellorum. ① ad vias et in agris passim. — Fl. fr. 4. p. 336. — J. Bauh. 3. p. 2. p. 338. f. 1. (v. v.)

Sect. II. *PETTORHAGIA*. *Ser. l. c.*

Calyces squamis scariosis 2-4 oppositis basi instructi.

4. *G. SAXIFRAGA* (Lin. spec. 584.) caulibus numerosis erectis rigidis, floribus

paniculatis terminalibus, calycibus squamis quaternis involucretis, foliis linearibus rigidis.  $\mathcal{U}$  in saxosis Galliæ australis, mediæ et occidentalis. — Fl. fr. 4. p. 737. — Barr. ic. 998. (v. v.)

## II. DIANTHUS. Lin. gen. n. 770.

Calyx tubulosus 5-dentatus basi squamis 2-4 oppositis imbricatis; petala 5 longè unguiculata; stamina 10; styli 2; capsula 1-locularis; semina compressa hinc convexa indè concava, peltata; embryo vix curvatus.

### Sect. I. ARMERIASTRUM. Ser. in DC. prod. 1. p. 355.

Flores capitati vel corymbosi, sessiles vel pedunculati.

#### §. 1. *Bracteis ovatis muticis.*

1. D. PROLIFER (Lin. spec. 587.) floribus aggregatis capitatis, squamis calycinis ovatis muticis tubum superantibus, foliis serrulatis. ① ad marginem sylvarum et agrorum. — Fl. fr. 4. p. 741. — J. Bauh. 3. p. 335. f. 1. (v. v.)

#### §. 2. *Bracteis lanceolatis acutis, calycibus striatis villosis.*

2. D. ARMERIA (Lin. sp. 586.) floribus laxè fasciculatis, squamis calycinis lanceolato-subulatis tubum æquantibus, foliis subulatis calycibusque hirsutis. ① in sterilibus. — Fl. fr. 4. p. 741. OEd. fl. dan. t. 230. (v. v.)

#### §. 3. *Bracteis ovatis vel lanceolatis, calycibus vix striatis glabris.*

3. D. BARBATUS (Lin. spec. 586.) floribus aggregatis fasciculatis, squamis calycinis ovato-subulatis tubum æquantibus, foliis lanceolatis.  $\mathcal{U}$  in sterilibus Galliæ australis, in hortis cultus. — Fl. fr. 4. p. 739. (v. v.)
  4. D. BALBISII (Ser. in DC. prod. 1. p. 356.) glaucescens, floribus subsessilibus capitato-fasciculatis, squamis calycinis lanceolatis patulis tubo paulò minoribus, foliis lanceolato-linearibus, caule angulato.  $\mathcal{U}$  circà Niceam. — Folia lanceolato-linearia basi non dilatata. (v. v.)
  5. D. FERRUGINEUS (Lin. mant. 563.) floribus aggregatis, involucreis squamisque scariosis rufis oblongis aristatis calyce paulò brevioribus, foliis linearibus longè connatis.  $\mathcal{U}$  in Pyrenæis Narbonensibus. — Fl. fr. 4. p. 740. — Barr. ic. t. 497. An sequentis var? (v. s.)
  6. D. CARTHUSIANORUM (Lin. spec. 586.) floribus aggregatis sessilibus pedunculatisque, squamis calycinis ovatis aristatis tubo brevioribus, involucreo oblongo aristato capitulo brevior, foliis linearibus trinerviis.  $\mathcal{U}$  in incultis et sterilibus. — Fl. fr. 4. p. 740. (v. v.)
- $\beta$ . *atrorubens* (Ser.) floribus aggregato-capitatis sessilibus 3-8. — In collibus agri Ruscinonensis, Ligericini. — D. atrorubens. Jacq. ic. rar. 467. Fl. fr. 4. p. 740. (v. s.)

### Sect. II. CARYOPHYLLUM. Ser. in DC. prod. 1. c.

Flores paniculati vel solitarii.

#### §. 1. *Petalis dentatis.*

7. D. ASPER (Willd. enum. 466.) floribus fasciculatis, squamis calycinis ovato-lanceolatis acuminatis tubo brevioribus, petalis acute dentatis, foliis linearilanceolatis 3-5-nerviis cauleque scabris apice serrulatis.  $\mathcal{U}$  in provinciis australioribus.

$\alpha$ . *angustifolius* (Ser.) pauciflorus, foliis linearibus acutis. D. serratus  $\beta$  DC. Fl. fr. 5. p. 601. (v. s.)

$\beta$ . *collinus* (Ser.) multiflorus, foliis linearilanceolatis. — In montibus Arvernæ et propè Mimatem. — D. Collinus W. et Kit. hung. 1. t. 38. (v. s.)

$\gamma$ . *serratus* (Ser.) caulibus 1-3-floris, floribus majoribus, foliis linearibus. — In Pyr. orient. D. serratus Lapeyr. et DC. fl. fr. 5. p. 601.  $\alpha$ . (v. s.)

8. *D. ALPESTRIS* (Balb. act. acad. taur. 7. p. 11. t. 1.) floribus subbinis, squamis calycinis 4, tubo brevioribus, 2 interioribus latis mucronatis, 2 exterioribus lanceolatis acuminatis, petalis emarginatis.  $\frac{1}{2}$  circa Monspe-  
linum et in Alpibus Galloprovinciæ inferioris? (v. s.)
9. *D. ATTENUATUS* (Smith. act. soc. Linn. 2. p. 301.) floribus solitariis, squamis calycinis subsevis brevibus lanceolatis acuminatis margine membranaceis, calyce longo apice attenuato, petalis crenatis unguibus calycem superanti-  
bus, foliis subulatis.  $\frac{1}{2}$  in maritimis agri Ruscinonensis, Galloprovinciæ et  
propè Mimatem. (cl. Prost.) — Fl. fr. 4. p. 742. *D. geminifloras*. Fl. fr. 5.  
p. 602. (v. v.)
10. *D. NIRTUS* (Vill. delph. 3. p. 593. t. 46.) floribus subsolitariis, squamis calycinis senis ovatis mucronatis calyce multò brevioribus, petalis crenatis,  
foliis subulatis margine scabris.  $\frac{1}{2}$  in collibus Delph. Galloprov., agri Lige-  
ricini (cl. Prost. et Pyren. orient. ex cl. Benth. — Ser. in DC. prod. 1. p. 358.  
(v. s.)
11. *D. CARYOPHYLLUS* (Lin. spec. 587.) caule ramoso, floribus solitariis, squa-  
mis calycinis subquaternis brevissimis ovatis submucronatis, petalis latis-  
simis imberbibus, foliis lineari-subulatis canaliculatis glaucis.  $\frac{1}{2}$  in Gallia  
australi et occidentali. — Fl. fr. 4. p. 741. — Lam. ill. t. 376. f. 1. (v. v.)
12. *D. SYLVESTRIS* (Jacq. coll. 1. p. 237. ic. rar. t. 82.) caule ramoso vel sim-  
plici, floribus solitariis, squamis calycinis brevissimis quaternis ovatis, exte-  
rioribus acutis, interioribus obtusissimis, petalis latissimis imberbibus den-  
tatis.  $\frac{1}{2}$  in saxosis montanis Galliz orientalis et montis Canigou ex Benth.  
— Fl. fr. 4. p. 742. Jacq. ic. rar. t. 82. (v. v.)
13. *D. PUGNONS* (Lin. mant. 240.) caule paucifloro, floribus solitariis, squamis calycinis brevissimis mucronatis patentibus, tubo gibbo, petalis integris,  
foliis cæspitosis subulatis.  $\frac{1}{2}$  in monte dicto la Clape propè Narbonam ex cl.  
Benth. (v. s. ex Hisp.)
14. *D. VIRGÆNEUS* (Lin. spec. 590.) caule subunifloro, squamis calycinis bre-  
vissimis obtusis mucronatis binis distantibus, petalis crenatis, foliis cæpi-  
tosis linearibus rigidis serrulatis.  $\frac{1}{2}$  in collibus agri Monspel., Narbonensis,  
Ruscinonensis. — Fl. fr. 4. p. 744. (v. v. DC.)
15. *D. subcaulis* (Ser.) caulibus cæspitosis brevissimis 1-floris, foliis floribus  
squamisque calycinis abbreviatis. — In rupibus montis Ventosi. — D.  
caulis. Lois. not. t. 6. f. 1. Fl. fr. 5. p. 603. (v. s.)
15. *D. DENTATUS* (Lin. sp. 588.) caulibus decumbentibus ramosis, floribus so-  
litariis, squamis calycinis ovato-lanceolatis acutis subbinis, foliis superiori-  
bus angustis acutis, inferioribus oblongis obtusis.  $\frac{1}{2}$  in arenosis sylvaticis  
Alsatiæ, Andegaviæ, Pyren., Cebenn. — Fl. fr. 4. p. 744. — J. Bauh. 3. p. 329.  
f. 4. (v. v.)
16. *D. CÆSIUS* (Smith act. soc. Linn. 2. p. 302. engl. bot. t. 62.) caulibus cæpi-  
tosis subunifloris, squamis calycinis acutis brevibus exterioribus lanceolatis  
interioribus subrotundis, petalis crenatis pubescentibus, foliis obtusiusculis  
margine scabris.  $\frac{1}{2}$  in saxosis montanis Jurassi, Arverniz. — Fl. fr. 4.  
p. 745. — Dill. elth. t. 298. f. 385. (v. v.)
17. *D. GLACIALIS* (Jacq. coll. 2. p. 84.) caulibus erectis cæspitosis bre-  
vibus subunifloris, squamis calycinis elongatis tubum æquantibus vel su-  
perantibus, calyce striato, petalis serratis, foliis linearibus acutis serrulatis.  
 $\frac{1}{2}$  in montanis Galloprov. et Delph. — Fl. fr. 5. p. 603. *D. Alpinus*. Fl. fr. 4.  
p. 746. non L. (v. s.)

### §. 2. *Petalis fimbriatis.*

18. *D. GALLICUS* (Pers. ench. 1. p. 495.) caulibus adscendentibus subunifloris,  
squamis calycinis brevibus ovatis submucronatis, petalis dentato-multifidis,  
foliis linearibus subciliatis.  $\frac{1}{2}$  in arenosis maritimis à Bayona ad Nannetes. —  
Fl. fr. 5. p. 602. *D. arenarius*. DC. syn. p. 588. ic. rar. t. 41. non L. (v. v.)
19. *D. PLUMARIUS* (Linn. spec. 589. hort. upsal. 105. n. 4.) glaucescens,

- caulibus 2-3 floris, dentibus obtusis, bracteis subovatis brevissimis mucronatis, petalis laciniato-multifidis barbatis, foliis linearibus margine scabris.  $\mathcal{L}$  in hortis cultus. — Fl. fr. 4. p. 745. (v. v. c.)
20. *D. SAXATILIS* (Pers. ench. 1. p. 494.) cespitosus subdecumbens, caule 2-3-floro, squamis ovatis mucrone longiusculo distinctis tubo brevioribus, petalis multifidis fauce glabris.  $\mathcal{L}$ ? ad rupes propè Clermont. D. Plumarius quorundam sed folia linearia non glauca, corolla fauce non pubescens, nec squamæ obtusæ. (Pers. l. c.) †.
21. *D. MONSPRESSULANUS* (Lin. amæn. 4. p. 313. spec. 588.) caule paniculato paucifloro, floribus solitariis, squamis calycinis subulatis rectis tubo dimidiò brevioribus, petalis digitato-multifidis fauce subglabris, foliis linearibus serrulatis.  $\mathcal{L}$  in Jureaso, Cebennis, Pyrenæis. — Fl. fr. 4. p. 745. — Clas. hist. 1. p. 284. f. 1. (v. v.)
22. *D. SUPRÆUS* (Lin. amæn. 4. p. 272. spec. p. 589.) caule paniculato multifloro, floribus subfastigiatis, squamis calycinis brevibus ovatis mucronatis, petalis ultrà medium pinnato-multifidis fauce pilosis.  $\mathcal{L}$  in nemoribus et pratis umbrosis montanis. — Fl. fr. 4. p. 744. — Clas. hist. 1. p. 284. f. 2. (v. v.)

### III. SAPONARIA. *Lin. gen. n.* 769.

Calyx tubulosus 5-dentatus basi nudus; petala unguiculata, ungue calycem æquante; stam. 10; styli 2; caps. 1-locularis.

#### Sect. I. VACCARIA. *Dod. pempt.* 104.

Flores paniculati, calyx angulosus glaber.

1. *S. VACCARIA* (Lin. spec. 585.) floribus paniculatis, calycibus pyramidalis quinquangularibus glabris, bracteis membranaceis acutis, foliis ovato-lanceolatis sessilibus.  $\odot$  inter segetes passim. — Fl. fr. 4. p. 737. — J. Bauh. 3. p. 2. p. 357. f. 2. (v. v.)

#### Sect. II. BOOTIA. *Neck. delic. gallo-belg.* 1. p. 193.

Flores fasciculato-paniculati, calyx teres sæpius pilosus.

2. *S. OFFICINALIS* (Lin. spec. 584.) floribus fasciculato-paniculatis, calycibus teretibus villosis luteolis, petalorum appendicibus linearibus, foliis ovato-lanceolatis acutis obtusisve.  $\mathcal{L}$  ad margines viarum et agrorum. — Fl. fr. 4. p. 737. Eng. bot. t. 1060. (v. v.)
3. *S. OXYMOIDES* (Lin. spec. 585.) caulibus procumbentibus dichotomis, floribus fasciculatis, calycibus teretibus villosis purpureis glanduloso-pilosis, foliis ovato-lanceolatis subuninerviis.  $\mathcal{L}$  in petrosis provinc. austral. et occidentaliun. — Fl. fr. 4. p. 738. — Lob. ic. t. 341. f. 2. (v. v.)

#### Sect. III. PROTEINIA. *Ser. in DC. prod.* 1. p. 366.

Flores solitarii axillares, calyx pilosus raro glaber.

4. *S. ORIENTALIS* (L. sp. 785.) caule dichotomo, ramis divaricatis, floribus axillaribus, pedunculis rigidis florem æquantibus calycibusq. hispidulis teretibus, laciniis acutis, foliis lineari spathulatis.  $\odot$  circà Collioure et Perpignan in agro Ruscionensi. — Fl. fr. 5. p. 601. — Dill. Elth. 205. t. 167. f. 204. (v. v. c.)

#### Sect. IV. BOLANTHUS. *Ser. in DC. prod.* 1. p. 366.

Caulis folisque cespitosi, flores aggregati, calyx teres villosus.

5. *S. CESPITOSA* (DC. rap. voy. 2. p. 78. 1808. fl. fr. 5. p. 601.) cespitosa, caulibus subnudis apice floriferis, floribus subumbellatis, calycibus cylindricis villosis profundè lobatis, lobis acutis, petalis apice emarginatis appendice bifido, lobis angustissimis, foliis linearibus glabris subradicalibus vix denticulatis.  $\mathcal{L}$  in sterilibus Pyren. editiorum. — S. elegans Lapeyr. Abr. pyr. 238. — Habitus *S. luteæ* sed flores rosei majores. (v. v. DC.)
6. *S. LUTEA* (Lin. spec. 585. excl. syn. Boeck.) cespitosa, caulibus diphyllis, flo-

ribus capitatis involucreatis, calycibus cylindricis lanatis, lobis obtosis brevibus, petalis obovatis integris audis, foliis linearibus basi ciliatis subradicalibus.  $\mathcal{L}$  in Pyrenæis? ex Lapeyr. — Fl. fr. 4. p. 758. All. ped. t. 23. f. 1. (v. s. ex Pedemontio.)

IV. CUCUBALUS. *Gärtn. fruct.* 1. p. 376. t. 77. f. 7.

Calyx campanulatus 5-dentatus nudus; petala 5 unguiculata, fauce sæpius coronata, limbo bifido; capsula carnea 1-locularis.

1. *C. RACCIFERUS* (Lin. spec. 591.) ramis divaricatis, foliis ovatis, calycibus campanulatis, petalis distantibus.  $\mathcal{L}$  in umbrosis. — Fl. fr. 4. p. 760. — Mill. ic. t. 112. (v. v.)

V. SILENE. *Lin. gen. n.* 772. *Gärtn. fruct.* 2. p. 233. t. 130. f. 8. *Outh. in DC. prod.* 1. p. 367.

Calyx tubulosus 5-dentatus nudus; petala 5 unguiculata, fauce sæpius coronata, limbo bifido; stamina 10; styli 3; capsulæ basi 3-loculares apice in 6 dentes dehiscentes.

§. 1. *Cespitosæ, caulibus subnullis.*

1. *S. ACAULIS* (L. sp. 603.) glabra, caulibus densis humilibus, foliis linearilanceolatis, pedunculis brevibus unifloris, calycibus campanulatis.  $\mathcal{L}$  in Alpibus et Pyrenæis. — Fl. fr. 4. p. 749. All. ped. t. 79. f. 1 et 2. (v. v.)

§. 2. *Caulescentes, floribus solitariis paniculatisve, calyce vesiculoso-inflato.*

2. *S. INFLATA* (Smith fl. brit. 467.) caulibus ramosis, floribus paniculatis, calycibus vesiculato-ovatis, petalis bifidis nudis, unguiculis cuneiformibus, stylis longissimis.  $\mathcal{L}$  in Galliâ frequens. — Fl. fr. 4. p. 746.

*a. vulgaris* (Otth.) glabra, erecta, foliis lanceolatis. — In agris. — Engl. bot. t. 164. (v. v.)

*β. uniflora* (Otth.) glabra, procumbens, foliis lanceolato-ovatis, floribus subsolitariis. — In Alpibus. *S. uniflora. β.* Fl. fr. 4. p. 747. — All. spec. t. 5. f. 3. (v. v.)

*γ. maritima* (DC.) piloso-pubescent, procumbens, foliis ovato-lanceolatis. — In maritimis. *S. uniflora. α.* Fl. fr. 4. p. 747. (v. v.)

*δ. fabaria* (Otth.) repens, foliis subspathulatis margine tenuissimè ciliato-serratis. — In arenosis maritimis Vasconie. — Cucubalus fabarius. Thore. S. Thorei. Dufour in Ann. sc. nat. 5. p. 184. (v. v.)

§. 3. *Caulescentes, floribus verticillato-spicatis.*

3. *S. OTITES* (Pers. ench. 1. p. 497.) caulibus erectis vix pubescentibus oligophyllis, foliis inferioribus numerosis spathulatis subcarnosis, superioribus lanceolatis, floribus dioicis parvis, calycibus femininis sphaericis, masculinis subclavatis, petalis linearibus indivisis.  $\mathcal{L}$  in sterilibus arenosis. — Fl. fr. 4. p. 752. — Clus. hist. 1. p. 295. f. 1. (v. v.)

*β. umbellata* (Otth.) caulibus humilibus, floribus subumbellatis. — In maritimis circa Nannetes. (v. v. DC.)

§. 4. *Caulescentes, calyce conoideo, fundo umbilicato dentibus longissimis.* (Otth.)

4. *S. CONICA* (L. sp. 598.) pubescens, foliis linearibus mollibus, floribus solitariis aut paniculatis, calycibus brevioribus conicis 30-striatis, petalis bipartitis, capsulâ ovatâ.  $\textcircled{1}$  in arenosis Galliæ mediæ, occidentalis et australis. — Fl. fr. 4. p. 759. Jacq. aust. t. 253. (v. v.)

5. *S. CONOIDES* (L. sp. 598.) caulibus pubescentibus, foliis lanceolato-linearibus ferè glabris, floribus solitariis aut paniculatis, calycibus longis conoidis 30-

striatis, petalis integris obovatis, capsulâ lageniformi. ① in arenosis Nicæ ex All. Monspelii ex Gou., Calvadosii ex cl. de Brébisson. — Fl. fr. 4. p. 760. — Cus. hist. 1. p. 288. f. 2. (v. v. c.)

§. 5. *Caulescentes, calycibus cylindricis campanulatisve.*

\* *Floribus spicatis.*

6. S. ANGELICA (L. sp. 594.) pilosa, caulibus ramosis, foliis lanceolatis acutis, calycibus ventricosus-cylindricis 10-striatis, dentibus longissimis acutis, petalis obcordatis parvis, appendicibus pyramidalibus. ① in provinciis occidentibus. — Fl. fr. 4. p. 757. Eng. bot. t. 1178. (v. a.)
7. S. TRIDENTATA (Desf. atl. 1. p. 349.) caulibus ramosis, foliis lineari-lanceolatis, spicâ secundâ, calycibus subcylindricis decemcostatis, dentibus longis, petalis 3-dentatis. ① Tarbæ ad pontem Aturis reperiit cl. Ramond. — Fl. fr. p. 758. — Cus. hist. 1. p. 290. f. 1. An à priori satis distincta species? (v. a.)
8. S. LUSITANICA (L. sp. 594.) pilosissima, caulibus ramosissimis, foliis inferioribus obovato-spathulatis, superioribus lanceolatis obtusis, spicâ distichâ, calycibus cylindrico-ventricosus 10-striatis, dentibus longioribus, petalis indivisis, limbo margine exteriori denticulato. ① Monspelii. — Fl. fr. 5. p. 606. — Dill. elth. p. 311. f. 401. Nimia cum sequente affinitas. (v. a.)
9. S. GALLICA (L. sp. 595.) hirsuta, caulibus ramosis, foliis inferioribus spathulatis, superioribus lanceolatis obtusis, spicâ secundâ, calycibus cylindrico-ventricosus 10-striatis, dentibus brevibus acutis, petalis obovatis coronatis. ① in Galliâ mediâ, occidentali et in Occitaniâ. — Fl. fr. 4. p. 757. S. stricta Lapeyr. abr. 246. ex autopsiâ cl. Arnott. — Vaill. bot. par. t. 16. f. 12. (v. v.)
10. S. QUINQUEVULNERA (L. sp. 795.) caulibus villosis, foliis pubescentibus inferioribus spathulatis, superioribus lanceolatis, spicâ distichâ, calycibus villosis cylindricis demum ventricosus 10-striatis, petalis obovatis retusis coronatis purpureis margine pallescentibus. ② in provinciis australioribus. — Fl. fr. 4. p. 758. Eng. bot. t. 86. (v. v.)  
*ℓ. cerastoides*, petalis pallidioribus. — S. cerastoides L. sp. 596. Vill. Dauph. 4. t. 48. Fl. fr. 4. p. 756. (v. v.)
11. S. NOCTURNA (L. sp. 595.) caule subramoso infra piloso, foliis pubescentibus basi longè ciliatis, inferioribus spathulatis, superioribus lineari-lanceolatis, spicis secundis densis, calycibus cylindricis subglabris reticulato-costatis, petalis bipartitis angustatis. ①? in provinciis mediterraneis et agro Syrtico ex Thore. — S. spicata DC. fl. fr. 4. p. 759. — Dill. elth. t. 310. f. 400. (v. v.)
12. S. BRACHYPETALA (Rob. et Cast. in DC. fl. fr. 5. p. 607.) caule simplici, foliis obovato-spathulatis obtusis hirsutis basi longè ciliatis, floribus secundis erectis, calycibus subcylindricis hirsutis dentibus longis, petalis bifidis minimis. ① Massiliæ. (v. v. D.)

\*\* *Floribus paniculatis solitariisve.*

13. S. MUTANS (L. sp. 596.) pubescens, caulibus basi foliosissimis, foliis radicalibus spathulatis, superioribus lanceolato-linearibus, floribus paniculatis secundis pendulis, calycibus cylindrico-ventricosus, petalis bipartitis reflexis longè coronatis. 2ℓ in vineis, ad muros et in nemoribus. — Fl. fr. 4. p. 753. Fl. dan. t. 292. S. paradoxa. Lapeyr. abr. 246. ex cl. Arnott. (v. v.)
14. S. NICAËNSIS (All. ped. t. 44. f. 2.) viscosissimo-villosa, caulibus ramosis, foliis linearibus obtusis, floribus paniculatis suberectis, calycibus cylindricis, petalis longè unguiculatis semibifidis, lobis ovatis planis, appendice bifidâ. ① in maritimis Corsicæ, Nicæ, Galloprov. orientalis. — Fl. fr. 4. p. 754. (v. v.)
15. S. SENOIDES (Jacq. coll. supp. t. 14. f. 1.) viscoso-pubescens, caulibus erectis ramosissimis, foliis subcarnosis inferioribus spathulatis, superioribus ovatis, floribus parvis terminalibus et in dichotomiâ, petalis obcordatis coro-

natis. ① in apricis reptantibus maritimis Massiliæ. — Fl. fr. 5. p. 605. — Bocc. sic. t. 12. f. 4. (v. v.)

16. *S. QUADRIDENTATA* (DC. fl. fr. 4. p. 748.) cespitosa, caulibus erectis gracilibus ramosis, foliis parvis linearibus angustissimis, floribus minimis paniculatis, calycibus campanulatis, petalis breviter 4-dentatis. ① ½ in Alpibus et Pyrenæis. — *S. quadrifida* L. sp. 602. — Seg. ver. t. 5. f. 1. (v. v.)

17. *S. RUPESTRIS* (L. sp. 602.) glabra, caulibus erectis ramosis, foliis ovato-lanceolatis, floribus paniculatis minimis, calycibus campanulatis, petalis obcordatis. ½ in saxosis umbrosis Voges. Alp. Cebenn. Pyrenæorum. — Fl. fr. 4. p. 748. — J. Bauh. 3. p. 360. f. 3. (v. v.)

§. 6. *Caulescentes, calycibus clavatis, floribus solitariis paniculatisve.*

18. *S. SAXIFRAGA* (L. sp. 602.) cespitosa, caulibus assurgentibus, foliis linearibus acutis, pedunculis longissimis, floribus terminalibus solitariis raro axillaribus, calycibus longè clavatis, petalis bipartitis, unguiculo ciliato, appendice bicuspidata. ½ in saxosis apricis Alpium, Cebenn., Pyren. — Fl. fl. 4. p. 749. — Wald. et Kit. hung. 2. t. 163. (v. v.)

19. *S. SICULOR* (Thore land. p. 174.) cespitosa glabra subviscosa, caulibus basi ramosis, foliis linearibus, floribus paniculatis, calycibus longè clavatis reticulatis, petalis bifidis, lobis lanceolatis. ① in Galhiâ occidentali. — DC. fl. fr. 4. p. 751. ic. rar. t. 42. (v. v.)

20. *S. CLANDESTINA* (Jacq. coll. supp. t. 3. f. 3.) pubescens, caule erecto ramossimo gracili, foliis inferioribus oblongis obtusis, superioribus lanceolatis subangustis, floribus laxè paniculatis, petalis brevibus bifidis, calycibus ventricosis, capsulis rugulosis in calyce subæssilibus. ① in prov. occidentalibus et australibus? — *S. rubella* Fl. fr. 5. p. 604. non L. *S. annulata* Thore land. (v. s.)

21. *S. INAPERTA* (L. sp. 600.) glabra, caulibus erectis ramosis, foliis acutis inferioribus lanceolatis, superioribus linearibus, floribus laxè paniculatis, calycibus oblongo-clavatis, petalis insperitis obcordatis, capsulâ lævi in calyce pedicellatâ. ① in Corsicâ et inter Narbonam et Perpignanum. — Fl. fr. 5. p. 604. *S. polyphylla* Fl. fr. 4. p. 750. — Dill. elth. t. 314. f. 407. (v. v.)

22. *S. PAUCIFLORA* (Oth. in DC. prod. 1. p. 383.) cespitosa pubescens, caulibus numerosis simplicibus gracilibus oligophyllis, foliis linearibus lanceolatis basi numerosis, floribus solitariis vel paniculatis paucis, calycibus angustis, petalis bipartitis reticulatis. ½ in Corsicâ. (v. s.)

23. *S. CILIATA* (Pourt. act. Toul. 3. p. 328.) pubescens, caulibus numerosis prostratis basi valdè foliosis, foliis linearibus setaceo-ciliatis, floribus paucis secundis, calycibus inflato-clavatis, petalis bipartitis, sinibus calycis deflexis. ½ in Pyrenæis et summo Arvernæ jugo Cantal. — Fl. fr. 4. p. 756. *S. stellata* Lapeyr. non Ait. *S. Pourretii* Poir. (v. v.)

24. *S. PARADOXA* (Lin. sp. 1678.) caulibus erectis pubescentibus, foliis scabriusculis vix ciliatis, inferioribus lanceolato-spathulatis, superioribus linearibus, floribus magnis paniculatis, calycibus longis cylindrico-clavatis, petalis 2-lobis, lobis latius obovatis, appendice bipartita. ½ in Corsicâ et circâ la Roche des Arnauds in Delphinatû ex Vill. — Fl. fr. 4. p. 754. Jacq. vind. 3. t. 84. (v. s.)

25. *S. MUSCIPULA* (L. sp. 601.) glabriuscula viscosa, caule erecto, ramis alternis longis, foliis inferioribus lanceolato-spathulatis, superioribus linearibus, floribus paniculatis, calycibus amplo-clavatis reticulatis, petalis bifidis. ①? circâ Dragnignan, Narbonam et in Corbariensibus. — Fl. fr. 4. p. 752. — Clus. hist. 1. p. 289. f. 1. (v. v. DC.)

26. *S. NOCTIFLORA* (Lin. sp. 599.) viscoso-pubescens, caulibus erectis ramosis, foliis magnis, inferioribus spathulatis, superioribus lanceolatis, floribus magnis paniculatis, calycibus ventricosis venosis, dentibus longissimis, petalis 2-partitis denticulatis. ① in Alsatia, Vogesis, Delph. ex Vill. Galloprov. ex Ger. — Fl. fr. 4. p. 755. Eng. bot. t. 291. (v. s.)

27. *S. SERICHA* (All. ped. t. 79. f. 3.) caulibus ramosis, foliis basi longè ciliatis, inferioribus spathulatis, superioribus linearibus oblongis, floribus magnis terminalibus seu paniculatis paucis, calycibus clavatis, petalis longè unguiculatis bipartitis coronatis. ① in Corsicâ et Galloprovinciâ maritimis. — Fl. fr. 4. p. 759. *S. pubescens* Lois. *S. velutina* Pourr. Otth. in DC. prod. 1. p. 373. (v. s.)
28. *S. DECUMBENS* (Bern. sicul. cent. 1. p. 75.) pubescens, caulibus numerosis decumbentibus diffusis ramosis, foliis parvis spathulato-lanceolatis, floribus inclinatis, spicâ dichotomâ, calycibus clavatis angustis, petalis longè unguiculatis bipartitis. ① in Corsicâ reperiit Ph. Thomas. An à præcedente satis distincta? (v. s.)
29. *S. VALLESA* (Lin. sp. 603.) caespitosa viscido-pubescens, radice lignosâ, caulibus humilibus assurgentibus parum ramosis, foliis lanceolatis, inferioribus spathulatis, floribus terminalibus rarè geminis, calycibus longissimis reticulatis, petalis bifidis. ② in Alpibus Delph. et Galloprovinciâ. — Fl. fr. 4. p. 756. All. ped. t. 34. f. 2. (v. v. DC.)
30. *S. CORSICA* (DC. fl. fr. 4. p. 756.) pubescens viscosissima, caulibus procumbentibus foliosis, foliis obovatis, floribus terminalibus erectis, calycibus clavatis, petalis longissimè unguiculatis bipartitis coronatis. ② in Corsicâ. — Bocc. mus. t. 54. (v. s.)
31. *S. ITALICA* (DC. fl. fr. 4. p. 753.) piloso-pubescens, caulibus ramosissimis, foliis inferioribus ovato-spathulatis, superioribus linearibus, floribus amplo-paniculatis, calycibus longis clavatis, petalis bilobis nudis. ② in lapideosis ad sepes in provinciis australibus et circâ Mimatem et Albi. — Jacq. obs. 4. t. 79. (v. v.)
- β. cana* (Otth.) parum ramosa mollissima, paniculâ densâ. — In Corsicâ et Galloprovinciâ. S. Salzmanni. Otth. in DC. prod. 1. p. 381. (v. s.)
32. *S. CONGESTA* (Sibt. et Sm. prod. fl. græc. 1. p. 300. ex Otth. in DC. prod. 1. p. 384.) pubescens, caule subramoso, foliis obovatis acuminatis petiolatis, floribus corymboso-congestis, calycibus longissimè clavatis, petalis bipartitis nudis. ③ in Pyrenæis propè Fontpedrouze reperiit Ph. Thomas. An species à præcedente distincta? (v. s.)
33. *S. REQUIENII* (Otth. in DC. prod. 1. p. 381.) pubescens, radice crassâ lignosâ, caule erecto subsimplici folioso præcipuè basi, foliis oblongo-ovatis acuminatis, floribus paucis paniculatis, calycibus longissimis. ② in Corsicâ. — *S. fruticosa* Fl. fr. 5. p. 606. An *L. sp.* 597? (v. s.)
34. *S. ARMERIA* (L. sp. 601.) glaberrima glauca viscosa, caule ramoso, foliis ovato-lanceolatis, floribus paniculato-corymbosis, calycibus longè clavatis petalis obcordatis coronatis. ① in Galliâ mediâ, occidentali et australi. — Fl. fr. 4. p. 751. — Clus. hist. 1. p. 288. f. 1. (v. v.)

VI. *LYCHNIS*. DC. fl. fr. 4. p. 761. — *Lychnis* et *Agrostemma*. Lin.

Calyx tubulosus, 5-dentatus, nudus; petala 5 unguiculata, fauce sæpius coronatâ; stamina 10; styli 5; capsula 1-5-locularis, anthophoro longo vel nullo.

#### Sect. I. *VISCARIA*. DC. fl. fr. 4. p. 761.

Calyx cylindrico-clavatus, capsula semiquinque locularis; anthophorum longum.

1. *L. VISCARIA* (Lin. spec. 625.) caule subgeniculis viscoso, limbo petalorum subintegro, foliis linearibus subspathulatis. ② in pratis siccis Galliæ borealis et mediæ. — Fl. fr. 4. p. 761. — Clus. hist. 1. p. 289. f. 2. (v. s.)

#### Sect. II. *EULYCHNIS*. DC. l. c.

Calyx cylindrico-clavatus, capsula unilocularis, petala fauce appendiculata, anthophorum longum aut subabbreviatum.



2. *L. CHALCEDONICA* (Lin. spec. 625.) glaberrima, floribus fasciculatis, calycibus cylindrico-clavatis costatis, petalis bilobis, anthophoro longo, foliis lanceolatis cordatis subpilosis amplexicaulibus. 2/ in hortis culta sub nom. Croix de Malte. — Fl. fr. 4. p. 762. — Clus. hist. 1. p. 292. f. 1. (v. v. c.)
3. *L. FLOS-JOVIS* (Lin. sp. 625.) tomentosa, floribus umbellato-capitatis, calycibus cylindrico-clavatis costatis, petalis bilobis, anthophoro brevi crasso, foliis lanceolato-linearibus amplexicaulibus sericeo-tomentosis. 2/ in saxosis apricis Delphinatûs ex Vill., Galloprovinciæ ex Ger. — Fl. fr. 4. p. 763. — Moris. hist. 2. s. 5. t. 36. n. 2. (v. v. in Helv.)
4. *L. CÆLI-ROSA* (Desrouss. in Lam. dict. 3. p. 644.) glabra, caule dichotomo paniculato erecto, floribus solitariis terminalibus, calycibus clavatis decemcostatis, lobis acutissimis, petalis lobatis, anthophoro longo, foliis linearibus acutis. ① circa Nicæam ex All. et in Corsicâ. — Fl. fr. 4. p. 764. — Moris. s. 5. t. 22. n. 32. (v. v. c.)

## Sect. III. AGROSTEMA. DC. fl. fr. 4. p. 763.

Calyx ovoides, dentibus brevibus; capsula 1-localeris; anthophorum brevissimum vel nullum.

5. *L. SYLVESTRIS* (Hope. cent. exs. 3. n. 33. DC. fl. fr. 4. p. 763.) floribus (rubellis) dichotomo-paniculatis subdioicis, petalis semibifidis, lobis angustis divergentibus, capsulis subrotundis, valvis recurvis, foliis ovatis vel lanceolatis. 2/ in umbrosis humidis. — Fl. fr. 4. p. 763. Eng. bot. t. 1579. Fl. inodori subseasiles. (v. v.)
6. *L. DIOICA* (Lin. spec. 626. var.  $\beta$ . DC. fl. fr. 4. p. 762.) floribus (albis) dichotomo-paniculatis dioicis, petalis semibifidis, lobis latis approximatis, capsulis conicis, apicibus rectis, foliis ovatis. 2/ ad vias et sepes. — Fl. fr. 4. p. 762. Eng. bot. t. 1580. Flores odorati pedunculati. (v. v.)
7. *L. LETA* (Ait. Kew. ed. 2. t. 3. p. 134.) caule simplici, floribus solitariis, calycibus 10-striatis, petalis 2-fidis, foliis lineari-lanceolatis. ① in arenosis paludosis agri Syrtici reperierunt Cl. Laterrade et Des Moulins. — Planta vix pallicaria, fl. cærulescentes? (v. s.)
8. *L. CORSICA* (Lois. not. 73.) caule erectiusculo ramoso subdichotomo, pedunculis elongatis unifloris, petalis oblongis subemarginatis, foliis lineari-lanceolatis glabris. 2/ in Corsicâ. — Fl. fr. 5. p. 607. Pedalis et ultra, fl. rubelli. An var. præcedentis? (v. s.)
9. *L. PYRÆNICA* (Berg. fl. bass. pyren. 2. p. 264.) glabra, caulibus cespitosius diffusis, floribus dichotomo-fasciculatis, flore longè pedunculato in dichotomiâ, calycibus campanulatis, lobis brevibus, petalis subemarginatis appendiculatis, foliis coriaceis, radicalibus spathulatis longè petiolatis, caulinis cordatis sessilibus. 2/ ad rupes in Pyrenæis occidentilibus. — Fl. fr. 5. p. 608. *L. nummularia* Lapeyr. abr. 263. (v. v. DC.)
10. *L. FLOS-CUCULI* (Lin. spec. 625.) caulibus adscendentibus glaberrimis, floribus dichotomo-fasciculatis, calycibus campanulatis decemcostatis, petalis laciniatis appendiculatis, foliis lanceolato-linearibus. 2/ in pratis humidis. — Fl. fr. 4. p. 762. — Clus. hist. 1. p. 292. f. 2. (v. v.)
11. *L. ALPINA* (Lin. spec. 626.) glabra, caulibus cespitosius strictis, floribus densè umbellato-capitatis, calycibus campanulatis, petalis bifidis, anthophoro nullo, fructibus globosis, foliis lineari-lanceolatis. 2/ in pratis editiorum Alpium et Pyrenæorum. — Fl. fr. 4. p. 762. — Hall. helv. t. 17. f. 4. (v. v.)
12. *L. CORONARIA* (Lam. dict. 3. p. 643.) tomentosa, caule dichotomo, pedunculis elongatis 1-floris, calycibus subcampanulatis costatis, petalis emarginatis coronatis serratis, foliis lanceolatis latissimis coriaceis. 2/ Alpibus Helvetiæ et Pedemontii indigena, in hortis culta. — Fl. fr. 4. p. 763. Curt. bot. mag. t. 24. (v. v. c.)

Sect. IV. GITHAGO. *Desf. cat.* 159.

Calyx cylindrico-campanulatus coriaceus, lacinii longissimis; capsulæ 1-loculares; anthophorum nullum.

13. *L. GITHAGO* (Lam. dict. 3. p. 643. DC. fl. fr. 4. p. 764.) hirsuta, caule dichotomo, floribus longè pedunculatis, foliis linearibus. ① abundè inter segetes — *Agrostemma Githago* L. sp. 624. — Fuchs. hist. 127. ic. (v. v.)

VII. VELEZIA. *Lin. gen. n.* 448. *Gærtn. fruct.* 2. t. 129. f. 12.

Calyx tubulosus, 5-6-dentatus; petala 5-6 brevina, unguibus filiformibus barbatis, limbo emarginato; stamina 5-6; styli 2; capsula 1-locularis.

1. *V. RIGIDA* (Lin. spec. 474.) calycibus filiformibus pubescentibus, petalis bifidis. ① in aridis Galloprovinciæ et Occitanicæ. — Fl. fr. 4. p. 765. — Barr. ic. 1017 et 1018. (v. s.)

Trib. II. *ALSINEÆ*. DC. fl. fr. 4. p. 766.

Calycis sepala 4-5 libera aut imâ basi vix subcoalita.

VIII. GOUFFEIA. *Robill. et Cast. diss. ined. DC. fl. fr.* 5. p. 609.

Calyx 5-partitus patulus; petala 5 integra; stamina 10; styli 2; capsula globosa 1-locularis 2-valvis 1-2-sperma.

1. *G. ARUMARIOIDES* (Rob. et Cast. ined. DC. l. c.) ramis divaricatis, floribus subcorymbosis, foliis linearibus brevibus deflexis. ① in rupestribus apricis Massiliæ. (v. v. D.)

IX. BUFFONIA. *Sauv. meth. fol.* 141. *Lin. gen. n.* 225. *Gærtn. fruct.* 2. t. 129. f. 1.

Calyx 4-sepalus; petala 4 integra; stamina 4; styli 2; capsula compressa, 1-locularis, 2-valvis, 2-sperma.

1. *B. ANNUA* (DC. fl. fr. 4. p. 768.) caule basi laxè paniculato, ramis divaricatis brevibus firmis, striis calycinis rectis parallelis, capsulis calycem vix æquantibus, foliis subulatis basi dilatatis. ① ad sepes et vias Galliæ australis, in montosis agri Ligericini, Arvernici et circà Divionem (Cordienne). — Fl. fr. 4. p. 768. — Lam. ill. t. 87. f. 1. (v. s.)
2. *B. PERENNIS* (Pourr. act. toul. 3. p. 219.) caule apice ramoso, ramis filiformibus elongatis, striis calycinis arcuatis convergentibus. ② in siccis agri Ruscinonensis, Narbonensis et Monsp. — Fl. fr. 4. p. 768. Lam. ill. t. 87. f. 2. (v. s.)

X. SAGINA. *Lin. gen. n.* 236. *Gærtn. fruct.* 2. t. 129. f. 10.

Calyx 4-5-partitus; petala 4-5- aut 0; stamina 4-5; capsula 4-5-valvis, 1-locularis, oo-sperma.

1. *S. PROCUMBENS* (Lin. spec. 185.) glabra ramis procumbentibus, foliis linearibus mucronatis, pedunculis fructiferis adscendentibus, petalis brevibus obtusis, lacinii calycinis subrotundis. ① ad muros et in arvis sabulosis. — Fl. fr. 4. p. 768. Eng. bot. t. 880. (v. v.)
2. *S. APETALA* (L. mant. p. 759.) hispido-pubescent, ramis erectiusculis dichotomis, foliis linearibus mucronatis, pedunculis fructiferis adscendentibus, petalis brevissimis subrotundis, lacinii calycinis lanceolatis obtusiusculis. ① in sabulosis passim. — Fl. fr. 4. p. 769. — Arduin. spec. 2. t. 8. f. 1. (v. s.)
- β. *agglomerata* (Ser.) floribus sessilibus pedicellatisque agglomeratis. Circà Agatham. (Ph. Thomas.) (v. s.)

3. *S. ERECTA* (Lin. spec. 185.) caulibus erectis subunifloris glabris, foliis linearibus acutis, pedunculis strictis, petalis brevibus, sepalis lanceolatis acutis. (1) in sterilibus glareosis. — Fl. fr. 4. p. 769. — Vaill. bot. par. t. 3. f. 2. *Moenchia glauca* Pers. (v. v.)

XI. *MOEHRINGIA*. Lin. gen. 264. Gärtn. fruct. 226, t. 129. f. 11.

Calyx 4-partitus; petala 4; stamina 8; styli 2; capsula 4 valvis, 1-locularis, ∞-sperma.

1. *M. MUSCOSA* (Lin. spec. p. 515.) foliis linearibus connatis, laciniis calycinis planis longitudine ferè internodiorum lanceolatis acutis. 2/ in humidis montosis. — Fl. fr. 4. p. 771. — Pluck. t. 74. f. 3. et t. 75. f. 1. (v. v.)

XII. *ELATINE*. Lin. gen. n. 685. Gärtn. fruct. 2. p. 142. t. 102. f. 2.

Calyx 3-4-partitus; petala 3-4 unguibus destituta; stamina petalis numero æqualia? aut dupla, ideò 3-4 aut 6-8; styli 4 apice capitati; capsula 4-valvis, 4-locularis, ∞-sperma; semina cylindracea.

1. *E. HYDROPHILA* (Lin. spec. 527.) foliis oppositis, floribus alternis pedicellatis octandris tetrapetalis. 2/ in Gallia occidentalis, borealis, orientalisve inundatis. — Fl. fr. 4. p. 771. var. a.; ic. rar. t. 43. f. 2. — Vaill. bot. t. 2. f. 2. (v. s.)  
2. *E. HEXANDRA* (DC. ic. pl. rar. 1. p. 14. t. 43. f. 1.) foliis oppositis, floribus alternis pedicellatis hexandris tripetalis. 2/ in inundatis Gallia occidentalis, mediæ et boreali-orientalis. — Fl. fr. 5. p. 609. — Vaill. bot. t. 2. f. 1. (v. v. circa Genevæ D.)  
3. *E. ALSTRAUTUM* (Lin. spec. 527.) foliis verticillatis, floribus sessilibus verticillatisque. 2/ in inundatis circa Argentoratam, Parisios, Lugdunum, etc. — Fl. fr. 4. p. 772. — Vaill. bot. par. t. f. 6. (v. v.)

XIII. *HOLOSTEUM*. Lin. gen. n. 136. Gärtn. fruct. t. 130. f. 5. excl. sp.

Calyx 5-sepalus; petala 5 dentata; stamina 5 aut abortu 3-4; styli 3; capsula 1-locularis apice in 6 dentes dehiscens; embryo intra albumen replicatus.

1. *H. UMBELLATUM* (L. sp. 130.) foliis radicalibus ellipticis glaucis glabris, caulibus majoribus ovatis, floribus umbellatis, pedicellis post anthesin deflexis, valvis circinnatis. (1) in muris et arvis passim. — *Alaine umbellata*. Fl. fr. 4. p. 770. Lam. ill. t. 51. f. 1. (v. v.)

XIII. *SPERGULA*. Lin. gen. n. 798. Gärtn. fruct. t. 130. f. 4.

Calyx 5-partitus; petala 5 integra; stamina 5-10; styli 5; capsula 1-locularis, 6 valvis, ∞-sperma.

§. 1. *Foliis verticillatis basi stipulatis.*

1. *S. ARVENSIS* (Lin. spec. 630.) foliis verticillatis basi stipulatis, floribus decandris, seminibus sphaeroidis hispidulis angustè marginatis nigris. (1) in arvis sabulosis. — Fl. fr. 4. p. 773. Lam. ill. t. 392. f. 1 (v. v.)  
2. *S. PENTANDRA* (Lin. spec. 630.) foliis verticillatis basi stipulatis, floribus subpentandris, seminibus lenticularibus margine punctatis alatisque. (1) in sylvaticis Gallia mediæ, occidentalis et australis. — Fl. fr. 4. p. 773. Lam. ill. t. 392. f. 2. (v. s.)

§. 2. *Foliis oppositis, junioribus in axillâ fasciculatis; stipulis nullis.*

3. *S. MODICA* (Lin. spec. 630.) foliis oppositis subulatis levibus obtusiusculis, infimis latè vaginantibus, caulibus cespitosis subsimplicibus paucifloris, petalis calyce duplò longioribus, seminibus subreniformibus rugosis. 2/ in paludo-

- sis Gallie occidentalis, media, boreali-orientalis. — Fl. fr. 4. p. 773. — J. Bauh. 3. p. 724. ic. (v. v.)
4. *S. SAGINOIDES* (L. spec. 631.) caule repente, foliis oppositis linearibus lævibus, pedunculis solitariis longissimis, petalis obtusissimis sepala obtusa vix æquantibus, seminibus subreniformibus punctatis. ① in Gallia occidentali et circa Valletriam ex Ramond. — Fl. fr. 4. p. 774. Eng. bot. t. 3105. (v. s.)
5. *S. SUBULATA* (Swartz act. holm. 1789. p. 45. t. 1. f. 3.) subpilosa, foliis oppositis subsecundis lineari-subulatis subaristatis, pedunculis solitariis longissimis, petalis calycis longitudine. ② ① in sabulosis Gallie occidentalis et circa Parisios. — Fl. fr. 4. p. 775. Eng. bot. t. 1082. (v. s.)
6. *S. FILIFERA* (DC. fl. fr. 4. p. 774.) foliis oppositis linearibus aristatis subrigidis glabris fasciculatis, caulibus repentibus ramosis cespitosus, pedunculis longissimis, petalis calyce duplò longioribus, semin. subovoides. — In montibus editioribus Corsicæ. (v. s.)
7. *S. GLABRA* (Wild. spec. 2. p. 821.) decumbens, foliis oppositis filiformibus glabris subacutis, floribus decandris, petalis calyce majoribus. ② in Alpibus Delph. Galloprov. Corsicæ. — Fl. fr. 4. p. 774. — All. ped. t. 64. f. 1. (v. v. DC.)

#### XIV. LABRÆA. St.-Hil. in mem. mus. 2. p. 287.

Calyx 5-fidus basi urceolatus; petala 5 bipartita perigyna; stamina 10 perigyna; styli 5; ovarium 1-loculare, oo-spermum, capsula apice 6-valvis.

1. *L. AQUATICA* (St.-Hil. l. c.) caulibus dichotomis, foliis lanceolatis superioribus sessilibus, floribus axillaribus longè pedunculatis post anthesin deflexis. ② in paludosis et ad fossas. — Ser. in DC. prod. 1. p. 393. excl. syn. L. et Sm. — *Stellaria aquatica* Poll. pal. 422. Fl. fr. 4. p. 795. — Eng. bot. t. 1074. (v. v.)

#### XV. STELLARIA. Lin. gen. n. 773. Gært. fruct. 2. t. 130. f. 3.

Calyx 5-partitus; petala 5 bifida; stamina 10 aut abortu 3-8; styli 3; capsula 1-locularis, apice 6-valvis, oo-sperma.

1. *S. NEMORUM* (Lin. spec. 603.) foliis inferioribus cordatis petiolatis, superioribus lanceolatis subsessilibus, paniculà dichotomà, petalis calyce duplò longioribus, seminibus orbiculatis compressis margine tuberculatis. ② in nemoribus et umbrosis. — Fl. fr. 4. p. 793. Eng. bot. t. 92. (v. v.)
2. *S. LATIFOLIA* (Pers. ench. 1. p. 561.) caulibus diffusis dichotomis basi subradicantibus, foliis ovatis, inferioribus petiolatis subcordatis, superioribus sessilibus, pedicellis axillaribus, junioribus umbellato-aggregatis, fructiferis reflexis, petalis calyce brevioribus, seminibus disciformibus rugosis. ① in aquaticis circa Monspelium. — Fl. fr. 5. p. 614. (v. s.)
3. *S. MEDIA* (Smith. engl. bot. t. 537. fl. brit. p. 473.) caulibus procumbentibus lineà laterali pilosà alternà, foliis lanceolatis tenerissimis, fructibus deflexis, capsulis profundè 6-valvibus calyce vix longioribus, seminibus subreniformibus rugosis. ① in cultis frequens. — *Alsine media* L. ap. 389. Lam. ill. t. 214. DC. fl. fr. 4. p. 770 (v. v.)
4. *S. VISCIDA* (Bieb. taur. 1. p. 342.) villosa-viscosa, foliis lineari-lanceolatis, caulibus dichotomis diffusis, petalis calyce longioribus, capsulis subcylindricis sepalis lanceolatis duplò longioribus. ① prope Metz. (el. Soleirol.) — Ser. in DC. prod. 1. p. 397. Waldst et Kit. hung. 1. t. 22. (v. s.)
5. *S. RUBRA* (Bast. suppl. p. 24. DC. fl. fr. 5. p. 614.) foliis linearibus glabris margine subciliatis, caule erecto, pedunculis erectis, sepalis trinerviis. ① in agro Nannetensi et Andegavensi. (v. s.)
6. *S. HOLOSTRA* (Lin. ap. 711.) foliis lanceolatis acuminatis serrulatis scabrisculis, superioribus latioribus brevioribusque, pedunculis filiformibus præ-

- longis, petalis semihifidis calyce lanceolato acuto longioribus.  $\frac{1}{2}$  ad sepes et in demetis. — Fl. fr. 4. p. 794. Eng. bot. t. 511. (v. v.)
7. *S. GRAMINEA* (Lin. spec. 604.) foliis linearibus margine lævibus, caulibus diffusis, floribus paniculatis divaricatis, petalis calyce longitudine.  $\frac{1}{2}$  ad margines sylvarum. — Fl. fr. 4. p. 795. Eng. bot. t. 803. (v. v.)
8. *S. GLAUCA* (With. bot. arrang. 1. p. 420.) glauca, foliis lineari-lanceolatis margine lævibus, floralibus scariosis, petalis calyce duplò longioribus, caule erectiusculo debili.  $\frac{1}{2}$  ad fossas et in pratis humidis Gallie boreali-occidentalis et circà Parisios. — Fl. fr. 4. p. 794. Eng. bot. t. 825. (v. s.)
9. *S. CERASTIODES* (Lin. spec. 604.) foliis oblongis pubescentibus, pedunculis unifloris geminatis, fructiferis deflexis, petalis calyce majoribus, capsulis oblongis sepalis obtusis ferè duplò longioribus, caulibus cespitoso-decumbentibus subdichotomis biflorisque.  $\frac{1}{2}$  in herbosis humidis propè nives in Alpibus et Pyrenæis. — Fl. fr. 4. p. 796. Cerastium trigynum Vill. dampf. 3. t. 46. (v. v.)
10. *S. RADICANS* (Lapeyr. fl. Pyr. t. 93. ex Abr. p. 250.) caulibus prostratis teretibus multifloris, foliis ellipticis obtusis lucidis, caulinis subsecundis, pedunculis divaricatis, sepalis linearibus obtusis corollà duplò longioribus, capsulis pyriformibus.  $\frac{1}{2}$  in Pyrenæis ex Lapeyr. Cum priori conjungit el. Benth. †.

XVI. ARENARIA. Lin. gen. n. 774. Gartn. fruct. 2. t. 130. f. 9.

Calyx 5-sepalus; petala 5 integra; stamina 10 aut abortu pauciora; styli 3; capsula 1-locularis, apice 6-3-valvis, oo - sperma.

Sect. I. SPERGULARIA. Pers. ench. 1. p. 504.

Capsula trivalvis; folia linearia, stipulis scariosis ad basin donata.

1. *A. SEGETALIS* (Lam. fl. fr. 3. p. 43.) glabra, caule erecto, foliis subulatis stipulatis subsecundis, sepalis scariosis medio lineà viridescente longitudinaliter notatis, petalis calyce brevioribus, pedunculis defloratis deflexis, seminibus subpyriformibus rugosis. ① inter segetes circà Parisios, Rothomagum, in Delphinatà et Andegavià. — Fl. fr. 4. p. 792. — Vail. bot. par. t. 3. f. 3. (v. s.)
2. *A. RUBRA* (Lin. spec. 606.) caulibus prostratis pilosis, foliis filiformibus internodio brevioribus stipulatis, sepalis lanceolatis obtusiusculis margine scariosis, pedunculis defloratis deflexis, seminibus angulato-subpyriformibus rugosis immarginatis. ① in Gallià passim. — Fl. fr. 4. p. 792.
- a. campestris* (L. sp. 606.) pilosa. — In arvis sabulosis. — J. Bauh. 3. p. 723. f. 3. (v. v.)
- β. marina* (L. sp. 606.) glabriuscula. — In maritimis salsis. (v. v.)
3. *A. MEDIA* (Lin. spec. 606.) caulibus prostratis, foliis semi-cylindricis carnosius internodia æquantibus stipulatis, sepalis lanceolatis obtusiusculis margine scariosis, pedunculis defloratis deflexis, seminibus alà membranaceà cinctis. ① in pratis et sabulosis maritimis. — A. marginata DC. fl. fr. 4. p. 793. ic. rar. t. 48. Affinis *A. rubræ* marinæ sed minus ramosa. (v. v.)

Sect. II. ARENARIUM. Ser. in DC. prod. 1. p. 401.

Folia graminea linearia lanceolata vel rotundata, exstipulata; capsula 3-6-valvis.

§. I. Foliis subulatis linearibusve.

4. *A. LARICIFOLIA* (L. sp. 607.) foliis subulatis denticulato-ciliatis, caulibus ascendentibus 3-6-floris subcævis, calycibus cylindricis, sepalis oblongis obtusiusculis triplinerviis hirtis corollà duplò brevioribus, capsulis 3-valvibus calycem superantibus.  $\frac{1}{2}$  in Alpibus et Pyrenæis. — Fl. fr. 5. p. 612.
- a. multiflora* (Ser.) calycibus pedunculisque tomentoso-hirtis. — A. larici-folia Vill. dampf. 4. t. 47. f. 5. (v. v.)

- β. striata* (Ser.) calycibus pedunculisque hirsuto-viscosis. — *A. striata* Vill. dauph. 4. t. 47. f. 6. (v. v.)
5. *A. AUSTRACA* (Jacq. aust. 3. t. 270.) foliis lineari-subulatis 3-nerviis, caule subpaniculato, pedunculis terminalibus longissimis binis puberulis, petalis obtusis emarginatis, sepalis 3-nerviis acutissimis patentibus.  $\gamma$  in Alpibus Delph. et Galloprovinciæ. — Fl. fr. 4. p. 736. — All. ped. t. 64. f. 2. *A. triflora* Vill. dauph. 4. t. 47. (v. s.)
6. *A. GRANDIFLORA* (L. sp. 608.) foliis subulatis latiusculis planis trinerviis ciliatis, radicalibus confertis, caulibus subunifloris, pedunculis longissimis pubescentibus, sepalis ovatis aristatis grosse 3-nerviis corollâ duplò minoribus, capsulis ovatis 6-valvibus calycem vix superantibus.  $\gamma$  in petrosis editioribus Jurassi, Alpium, montium Occitanie, Pyrenæorum. — Fl. fr. 4. p. 787. All. ped. t. 10. f. 1. *A. mixta* Lapeyr. abr. p. 255. (v. v.)
- β. triflora* (Ser.) caulibus 2-4-floris, foliis angustioribus recurvis. In montosis sabulosis Fontis-Bleaudi, Jurassi, montis Ventosi, Pyrenæorum. (v. v.)
- γ. stolomifera* (Ser.) caulibus ramosissimis, foliis inferioribus latiusculis brevibus densis, superioribus sepalisque angustissimis longissimis distantibus. In Delphinatâ. (v. s.)
7. *A. VERNA* (L. mant. 72.) cespitosa multicaulis, foliis subulatis obtusiusculis nervosis, caulibus subpaniculatis elongatis, sepalis ovato-lanceolatis acuminatis striatis, petalis obovatis calycem vix æquantibus, capsulis cylindricis 3-valvibus.  $\gamma$  in collibus sabulosis montanis. — Fl. fr. 4. p. 788. Vaill. bot. t. 2. f. 3. (v. v.)
- β. cespitosa* (Ser.) caulibus valdè foliosis, calycibus pedunculisque glabrisculis. — *A. cespitosa* Ehrh. Fl. fr. 5. p. 613. (v. v.)
8. *A. GERARDI* (W. sp. 2. p. 729.) erecta ramosa, foliis lineari-subulatis 3-nerviis, pedunculis geminis terminalibus 1-floris, sepalis acuminatis margine membranaceis 3-nerviis.  $\gamma$  in pratis sterilibus Alpium Galloprovinciæ ex Ger? Delphinatâ ex Vill. — Fl. fr. 4. p. 788. — Ger. Gallop. t. 15. f. 1. An præcedentis var? (v. s. ex Pedem.)
9. *A. HERPINA* (L. sp. 608.) cespitosa erecta multicaulis piloso-hispida, peniculâ dichotomâ, foliis subulatis sepalisque subtus hispidis vix striatis, corollâ calycem superante, capsulis subglobosis trivalvibus calyce vix longioribus.  $\gamma$  circa Monspelium. — Fl. fr. 4. p. 789. Habitus *Spergularæ*, caules simplices pilis raris adpersi. (v. v. DC.)
10. *A. TENUIFOLIA* (L. sp. 607.) foliis subulatis setaceis, caule ramosissimo dichotomo, sepalis subulatis striatis petalis multò longioribus, capsulâ 3-valvi calycem vix superante. ① ad muros et in sabulosis. — Fl. fr. 4. p. 789.
- α. Vaillantii* (Ser.) caulibus cespitosis strictis glabris supernè ramosis. — Vaill. bot. t. 3. f. 1. (v. v.)
- β. Barrelieri* (DC.) caulibus ramosissimis glabris subprocumbentibus. — *A. Barrelieri* Vill. — Barr. ic. t. 580. (v. s.)
- γ. hybrida* (DC.) caule supernè ramoso glabro striato, calycibus piloso-glandulosis viscidis. — *A. hybrida* Vill. dauph. 4. t. 47. *A. pentandra* Duf. ann. gen. 7. p. 292. (v. v.)
- δ. viscidula* (DC.) caule erecto supernè ramoso calycibusque hispido-viscidis. — *A. viscidula* Thuil. *A. viscosa* Pers.
11. *A. RECURVA* (All. ped. t. 89. f. 3.) foliis radicalibus congestis recurvis subulatis subsecundis, caulibus cespitosis simplicibus subtrifloris, sepalis ovato-lanceolatis striatis pedunculisque piloso-glandulosis, petalis ovatis calycem æquantibus, capsulis ovatis 3-valvibus calyce brevioribus, seminibus subreniformibus vix punctato-rugosis.  $\gamma$  in pratis saxosis editiorum Alpium et Pyrenæorum. — Fl. fr. 4. p. 790. Jacq. coll. 2. t. 16. f. 1. sed in icône fl. nimis magni. (v. v.)
12. *A. SETACKA* (Thuil. fl. par. ed. 2. p. 220.) caule ramosissimo, floribus

paniculato-fastigiatis, foliis setaceis fasciculatis secundis basi ciliatis, calycibus ovato-acutis margine albo-membranaceis petala subæquantibus, capsulis 3-valvibus calyce superantibus.  $\frac{1}{2}$  ad saxa in collibus circè Parisios et Fontem-Bleaudi. — Fl. fr. 4. p. 790. A. heteromalla Pers. (v. s.)

13. A. FASCICULATA (Gou. ill. 30.) foliis subulatis fasciculatis setaceis, caulibus erectis strictis subimplexicibus, floribus fasciculatis, pedicellis folio brevioribus, sepalis acuminatis submembranaceis 2-lineatis petalis subduplò longioribus, capsulis 3-valvibus calyce brevioribus, seminibus reniformibus rugosis margine subserratis. ① ② in apricis montanis Jurassi, Delphinatús, et agri Ligericini. — Fl. fr. 4. p. 791. — Hall. helv. t. 17. (v. v.)

14. A. MUCRONATA (DC. fl. fr. 4. p. 791.) foliis setaceis basi non ciliatis, caulibus cespitosi basi prostratis, paniculà subdichotomà, pedunculis folio longioribus, sepalis acuminatis submembranaceis bilineatis aristatis petala capsulasque trivalves superantibus. ①? in saxosis montanis prov. austr., Pyrenæorum et agri Mimatensis. — A. mutabilis Lapeyr. abr. 256. An var. præcedentis. (v. v.)

15. A. ULIGINOSA (DC. fl. fr. 4. p. 786. ic. rar. t. 46.) erecta basi ramosa glabra, foliis linearibus obtusiusculis, ramis supernè nudis, pedunculis geminis unifloris subterminalibus basi bibracteatis, sepalis lanceolatis enerviis petalis ovatis vix longioribus, capsulis ovatis calycem æquantibus trivalvibus, seminibus subreniformibus.  $\frac{1}{2}$  in paludosis turfosis Jurassi. (v. s.)

§. 2. *Foliis lanceolatis ovalibus rotundatisque.*

16. A. TETRAQUETRA (L. sp. 605.) caulibus cespitosi, foliis oblongis coriaceis infernè margine ciliatis, sepalis ovato-lanceolatis coriaceis rigidis acutis trinerviis corollam subæquantibus, capsulis ovatis truncatis calycem æquantibus, seminibus reniformibus tuberculatis.  $\frac{1}{2}$  in sterilibus montosis Gallie Mediterraneæ. — Fl. fr. 4. p. 781. Gay. in ann. se. nat. 3. p. 27.

a. *uniflora* (Gay.) caulibus floriferis brevissimis, foliis dense imbricatis, floribus solitariis, sepalis petalisq. 4, stamin. 8. In Pyrenæis editioribus. (v. v.)

ß. *aggregata* (Gay.) caulibus floriferis elongatis, foliis supernè laxis, floribus 5-10 aggregatis, sepalis petalisque 5, staminibus 10. — Barr. ic. t. 595. (v. v.)

17. A. LANCEOLATA (All. ped. t. 26. f. 5.) cespitosa villosula, ramis ascendentibus, foliis lanceolatis angustis acutis rigidis nervosis, pedicellis foliis duplò longioribus, sepalis lanceolatis acutis nervosis corollà vix longioribus, capsulis ovatis longitudine calycis 3-valvibus, valvis apice callosis.  $\frac{1}{2}$  In Alpibus Galloprovinciæ et Delphinatús. — Fl. fr. 4. p. 785. A. cherlerioides Vill. dauph. 4. p. 47. t. 47. Folia imbricata. (v. v. DC.)

18. A. PURPURESCENS (Ram. in DC. fl. fr. 4. p. 785.) cespitoso-decumbens, ramis erectis 2-3-floris, foliis ovato-lanceolatis acuminatis glabris, pedicellis tomentosis folia vix superantibus, sepalis lanceolatis lævibus margine scariosis corollà longioribus, capsulis ovato-cylindricis semi-sexvalvibus calycem superantibus, valvis acutis, seminibus reniformibus.  $\frac{1}{2}$  in Pyrenæis editissimis. — Fl. fr. 6. p. 611. ic. rar. t. 45. A. cerastoides Lap. (v. v.)

19. A. MONTANA (L. sp. 606.) pubescens, foliis lanceolato-linearibus, caulibus sterilibus longissimis procumbentibus, pedunculis terminalibus longis unifloris, fructiferis nutantibus, sepalis lanceolatis acuminatis corollà multò brevioribus, capsulis ovato-globosis 6-valvibus calyce æquantibus, valvis obtusiusculis, seminibus reniformibus rugosis.  $\frac{1}{2}$  in sabulosis Gallie occidentalis, in montanis circè Mimatem, Mompelinum, et Pyren. orient. ex Benth. — Fl. fr. 4. p. 784. Vent. cels. t. 34. A. linearifolia Poir. (v. v.)

20. A. SERPYLLIIFOLIA (L. sp. 606.) foliis ovatis acutis sessilibus ciliatis, caulibus ramosissimis, sepalis lanceolatis acutis trinerviis corollà ferè duplò majoribus, capsulis ovatis apice 6-valvibus calycem æquantibus, seminibus reniformibus rugosis. ① ad muros et in arvis sabulosis. — Fl. fr. 4. p. 784. Fuchs. hist. 23. ic. (v. v.)

21. *A. CINEREA* (DC. fl. fr. 5. p. 811.) piloso-cinerea, foliis ovato-lanceolatis acutis ciliatis, superioribus distantibus lanceolatis acutis, floribus paniculato-dichotomis longè pedunculatis, sepalis lanceolatis acutis subcarinatis corollâ ferè duplò minoribus, capsulis ovatis calycem æquantibus sexvalvibus, valvis callosis.  $\frac{1}{2}$  in Galloprovinciæ petrosis. (v. s.)

22. *A. CILIATA* (L. sp. 608.) foliis ovatis obovatisve nervosis ciliatis, caulibus procumbentibus, sepalis lanceolatis acutis nervosis corollâ ferè duplò brevioribus, capsulis ovatis 6-valvibus calycem æquantibus.  $\frac{1}{2}$  in Alpibus Delphinatûs, Galloprov., Arvernæ, Pyrenæis. — Fl. fr. 4. p. 783. — Hall. hist. t. 17. f. 3. (v. v.)

$\beta$ . *multicaulis* (DC.) foliis subcoriaceis sepalisque vix nervosis. — *A. multicaulis* Wulf. in Jacq. coll. 1. t. 17. f. 1. (v. v.)

23. *A. TRINERVIA* (L. sp. 605.) caule gracili ramoso, foliis ovatis acutis petiolatis ciliatis nervosis, pedunculis longis post anthesin deflexis, sepalis lanceolatis acutis margine membranaceis corollâ multò longioribus, capsulis ovatis calycem æquantibus 6-valvibus, valvis apice circinnatis (I) ? in nemorosis. — Fl. fr. 4. p. 788. Fl. dan. t. 429. (v. v.)

24. *A. CERASTIFOLIA* (Ram. in DC. fl. fr. 4. p. 783.) cespitosa sublignosa ramosa, foliis ovato-lanceolatis trinerviis pubescentibus, pedunculis terminalibus unifloris piloso-glandulosis, sepalis lanceolatis obtusiusculis nervosis corollâ brevioribus, capsulis ovatis calycem vix superantibus trivalvibus.  $\frac{1}{2}$  in rimis saxorum in editioribus Pyrenæis. — *A. Ramondi* Poir. (v. v. DC.)

25. *A. BALNEARICA* (L. syst. nat. ed. 12. app. 230.) cespitosa repens, foliis ovatis lucidis subcarnosis ciliatis, pedunculis elongatis 1-floris, floribus cernuis, sepalis ovatis obtusiusculis enerviis corollâ multò brevioribus, capsulâ ovatâ calycem vix superante 6-valvi, valvis apice subcircinnatis.  $\frac{1}{2}$  in Corsicâ. — Fl. fr. 4. p. 782. L'Her. stirp. 1. t. 15. (v. s.)

26. *A. NIFLORA* (L. mant. 71.) foliis ovato-subrotundis glabris subimbricatis, caulibus procumbentibus ramosissimis, pedunculis bifloris lateralibus, sepalis ovatis obtusis corollâ brevioribus, capsulis ovatis longitudine calycis 3-valvibus.  $\frac{1}{2}$  in Alpibus editissimis Delphinatûs propè nives. — Fl. fr. 4. p. 782. All. ped. t. 44. f. 1. et t. 64. f. 3. (v. v. in Pedem.)

$\beta$ . *apetala* (DC.) foliis imbricatis ciliatis, floribus apetalis. — *A. apetala* Vill. dauph. t. 48. (v. s.)

XVII. ADENARIUM. *Rafin. journ. phys.* 1818. DC. in *prod.* t. 3. ined. — Honckeney, *Ehr. beytr.* 2. p. 181. — *Arenariæ sp. auct.*

Calyx 5-partitus; petala 5, imò calyci inserta integra; stam. 10, cum petalis inserta; styli 3-5; glandulæ 10, in toro circâ ovarium; capsula 3-5-valvis 1-locularis; semina pauca.

1. *A. PEPLOIDES* (Rafin. l. c.) foliis ovatis carnosis, floribus solitariis, seminibus maximis subpyriformibus.  $\frac{1}{2}$  in arenosis maritimis secûs Oceani ripas. — *Arenaria peploides*. L. sp. 605. Ser. in DC. prod. 1. p. 413. Fl. fr. 4. p. 781. fl. dan. t. 624. non Lapeyr. (v. v.)

#### XVIII. CERASTIUM. L. gen. 797.

Calyx 5-partitus; petala 5 bifida; stamina 10; styli 5; capsula 1-locularis cylindrica aut globosa, apice dehiscens dentibus 10 circinnatis vel ascendentibus.

##### §. 1. Calyce petala æquante vel superante.

1. *C. VULGATUM* (L. sp. 627.) hirsutum, caulibus erectis, foliis ovatis obtusissimis hirsutis, floribus dichotomis subumbellatis pedunculis longioribus, petalis calycem æquantibus, capsulis oblongis attenuatis calyce duplò longioribus. (I) (2) in raderatis. — Fl. fr. 4. p. 775. — Vaill. bot. t. 30. f. 3. (v. v.)

$\beta$ . *glomeratum* (DC.) foliis obtusissimis, floribus umbellato-glomeratis. C. ovale. Pers. (v. v.)



2. *C. viscosum* (L. sp. 627.) hirsuto-viscosum, caulis erectis, foliis lanceolato-oblongis, floribus dichotomo-subumbellatis, pedunculis petalisque calyce æquantibus, capsulis subpendulis teretibus calyce duplò longioribus. (1) in ruderalis. — Fl. fr. 4. p. 776. *C. obscurum* Chaub. in St.-Am. agen. et bouq. t. 4. f. 1. — Vaill. bot. t. 3o. f. 1. (v. v.)
3. *C. sennecianum* (L. sp. 607.) hirsuto-viscidum, caulis erectis, foliis ovato-lanceolatis, floribus pentandris dichotomo subumbellatis, pedunculis calyce longioribus, capsulis teretibus post florescentiam deflexis calyce duplò longioribus. (1) in sabulosis et ad margines agrorum. — Fl. fr. 4. p. 776. — Vaill. bot. t. 3o. f. 2. (v. s.)
  - β. *albidoides* (DC.) lobis calycinis margine spiceoque scariosis. — *C. alsinoides* Pers. — Circà Monopolium et Massiliam (Solier). (v. v.)
  - γ. *pellucidum*, lobis calycinis bracteisque spice scarioso-pellucidis. — In arena versus Agianum. — *C. pellucidum* Chaub. in St.-Am. agen. et bouq. t. 4. f. 2. Ser. in DC. prod. 1. p. 416. (v. s. D.)
4. *C. androsacrum* (Ser. in DC. prod. 1. p. 416.) pilosissimum, caule dichotomo divaricato, foliis ovatis obtusis, floribus solitariis aut subcapitatis pedunculatis basi subinvolutis, sepalis linearibus angustis acutissimis, petalis ovato-lanceolatis obtusis calyce triplò brevioribus, capsulà oblongà calyce paulò brevioris. (1) ad vias Corsicæ propè Calvi detexit al. Soleirol. (v. v.)
5. *C. brachystylum* (Desp. in Pers. ench. 1. p. 52o.) caule erecto tomentoso dichotomo, foliis ovatis, floribus paniculatis, pedunculis flore longioribus, calyce villosa petalis longiore, capsulis calycem vix superantibus. (1) in agro Mimattensi, Aginensi, Cænomanensi, Aureliensi et Argentor., probabiliter in omni Gallia boreali et occidentali. — Fl. fr. 4. p. 777. ic. rar. t. 44. Caulis folisque pilosissima. (v. s.)
6. *C. murali* (Desp. in DC. fl. fr. 5. p. 609.) hirsutum suberectum rigidum, foliis ovato-oblongis acutis numerosis, floribus fasciculatis longitudine pedicelli, petalis emarginatis calycem æquantibus, capsulis oblongis calycem vix superantibus. 2/ ? circà Cænomanum et Beauvais ex D. Gravet. (v. s.)

### §. 2. *Petalis calycem superantibus.*

7. *C. aquaticum* (L. sp. 609.) foliis cordatis sessilibus, floribus laxè dichotomo-paniculatis, petalis bifidis calyce vix longioribus, capsulis pendulis subglobosis calycem subsequantibus. 2/ in fossis aquosis prope lacus. — Fl. fr. 4. p. 780. Eng. bot. t. 538. (v. v.)
8. *C. Soleirolii* (Ser. ined.) hirsuto-viscosum, caulis prostratis ascendentibus 1-2-floris, foliis inferioribus ovatis confertis, superioribus lanceolatis, floribus erectis, pedunculis calycem subsequantibus, sepalis lanceolatis apice subscariosis, capsulis calycem æquantibus? 2/ in Corsicà reperiit Soleirol. (v. s.)
9. *C. romerosum* (L. sp. 629. var. β) caulis repentibus diffusis foliisque oblongis spatulatis tomentoso-canis, pedunculis erectis dichotomis, sepalis oblongis tomentoso-canis margine scariosis, capsulà subcylindracè calyce longiore. 2/ in montibus Galloprovinciæ, in hortis vulgatissimum. — Fl. fr. 4. p. 777. — Col. phytob. ed. 1744. p. 115. t. 31. (v. v.)
10. *C. lanatum* (Lam. dict. 1. p. 68o.) caulis prostratis cespitosus, foliis densè lanatis inferioribus subrotandis, superioribus ovatis, floribus dichotomis, sepalis lanceolatis margine scariosis, capsulà ovato-cylindricà calyce subduplò longiore. 2/ in Pyrenæis, Alpibus et Arvernæ summo jugo Cantal. — Fl. fr. 4. p. 778. (v. v.)
  - β. *Thomasianum* (Ser.) caulis foliis calycibusque glabris. — in valle Pyren. orient. dictà Eynes. (v. s.)
11. *C. alpinum* (L. sp. 628.) caulis prostratis, foliis ovato-lanceolatis pilosis nudisve, floribus dichotomis parvis longè pedunculatis, sepalis oblongis

obtusiusculis margine scariosis pilosis, petalis calyce duplò longioribus, capsulâ oblongâ calyce subduplò longiore.  $\frac{1}{2}$  in herbosis humidis Alpium, Pyrenæorum. — Fl. fr. fr. 4. p. 779. — Rai. syn. p. 349. t. 15. (v. v.)

12. *C. LATIFOLIUM* (L. sp. 627.) hirsuto - subviscosum, caulibus prostratis, foliis ovatis r-rarò 3-floris, floribus terminalibus, pedunculo florem æquante, sepalis ovatis margine scariosis, petalis calyce duplò longioribus, capsulâ ovatâ turgidâ calycem superante.  $\frac{1}{2}$  in Alpibus Delph. et Galloprovinciæ. — Fl. fr. 4. p. 778. Jac. coll. 1. t. 20. (v. v. in Pedem.)

13. *C. ARVENSE* (Lin. sp. 628.) caulibus declinatis, foliis lineari-lanceolatis obtusiusculis subpilosis, floribus dichotomo-paniculatis, pedunculis deflexo-pubescentibus, petalis calyce duplò majoribus, capsulâ oblongo-cylindricâ calyce duplò longiore.  $\frac{1}{2}$  ad vias et in versuris agrorum. — Fl. fr. 4. p. 778. — Vaill. bot. par. t. 30. f. 4. 5. (v. v.)

14. *C. STICTUM* (L. sp. 629? DC. fl. fr. 5. p. 610.) caulibus declinatis, foliis sublinearibus acuminatis glabris aut subhirsutis, pedunculis glanduloso-pubescentibus, petalis calyce duplò majoribus, capsulâ oblongâ.  $\frac{1}{2}$  in Alpibus et Galloprovinciæ montanis. An *C. arvensis* var.?

*a. suffruticosum* (DC.) foliis angustissimis glabrisculis. — *C. suffruticosum*. L. Fl. fr. 4. p. 780. Benth. cat. 70. (v. v.)

*β. lineare* (DC.) foliis lineari-lanceolatis elongatis acuminatis glabris. — *C. lineare* All. ped. t. 88. f. 5. (v. v.)

*γ. commune* (DC.) foliis linearibus obtusiusculis. — Scop. carn. 1. t. 19 f. 1. (v. v.)

#### XIX. CHERLERIA. Hall. itin. helv. 1. L. gen. n. 775.

Calyx 5-sepalus; petala 5 minima emarginata; stamina 10, styli 3, capsula 3-locularis? 3-valvis, loculis 2-spermis.

1. *C. SEDOTES* (L. sp. 608.) caespitosa parvula, foliis triquetris obtusiusculis nervosis patulis, capsulæ valvis obtusiusculis apice callosis calyce duplò longioribus.  $\frac{1}{2}$  prope nives æternas in editioribus Alpibus et Pyrenæis. — Fl. fr. 4. p. 781. Lam. ill. t. 379. (v. v.)

#### Ordo XV. LINEÆ. DC. theor. ed. 1. p. 217. prod. 1. p. 423.

Cal. sejala 3, 4, aut sæpiùs 5, basi vix imâ subcohærentia, per æstivationem imbricata cum pedunculo continua et ideò persistentia. Petala tot quot sejala, hypogyna, basi unguiculata, cum staminum annulo et interdum inter se subcoalita, sepalis alterna, per æstivationem contorta. Stam. petalorum numero æqualia, petalis alterna, basi in annulum monadelpha. Anthæræ basi insertæ, biloculares, birimosæ. Ovarium subglobosum loculis sæpiùs intus tot quot sunt sejala, rarò paucioribus. Styli tot quot loculi. Capsula globosa, constans carpellis marginibus induplicatis, apice bivalvi dehiscentibus, intus dissepimento incompleto è centro orto in concamerationes duas divisus. Semina in quoque carpello 2, inversa. Albumen sæpiùs nullum at hujus loco endoplevra tumida carnosâ. Embryo rectus, radiculâ ad hylum sursus tendente. — Herbæ aut suffrutices, foliis integris exstipulatis.

I. LINUM. *C. Bauh. Smich. Fl. brie. 1. p. 342. DC. prod. 1. p. 423.*

Flores proportione partium quinarîâ; sepala integra; styli rarissimè 3, cum petalis staminibus sepalisque 5.

§. 1. *Floribus flavis.*

1. *L. CALLICUR* (L. sp. 401.) glabrum, sæpius multicaule, foliis alternis lineari-lanceolatis, floribus laxè paniculato-corymbosis, pedicellis calycis longitudine, sepalis basi ciliatis apice subulatis, petalis obtusis calyce duplò longioribus. ① in arvis et sterilibus Gallie australis et occidentalis. — Fl. fr. 4. p. 796. Ger. Gallop. t. 16. f. 1. (v. v.)

2. *L. STRICTUM* (L. sp. 400.) glabrum erectum, foliis lineari-lanceolatis strictis sepalisque acuminatissimis ciliato-scabris, paniculâ corymbosâ confertâ, pedicellis brevissimis, petalis calycis longitudine. ① in agris et arenosis Gallie australis et occidentalis. — Fl. fr. 4. p. 798. — Lob. ic. t. 411. f. 2. (v. v.)

β. *alternum* (Pers.) caule supernè evidentius corymboso. In Corsicâ et Occitaniâ maritimâ ex cl. Benth. (v. s.)

3. *L. MARITIMUM* (L. sp. 400.) glabrum erectum, foliis lanceolatis trinerviis alternis, infimis oppositis obtusis, paniculâ laxâ subcorymbosâ, sepalis ovatis breviter mucronatis, petalis calyce triplò longioribus. ♀ in herbosis submaritimis Gallie australis. — Fl. fr. 4. p. 797. — Lob. ic. t. 412. f. 2. (v. v.)

4. *L. GLANDULOSUM* (Moench. meth. 287.) glabrum erectum basi suffruticosum, foliis basi biglandulosis lanceolatis alternis, infimis subpathulatis obtusis, sepalis acuminatis, petalis calyce triplò longioribus. ♀ §. *L. campanulatum*. Fl. fr. 4. p. 797.

α. *campanulatum* (L. sp. 400.) basi fruticosum glaucum, floribus corymbosis, sepalis lineari-lanceolatis. § in montanis lapideis Gallie australis. — Lob. ic. 414. f. 2. (v. v.)

β. *flavum* (L. sp. 399.) basi suffruticosum virens, floribus corymbosis, sepalis subciliatis. ♀ § in Gallopr. ? (v. s. ex Genuâ.)

§. 2. *Floribus cæruleis roseisve rarò albis.*

5. *L. HIRSUM* (L. sp. 398.) foliis lanceolatis 3-5-nerviis alternis suboppositisque cauleque hirsutis, summis sepalisque margine pilos glanduloso-capitato gerentibus, stylis stamina ferè duplò superantibus. ♀ in agro Nicœnsi et Pyrenæis orientalibus ex Lapeyr. — Fl. fr. 4. p. 800 et 5. p. 616. Jacq. fl. aust. t. 31. (v. s.)

6. *L. NARBONENSE* (L. sp. 398.) glabrum erectum, foliis lanceolato-linearibus acutissimis rigidiusculis, paniculâ subcorymbosâ, calycibus acuminatissimis basi ad marginem scariosis. ♀ in collibus sterilibus Gallie australis. — Fl. fr. 4. p. 799. — Barr. ic. t. 1007. (v. v.)

7. *L. USITATISSIMUM* (L. sp. 397.) glabrum erectum, foliis lanceolatis linearibusve, paniculâ corymbosâ, sepalis ovatis acutis margine membranaceis, petalis subcrenatis calyce triplò majoribus. ① in arvis. — Fl. fr. 4. p. 798. Black. herb. t. 180. (v. v.)

8. *L. ANGUSTIFOLIUM* (Huds. ang. 134.) glabrum multicaule suberectum, foliis lineari-lanceolatis acutis sub-3-nerviis, sepalis ellipticis sub-3-nerviis acutiusculis, petalis calyce vix duplò majoribus. ♀ in pascuis Gallie mediæ, occidentalis et australis. — Fl. fr. 4. p. 799. Eng. bot. t. 381. Flores pallide cærulei. (v. v.)

9. *L. MONTANUM* (Schleich. cat. DC. prod. 1. p. 427.) glabrum erectiusculum multicaule, foliis linearibus lanceolatisve acutis erectiusculis, floribus paniculato-corymbosis, pedicellis fructiferis erectis, sepalis ovalibus basi trinerviis, ext. acutis, int. obtusis margine membranaceis, petalis obtusis calyce triplò majoribus. ♀ in herbosis Jurassi, Alpium et Pyrenæorum. — L. Anas

- triacum Fl. fr. 5. p. 615. non Lin. L. *Alpinum* Fl. fr. 4. p. 799. non L.—Scop. carn. ed. 2. t. 11. (v. v.)
10. L. *ALPINUM* (L. sp. 1672.) glabrum multicaule decumbens, foliis linearisubulatis erectis, floribus paucis subcorymbosis, pedicellis fructiferis erectis, sepalis ovalibus basi trinerviis, exter. subacutis, inter. obtusis margine membranaceis, petalis obtusis calyce triplò majoribus.  $\frac{2}{3}$  in siccis montanis Galloprov., Arvernæ, Pyrenæorum.—Fl. fr. 5. p. 615. non 4. p. 799. Jacq. aust. t. 321. Flores cærulei. (v. v.)
11. L. *TEXENSOLIMUM* (L. sp. 398.) caulibus a basi ramosis erectis glabris, foliis linearis-etaceis glabriusculis, sepalis lanceolatis acuminatis medio glanduloso-ciliatis capsulam superantibus, petalis calyce triplò longioribus.  $\frac{2}{3}$  in collibus aridis Gallie occidentalis, australis, medise et orientalis passim. — Fl. fr. 4. p. 800. — Clus. hist. 1. p. 318. f. 2. Petala sordidè albo-carnæa. (v. v.)
12. L. *SALSOLOIDES* (Lam. dict. 3. p. 521.) caulibus basi fruticosis subtertosis ramosis, ramis ascendentibus, sterilibus brevioribus, foliis linearis-etaceis glabriusculis, sepalis ovatis acuminatis medio glanduloso-ciliatis capsulâ subbrevioribus, petalis calyce triplò longioribus.  $\frac{2}{3}$  in aridis Gallopr., Occitanie, Vasconie et Pictavie ex cl. Bast. — L. suffruticosum. Fl. fr. 5. p. 826. non Cav. — Barr. ic. t. 795? Corolla præcedentia. (v. v.)

§. 3. *Floribus albis, foliis oppositis.*

13. L. *CATHARTICUM* (L. sp. 401.) glabrum erectum, foliis oppositis obovato-lanceolatis, caule superne dichotomo. ① in pascuis et arvis.—Fl. fr. 4. p. 801. — Barr. ic. t. 1165. f. 1. (v. v.)

II. *RADIOLA*. Gm. *yst.* 1. p. 289.

Flores proportionem partium quaternariâ; sepala ad medium fere coalita, apice trifida.

1. R. *LUTIDUS* (Gmel. l. c.) ① in sabulosis humidis. — L. *Radiola* L. sp. 402. Fl. fr. 4. p. 801. — Vaill. bot. t. 4. f. 6. Herba passila floribus numerosis minimis. (v. v.)

Ordo XVI. *MALVACEÆ*. Brown. *cong.* p. 8. DC. *prod.* 1. p. 429. — *Malvaceæ* s. 1. 2. 3. *Juss. gen.* 271.

Calyx constans sepalis 5, rarissimè 3-4, basi plus minùs coalitis per æstivationem valvatis, sæpè extus gerens bracteas seu sepala exteriora involucrum calycemve exteriorem constituentes. Petala tot quot sepala et iis alterna, hypogyna, inter se æqualia, per æstivationem spiraliter contorta; nunc omninò distincta, sæpiùs imò staminum tubo adnata. Stamina numero petalorum æqualia aut sæpiùs multiplicia, definita aut sæpiùs indefinita, hypogyna; filamentis in tubum coalitis inæqualibus, exterioribus brevioribus; antheris unilocularibus rimâ transversali dehiscentibus. Ovarium carpellis plurimis sæpiùs circâ axim verticillatis, interdum congestis, rariùs liberis, fere semper coalitis constans; styli tot quot carpella nunc distincti, nunc in unum coaliti; stigmata tot quot carpella plus minùs distincta. Carpella nunc 1-2-sperma intus rimâ dehiscentia,

nunc polysperma loculicido-dehiscencia aut medio septifera, septo ad latus interius semina gerente, nunc sublibera, nunc in capsulam multilocularem, nunc in baccam anomalam conata. Albumen nullum. Embryo rectus, radicula tereti, cotyledonibus chrysaloïdeo-contortuplicatis. — Herbæ, frutices aut arbores, foliis alternis sæpius petiolatis stipulatis.

I. MALOPE. *L. gen. n. 843.*

Calyx cinctus involucri 3-phylo, foliolis cordatis; carpella plurima 1-sperma in capitulum aggregata.

1. *M. MALACOÏDES* (L. sp. 974.) foliis ovatis crematis, stipulis oblongo-Huearibus, pedunculis axillaribus 1-floris. ① in pratis Galloprovinciæ circa Grasse et Cannes. — Fl. fr. 4. p. 827. — Cav. diss. t. 27. f. 1. — Barr. ic. t. 1189. (v. s.)

II. MALVA. *L. gen. n. 841.*

Calyx cinctus involucri triphylo rariùs 5-6-phylo, bracteolis oblongis setaceis; carpella capsularia plurima in orbem disposita.

Sect. I. *Bismalvæ.* (Medik.)

Pedicellis axillaribus solitariis 1-floris, involucri 3-phylo, foliis multipartitis, caule herbaceo.

1. *M. TOURNEFORTIANA* (L. sp. 971.) foliis multipartitis, lobis trifidia linearibus, caule decumbente, pilis subnullis fasciculato-stellatis. ① in maritimis Galloprovinciæ circa Forum Julii, et agri Ruscinonensis ex cl. Arnott. — Fl. fr. 4. p. 830. Cav. diss. 2. t. 17. f. 3. (v. s.)
2. *M. ALCA* (L. sp. 971.) foliis inferioribus angulatis, superioribus 5-partitis incis, caulibus calycibusque pube stellatâ fasciculatâ subvelutino-scabris. ② ad sylvarum oras, in incultis et umbrosis. — Fl. fr. 4. p. 829. Cav. diss. 2. t. 17. f. 2. — Fuchs. hist. 80. (v. v.)
3. *M. MOSCHATA* (L. sp. 971.) foliis inferioribus reniformibus incis, caulibus multipartitis, laciniis linearibus, caulibus calycibusque pilis simplicibus scabris. ② in siccis et sterilibus. — Fl. fr. 4. p. 830. Cav. diss. 2. t. 18. f. 1. (v. v.)

β. *laciniata* (Desrouss.) foliis etiam inferioribus multifidis, laciniis linearibus. — Moris. s. 5. t. 18. n. 4. Variat etiam omninò glabra. (v. s.)

4. *M. FASTIGIATA* (Cav. diss. 2. t. 23. f. 2.) foliis cordatis 5-lobis acuminatis inæqualiter dentatis medio productiore cauleque erecto pube stellatâ tomentosis, pedicellis axillaribus et in corymbum terminalem approximatis. ③ in Arverniâ et propè Aginnum ex St.-Am. — Fl. fr. 5. p. 625. (v. v. DC.)

Sect. II. *Fasciculatæ.* (DC.)

Foliis angulatis cordatis 5 nerviis, pedicellis axillaribus plurimis 1-floris, calyce exteriori 3-phylo, caule herbaceo.

5. *M. SYLVESTRIS* (L. sp. 969.) caule erecto, foliis 5-7-lobatis acutis, pedicellis petiolisque pilosis. ② in incultis et ad sepes frequens. — Cav. diss. 2. t. 26. f. 2. Fl. fr. 4. p. 829. — Lob. ic. t. 650. f. 2. (v. v.)
6. *M. ROTUNDIFOLIA* (L. sp. 969.) caule prostrato, foliis cordato-orbiculatis obtusissimè 5-lobatis, pedicellis fructiferis declinatis petiolisque pubescentibus, corollis calyce duplò majoribus. ② in incultis et ad sepes. — Fl. fr. 4. p. 829. Cav. diss. 2. t. 26. f. 3. — Lob. ic. t. 651. f. 1. (v. v.)
7. *M. NICÆENSIS* (All. ped. n. 1416.) caule prostrato, foliis angulatis acutis, pedicellis 3-4 erectis petiolo quadruplò brevioribus, calycibus pilosis corollâ brevioribus. ① in ruderatis circa Nicæam, Mompelium et Burdigalam (Dea Moulins). — Fl. fr. 4. p. 828. Cav. diss. 2. t. 25. f. 1. (v. v. DC.)

8. *M. PARVIFLORA* (Lin. amœn. 3. p. 416.) caule patulo, foliis subrotundis obtuse angulatis crenatis glabrisculis, floribus axillaribus sessilibus glomeratis, corollis vix calyce longioribus. ② circa Niceam, Massiliam (Solier.), Montpellierum et in Pyren. orient. — Fl. fr. 4. p. 828. Cav. diss. 2. t. 26. f. 1. Carpella supernè rugosa margine dentata. (v. s.)
9. *M. CRISPA* (L. sp. 970.) caule erecto, foliis angulatis dentatis crispis glabris, floribus axillaribus sessilibus. ① ex Syria orta, in cultis et hortis subspontaneas. — Fl. fr. 4. p. 829. Cav. diss. 2. t. 23. f. 1. (v. v.)

### III. ALTHEA. Cav. diss. p. 91.

Calyx cinctus involucello 6-9-fido; carpella capsularia 1-sperma in orbem disposita.

#### SECT. I. ALTHÆASTRUM. DC. prod. 1. p. 436.

Carpella emarginata, margine membranaceo destituta; involucellum sæpius 8-9-fidum.

1. *A. OFFICINALIS* (L. sp. 966.) foliis utrinquè molliter tomentosis cordatis ovatisve dentatis indivisis subtrilobisve, pedunculis axillaribus multifloris folio multò brevioribus. ② in humidis passim, præsertim in provinciis occident. borealibusque. — Fl. fr. 4. p. 831. Cav. diss. 2. t. 30. f. 2. — Fuchs. hist. p. 15. ic. (v. v.)
2. *A. NARBONENSIS* (Pourr. in Cav. diss. 2. t. 29. f. 2.) foliis pubescentibus subtomentosis inferioribus 5-7-partitis, superioribus 3-fidis, pedunculis multifloris folio longioribus laxiusculis. ② in agro Narbonensi, et maritimis Montpellier ex cl. Benth. — Fl. fr. 4. p. 832. Jacq. ic. rar. 1. t. 138. (v. s.)
3. *A. CANNABINA* (L. sp. 966.) foliis pubescentibus subtus subincanis, inferioribus palmato-partitis, superioribus 3-partitis, lobis angustis grossè dentatis, pedunculis axillaribus multifloris laxis folio longioribus. ② ad sepes et in vineis Gallie australis occidentalisque passim. — Fl. fr. 4. p. 832. Cav. 2. t. 30. f. 1. — Loh. ic. t. 656. f. 1. (v. v.)
4. *A. HIRSUTA* (L. sp. 965.) foliis cordatis piloso-scabris suprâ glabris, infimis obtusis, superioribus 5-lobis, caule hispido, pedicellis 1-floris folio longioribus. ① ad sepes et in incultis Gallie mediterranæ, occidentalis et circa Parisios ex cl. Merat. — Fl. fr. 4. p. 832. Cav. diss. 2. t. 29. f. 1. (v. v.)

#### SECT. II. ALCEA. L. gen. n. 840.

Carpella margine membranaceo sulcato circumdata; involucellum 6-7-fidum.

5. *A. ROSA* (Cav. diss. 2. t. 29. f. 3.) caule stricto hirsuto, foliis cordatis 5-7-angulatis crenatis rugosis, floribus axillaribus sessilibus ad apicem subapicatis, petalis subcrenatis, unguibus villosis. ② Ex Oriente indigena, in hortis culta. — Fl. fr. 4. p. 821. Alcea rosea. L. (v. v. c.)

### IV. LAVATERA. L. gen. n. 842.

Calyx cinctus involucello 3-6-fido, foliolis nempè ad medium coactis; carpella capsularia, 1-sperma, in orbem circa axim variè dilatatum disposita.

#### SECT. I. STEGIA. Fl. fr. 4. p. 835.

Receptaculum seu axis fructus apice expansus in discum ovaria tegentem.

1. *L. TRIMESTRIS* (L. sp. 974.) caule herbaceo, foliis glabrisculis subrotundis cordatis, superioribus angulatis, pedicellis solitariis, orbiculo carpella omnino tegente. ① circa Niceam ex Allioni, Massiliam (Solier.), et in agro Ruscinonensi ex Lapeyr. — Cav. diss. 2. t. 3. f. 1. Stegia Lavatera Fl. fr. 4. p. 836. (v. s.)

#### SECT. II. OLBIA. Medik. malv. p. 41.

Receptaculum fructus centrale conicum exsertum.

2. *L. OLIVA* (L. sp. 972.) caule fruticoso pilis subs fasciculatis distantibus subscabro, foliis molliter cano-tomentosis 5-lobis, lobo medio elongato, summis oblongis subdivisis, floribus solitariis sessilibus. § in sepihus rupestribusque circa Olbiam, in insulâ Margaritæ (Stire), et agro Ruscinonensi propè Bagnols ex cl. Benth. — Fl. fr. 4. p. 833. — Lob. ic. t. 653. f. 2. (v. v.)
3. *L. PUNCTATA* (All. auct. p. 26.) caule herbaceo pube stellatâ subscabro, foliis subtomentosis, inferioribus orbiculato-cordatis, summis trilobis, pedicellis solitariis 1-floris folio longioribus. ① in agris et culis Nicæ, Gallu-provinciæ, Corsicæ. — Fl. fr. 4. p. 835. (v. v.)

Sect. III. *AXOLOPHA*. DC. prod. 1. p. 439.

Receptaculum apice truncatum in cristas tot quot carpella membranaceas laterales verticales expansum.

4. *L. MARITIMA* (Gou. ill. t. 11. f. 2.) caule fruticoso foliisque tomentosis subrotundis obtusè angulatis crenatis, pedicellis axillaribus solitariis. § in rupestribus maritimis provinc. mediterræ. — Fl. fr. 4. p. 834. Cav. diss. 2. t. 32. f. 3. (v. v.)

Sect. IV. *ANTHEMA*. Med. malv. p. 42.

Receptaculum parvum foveolatum, nec exsertum, nec in cristas expansum.

5. *L. ARBOREA* (L. sp. 972.) caule arboreo, foliis 7-angularibus subtomentosis plicatis, pedicellis axillaribus 1-floris confertis petiolo multò brevioribus. ② in Corsicâ et Caloneso (Belle-Ile en mer.) ex cl. Bory in herb. DC. — Fl. fr. 4. p. 834. Cav. diss. 5. t. 139. f. 2. (v. v. c.)
6. *L. NEAPOLITANA* (Ten. cat. 1819. n. 125.) caule herbaceo scabro erecto, foliis subrotundis 7-nerviis obtusissimè 7-lobatis crenatis, pedunculis axillaribus aggregatis, involucello calyce brevioris, calycis lobis acuminatis. 2/ circa Bonifacio ex cl. Viviani aora. 11. (v. s. ex Neap.)
7. *L. CRETICA* (L. sp. 973.) caule herbaceo hispido-scabro, foliis 5-lobis acutis, pedicellis axillaribus 1-floris aggregatis petiolo multò brevioribus. ① in Corsicâ ex cl. Viviani. — Cav. diss. 2. t. 32. f. 1. (v. s. ex Cret.)

V. *HIBISCUS*. L. gen. n. 846.

Calyx cinctus involucello sæpius polyphylo, rariùs foliolis paucis aut inter se coalitis; stigmata 5; carpella in capsulam 5-locularem coalita, valvis intùs medio septiferis, loculis polyspermis aut rariùs 1-spermis.

1. *H. SYRIACUS* (L. sp. 978.) caule inermi arboreo, foliis cuneiformi-ovatis trilobis dentatis, pedicellis petiolo vix longioribus, involucello 6-7-phylo, carpellis polyspermis. 5 ex Syriâ et Carniolâ ortus, in hortis cultus. — Fl. fr. 4. p. 837. Cav. diss. 3. p. 69. f. 1. (v. v. c.)
2. *H. ROSA* (Thore in Lois. gall. p. 434.) foliis cordatis dentatis subtrilobis subtùs cano-tomentosis, pedicellis axillaribus petiolo liberis 1-floris supra medium articulatis, involucello 8-15-phylo, carpellis polyspermis. 2/ in agro Syrtico propè Dax ad ripas fluminis Adour. — Fl. fr. 5. p. 626. H. palustris Fl. fr. 4. p. 837. non L. (v. v.)

VI. *SIDA*. Cav. diss. p. 5.

Calyx nudus 5-fidus; stylus apice multifidus; carpella 6-30 circa axim verticillata plus minùsve inter se coalita 1-locularia mono aut oligosperma.

1. *S. ANATILON* (L. sp. 963.) foliis subrotundo-cordatis acuminatis dentatis tomentosis, pedunculis petiolo brevioribus, carpellis 15 truncatis birostratis pilosis. ① in paludosis propè Vindomagus. — Fl. fr. 4. p. 836. (v. s.)

Ordo XVII. TILIACEÆ. *Juss. gen. p. 290. s. 2. excl. gen. DC. prod. 1. p. 503.*

Calyx extus nudus 4-5-sepalus, sepalis per æstivationem valvatis. Petala tot quot sepala, iis alterna sæpè ad unguem foveolata, integra, rarissimè nulla. Stam. hypogyna, libera, numero sæpiùs indefinita, rarò definita. Antheræ biloculares rimâ duplici longitudinali dehiscentes. Glandulæ tot quot petala, iis oppositæ, stipiti ovariifero adnatæ. Ovarium 1 carpellis 4-10 coalitis constans. Styli totidem in 1 coaliti, stigmatibus sæpiùs liberis. Capsula plurilocularis; semina in loculis plurima. Albumen carnosum. Embryo erectus, cotyledonibus planis foliaceis. — Arbores fruticesve, foliis simplicibus bistipulatis.

I. TILIA. *L. gen. n. 660.*

Calyx 5-partitus deciduus; petala 5 intus aut nuda aut squamulâ sucta; stam. ∞ filamentis liberis aut subpolyadelphis; ovarium globosum villosum 1-stylum 5-locul., loculis 2-spermis; nux coriacea, abortu 1-locularis 1-2-sperma.

1. *T. MICROPHYLLA* (Vent. diss. p. 4. t. 1. f. 1.) foliis cordato-subrotundis acuminatis argutè serratis glabris ad axillas venarum subtus barbatis petiolo vix longioribus, fructibus subglobosis vix costulatis tenuissimis fragilibus. *♂* in sylvis pasalm. — Fl. fr. 4. p. 825. (v. v.)
2. *T. PLATYPHYLLA* (Scop. carn. n. 641.) foliis cordato-subrotundis acuminatis argutè serratis subtus puberulis, fructibus turbinatis prominulo-costatis lignosis crassis. *♂* in sylvaticis submontanis. — Fl. fr. 4. p. 826. Vent. l. c. p. 6. t. 1. f. 2. (v. v.)

Ordo XVIII. AURANTIACEÆ. *Corr. ann. mus. 6. p. 376. DC. prod. 1. p. 535.*

Calyx urceolatus campanulatusve, disco subadnatus, brevis, 3-5-dentatus, marcescens. Petala 3-5 basi lata, nunc libera, nunc inter se subcoalita, disco hypogyno extus inserta, per æstivationem marginibus leviter imbricata. Stam. numero petalorum æqualia dupla aut multiplicia, disco hypogyno inserta; filamenta basi complanata nunc libera, nunc inter se variè adnexa polyadelphe, nunc longè monadelphæ, semper apice libera subulata; antheræ terminales basi insertæ erectæ. Ovarium ovatum multiloculare. Stylus 1; stigma subdivisum. Fructus (Aurantium DC.) constans 1°. indusio crasso glanduloso evalvi indehiscente; 2°. carpellis plurimis circa axim rationalem verticillatis membranaceis, nunc semina tantum



foventibus, nunc intus pulposis carnosive. Semina ad carpel-  
lorum angulum interiorem adfixa, nunc plurima nunc solita-  
ria, exalbuminosa, sæpius pendentia. Embryo rectus, radicula  
retracta superâ ad hilum versa, cotyledonibus magnis crassis.  
— Arbores aut frutices, in foliis, calycibus, petalis, stam.  
filamentis, fructus, indusio et cotyledonibus glandulis vesicu-  
laribus oleo volatili fætis instructæ, foliis alternis super cau-  
lem articulatis.

I. CITRUS. *L. gen. n. 1218.*

Flores proportionem partium sæpius quinariâ; calyx urceolatus 3-5-  
fidus; petala 5-8; stam. 20-60, filamentis compressis plus minusve  
basi coalitis polyadelphis, antheris oblongis; stigma hemisphæricum;  
fructus baccatus 7-12-locul. loculis polyspermis pulposis; spermo-  
dermis membranacea. — Folia 1-foliolata petiolo sæpè alato.

1. *C. MEDICA* (Risso ann. mus. 20. t. 2. f. 2.) petiolis nudis, foliis oblongis  
acutis, floribus 40-andris sæpè agynis, fructuum oblongorum rugosorum  
cortice crasso, pulpâ acidulâ. 5 in Asiâ, culta Nicæ et in Galloprovinciâ.  
— Ferr. hesp. t. 59. 60. 63. *C. medica* α Fl. fr. 4. p. 859. Petala extus purpu-  
rascentia. (v. v.)
2. *C. LIMONIUM* (Risso ann. mus. 20. p. 201.) petiolis subulatis, foliis oblongis  
acutis dentatis, floribus 35-andris sæpè agynis, fructuum oblongorum cortice  
tenuissimo, pulpâ acidissimâ. 5 in Asiâ, nunc in Galloprovinciâ et Nicæ  
cultâ. — *C. medica* β Fl. fr. 4. p. 860. — Ferr. hesp. t. 247, 211, etc. Petala  
extus purpurascens. (v. v.)
3. *C. AURANTIUM* (Risso ann. mus. 20. t. 1. f. 1 et 2.) petiolis subnudis, foliis  
ovato-oblongis acutis, floribus 20-andris, fructuum globosorum cortice tenui,  
pulpâ dulci. 5 in Asiâ orientali, nunc in provinciis australioribus culta. —  
*C. Aurantium* α Fl. fr. 4. p. 860. — Ferr. hesp. t. 427. 385, etc. Petala alba.  
(v. v.)
4. *C. VULGARIS* (Risso ann. mus. 20. p. 190.) petiolis alatis, foliis ellipticis  
acutis crenulatis, floribus 20-andris, fructuum globosorum cortice tenui  
scabroso, pulpâ acri amarâ. 5 in Asiâ, nunc in provinciis australioribus  
cultâ. — *C. Aurantium* β Fl. fr. 4. p. 860. — Ferr. hesp. t. 409, 433. Petala  
alba. (v. v.)

Ordo XIX. HYPERICINÆ. DC. *fl. fr.* 4. p. 860. *Chois. prod. hyp.* 32.

Calyx monosepalus 4-5-partitus seu 4-5-sepalus, persistens, sæpè inæqualis, nempe 2 sepalis minoribus externis, 2 aut 3 majoribus. Petala 4-5 hypogyna, calycis lobis alterna, contortè æstivata. Stamina numerosa, sæpius indefinita, basi polyadelphe rarò libera aut monadelphe; filamenta longa; antheræ oscillantes minutæ. Ovarium unicum, liberum. Styli plurimi longi aliquando in unum coaliti. Stigmata simplicia, rarò capitata. Capsula baccave multivalvis multilocularis; loculi numero stylos æquantes. Placenta integra centralis seu multipartita margine introflexo valvularum adfixa. Semina numerosissima, vulgò teretia rarò complanata. Embryo rectus; radícula infera; albumen nullum. — Herbæ aut frutices variis glandulis obsitæ, foliis oppositis rarissimè alternis sæpè punctatis.

I. ANDROSÆMUM. *All. ped. n.* 1440. *Chois. prod. hyp.* 37.

Capsula baccata sub-1-locularis; calyx 5-partitus, lobis inæqualibus; petala 5; styli 3; stamina  $\infty$  basi coalita.

1. *A. OFFICINALE* (All. l. c.) foliis sessilibus, floribus terminalibus pedunculatis.  $\zeta$  in humidis provinciarum occidentalium, australium rarior, et propè Fontem-Bleaudi ex Merat. — *Fl. fr.* 4. p. 861. *Black. herb. t.* 94. (v. v.)

II. HYPERICUM. *L. gen. n.* 902.

Capsula membranacea; styli 3-5 in quibusdam numero variabiles; stamina  $\infty$  basi polyadelphe, rarò numero subdefinita; petala 5; sepala 5 basi plus minusve coalita et inæqualia.

Sect. I. ASCYRRIA. *Chois. l. c.* 38.

Sepala basi coalita et inæqualia; styli 3-5. — Suffrutices, floribus vulgò terminalibus magnis paucis.

1. *H. HIRCINUM* (L. sp. 1103.) ramis alatis, foliis basi submarginatis dilatatis sessilibus apice acutis ovato-lanceolatis margine glandulosis, pedunculis 2-bracteatis, staminibus corollam superantibus, seminibus 2 appendiculatis.  $\zeta$  in Corsicâ. — *Schk. bot. handb.* 3. t. 213. 3. (v. s.)

Sect. II. PERFORARIA. *Chois. l. c.* 44.

Calyx 5-sepalus, sepalis æqualibus, styli vulgò 3. — Herbæ aut suffrutices, floribus axillaribus paniculatisve.

\* *Sepalis integris.*

2. *H. QUADRANGULUM* (L. sp. 1104.) caule herbaceo quadrangulo recto ramoso, foliis ovatis obtusis pellucido-punctatis margine nigro-punctatis, florum paniculis multifloris, calyce lanceolato, antheris nigro-punctatis.  $\mathcal{U}$  in paludosis fossisque humidis. — *Fl. fr.* 4. p. 862. — *Cam. epit.* 676. 10. (v. v.)

*ß. dubium* (Chois. in DC. *prod.* 1. p. 548.) caule minus ramoso, calyce

vix punctato, corollâ nigro-punctatâ, foliis vix pellacido-punctatis minus obtusis. — In Pyren. et montosis Gallie ab Argentorato ad Olbias. — *H. dubium* Leers. Fl. fr. 4. p. 862. Vill. dauph. 3. t. 44. Benth. cat. gr. (v. v.)

3. *H. REPENS* (L. sp. 1103.) caule suffruticoso tereti purpurascente ascendente et prostrato, foliis ovato-linearibus obtusis approximatis vix perforatis, corymbo paucifloro, calyce ovato-lanceolato acuto nigro-punctato.  $\mathcal{L}$  in Corsicâ et Galloprovinciâ inter Cannas et Antipolim. (v. s.)
4. *H. CATERUM* (L. mant. 106.) caule tereti ramosissimo, foliis sessilibus lanceolatis basi undulato-sinuatis pellucido-punctatis, calyce minimo obtuso.  $\mathcal{L}$  in sterilibus Monspelii propè Pont-Juvenal, forsan in lauis extraneis allatum. — Fl. fr. 4. p. 863. — Bocc. mus. 2. t. 12. (v. v.)
5. *H. HUMIFUSUM* (L. sp. 1103.) caule herbaceo humifuso, foliis oblongis obtusis tenuissimè pellucido-et nigro-punctatis, floribus cynosis terminatis, calyce lineari-lanceolato corollam superante, staminibus 15-20.  $\mathcal{L}$  in arenosis et agris. — Fl. fr. 4. p. 863. — Clus. hist. 2. p. 181. f. 3. (v. v.)  
 $\beta$ . *Liottardi* (DC.) caule humili erectiore bisannuo, floribus aliquandò 4-partitis, calyce longiore.  $\mathcal{L}$  in agris post massem circâ Gratianopolim. — *H. Liottardi* Vill. dauph. t. 44. (v. s.)
6. *H. PERFORATUM* (L. sp. 1105.) caule ancipiti, foliis obtusis ovato-ellipticis pellucido-punctatis, floribus paniculatis, calyce lanceolato pellucido-punctato, antheris nigro-punctatis, stylis divergentibus.  $\mathcal{L}$  in sylvis, incultis et ad sepes frequens. — Fl. fr. 4. p. 862. — Fuchs. hist. p. 835. ic. (v. v.)  
 $\beta$ . *elatum* (Chois.) caule elatiore, internodiis distantioribus, foliis minoribus margine revolutis, paniculis laxis paucifloris. (v. s.)
7. *mycrophylum* (DC.) omnibus partibus minoribus confertioribus, paniculis strictis multifloris. (v. s.)

\* \* *Sepalis dentatis sæpè dentato-glandulosis.*

7. *H. ELONUM* (Lin. sp. 1106.) caule herbaceo villosa tereti, foliis ambrotundis ovatis sessilibus suprâ emarginatis, paniculis laxis paucifloris, calyce serrato-glanduloso linearis acuto, staminibus paucis, stigmatibus capitatis.  $\mathcal{L}$  in paludosis Gallie mediæ et occidentalis. — Fl. fr. 4. p. 866. — Ments. pug. t. 7. f. 3. *Chironia uliginosa* Lapeyr. abr. supp. 39. (v. v.)
8. *H. TOMENTOSUM* (L. sp. 1106.) caule herbaceo tomentoso tereti ascendente, foliis ovatis obtusis subamplexicaulibus margine nigro-punctatis, paniculis laxis dichotomis, calyce acuminato, stigmatibus simplicibus.  $\mathcal{L}$  in pratis humidiusculis Gallie mediterranee. — Fl. fr. 4. p. 865. — Moris. 2. s. 5. t. 6. n. 5. (v. v.)
9. *H. HIRsutum* (L. sp. 1105.) caule tereti suffruticoso hirsuto recto, foliis ovato-oblongis nervatis pellucido-punctatis, paniculâ longâ racemosâ, calyce lineari subacuto, stylis divergentibus.  $\mathcal{L}$  in sylvis montosis passim — Fl. fr. 4. p. 865. — Moris. hist. 2. s. 5. t. 6. n. 11. (v. v.)
10. *H. NUMMULARIUM* (L. sp. 1106.) caule tereti ascendente herbaceo, foliis orbiculatis petiolatis, calyce ovato obtuso.  $\mathcal{L}$  in saxosis Pyrenæorum et Alpium Delphinatûs. — Fl. fr. 4. p. 866. Lam. ill. t. 643. f. 3. (v. v.)
11. *H. PULCRUM* (L. sp. 1106.) caule herbaceo recto tereti, foliis amplexicaulibus cordatis obtusis pellucido-punctatis margine subrevolutis, calyce ovato obtuso, corollâ glandulosâ.  $\mathcal{L}$  in umbrosis siccis. — Fl. fr. 4. p. 865. Lam. ill. t. 643. f. 4. (v. v.)
12. *H. DENTATUM* (Lois. gall. p. 499. t. 17.) caule suffruticoso tereti ascendente, foliis amplexicaulibus oblongis subobtusis pellucido-punctatis, superioribus aliquandò dentatis, calyce lanceolato acuto pilis glandulosis ciliato, corollâ antherisque nigro-punctatis.  $\mathcal{L}$  in insulis Stæchadum, prope Olbias et Telonem. — Fl. fr. 5. p. 631. (v. v. D.)
13. *H. MONTANUM* (L. sp. 1105.) caule herbaceo tereti recto, foliis amplexicaul-

- libus ovatis obtusis tenuissimè pellucido-punctatis margine nigro-punctatis, internodiis superioribus longioribus, calyce lineari-lanceolato acuto, antheris nigro-punctatis.  $\mathcal{L}$  in montanis. — Fl. fr. 4. p. 864. — J. Bauh. 3. p. 283. t. 2. (v. v.)
14. *H. vimbratum* (Lam. dict. 4. p. 148.) caule herbaceo tereti purpurascente simplici, foliis amplexicaulibus ovatis margine nigro-punctatis non pellucido-punctatis, calyce ovato acuto ciliato, stylis 3-5.  $\mathcal{L}$  in declivibus Jurassi, Alpium, Pyrenæorum. — Fl. fr. 4. p. 863. H. Richeri Vill. dauph. 3. t. 44. (v. v.)
- $\beta$ . *androsæmifolium* (Chois.) caule magis repente, foliis magis ovatis. Vill. dauph. 3. t. 44. (v. s.)
- $\gamma$ . *Burseri* (DC. fl. fr. 5. p. 630.) caule altiore, foliis obtusioribus, bracteis longioribus, minùs ciliatis, calyce breviori minùs ciliato. Pluck. t. 93. f. 6. (v. s.)
15. *H. hyssopifolium* (Vill. dauph. 3. t. 44.) caule suffruticoso tereti ascendente, foliis oblongo-lanceolatis subobtusis utrinquè attenuatis pellucido-punctatis in axillâ fasciculatis, calyce subobtusos, stylis 3-4.  $\mathcal{L}$  in montosis Galloprovinciæ, Cebennarum (cl. Prost), Pyren. orient. — H. diversifolium Fl. fr. 5. p. 631. (v. s.)
16. *H. linearifolium* (Vahl. symb. 1. p. 65.) caule herbaceo recto tereti, foliis linearibus obtusis margine nigro-punctatis non pellucido-punctatis, calyce lanceolato nigro-punctato.  $\mathcal{L}$  in ericetis provinc. occidentaliû, circâ Mimatem (cl. Prost). — Fl. fr. 5. p. 631. (v. s.)
17. *H. Comis* (L. sp. 1107.) caule suffruticoso ascendente tereti, foliis verticillatis linearibus margine revolutis, calyce subobtusosq.  $\mathcal{L}$  in collibus aridis Galloprovinciæ, Nicææ. — Fl. fr. 4. p. 867. bot. mag. 178. (v. v.)

Ordo XX. ACERINEÆ. DC. theor. ed. 2. p. 244. prod. 1. p. 593.

Cal. 5-rariùs 4-9-partitus. Petala totidem circâ discum hypogynum inserta, lobis calycinis alterna, sæpiùs concolora, rariùs nulla. Stam. disco hypogyno inserta, sæpiùs 8 rariùs 5-12. Antheræ oblongæ. Ovarium didymum. Stylus 1. Stigmata 2. Fructus constans carpellis 2 demùm separabilibus indehiscen-  
tibus samaroïdeis, nempè 1-locul. 1-2-spermis compressis superiùs desinentibus in alam membranaceam divergentem ad  
latus inferiùs incrassatam. Semina basi loculamentorum adfixa oblonga. Endopleurum subcarnosum. Albumen nullum. Embryo curvatus convolutusve, cotyledonibus foliaceis, radiculâ ad basin loculi directâ. — Arbores, foliis oppositis, (in nostris) simplicibus.

I. ACER. Mæsch. met. 334. DC. prod. 1. p. 593.

Flores polygami; calyx 5-lobus interdùm 5-partitus; stam. rariùs 5, sæpiùs 7-9.

§. 1. *Floribus racemosis.*

1. *A. PSEUDO-PLATANUS* (L. sp. 1496.) foliis cordatis 5-lobis glabris, lobis acu-

minatis inæqualiter dentatis, racemis pendulis subcompositis rachi staminumque filamentis hirsutis, fructibus glabris, alis subdivergentibus.  $\S$  in sylvis montosis. — Fl. fr. 4. p. 86. Duham. arb. 1. t. 368. (v. v.)

2. *A. CAMPESTRIS* (L. sp. 1497.) foliis cordatis 5-lobis, lobis grossè subdentatis, racemis erectis, fructuum alis divaricatissimis.  $\S$  in sylvis et sepibus. — Fl. fr. 4. p. 86g. J. Bauh. h. 1. p. 2. p. 166. ic.

a. *hebecarpum* (DC.) fructibus velutino-pubescentibus. In Galliâ orientali et australi. (v. v.)

$\beta$ . *collinum* (Wallr.) fructibus glabris, foliorum lobis obtusis, floribus minoribus. (v. v.)

$\S$ . 2. *Floribus corymbosis fasciculatisve.*

3. *A. OPULIFOLIUM* (Vill. danph. 4. p. 802.) foliis cordatis subrotundis 5 lobis, lobis obtusis obtusè grossèque dentatis, corymbis subsessilibus, ovariiis fructibusque glabris, alis subdivergentibus.  $\S$  in lapidosis apriciis Delphin., Galloprov., Cebennarum. — Fl. fr. 4. p. 86g. (v. v.)

4. *A. MONARZSULANUM* (L. sp. 1497.) foliis annuis cordatis trilobis, lobis subintegerrimis aequalibus, corymbis paucifloris erectis, fructibus glabris, alis vix divergentibus,  $\S$  —  $\S$ . In apriciis lapidosis Galliæ orientalis, australis et rariùs occidentalis. — Fl. fr. 4. p. 86g. — Pluck. t. 251. f. 3. (v. v.)

5. *A. PLATANOIDES* (L. sp. 1496.) foliis cordatis glabris 5-lobis, lobis acuminatis, dentibus paucis grossis acutis, corymbis pedunculatis erectiusculis fructibusque glabris, alis divaricatis.  $\S$  in sylvis montosis Arvernæ, Cebennarum, Alpium. — Fl. fr. 4. p. 868. Duham. arb. 1. t. 10. f. 1. — Cam. epit. 63. ic. (v. v. c.)

Ordo XXI. HIPPOCASTANÆ. DC. *theor. elem. ed.* 2. p. 244.  
*prod.* 1. p. 597.

Cal. campanulatus 5-lobus. Petala 5 aut abortu 4, inæqualia, hypogyna. Stam. 7-8 disco hypogyno inserta, libera, inæqualia. Antheræ subincumbentes. Stylus 1. Caps. junior 3-locul. 3-valvis, loculis biovulatis, valvis medio septiferis, ovulis septis adfixis; adulta coriacea subglobosa abortu 2-3-valvis, 2-3-locularis, 2-4-sperma. Semina grossa, subglobosa, variè compressa et angulata, testâ nitidâ glaberrimâ, hilo latò discolore basilari. Albumen nullum. Embryo curvatus inversus, cotyledonibus carnosissimis per germinationem intrâ spermodermium hypogæis, radiculâ conicâ curvatâ ad hilum directâ. — Arbores fruticesve, foliis oppositis compositis palmatis.

I. *ÆSCULUS*. L. *gen. n.* 462. *excl. sp.* DC. *prod.* 1. c.

Calyx campanulatus; petala 4-5 expansa, limbo ovato; stam. filamenta intorsum recurva; capsula echinata.

1. *Æ. HIPPOCASTANUM* (L. sp. 488.) floribus 5-petalis 7-andris, foliis 7 obovato-cuneatis acutis dentatis.  $\S$  in Indiâ boreali, in Galliâ nunc subspontanea. — Fl. fr. 4. p. 870. — Riv. pentap. irr. t. 123. (v. v.)

Ordo XXII. MELIACEÆ. *Juss. gen.* 263. *mem. mus.* 3. p. 436.

Cal. sepalis 4-5 plus minùs inter se in calycem 1-phyllum coalitis. Petala totidem sepalis alterna, hypogyna, ungue lato sæpiùs basi conniventia aut coalita, æstivatione sæpiùs valvatâ. Stam. numero petalorum sæpiùs dupla; filamenta in tubum apice dentatum longè coalita, antheris ad faucem tubi sessilibus intùs adnatis. Ovarium 1. Stylus 1. Fructus varius, (in nostris) drupaceus multilocularis aut abortu 1-locularis, valvis (dùm adsunt) medio septiferis. Semina nunc albuminosa nunc exalbuminosa, embryone formâ vario. — Arbores aut frutices foliis alternis exstipulatis.

I. MELIA. *L. gen. n.* 576.

Calyx parvus 5-fidus; petala 5 oblongo-linearia patentia; stam. 10; ovarium toro subelevato insidens; stylus filiformis; drupa ovata, nucleo 5-sulcato 5-locul. loculis 1-spermis; albumen carnosum.

1. *M. AZEDARACH* (L. sp. 550.) foliis bipinnatis, foliolis lævibus incisive subquinnatis. *h* ex Syriâ orta, nunc in provinciis australibus quasi spontanea. — *Fl. fr.* 4. p. 858. *Comm. hort.* 1, t. 70. (v. v. o.)

Ordo XXIII. AMPELIDEÆ. *H. B. et Kunth. nov. gen.* 5. p. 223. — *Vites. Juss. gen.* 267.

Calyx parvus margine integer aut subdentatus. Petala 4-5 dentibus calycinis alterna, disco ovarium ambiente extùs inserta et idè inter hypogyna et perigyna ambigua, basi latiora, rariùs in corollam 1-petalam coalita, per æstivationem intùs apice inflexa subvalvata. Stamina tot quot petala, antè petala disco inserta, interdùm abortu sterilia; filamenta libera aut basi subcoalita; antheræ ovatæ birimosæ dorso insertæ oscillatoriæ. Ovarium globosum liberum. Stylus 1 brevissimus aut subnullus. Stigma simplex. Bacca globosa, junior 2-locularis loculis 2-spermis, adultâ sæpè dissepimentis evanidis 1-locularis, aquosa aut subcarnosa, epicarpio non secedente. Semina 4-5 (aut abortu nulla) axi centrali funiculo brevi adfixa erecta, ossea. Albumen carnosum durum. Embryo erectus, radiculâ teretiùsculâ inferâ. — Frutices sarmentosi scandentes, foliis basi stipulaceis, inferioribus oppositis, superioribus alternis pedunculis oppositis, pedunculis interdùm in cirrhos evadentibus.

I. VITIS. *L. gen. n. 284.*

Cal. sub-5-dentatus; petala 5 apice cohærentia et instar calyptræ basi abscedentia simulque decidua; stam. 5; stylus 0; bacca 2-locularis 4-sperma, loculis seminibusve sæpè abortivis.

1. *V. VINIFERA* (*L. sp. 293.*) foliis lobatis sinuato-dentatis nudis aut tomentosis. 3 in Asia australi indigena, nunc exceptis provinciis boreali-occidentalibus ubique culta. — *Fl. fr. 4. p. 857. Blackw. herb. t. 154. Admodum varians.* (*v. v.*)
2. *V. LACINIOSA* (*L. sp. 293.*) foliis 5-sectis, segmentis petiolulatis multifidis. 5 passim culta. — *Corn. can. t. 183. Potius prioris mera varietas ex DC. prod.* (*v. v.*)

Ordo XXIV. GERANIACEÆ. *DC. fl. fr. 4. p. 838. prod. 1. p. 637.* — *Gerania. Juss. gen. 268.*

Calyx persistens 5-sepalus, sepalis plūs minūs inæqualibus æstivatione imbricatis, interdum uno, basi in calcar cavum cum pedunculo arcuè connatum, producto. Petala 5 sepalis alterna, unguiculata, æqualia aut inæqualia, in prioribus hypogyna, in posterioribus sæpius calyci inserta aut cum eo connexa. Stamina filamenta rarissimè libera, ferè semper basi plus minūs monadelpha, hypogyna aut perigyna, numero petalorum dupla, interdum nonnulla sterilia, æqualia aut inæqualia. Ovarium primà fronte 5-loculare in stylum longum crassum et stigmatibus 5 coronatum desinens, sed revera torus elongatus in axim subpentagonum gracilem. Carpella 5, submembranacea indehiscentia 1-locularia biovulata ad basim tori adpressa, apice gerentia stylos filiformes totidem secus tori sulcos arcuè adnatos et apice summo in stigmata 5 brevia acuta simplicia desinentes, post anthesim styli à basi ad apicem variis modis tortiles carpella è calyce extrahentes et cum iis demum è toro secedentes. Semina in carpellis solitaria, pendula, exalbuminosa; embryo curvus, radicula deflexa ad fundum carpelli directà, cotyledonibus convoluto-aut flexuoso-plicatis. — Herbæ aut suffrutices, foliis ad nodos inferiores oppositis, aut alternis pedunculis tunc folio oppositis.

I. GERANIUM. *L'Hér. ger. DC. fl. fr. 4. p. 844.*

Sepala 5 æqualia; pet. 5 æqualia; stam. 10 fertilia alterna majora; glandulæ nectariferæ ad basin stam. majorum; carpellorum aristæ intus glabræ, demum elasticè a basi ad axeos apicem circinnatim revolutæ.

\* *Perennia, pedunculis 1-floris.*

1. *G. SANGUINEUM* (*L. sp. 958.*) caule erecto aut diffuso ramoso, pedunculis axillaribus petiolo multò longioribus medio hibracteolatis, foliis oppositis

5-partitis, lobis 3-fidis, lobulis linearibus.  $\mathcal{Z}$  in pratis siccis sylvaticis. — Fl. fr. 4. p. 845. Cav. diss. 4. t. 76. f. 1. Flores sanguinei ampli. (v. v.)

\* \* Perennia, pedunculis bifloris.

2. *G. ARGENTRUM* (L. smæn. 4. p. 324.) caule brevissimo, foliis subradicalibus longè petiolatis utrinque sericeo-canis 5-7-partitis, lobis 3-fidis, lobulis linearibus, pedunculis subradicalibus, petalis emarginatis.  $\mathcal{Z}$  in Alpibus Galloprovinciæ. — Vill. dauph. 3. t. 40. Cav. diss. 4. t. 77. f. 3. Fl. fr. 4. p. 849. (v. s.)

3. *G. CINERUM* (Cav. diss. 4. t. 89. f. 1.) subcaule, foliis subradicalibus petiolatis glauco-pubescentibus 5-partitis, lobis cuneiformibus trifidis, pedunculis subradicalibus, petalis emarginatis.  $\mathcal{Z}$  in Pyrenæis editioribus. — Fl. fr. 4. p. 849. (v. v.)

4. *G. TUBEROSUM* (L. sp. 953.) radice subglobosâ, caule à basi ad bifurcationem nudo, foliis 5-partitis, lobis linearibus pinnatim inciso-serratis.  $\mathcal{Z}$  circa Massiliam, et Agatham ex Benth. — Fl. fr. 5. p. 628. — Lob. ic. t. 661. f. 2. Moris. hist. s. 5. t. 16. n. 21. (v. v. D.)

5. *G. NODOSUM* (L. sp. 953.) caule tetragono, foliis inferioribus 5-lobis, superioribus 3-lobis, lobis oblongis acuminatis serratis, petalis emarginatis.  $\mathcal{Z}$  in sylvis montosis Galloprov., Arverniæ, Mimatis, Aggini, Pyrenæorum. — Fl. fr. 4. p. 847. Cav. diss. 4. t. 80. f. 1. Carpella undiquè pilosa; folia subhis lucida. (v. v.)

6. *G. PHÆUM* (Lam. dict. 2. p. 658.) caule tereti, foliis 5-lobis inciso-dentatis, summis sessilibus, petalis patulis integris, staminum filamentis basi pilosis, carpellis transversè plicatis.  $\mathcal{Z}$  in Alpibus et Pyrenæis. — Fl. fr. 4. p. 846.

*a. vulgatus* (DC.) petalis livide rubro-fuscis basi albo-maculatis. — *G. phæum* L. sp. 933. Cav. diss. 4. t. 89. f. 2. (v. v.)

*β. lividum* (L'Hér. ger. t. 39.) petalis roseo-violaceis subemarginatis. (v. s.)

7. *G. SYLVATICUM* (L. sp. 954.) caule tereti erecto glabro, foliis sub-7-lobis, lobis oblongis inciso-serratis, pedunculis subcorymbosis, petalis subemarginatis, staminum filamentis subulatis ad medium ciliatis.  $\mathcal{Z}$  in sylvis et pascuis montosis. — Fl. fr. 4. p. 847. eng. bot. t. 121. Carpella pilosa. (v. v.)

*β. batrachioïdes* (Cav. diss. 4. t. 85. f. 1.) floribus majoribus, petalis integris. (v. s.)

8. *G. FRATENSE* (L. sp. 954.) caule tereti erecto tomentoso, foliis sub-7-lobis, lobis lineari-oblongis inciso-serratis, pedunculis subcorymbosis, petalis integris, staminum filamentis glabris basi valde dilatatis.  $\mathcal{Z}$  in pratis humidis montanis Pyren., agri Ligericini (Prost), Delph., etc. — Fl. fr. 4. p. 848. Cav. 4. t. 87. f. 1. Flores ampli cærulei (v. v.)

9. *G. PALUSTRE* (L. smæn. 4. p. 323.) caule decumbente pilis patulis subhispido, foliis 5-7-lobis, lobis inciso-dentatis, pedunculis longissimis pilosulis, pedicellis declinatis, petalis integris, staminibus subulatis glabris.  $\mathcal{Z}$  in humidis Alpium Delphinatûs? — Fl. fr. 4. p. 848. Cav. diss. 4. t. 87. f. 2. — Oed. fl. dan. t. 596. (v. v. D. in Helv.)

10. *G. ACONITIFOLIUM* (L'Hér. ger. t. 40.) caule subascendente glabriusculo, foliis subpeltato-7-partitis lobis laciniatis, pedunculis calycibusque villosis, petalis integris, staminibus subulatis glabriusculis.  $\mathcal{Z}$  in Alpibus Delphinatûs. — Fl. fr. 4. p. 848. *G. rivulare* Vill. dauph. 3. t. 40. Fl. albi lineati. (v. v. DC.)

11. *G. PYRÆNICUM* (L. mant. 97.) caule erecto ramoso, foliis reniformibus 7-lobis, lobis oblongis obtusis 3-fidis, lobulis 3-dentatis, floralibus 3-partitis, petalis emarginatis calyce paulò majoribus, carpellis ecristatis pubescentibus.  $\mathcal{Z}$  in montibus Delph., Arverniæ, Pyren. et circa Angers. (Bast.) — Fl. fr. 4. p. 850. eng. bot. t. 405. Cav. diss. 4. t. 99. f. 2. Variat flore albo et palidè purpureo. (v. v.)

\* \* \* Annuâ, pedunculis bifloris.

12. *G. MOLLE* (L. sp. 955.) foliis reniformibus; radicalibus 9-lobis, caulinis



- 7-lobis, lobis trifidis, petalis bifidis longitudine calycis mutici, carpellis glabris rugosis, seminibus lævibus. ① in ruderatis. — Fl. fr. 4. p. 850. — Vaill. t. 15. f. 3. (v. v.)
13. *G. FUSILLUM* (L. sp. 957.) foliis subreniformibus 7-lobis, lobis 3-fidis, petalis emarginatis longitudine calycis mutici, carpellis pubescentibus non rugosis, seminibus lævibus. ① in ruderatis. — Fl. fr. 4. p. 852. — Vaill. bot. t. 15. f. 1. (v. v.)
- ℓ. humile* (Cav. diss. 4. t. 83. f. 2.) foliis tenuius incisus, caule humiliore (v. v.)
14. *G. ROTUNDIFOLIUM* (L. sp. 957.) foliis radicalibus reniformibus 7-lobis, caulinis subrotundis basi truncatis 5-lobis, lobis 3-fidis, petalis integris longitudine calycis aristati, carpellis hirsutis, seminibus reticulatis. ① in paucis et pratis. — Fl. fr. 4. p. 852. Cav. diss. 4. t. 93. f. 2. Flores parvi rosei. (v. v.)
15. *G. COLUMBINUM* (L. sp. 956.) foliis 5-partitis, lobis multifidis linearibus, petalis emarginatis longitudine calycis aristati, carpellis glabris lævibus, seminibus reticulatis. ① in arvis. — Fl. fr. 4. p. 851. Cav. diss. 4. t. 82. f. 1. (v. v.)
16. *G. DISSECTUM* (L. sp. 956.) foliis 5-partitis, lobis 3-fidis linearibus, petalis emarginatis longitudine calycis aristati, carpellis pilosis, seminibus reticulatis. ① in arvis. — Fl. fr. 4. p. 851. Cav. diss. 4. t. 78. f. 2. Vaill. bot. t. 15. f. 2. Petala purpurea. (v. v.)
17. *G. LUCIDUM* (L. sp. 955.) glaberrimum, foliis rotundatis 5-lobis, calycibus pyramidato-angulatis transversè corrugatis, carpellis dorso sulcato-muricatis. ① in rupibus umbrosis. — Fl. fr. 4. p. 850. Cav. diss. 4. t. 80. f. 2. Flores rosei. (v. v.)
18. *G. ROBERTIANUM* (L. sp. 955.) foliis 3-5-partitis, lobis trifido-pinnatifidis, petalis integris calyce angulato aristato duplò longioribus, carpellis glabris reticulato-rugosis, seminibus lævibus. ① in rupibus, sepibus, incultis frequentis. — Fl. fr. 4. p. 855. Cav. diss. 4. t. 86. f. 1. (v. v.)
- ℓ. purpureum*. (Vill. dauph. 3. t. 40.) petalis calyce paulo majoribus. (v. s.)

II. ERODIUM. *L'Hér. DC. fl. fr. 4. p. 838.*

Sepala 5 æqualia; petala 5 regularia aut irregularia; stam. 10 filamentis basi monadelphis, 5 antheriferis, 3 alternis sterilibus; glandulæ 5 ad basin staminum steriliū; carpellorum aristæ intus barbatae demum elasticè spiraliter tortæ.

\* *Foliis pinnati-sectis aut pinnati-partitis.*

1. *E. PETRÆUM* (W. sp. 3. p. 625.) acaule, pedunculis multifloris, foliis glabrisculis pinnati-sectis, segmentis pinnatifidis, lobis lanceolato-linearibus, rachi inter segmenta dentatâ, petalis retusis calyce duplò longioribus. ② in Occitanis, agri Roseinonensis et Pyren. orientalium rupibus apricis. — Fl. fr. 4. p. 839. Gou. ill. t. 21. f. 1. Flores purpurascens. (v. v.)
2. *E. GLANDULOSUM* (W. sp. 3. p. 628.) acaule, pedunculis multifloris, foliis pubescenti-glandulosis pinnati-sectis, segmentis pinnatifidis, lobis lanceolato-linearibus, rachi inter segmenta dentatâ, petalis subæqualibus acutis calyce duplò longioribus. ② in editioribus Pyrenæis. — Cav. diss. 5. t. 125. f. 2. *E. graveolens* Lapeyr. abr. 390. An à præcedente satis distinctum? (v. v. DC.)
3. *E. CICONIUM* (W. sp. 3. p. 629.) caule ascendente folisque subvillosis pinnati-sectis, segmentis obtusis pinnatifidis dentatis, rachi inter segmenta dentatâ, pedunculis multifloris, petalis calycis longitudine 2 subemarginatis. ① in maritimis Occitanis. — Fl. fr. 4. p. 841. Cav. diss. 4. t. 95. f. 2. Calycis sepala longè mucronata. (v. s.)
4. *E. CICTARIUM* (Léman. in DC. fl. fr. 4. p. 840.) caule prostrato aut diffuso

hirto, foliis pinnati-sectis, segmentis sessilibus pinnatifidis incisive acutisve, pedunculis multifloris, petalis inæqualibus. ① in ruderatis frequens. — Fl. fr. 4. p. 840.

α. *præcox* (Cav. diss. 5. t. 126. f. 2.) acaule, foliis patulis, segmentis subincisis, petalis calyce majoribus. — In apricis. (v. v.)

β. *pimpinellæfolium* (Cav. diss. 4. t. 126. f. 1.) caulescens demum suberectum, foliis longè petiolatis, segmentis acutè incisis, petalis calyci subæqualibus. — In pratis. (v. v.)

γ. *Cherophyllum* (Cav. diss. 4. t. 95. f. 1.) multicaule subprostratum, segmentis tenuiter pinnati-partitis, floribus pallidis aut albis. — In siccis lapidosis. (v. s.)

δ. *pilosum*. (Thuil. fl. par. ed. 2. p. 347.) multicaule subprostratum undique pilosum, segmentis tenuiter pinnati-partitis, floribus intensè purpureis. — In arenosis. (v. v.)

ε. *cicutæfolium* (Thuil. l. c. sub cicutario.) subcaulescens, segmentis oblongis obtusè incisis, floribus pallidis roseisve. — In apricis. (v. v.)

5. *E. ROMANUM* (W. sp. 3. p. 630.) subacaule, foliis pinnati-sectis, segmentis ovatis pinnatifidis, pedunculis multifloris, petalis calyce majoribus æqualibus. 2/2 secus vias Occitanicæ, agri Ruscinonensis, et Tolosæ ex Benth. — Fl. fr. 5. p. 628. Cav. diss. 4. t. 94. f. 2. — Barr. ic. t. 1245. (v. v.)

6. *E. MOSCHATUM* (W. sp. 3. p. 631.) caule procumbente, foliis pinnati-sectis, segmentis petiolulatis ovatis inæqualiter serrato-incisis, pedunculis multifloris pubescenti-subglandulosis. ① in siccis Nicææ, Occitanicæ, agri Ruscinonensis; circà Burdigalam ex Laterrade, Abbatisvillam ex Boncher. — Fl. fr. 4. p. 841. Cav. diss. t. 94. f. 1. — Lob. ic. t. 658. f. 2. Variat glabrum et in caule costisque glanduloso-pubescent. (v. v. c.)

7. *E. BOTRYS* (Bert. amæn. it. p. 35.) caulescens setis substrigosis hispidum, foliis sinuato-pinnatifidis, lobis obtusis dentatis, pedunculis 2-4-floris, calycibus pubescentibus breviter mucronatis. ① in Corsicâ. — Ger. botrys Cav. diss. 4. t. 90. f. 2. (v. s.)

8. *E. LITTORÆUM* (Léman in DC. fl. fr. 4. p. 842.) caulescens diffuso-prostratum glabriusculum, foliis sinuato-pinnatifidis, lobis 3-7 obtusis crenatis, pedunculis multifloris elongatis, aristis barbatis. 2/2 in Corsicâ et circà Narbonam. (v. s.)

\*\* *Foliis subtrilobatis aut indivisis.*

9. *E. MALACHOIDES* (W. sp. 3. p. 639.) caule herbaceo ramoso pilosiusculo, foliis cordatis indivisis trilobisve obtusis dentatis, pedunculis multifloris, petalis calycis longitudine. ① in provinciis mediterraneis et circà Burdigalam ex Laterr. — Fl. fr. 4. p. 842. Cav. diss. 4. t. 91. f. 1. — J. Bauh. hist. 3. p. 472. t. 1. (v. v.)

10. *E. MALIPOIDES* (W. sp. 3. p. 640.) caulescens erectum aut subacaule totum molliter cinereo-velutinum, foliis ovatis subcordatis crenatis interdum sublobatis, pedunculis subbifloris, aristis barbatis. 2/2 in Corsicâ rupibus maritimis. — E. malopoides β. DC. prod. 1. p. 648. E. Corsicum. Lem. in fl. fr. 4. p. 842. — Cav. diss. 4. t. 90. f. 1. (v. s.)

11. *E. MARITIMUM* (Smith. fl. brit. 2. p. 728.) caulescens diffuso-prostratum glabriusculum, foliis cordatis ovatis inciso-crenatis pubescentibus, pedunculis uni aut sæpius bifloris, aristis imberbibus. 2/2 in Gallicæ Narbonensis, Armoracæ et Picardiæ maritimis. — Fl. fr. 4. p. 843. Cav. 4. t. 88. f. 1. (v. s.)

β. *Bocconi* (DC.) subcaulescens cespitosum pumilum, foliis ovatis inciso-crenatis, pedunculis 1-rarius-2-floris, petalis vix calycis longitudine. 2/2 in Corsicâ montibus. — E. Bocconi. Viv. cors. app. 5. — Bocc. mus. 2. p. 160. f. 128. (v. s.)

Ordo XXV. TROPÆOLEÆ. *Juss. mem. mus. 3. p. 447. DC. prod. 1. p. 683.*

Calyx 5-partitus coloratus, laciniâ superiore basi calcaratâ, calcare libero intrâ florem hinc hiantè extrâ stamina, lobis nunc liberis, nunc inter se variè coalitis. Petala 5 calyci inserta, ejusdem lobis alterna, inæqualia, irregularia, 2 superiora sessilia remota, calcaris faucibus inserta, 3 inferiora unguiculata minora interdum abortiva. Stamina 8, filamentis liberis disco insertis, antheris 2-ocularibus rimâ duplici dehiscentibus. Carpella 3 arcuè connata in ovarium 1 trigonum. Styli 3 in 1 connexi. Stigmata 3. Carpella 3, 1-ocularia, 1-sperma. Semina magna, exalbuminosa. Embryo magnus, cotyledonibus rectis, junioribus distinctis, dein arcuè conferruminatis et etiam cum spermodermate adhærentibus, imâ basi subdistinctis; radicula intrâ cotyledonum processus latente, tubercula 4 mox radicellas proferentia gerente. — Herbæ foliis alternis non stipulaceis peltinerviis.

I. TROPÆOLUM. *L. gen. n. 466.*

Cal. 5-partitus, lobo superiore calcarato; petala 5 inæqualia, 3 inferiora minora aut evanida; stam. 8 ab ipsâ basi libera; carpella 3 suberosa reniformia indehiscentia hinc sulcata rotundata.

1. *T. minus* (*L. sp. 460.*) foliis peltinerviis orbiculatis subrepandis, nervis apice mucronatis, petalis acuminato-setaceis. ① sub dio. ② in cald. ex Peruvia ortum in hortis cultum. — *Curt. bot. mag. t. 98. (v. v. c.)*
2. *T. majus* (*Lin. sp. 490.*) foliis peltinerviis orbiculatis subquinelobis, nervis apice non exsertis, petalis obtusis. ① ② in Peruvia undè in Europam allatum et in hortis cultum. — *Fl. fr. 4. p. 854. Curt. bot. mag. t. 23. (v. v. c.)*

Ordo XXVI. BALSAMINÆÆ. *A. Rich. dict. class. 2. p. 173. DC. prod. 1. p. 685.*

Calyx 2-sepalus, sepalis parvis deciduis oppositis sæpè mucronatis per æstivationem imbricatis. Petala 4 hypogyna cruciata; 2 exteriora sepalis alterna apice callosa; superius fornicatum emarginatum, inferius integrum basi in calcar productum; 2 interiora prioribus alterna, magis petaloidea, inter se æqualia, sæpè 2-fida seu appendiculata. Stamina 5 hypogyna, filamentis brevibus apice incrassatis, antheris subconnatis, 3 inferiora petalis opposita antheras constanter 2-oculares gerentia, 2 superiora antè petalum supremum orta antheras nunc 2-oculares nunc 1-oculares gerentia. Antheræ rimâ

longitudinali dehiscentes. Ovarium 1. Stylus 0. Stigmata 5 aut in unum coalita sessilia. Capsula 5-valvis, valvulis elasticè dissilientibus, placenta centrali, 5-angulari, angulis membranaceis suturas inter valvulares attingentibus et ideò capsula junior 5-locularis. Semina pendula, in quoque loculo plurima, exalbuminosa. Embryo rectus, radicula superà. — *Herbæ teneræ, foliis exstipulatis penninerviis.*

I. BALSAMINA. *Riv. irr. tetr. ic.*

Antheræ 5 biloculares; stigmata 5 distincta; capsula ovata; valvis ad maturitatem introrsum apice elasticè inflexis; cotyledones crassæ.

1. *B. hortensis* (Desp. dict. sc. nat. 3. p. 485.) pedicellis adgregatis, foliis lanceolatis, superioribus alternis, calcaribus flore brevioribus. ① in Indiâ orientali, in hortis culta. — *Impatiens Balsamina* L. Fl. fr. 4. p. 854. Black. t. 583. (v. v. c.)

II. IMPATIENS. *Riv. tetr. ic.*

Antheræ 5, nempe 3 biloculares, 2 antè petalum superius 1-loculares; stigmata 5 coalita; capsula prismatico-teretiuscula elongata, valvis à basi ad apicem extrorsum revolutis; cotyledones planiusculæ.

1. *I. NOLI-TANGERE* (L. sp. 1328.) pedunculis 3-4-floris folio brevioribus sub folio patulis, floribus pendulis, calcare apice recurvo, foliis ovatis grosse dentatis, geniculis caulinis tumentibus. ① in nemoribus passim. — Fl. fr. 4. p. 855. Barr. ic. t. 1197. (v. v.)

Ordo XXVII. OXALIDÆ. *DC. prod. 1. p. 689.*

Cal. 5-sepalus 5-partitusve persistens æqualis. Pet. 5 hypogyna æqualia, interdum infernè inter se cohærentia, per æstivationem spiraliter contorta. Stam. 10, filamentis basi sæpius monadelphis, 5 sepalis opposita exteriora breviora, 5 petalis opposita interiora longiora. Anth. 2-loculares non adnatæ. Ovarium liberum 5-loculare. Styli 5. Stigmata penicilliformia, capitata aut subbifida. Capsula membranacea 5-locul. 5-10-valvis, ad angulos longitudinaliter dehiscentem. Semina pauca, angulo loculamentorum centrali adfixa, juniora inclusa intra arillum carnosum demum ab apice elasticè dehiscentem et semen projicientem. Albumen cartilagineo-carnosum. Embryo inversus, radicula superà longâ. — *Herbæ foliis alternis rariùs oppositis.*

I. OXALIS. *L. gen. 582.*

Cal. 5-sepalus, sepalis liberis aut basi coalitis; pet 5; stam. 10, filamentis basi breviter monadelphis; capsula pentagona oblonga aut cylindræa.

1. *O. STRICTA* (L. sp. 624.) caule erecto folioso, pedunculis umbelliferis 3-6-floris folii circiter longitudine, foliolis obcordatis, petalis integris, stylis staminum interiorum longitudine. (1) in ruderatis, sepibus passim. — Fl. fr. 4. p. 856. — Fl. dan. t. 8-3. Flores flavi. (v. v.)
2. *O. CORNICULATA* (L. sp. 624.) caule decumbente ramoso radicante, pedunculis subumbellatis petiolo brevioribus, foliolis obcordatis, petalis emarginatis, stylis longitudine staminum interiorum. 2½ in cultis passim. — Fl. fr. 4. p. 856. Jacq. ox. t. 5. Flores flavi. (v. v.)  
*β. villosa*, undique hirsuta, pedunculis petiolo sublongioribus. — In arvis Corsicæ propè Bastia (cl. Soleirol.) — *O. villosa*, Bieb. taur. t. p. 355. DC. prod. 1. p. 692. (v. s.)
3. *O. ACROSZELLA* (Lin. sp. 620.) acaulis, radice dentatâ repente, foliolis obcordatis puberulis, scapo foliis longiore suprâ medium hibracteolato, petalis ovalibus obtusis, stylis stamina interiora æquantibus aut superantibus. 2½ in umbrosis. — Fl. fr. 4. p. 855. — Lob. ic. 2. p. 32. f. 1. (v. v.)

Ordo XXVIII. ZYGOPHYLLÆÆ. *Brown. gen. rem. p. 13.*  
*DC. prod. 1. p. 703.*

Calyx sepalis 5 distinctis aut vix basi subconnexis. Petala 5 sepalis alterna receptaculo inserta. Stam. 10 distincta, hypogyna, 5 sepalis, 5 petalis opposita. Ovarium unicum 5-locul. Styli 5 in unum coaliti, interdum apice summo subdistincti. Capsula carpellis 5 plus minus inter se et axi centrali adnatis constans, loculis ad angulum superius dehiscentibus, sæpius polyspermis aut 1-spermis, nec cocculiferis nec arilliferis. Semina albuminosa aut sæpius exalbuminosa, embryone recto, radicula superâ. — Herbæ fruticesve, foliis basi stipulaceis, in nostris oppositis.

I. TRIBULUS. *Tourn. inst. t. 141. L. gen. n. 532.*

Calyx 5-sepalus deciduus; petala 5 patentia; stamina 10; stylus 0, carpella 5 axi adnexa triangularia indehiscentia dura extus in spinas tubercula aut alas exserta, intus transverse plurilocularia, rarius 1-locularia; semina in loculis solitaria horizontalia exalbuminosa.

1. *T. TERRESTRIS* (L. sp. 554.) foliis 6-jugis, foliolis subæqualibus, pedicellis petiolo brevioribus, carpellis 4-cornutis. (1) in arvis et ad vias Galliarum australis; circâ Burdigalam ex Laterr. et in insulâ Nigri-Monasterii. — Fl. fr. 4. p. 731. — Barr. ic. t. 558. (v. v.)

Ordo XXIX. RUTACEÆ. *Juss. gen. 296. excl. sect. prima.*  
*DC. prod. 1. p. 709.*

Calyx 3-5-sepalus, sepalis basi plus minusve inter se coalitis et ideo dentatus fissus aut partitus. Petala rarissimè nulla, sæpiùs tot quot sepala, sæpè unguiculata, distincta. Discus carnosus-glandulosus ovarium cingens, è receptaculo ortus, extùs petala, superiùs stamina gerens. Stam. sæpiùs sepalorum numero dupla et tunc aut omnia fertilia aut alterna sessilia, rariùs sepalis tripla. Carpella tot quot sepala, interdum abortu pauciora, nunc distincta, nunc basi coalita, nunc omnino connata. Stylus è centro ovariorum ortus, unicus in stigmata tot quot ovaria divisus et sic stylis ex ovario quoque ortis et concretis compositus. Carpella in fructu sæpiùs distincta 1-locularia, dehiscentia, bivalvia, intùs cocculosa. Semina angulo interno adfixa inversa. Embryo rectus compressus, radicula superà. — Herbæ aut frutices, foliis alternis oppositisve exstipulatis.

I. RUTA. *Tourn. inst. t. 133. L. gen. n. 523.*

Cal. persistens 4-partitus rariùs 3-5-partitus; pet. totidem unguiculata subcochleata; stam. petalorum numero dupla; pori nectariferi ad basin ovarii tot quot stamina; ovarium stipite crasso brevi impositum; capsula subglobosa divisa in tot loculos quot sunt petala; albumen carnosum; embryo arcuatus.

1. *R. MONTANA* (Clus. hist. 2. p. 136.) foliis suprà-decompositis, lobis omnibus linearibus, petalis integris.  $\mathcal{L}$   $\mathcal{F}$  in siccis collinis lapideis Gallie australis et in Andegaviâ (Bast.). — Fl. fr. 4. p. 732. — Jacq. ic. rar. 1. t. 76. (v. v.)
2. *R. GRAVEOLENS* (L. sp. 548. excl. var.  $\alpha$  et  $\beta$ .) foliis suprà-decompositis, lobis oblongis, terminali obovato, petalis integris aut subdentatis.  $\mathcal{L}$   $\mathcal{F}$  in siccis Gallie australis et in hortis frequens. — Fl. fr. 4. p. 732. — Duham. arb. 2. t. 61. (v. v.)
3. *R. BRacteosa* (DC. prod. 1. p. 710.) foliis suprà-decompositis latitudine vix triplo longioribus, lobis oblongo-cuneatis subæqualibus, bracteis maximis subcordatis, petalis ciliatis.  $\mathcal{F}$  circà Niceam. (v. s.)
4. *R. ANGUSTIFOLIA* (Pers. ench. 1. p. 464.) foliis suprà-decompositis latitudine quadruplò longioribus, lobis oblongo-cuneatis subæqualibus, bracteis minimis ovatis, petalis ciliatis.  $\mathcal{L}$   $\mathcal{F}$  in apricis lapideis Gallie australis et in muris veteribus Aquarum Tarbellicarum. (Dax.) — *R. Chalepensis*.  $\mathcal{F}$  Fl. fr. 4. p. 733. — Moris. hist. s. 5. t. 35. n. 8. Folia angusta valdè glauca. (v. v.)
5. *R. Corsica* (DC. prod. 1. p. 710.) foliis suprà-decompositis, lobis obovatis subæqualibus, racemis subsimplicibus, pedicellis elongatis 1-floris rigidè divaricatis, petalis integris.  $\mathcal{L}$   $\mathcal{F}$  in Corsicæ montibus. — Bocc. mus. p. 70. t. 79. ic. (v. s.)

II. DICTAMNUS. *L. gen. n. 522.*

Cal. deciduus 5-partitus; pet. 5 subinæqualia; stam. 10 declinata filam. apice glanduloso-tuberculosis; stigma simplex; capsula carpellis 5 interne connexis compressis 2-spermis.

1. *D. FRAZINELLA* (Pers. ench. 1. p. 464.)  $\mathcal{L}$  in sylvaticis Gallie australis circà Narbonam, etc., etc. in hortis culta. — *D. albus* L. sp. 548. Fl. fr. 4. p. 734. — Clus. hist. 1. p. 99. f. 2. — Variat floribus purpureis et albis. (v. v. c.)

Subclassis II. CALYCIFLORÆ.

Calyx gamosepalus, sepalis nempè basi plus minùs inter se coalitis. Torus (1) calyci intùs ad basin plus minùs adnatus. Petala et stamina è parte tori calyci adnatâ exserta et ideò orta è calyce vulgò dicta. Petala inter se libera aut coalita. Ovarium liberum aut calyci adnatum.

---

Ordo XXX. CELASTRINEÆ. *R. Brown. gen. rem. p. 22. DC. prod. 2. p. 2.*

Calycis sepala 4-5 imâ basi inter se coalita, ab ovario libera, æstivatione imbricata. Petala tot quot sepala, iis alterna, rarissimè nulla. Stamina tot quot sepala, petalis alterna, nempè sepalis opposita, insertione ambigüe perigyna. Antheræ biloculares. Ovarium liberum, disco subcarnoso cinctum, 2-3-4-loculare, loculis mono-aut-polyspermis, ovulis erectis rarò pendulis. Stylus 1 aut 0. Stigma 2-4-fidum. Pericarpium capsulare baccatum drupaceum samaroïdeumve, formâ varium, abortu loculorum sæpè deformatum. Semina in pluribus, præsertim capsularibus, arillatum. Albumen 0 aut carnosum. Embryo rectus, axilis. — Frutices vel arbores, foliis sæpè stipulatis, alternis aut oppositis.

Trib. I. STAPHYLEACEÆ. DC. l. c.

Semina ossea, ad hylum truncata, exarillata. Albumen 0 aut tenue. Cotyledones crassæ. — Folia composita.

I. STAPHYLEA. *L. gen. n. 374.*

Calyx 5-partitus, disco urceolato basi tectus, lobis oblongis concavis coloratis; petala 5 sepalis alterna; stamina 5 petalis alterna; ovarium 2-3-lobum; capsula 2-3-locularis, loculis membranaceis intùs dehiscentibus oligospermis.

1. *S. PINNATA* (*L. sp. 386.*) foliis pinnatis, petiolo eglanduloso, foliolis 5-7 oblongo-lanceolatis glaberrimis serratis, floribus racemosis, capsulis membranaceis vesicariis. ♂ in sylvis Alsatiæ, in hortis colina. — *Fl. fr. 4. p. 620.* — *Cam. epit. 171. ic. (v. v. c.)*

Trib. II. EVONYMEÆ. DC. l. c. p. 3.

Semina arillata, ad hylum non truncata. Embryo erectus in axi albuminis carnosi. Cotyledones foliaceæ. — Folia simplicia.

---

(1) Torus est pedunculi dilatatio è quâ originem ducunt organa floris mascula, nempè stamina seu antherifera, sen sterilia, seu in petala conversa.

II. EVONYMUS. *Tourn. inst. t. 388. L. gen. 271.*

Calyx 4-6-lobus planus, basi disco peltato tectus; petala 4-6 patentia disco inserta, stamina 4-6, glandulis suprâ discum prominulis inserta, petalis alterna; stylus 1; capsula 3-5-localaris, 3-5-angularis, valvis medio septiferis; seminibus in loculis 1-4; embryo viridis.

1. E. EUROPEUS (L. sp. 286. var. *a.* excl. *Clus. syn.*) ramis lævibus, foliis lanceolato-ovatis tenuissimè serratis, pedunculis subtrifloris, petalis oblongis subacutis, lobis capsularum obtusis. ♂ in sepibus. — Fl. fr. 4. p. 620. *Bull. herb. t. 135. (v. v.)*
2. E. LATIFOLIUS (Bauh. pin. 248.) ramis lævibus, foliis lato-ovatis denticulatis, pedunculis trichotomis multifloris, petalis ovalibus, capsularum lobis acutè angulatis alæformibus. ♂ in nemoribus Delph., Galloprovinciæ. — Fl. fr. 4. p. 621. — *Clus. hist. 1. p. 56. f. 2. (v. s.)*

Trib. III. *AQUIFOLIACEÆ*. DC. prod. 2. p. 11.

Petala basi latiora, interdum inter se coalita. Fructus indehiscens. Seminis embryo rectus in axi albuminis carnosus. — Folia simplicia.

III. ILEX. *L. gen. n. 172.*

Calyx 4-5-dentatus persistens, petala 4-5, sepalis alterna, nunc libera, nunc basi in corollam rotatam subcoalita; stamina 4-5, petalis alterna, hypogyna; ovarium sessile, 4-loculare, stigmatibus subsessilibus 4-5 nunc distinctis nunc in 1 coalitis coronatum; bacca 4-5-pyrena, nucleis oblongis apice umbilicatis 1-spermis.

1. I. AQUIFOLIUM (L. sp. 181.) foliis ovatis acutis nitidis undulatis dentibus et apice spinosis, pedunculis axillaribus brevibus multifloris, floribus subumbellatis. ♂ ad sepes et in sylvis. — Fl. fr. 4. p. 621. *eng. bot. t. 496. (v. v.)*

Ordo XXXI. RHAMNEÆ. *R. Brown. gen. rem. p. 22. DC. prod. 2. p. 19.*

Calycis tubus ovario adhærens, lobi æstivatione valvati, numero definiti 4-5. Petala totidem lobis calycinis alterna, sæpè squamiformia limbo concavo. Stamina tot quot petala iis opposita. Antheræ 2-loculares. Ovarium nunc totum calyci adnatum, nunc imâ basi aut ad medium adhærens, 2-4-loculare, loculis 1-ovulatis. Stylus 1. Stigmata 2-4. Pericarpium sæpiùs indehiscens baccatum drupaceum samaroideumve, rariùs capsulare; semina erecta, arillo destituta. Albumen nullum aut sæpiùs carnosum. Embryo rectus, axilis, radiculâ inferâ. — Frutices aut arbusculæ, foliis simplicibus alternis rarissimè oppositis, sæpè stipulaceis.



I. ZIZYPHUS. *Tourn. inst. t. 403. Gærtn. fruct. 1. t. 43.*

Calyx patens, 5-fidus, medio tubi post anthesin circumscissus, basi sub fructu persistente et cum eo subadhærente; discus glandulosus calyci adnatus petala 5 gerens; stamina 5 antè petala sita; styli 2-3 simplices; drupa ovoidea nucleo evalvi biloculari 2-spermo (rarius 1-3-loc. 1-3-spermo); semina suborbiculata, compressa, sulco destituta, ferè exalbuminosa.

1. *Z. VULGARIS* (Lam. ill. t. 185. f. 1.) foliis ovatis retusis denticulatis ramulisque glabris, aculeis nullis aut geminis, altero recurvo, drupis ovato-oblongis. 5 ex Syriâ orta, in regionibus mediterraneis nunc subspontanea. — Fl. fr. 4. p. 626. — Cam. epit. 167. ic. (v. v.)

II. PALIURUS. *Tourn. inst. t. 387. DC. prod. 2. p. 22.*

Flores Zizyphi, sed styli 3, fructus siccus indehiscens orbicularis alâ latâ membranaceâ cinctus, intus 3-locul. 3-spermus; semina ovata sulco destituta, albumine nullo aut parcissimo.

1. *P. ACULEATUS* (Lam. ill. t. 210.) ramulis pubescentibus, foliis basi ovatis acutis 3-nerviis, fructibus alâ erenulata. 5 in Galliæ mediterraneæ sterilibus. — Fl. fr. 4. p. 626. — Cam. epit. 83. ic. (v. v.)

III. RHAMNUS. *Lam. dict. 4. p. 461. ill. t. 128. DC. l. c. p. 23.*

Calyx 4-5-fidus, post anthesin medio sæpè circumscissus, basi sub fructu persistente et cum eo adhærente; petala lobis calycis alterna aut 0; stamina antè petala inserta; stylus 2-4-fidus; fructus subexsuccus aut baccatus 2-4-locul., loculis coccisve in exsiccis separabilibus 1-spermis (rarissimè 2-spermis), rimâ longit. intus dehiscen-  
tibus; semina oblonga, ad latus exterius sulco profundo basi latiore notata.

Sect. I. ALATERNUS. *Tourn. inst. t. 366.*

Flores 5-fidi 5-andri, fructus baccatus; seminum cicatricula non exserta.

1. *R. ALATERNUS* (L. sp. 281.) foliis ovatis ellipticis lanceolatisve coriaceis glaberrimis serratis, floribus dioicis breviter racemosis. 5 in provinciis australibus et circâ Nannetes. — Fl. fr. 4. p. 624. Duham. ed. nov. 3. p. 42. t. 14.

α. *vulgaris* (DC.) foliis ovatis serratis. (v. v.)

β. *angustifolius* (DC.) foliis lanceolatis sublinearibusve. (v. v.)

Sect. II. CERVISPINA. *Dill. nov. gen. t. 8. DC. l. c. p. 24.*

Flores 4-fidi 4-andri; fructus subbaccatus 4-spermus, seminum cicatricula non exserta.

\* *Ramis apice spinosis.*

2. *R. CATHARTICUS* (L. sp. 280.) erectus, foliis ovatis dentatis, floribus fasciculatis polygamo-dioicis, baccis 4-spermis subglobosis. 5 in sylvis, sepibus et incultis passim. — Fl. fr. 4. p. 622. eng. bot. t. 1629. (v. v.)
3. *R. INFECTORIUS* (L. mant. 49.) procumbens diffususve, foliis ovato-lanceolatis serrulatis glabresculis, floribus dioicis, utriusque sexus petaliferis. 5 in Galliæ australis asperis. — Fl. fr. 4. p. 623. — Clus. hist. 1. p. 111. ic. (v. v.)
4. *R. SAXATILIS* (L. sp. 1671.) procumbens suberectusve, foliis ovato-lanceolatis serrulatis glabresculis, floribus dioicis, femineis apetalis. 5 in montibus Delphinatûs ex Villars. — Fl. fr. 4. p. 623. — Clus. hist. 1. p. 112. f. 1. An prioris var? (v. s. ex Italiâ boreali.)

5. *R. PUBESCENS* (Poir. dict. 4. p. 464.) diffusus, foliis ovatis integerrimis coriaceis pubescentibus. § inter Caunas et Carcassonam ex Dom. Fourmeault. — *R. oleoides* Fl. fr. 4. p. 623. non L. An *R. oleoidis* var? (v. s. c.)

\* \* *Ramis apice non spinosis.*

6. *R. RUMILUS* (L. mant. 49.) procumbens ramosissimus, foliis ovatis serratis glabris, floribus hermaphroditis. § in Alpibus Galloprovinciæ, Delph., Pyren. — Fl. fr. 4. p. 625. *R. rupestris* Scop. carn. ed. 2. t. 5 (v. v. DC.)

β. *rupestris* (DC.) foliis integerrimis. In Delphinatū.

7. *R. ALPINUS* (L. sp. 280.) erecto-tortuosus, foliis ovali-lanceolatis crenato-serratis glabris nervulis lateralibus plurimis lineatis, floribus dioicis, femineorum stigmate 4-fido. § in Pyrenæis, Alpibus Delph., Gallop., Jurasso. — Fl. fr. 4. p. 624. Hall. helv. t. 40. (v. v.)

β. *Corsicus*, foliis rotundatis obtusis pubescentibus, petiolis ramisque superioribus velutinis. In Corsicā reperiit Ph. Thomas. (v. s.)

Sect. III. FRANGULA. *Tourn. inst. t. 388. DC. l. c. p. 26.*

Flores 5-fidi 5-andri, seminum cicatricula exserta.

8. *R. FRANGULA* (L. sp. 280.) foliis ovalibus integerrimis nervis lateralibus 10-12 lineatis calycibusque glabris, floribus hermaphroditis. § in dumetis, sylvalis sepiusque. — Fl. fr. 4. p. 624. Lam. ill. t. 128. f. 1. Baccæ 2-4-spermæ. (v. v.)

## Ordo XXXII. TEREBINTHACEÆ. *Juss. gen. 368.*

Flores hermaphroditi polygami aut dioici. Calyx sepalis 3-5 basi plus minus coalitis æstivatione imbricatis constans, ovario rarissimè adhærens. Petala rariùs 0, sæpissimè sepalorum numerum æquantia et iis alterna distincta, in paucissimis basi coalita, æstivatione imbricatâ aut valvatâ. Stamina cum petalis ex imo calyce aut è disco calycino, aut rariùs è toro ovarium ambiente orta, nunc petalis numero æqualia iisque alterna, nunc petalis numero dupla. Carpella nunc plurima distincta monostyla, nunc plurima ovariis coalita, nonnulla ex iis in utroque casu sæpè abortiva, et indè carpella in plurimis solitaria 1-ocularia evadunt, sed tunc stylorum stigmatumve numerus abortionem sæpè indicat. Fructus capsularis aut drupaceus. Semina pauca sæpiùs solitaria sæpiùs exalbuminosa. Embryo rectus curvatus arcuatus replicatusve, radiculâ sæpiùs superâ. — Arbores fruticesve, foliis alternis non stipulaceis, floribus parvis.

Trib. I. *ANACARDIÆ*. DC. prod. 2. p. 62.

Petala et stamina disco calycino aut calyci inserta. Ovarium 1, 1-loc. 1-ovulatum. Semen funiculo è fundo loculi orto apice inflexo sustentum, exalbuminosum. Cotyledones crassæ super radiculam replicatæ.

## TEREBINTHACEÆ.

113

### I. PISTACIA. *L. gen.* 1108.

Flores dioici spetali. *Masc.* Racemus amentaceus squamis 1-floris; calyx 5-fidus; stam. 5, antheris subsessilibus 4-gonis. *Fem.* Racemus laxior; calyx 3-4-fidus; ovarium 1-3-loculare; stigmata 3 crassiuscula; drupa sicca ovata nuce subossæ sæpius 1-locularis 1-sperma, semina in loculis solitaria. — Arbores foliis pinnatis.

1. *P. vera* (L. sp. 1454.) foliis impari-pinnatis, foliolis ovatis basi subattenuatis apice submacronatis quinis ternis aut solitariis. ♀ ex Syria orta, nunc in provinciis mediterraneis subspontanea. — Fl. fr. 4. p. 616. — Blackw. t. 461. — Boec. mus. 2. t. 92. (v. v.)
  2. *P. terebinthus* (L. sp. 1455.) foliis impari-pinnatis, foliis subsæptenis ovato-lanceolatis basi rotundatis acutis mucronatis. ♀ in provinciis austriacioribus. — Fl. fr. 4. p. 616. — Blackw. t. 478. (v. v.)
  3. *P. lentiscus* (L. sp. 1455.) foliis abrupte pinnatis, foliolis octonis lanceolatis, petiolo alato. ♀ in saxosis maritimis Nicæ, Galloprovinciæ, Occitanis. — Fl. fr. 4. p. 617. Blackw. t. 195. (v. v.)
- γ. *angustifolia* (DC.) foliolis sublinearibus.

### Trib. II. *SUMACHINEÆ*. DC. prod. 1. p. 66.

Petala et stamina disco calycino aut calyci inserta. Ovarium 1, 1-loc. 1-ovulatum. Semen funiculo è basi orto sustentum, pendulum, exalbuminosum. Cotyledones foliaceæ. Radicula superæuræ commissuram inflexa.

### II. RHUS. *L. gen.* n. 369.

Calyx parvus 5-partitus persistens; petala 5 ovata patentia; stamina 5, ovarium 1 subglobosum, styli breves 3 aut stigmata 3 sessilia; drupa ferè exsucca 1-locularis, nucleo osseo (abortu?) 1-spermo et interdum 2-3-spermo. — Frutices.

1. *R. cotinus* (L. sp. 383.) foliis obovatis simplicibus, floribus laxè paniculatis, hermaphroditis; drupâ semicordatâ venosâ glabrâ. ♀ in collibus Gallicis austro-orientalis. — Fl. fr. 4. p. 614. Jacq. aust. t. 238. (v. v.)
2. *R. cornaria* (L. sp. 379.) foliis impari-pinnatis 5-7-jugis villosis, floribus paniculatis, drupâ ovato-rotundâ villosâ. ♀ in aridis provinciis austriaciorum et aliquandò occidentalis. — Fl. fr. 4. p. 615. — Blackw. t. 486. (v. v.)
3. *R. radicans* (L. sp. 381.) foliis impari-pinnatis 3-foliolatis, foliolis glabris integris, floribus paniculatis, drupâ ovato-rotundâ. ♀ ex America boreali, nunc ex el. Le Prévost in sylvis paludosis circâ Montauré propè Louviers frequens. — bot. mag. t. 1806. Taeta venenosa. (v. v. v.)

### Trib. III. *PTELEACEÆ*. Kunth. gen. tereb. p. 22.

Petala 3-5 hypogyna sessilia, per æstivationem imbricata, rariùs valvata. Stamina petalorum numero æqualia. Torus disciformis. Ovarium 2-5-loculare. Fructus indehiscens 1-5-locul., loculis 1-2-spermis. Semina sæpè crustacea, albumine carnoso, cotyledonibus planiusculis, radicula superâ.

### III. CNEORUM. *L. gen.* n. 48.

Flores hermaphroditi; calyx 3-4-dentatus parvus persistens; petala 3-4-æqualia; torus subglobosus; stamina 3-4; stigmata 3-4; drupæ baccatæ 3-4 ad axem conjunctæ, putamine 2-loculari, loculis 1-spermis. — Suffrutices.

1. *C. ruscocoon* (L. sp. 49.) glabrum, floribus axillaribus, pedicellis à folio florali liberis. 3 in Galliæ australis glareosis rupestribus. — Fl. fr. 4. p. 615. — Cam. epit. 973. ic. (v. v.)

Ordo XXXIII. LEGUMINOSÆ. *Juss. gen.* 345. *DC. prod.* 2. p. 93.

**Calyx** constans sepalis 5 (rarissimè 4) basi plus minùs coalitis et ideò 5-dentatus 5-fidus 5-partitusve; sepala sæpiùs inæqualia nunc subæqualiter coalita, nunc concreta in labia 2, superius constans sepalis 2 apice liberis coalitisve, inferius sepalis 3, sæpiùs apice distinctis. **Petala** 5 aut abortu 4, 3, 2, 1 aut 0, sæpiùs inæqualia, imò calyci sæpiùs, toro rariùs inserta, per æstivationem sæpiùs imbricata, rariùs valvata, fere semper libera, interdum in corollam gamopetalam concreta. **Stamina** cum petalis inserta, petalorum numero dupla, rariùs aut tripla quadrupla aut pauciora; nunc omninò libera, nunc filamentis variè connexis aut monadelphis vaginà integrâ seu supernè fissâ, aut diadelphis 9 et 1, seu 5 et 5, nunc rarissimè triadelphe. **Antheræ** biloculares. **Carpellum** sæpiùs 1, cæteris nempe abortivis. **Ovarium** sessile aut stipitatum, liberum. **Stylus** 1 filiformis è suturâ superiore ortus. **Stigma** terminale aut laterale. **Legumen** bivalve membranaceum coriaceum, dehiscens aut indehiscens, 1-loculare, aut suturâ alterâ introflexâ longitudinaliter biloculare, aut isthmis articulationibusve transversè multiloculare. **Semina** 2 aut plura aut solitaria, suturæ superiori adfixa, alternatim cuique valvulæ inserta, sæpiùs ovalia aut reniformia; funiculus varius, rarè in arillum expansus. **Testa** lævis. **Endoplevra** sæpè tumida albumen mentiens. **Embryo** nunc rectus, nunc homotropus seu pleurorhizæus, radiculâ nempe inflexâ, super lorum commissuram pronâ, sed in utroque casu ad hytûm directâ. **Cotyledones** foliaceæ aut carnosæ; aliæ exsertæ, aliæ, intrâ spermodermium sub terrâ inclusæ. — **Arbores** frutices aut herbæ, foliis sæpissimè alternis, bistipulatis, simplicibus aut variè compositis, petiolatis.

Subordo I. PAPILIONACEÆ. *L. ord. nat.* 32. *DC. prod.* 2. p. 94.

**Embryo** homotropus pleurorhizæus nempe radiculâ super lorum commissuram pronâ. **Sepala** per æstivationem imbricata aut subvalvata. **Petala** in corollam papilionaceam disposita, semper ideò irregularia, stamina imò calyci inserta seu perigyna.

Trib. I. *SOPHOREÆ*. Spreng. DC. prod. 2. p. 94.

Stamina libera. Legumen continuum nec articulatum. Cotyledones planæ foliaceæ.

I. *ANAGYRIS*. Tourn. inst. t. 415. L. gen. n. 509.

Calyx 5-dentatus subbilabiatus; alæ vexillo longiores et carinâ dipetalâ breviores; legumen breviter stipitatum compressum polyspermum bivalve isthmis irregulariter subinterruptum.

1. *A. VETIDA* (L. sp. 534.) foliis trifoliolatis, foliolis integerrimis lanceolatis acutis, leguminibus acuminatis. § in collibus saxosis submontanis Gallias mediterraneæ circâ Niceam, Massiliam, Arelatem, Mompelium. — Fl. fr. 4. p. 491. — Clus. hist. 1. p. 93. ic. (v. s.)

Trib. II. *LOTEÆ*. DC. prod. 2. p. 115. leg. mem. vi.

Embryo homotropus. Stamina aut monadelphæ aut diadelphæ (9 et 1). Legumen continuum 1-loculare, aut rariùs suturâ alterâ introflexâ 2-loculare. Cotyledones planiusculæ per germinationem in folium stomatibus donatum conversæ.

Subtribus 1. *GENISTEÆ*. Bronn. DC. 1. c.

Legumen 1-loculare. Stamina sæpissimè monadelphæ. Folia simplicia aut palmatim trifoliolata rariùs pinnata. Caules sæpissimè fruticosi.

II. *ULEX*. L. gen. n. 881.

Calyx hibracteolatus, bipartitus, labio altero 3, altero 2-dentato; stam. omnia connexa; legumen ovali-oblongum turgidum vix calyce longius pluri-ovulatum oligospermum. — Frutices ramosissimi.

1. *U. EUROPEUS* (L. sp. 1045. var. æ.) erectus, foliis lanceolato-linearibus ramulisque subvillosis, bracteis ovatis laxis, calycis pubescentis dentibus obsoletis conniventibus. § in sterilibus et sepibus præsertim provinciarum occidentaliûm borealiûmque. — Fl. fr. 4. p. 492. eng. bot. t. 742. (v. v.)

2. *U. NAUUS* (Smith. fl. brit. 757. eng. bot. t. 743.) decumbens, foliis linearibus ramulisque glabris, bracteis minutis adpressis, calycis glabriusculi dentibus lanceolatis distantibus. § in sterilibus et ericetis Gallias mediæ et occidentalis. — Fl. fr. 4. p. 492. (v. v.)

3. *U. PROVINCIALIS* (Lois. not. 105. t. 6. f. 2.) erectus, foliis lanceolato-linearibus ramulisque glabriusculis, bracteis minutis adpressis, calycis subpubescentis corollamque vix superantis dentibus lanceolatis distantibus. § in sterilibus Galloprovinc. Occitaniz, Andegaviz. — Fl. fr. 5. p. 546. (v. v.)

III. *SPARTIUM*. DC. prod. 2. p. 145.

Cal. membranaceus spathaceus supernè fissus apice 5-dentatus subbilabiatus; corollæ vexillum subrotundum complicatum, carinâ acuminatâ et petalis parum adglutinatâ bipartibilis; stam. monadelphæ; legumen plano-compressum polyspermum eglandulosum.

1. *S. JUNCUM* (L. sp. 995.) glabrum, foliis paucis lanceolatis, floribus in racemos terminales dispositis. § in asperis apriciis Delphir. super. ex Villars, Galloprov., Occitaniz, circâ Montem-Albanum, Tolosam, Tarbam, Agiumum. — Gen. juncea. Fl. fr. 4. p. 495. (v. s.)

IV. *GENISTA*. Lam. dict. 2. p. 616. ill. t. 619. DC. leg. mem. vi. p. 204.

Calyx 2-labiatus, labio superiore 2-partito, inferiore 3-dentato; aut 5-lobus, lobis 3 inferioribus ad apicem serè coalitis; vexillum oblongo-ovale; carina oblonga recta genitalia non omninò continens; stamina

monadelphæ, legumen plano compressum aut rariùs subturgidum, polypermum, rariùs oligospermum, eglandulosum. — Frutices.

\* *Inermes, foliis omnibus aut plerisque trifoliolatis.*

1. *G. CANDICANS* (Lin. amæn. 4. p. 284.) foliis petiolatis 3-foliolatis, foliolis obovatis pube adpressâ pubescentibus, ramis angulatis, capitulis terminalibus paucifloris, leguminibus hirsuto-villosis. § in collibus Corsicæ, Nicæ, insularum Stæchadum, Monopelii, agri Ruscinonensis. — *Cytisus candicans* L. sp. 740. Fl. fr. 4. p. 504. (v. v. DC.)
2. *G. LINIFOLIA* (L. sp. 405.) foliis sessilibus 3-foliolatis, foliolis linearibus subtus sericeis margine demùm revolutis, racemis terminalibus confertis, leguminibus hirsutis, ramis teretibus sulcatis. § in insulis Stæchadum. — *Cytisus linifolius* Lam. DC. Fl. fr. 4. p. 504. — Desf. atl. 2. t. 181. (v. s.)

\*\* *Spinosæ, foliis aut omnibus aut nonnullis trifoliolatis.*

3. *G. HORRIDA* (DC. fl. fr. 4. p. 500.) foliis 3-foliolatis petiolatis oppositis, foliolis linearibus complicatis subsericeis, ramis angulatis spinosis confertis oppositis, floribus subterminalibus paucis, calyce pubescente. § in aridis Pyren. et agri Lugdunensis. — *Spartium horridum* Valh. *G. erinacea* Gilib. bot. prat. 2. p. 239. ic. (v. v. DC.)
4. *G. LOBELII* (DC. fl. fr. 4. p. 499.) foliis paucis, imis sessilibus 3-foliolatis, cæteris ferè omnibus simplicibus sparsis lineari-oblongis subsericeis, ramis confertis spinosis striatis hinc inde tuberculatis, floribus secus ramulos paucis solitariis subracemosis pedicellatis, calyce pubescente. § in aridis Corsicæ, Galloprov. — Lob. adv. p. 409. ic. (v. v. DC.)
5. *G. SALEMAINI* (DC. prod. 2. p. 147.) foliis sessilibus 3-foliolatis simplicibusve oblongis obtusis adpressè pubescentibus, ramis demùm striatis laxis spinosis, floribus secus ramos subracemosi geminis pedicellatis adpressè sericeo-pubescentibus, calycis lobis 3 infer. superiores subæquantibus angustioribus. § in petrosis Corsicæ propè Corte. DC. mém. leg. v. p. 211. — A præcedente vix diversa. (v. s.)

\*\*\* *Spinosæ, foliis æntibus simplicibus.*

6. *G. CORSICA* (DC. fl. fr. 5. p. 548.) tota glaberrima, foliis lineari-lanceolatis, stipulis subspinescentibus, spinis simplicibus rigidis, floribus axillaribus solitariis pedicellatis, carinâ vexilli glabri longitudine, legumine 4-8-spermo glaberrimo. § in Corsicâ circâ Bastia, Bonifacio, etc. (v. s.)  
*l. pubescens* (DC.) ramulis foliisque junioribus adpressè pubescentibus. § in Corsicâ propè Calvi. (v. s.)
7. *G. SCORPIUS* (DC. fl. fr. 4. p. 498.) foliis paucissimis oblongis subsericeis, spinis ramosis patentibus striatis glabris, floribus fasciculatis subracemosis breviter pedicellatis glabris, carinâ vexilli longitudine, legumine 2-4-spermo. § in sterilibus montosis Delphin. austr., Galloprov., Occit., Cebenn., Aquitanicæ. — *Spartium Scorpius* L. sp. 995. — Clus. hist. 1. p. 106. f. 1. (v. v.)
8. *G. HISPANICA* (L. sp. 999.) foliis lanceolatis villosis, spinis ramosis rigidis, ramis floriferis inermibus, racemo terminali subcapitato, carinâ villosâ vexilli glabri longitudine, legumine ovali 2-4-spermo subbispido, maturo glabrinaculo. § in aridis montosis provinz. australium à Nicæ ad Albi. — Fl. fr. 4. p. 499. — J. Bauh. 1. p. 2. p. 400. f. 1. (v. v.)
9. *G. ANGOLICA* (L. sp. 999.) tota glabra, foliis ovato-lanceolatis, spinis simplicibus, ramis floriferis inermibus, racemis paucifloris terminalibus, carinâ vexillo et alia longiore, legumine ovato-cylindraceo polyspermo. § in sylvis turfosis et ericetis Gallicæ occidentalis, mediæ et australis. — Fl. fr. 4. p. 498. — Lob. ic. 2. p. 93. f. 2. (v. v.)
10. *G. GERMANICA* (L. sp. 995.) foliis lanceolatis hirsutis, spinis simplicibus aut ramosis, ramis floriferis inermibus, racemis terminalibus, floribus subvillosis, carinâ vexillo et alia longiore, legumine ovato hirsutato 2-4-spermo.

♂ in sylvis et ericetis Gallie orientalis, medise et australis. — Fl. fr. 4. p. 499.  
— Fuchs. hist. 220. ic. (v. v.)

\*\*\*\* *Inermes, foliis omnibus simplicibus.*

11. *G. PURGANS* (L. sp. 999.) erecta ramosissima, ramis teretibus striatis, foliis paucissimis lanceolatis subsessilibus subsericeis, floribus axillaribus solitariis subpedicellatis, petalis inter se æqualibus glabris, legumine juniore adpressè pubescente. ♂ in siccis sterilibus montosis agri Lugdun., Arvern., Cebenn., provinc. mediter., Pyr. or. et Aurelie ex Dub. — Fl. fr. 4. p. 494. Bull. herb. 115. (v. v.)

12. *G. CINEREA* (DC. fl. fr. 4. p. 494.) erecta ramosissima, ramis striatis, foliis lanceolatis adpressè pubescentibus, floribus secus ramos subsessilibus solitariis, petalis sericeis longitudine subæqualibus, leguminibus adpressè villosis 4-6-spermis. ♂ in collibus aridis montosisque à Nicæa ad Pyr. orient. secus limites olearum. — Fl. fr. 4. p. 494. (v. v.)

13. *G. HUMIFUSA* (L. sp. 998? Vill. dauph. 3. p. 421. t. 44.) procumbens, ramis tortuosis tuberculosis apice spinoscenti-rigidis foliisque lineari-lanceolatis piloso-hirsutis, floribus axillaribus solitariis subsessilibus, petalis sericeis inter se subæqualibus, lobis calycinis ovatis subacutis. ♂ in Delphinatùs montosis propè Gap. — Fl. fr. 4. p. 496. (v. v.)

14. *G. TINCORIA* (L. sp. 998.) radice repente, caulibus suberectis, ramis teretibus striatis erectis, foliis lanceolatis glabriusculis, floribus apicato-racemosis, leguminibusque glabris. ♂ frequens in pascuis et sylvulis. — Fl. fr. 4. p. 495. — Fuchs. hist. 809. ic. (v. v.)

♂. *latifolia* (DC.) foliis lato-lanceolatis. In Arvernise Monte-Aureo. (v. v. DC.)

15. *G. SAGITTALIS* (L. sp. 998.) caule prostrato, ramis herbaceis ascendentibus ancipitibus membranaceis subarticulatis, foliis ovato-lanceolatis, floribus in apicem ovatam terminalem aphyllam digestis, corollis glabris cum carinâ ad dorsum lineâ villosâ. 2♂ in pratis pascuisque submontosis. — Fl. fr. 4. p. 497. — J. Bauh. 1. p. 2. p. 393. f. 3. (v. v.)

♂. *minor* (DC.) fruticulosa minor, ramis ad apicem foliisque adpressè pubescenti-sericeis. In siccioribus Gallie australis. (v. v.)

16. *G. PROSTRATA* (Lam. dict. 2. p. 618.) caulibus diffuso-prostratis, ramis angulatis striatis villosulis, foliis obovatis oblongis subtus subvillosis, floribus axillaribus longè pedicellatis erectis, corollis glabris, leguminibus villosis 3-4-spermis. ♂ in asperis Burgundiæ, Jurassi et ad urbem Velaunorum (le Puy en Velay). — Fl. fr. 4. p. 496. G. Halleri Reyn. mem. 1. p. 211. ic. (v. v.)

17. *G. PILOSA* (L. sp. 999.) caulibus procumbentibus striatis ramosis tuberculatis, foliis lanceolatis complicatis subtus adpressè sericeis, floribus axillaribus breviter pedicellatis, corollis sericeis, leguminibus pubescentibus polyspermis. ♂ in asperis apricis submontosis passim. — Fl. fr. 4. p. 495. — Cus. hist. 1. p. 103. f. 2. (v. v.)

V. *CYTISUS*. DC. fl. fr. 4. p. 501. leg. mem. vi. — *Cytisus* et *Spartii* sp. L. Lam. etc.

Calyx bilabiatus, labio superiore sæpius integro, inferiore subtridentato; vexillum ovatum amplum; carina obtusissima genitalia includens; stamina monadelpha; legumen plano-compressum polyspermum eglandulosum. — Frutices.

Sect. 1. *LABURNUM*. DC. prod. 2. p. 153.

Calyx campanulatus; legumina polysperma ad suturam superiorem non dilatata. — Rami inermes foliosi.

1. *C. LABURNUM* (L. sp. 1041.) ramis teretibus incanescuntibus, foliis petiolatis, foliolis ovato-lanceolatis subtus pubescentibus, ramis pendulis ampli-

eibus pedicellis calycibusque adpressissime pubescentibus, leguminibus linearibus polyspermis adpressè pubescentibus.  $\frac{3}{5}$  in lapidosis demissiorum Alpium, Jurassi, collium Bressæ, Bengesæ, Burgundiæ, Pyrenæorum ex Dum. Cours. — Fl. fr. 4. p. 501. excl. var.  $\beta$ . — J. Bauh. hist. 1. p. 2. p. 361. ic. (v. v.)

2. *C. ALPINUS* (Mill. dict. n. 2.) glabra, ramis teretibus, foliis petiolatis, foliolis ovato-lanceolatis basi rotundatis, racemis pendulis, pedicellis calyceque hirsuto-puberulis, leguminibus glabris oligospermis marginatis.  $\frac{3}{5}$  in nemorosis montibus Jurassi australis. — Waldst. et Kit. hung. 3. p. 288. t. 260. *C. laburnum*  $\beta$ . Fl. fr. 1. c. (v. v. D.)
3. *C. sessilifolius* (L. sp. 1041.) glaberrimus, ramis teretibus, foliis floralibus subsessilibus, foliolis ovatis, racemis terminalibus erectis abbreviatis, bractea triphyllâ sub calyce.  $\frac{3}{5}$  in asperis montosis Delph., Niceæ, Gallop., Occit., agri Ligericini et Ruscimonensis. — Fl. fr. 4. p. 502. — J. Bauh. hist. 1. p. 2. p. 373. f. 2. (v. v.)
4. *C. triflorus* (L'Her. stirp. 184.) hirsuta, ramis teretibus, foliis petiolatis, foliolis obovato-ellipticis, floribus axillaribus pedicellatis ternis apice ramorum subracemosis.  $\frac{3}{5}$  in Corsicâ, circâ Narbonam, Olbias, Antipolim. — Fl. fr. 4. p. 505. — Clus. hist. 1. p. 94. f. 3. (v. e.)
5. *C. scoparius* (Link. enum. 2. p. 241.) ramis angulatis glabratiss, foliis trifoliolatis petiolatis, summis simplicibus foliolisque oblongis, floribus axillaribus pedicellatis solitariis, leguminibus ad margines pilosis.  $\frac{3}{5}$  in arenosis et sylvaticis Galliæ præsertim occidentalis. — Genista scoparia. Fl. fr. 4. p. 497. non Vill. — Duham. arb. t. 84. (v. v.)

#### Sect. II. CALYCOTOME. Link. DC. l. c. p. 154.

Calyx campanulatus subbilabatus dein circumscissus truncatus; legumina ad suturam superiorem incrassata. — Rami spinosi.

6. *C. spinosus* (Lam. dict. 2. p. 247.) ramis angulatis spinosis, foliis trifoliolatis, foliolis obovato-oblongis, leguminibus glaberrimis.  $\frac{3}{5}$  in collibus et asperis Corsicæ et Galliæ merid. à Perpignano ad Niceam passim. — Fl. fr. 4. p. 503. — Lob. ic. 2. t. 95. (v. v. DC.)
7. *C. lanigerus* (DC. fl. fr. 4. p. 504.) ramis striatis spinosis, foliis 3-foliolatis, foliolis obovato-ellipticis, leguminibus lanato-hirsutissimis.  $\frac{3}{5}$  in Corsicæ collibus asperis. — Spartium lanigerum Desf. (v. e.)

#### Sect. III. TUBOCYTISUS. DC. l. c.

Calyx tubulosus apice bilabatus. — Frutices inermes.

8. *C. supinus* (Jacq. aust. 1. t. 20.) caulibus decumbentibus ramosis, ramis teretibus junioribus hirsutulis adultis glabratiss, foliolis obovato-obtusis subtus hirsutis, floribus 2-4 subterminalibus pedunculatis, calycibus leguminibusque parè villosis.  $\frac{3}{5}$  in collibus herbosis apricis subdomosis Alpium Delphinensium et propè Fontem-Bleandi ex Merat. — Fl. fr. 5. p. 549. — Clus. hist. p. 96. n. VII. ic. (v. v. DC.)
9. *C. capitatus* (Jacq. aust. t. 33.) caulibus erectis, ramis strictis hispidis, foliolis ovato-ellipticis villosis, floribus ad apices ramorum plurimis capitatis, calycibus leguminibusque villosis-hispidis.  $\frac{3}{5}$  ad margines sylvarum Burgundiæ orient., Lugduni, agri Ruscini. ex Benth. Andegaviæ ex Bast., Bardigaliæ ex Laterr. — Fl. fr. 4. p. 505. — Clus. hist. 1. p. 96. f. 1 et 3. (v. v. DC.)

#### Sect. IV. LOTOIDES. DC. l. c.

Calycis tubus brevis obconicus, labiq. superius bipartitum, inferius tridentatum; corolla vix calyce longior.

10. *C. argentus* (L. sp. 1043.) caulibus decumbentibus foliis calycibus corollis leguminibusque adpressè sericeis, foliis petiolatis trifoliolatis, foliolis oblongo-lanceolatis, floribus 3-4 terminalibus.  $\frac{3}{5}$   $\frac{2}{4}$  in petrosis Delphinatiss,



## LEGUMINOSÆ.

119

prov. austr. usque ad Tolosam. — Fl. fr. 4. p. 506. — Lqh. ic. 2. p. 41. f. 2. (v. v.)

### VI. ADENOCARPUS. DC. fl. fr. 5. p. 549. leg. mem. vi.

Calyx obconicus sæpè glandulosus bilabiatus, labio super. bipartito, inferiore productiore trifido; carina obtusa genitalia fovens; stam. monadelphæ; legumen oblongum plano-compressum glandulas subpedicellatas undiquè gerens. — Frutices.

1. *A. PARVIFOLIUS* (DC. l. c.) calyce glanduloso-subpubescente, labii inferioris laciniâ mediâ lateralibus longiore labium super. longè superante, ramis glabris, floribus remotis. ♂ in ericetis apricis Galliarum occidentalia. — Cytisus complicatus. Fl. fr. 4. p. 503. — Clus. hist. 1. p. 94. f. 1. (v. v.)
2. *A. TELONENSIS* (DC. l. c.) calyce eglanduloso pubescente, labii inferioris laciniis subæqualibus labium superius paulò superantibus, ramulis glabresculis, floribus remotis. ♂ in sterilibus et ericetis in Pyrenæis, Cebennis, montibus Galloprovinciæ. — Cyt. Telonensis Lois. in Duham. arb. ed. nov. 5. t. 47. f. 2. (v. v. DC.)

### VII. ONONIS. L. gen. n. 863.

Calyx campanulatus 5-fidus, laciniis linearibus; vexillum magnum striatum; stamina monadelphæ, decimo interdum sublibero; legumen sæpius turgidum sessile oligospermum.

\* *Floribus pedunculatis flavis.*

1. *O. ARACHNOÏDEA* (Lapeyr. abr. 409.) suffruticosa villosa non viscosa, foliis pinnatum 3-foliolatis, foliolis ovali-oblongis apice serratis, summis interdum simplicibus, pedicellis 1-floris aristatis. ♂ ♀ in Pyrenæis orientalibus secus torrentes et propè Perpignanum. — Fl. fr. 5. p. 552. Vexillum rubro striatum. Rami villis non viscosis molliter hirsuti. (v. v.)
2. *O. NATRIX* (DC. fl. fr. 4. p. 514.) suffruticosa viscoso-pubescentis, stipulis ovatis acuminatis, pedicellis 1-floris aristatis aristâ parte pedunculi superiore longiore, calycis hirsuto-glandulosi laciniis lanceolato-linearibus. ♂ ♀ in apricis. — Fl. fr. 4. p. 514.
  - α. vexillo flavo. — *O. Natrix* L. sp. 1008. — Cam. epit. 444. ic. (v. v.)
  - β. vexillo rubro striato. — *O. pinguis* L. sp. 1009. (v. v.)
3. *O. RAMOSISSIMA* (Desf. atl. 2. t. 186.) basi suffruticulosa ramosissima pubescenti-viscosa, stipulis lanceolatis acuminatis, pedunculis aristatis aristâ parte superiore pedunculi longiore, calycis viscido-hirsuti laciniis linearibus. ♂ ♀ in maritimis arenosis Nicæ. — Fl. fr. 4. p. 515. Vexillum venis purpureis striatum. An satis ab. *O. Natrice* β. distincta? (v. v.)
4. *O. ARENARIA* (DC. cat. Monsp. 128.) basi suffruticulosa ramosissima pubescenti-viscosa, stipulis lineari-lanceolatis longè acuminatis, pedicellis 1-floris folio brevioribus breviter aristatis. ♂ ♀ in arenosis maritimis Mompeli. — Fl. fr. 5. p. 551. Vexillum flavum non striatum. Flores minores quam in 2 præcedentibus. (v. v. DC.)
5. *O. VISCOSA* (L. sp. 1009. var. α.) herbacea pubescens viscosa, foliis superioribus simplicibus, inferioribus 3-foliolatis foliolo medio majore, stipulis petiolum subæquantibus, pedunculis unifloris aristatis folio longioribus, corollâ calycinis lobis longiore. ① circâ Olbias et Mompelium. — Fl. fr. 4. p. 513. — Barr. ic. t. 1239. Flores flavi vexillo subpurpureo. (v. α.)
6. *O. BREVIFLORA* (DC. prod. 2. p. 160.) herbacea pilosa subviscosa, foliis superioribus simplicibus, infer. 3-foliolatis, foliolis ellipticis obtusis serratis, medio majore, stipulis serratis petiolum subæquantibus, pedunculis 1-floris longè aristatis folii longitudine, corollâ lobis calycinis brevioribus. ① ex Foro-Julii (Fréjus) misit. cl. Solier. — *O. viscosa* β. L. DC. fl. fr. 4. p. 515. Legumen calyce duplò longius. (v. α.)

7. *O. PUNASCINA* (L. mant. 267.) herbacea erecta viscoso-pubescentia, foliis superioribus simplicibus, cæteris 3-foliolatis, foliolis ovali-oblongis serratis, stipulis amplis acuminatis integris, calycibus latis striatis, pedunculis 1-floris muticis folio brevioribus. ① in apricis agri Monspelienensis. — Fl. fr. 5. p. 557. Vexillum subpubescens. (v. s.)
8. *O. ORNITHOPODIÖIDES* (L. sp. 1009.) herbacea viscoso-pubescentia, foliolis trifoliolatis, foliolis oblongis, pedunculis 1-2-floris aristatis, leguminibus linearibus cernuis inter semina contractis. ① in Corsicâ (cl. Soleirol.) — Cav. ic. 2. t. 192. (v. s.)

\* \* *Floribus pedunculatis purpurascensibus rariis albis.*

9. *O. ROTUNDIFOLIA* (L. sp. ed. 1. p. 719. non ed. 2.) fruticosa, foliis trifoliolatis, foliolis ovatis dentatis, pedunculis trifloris ebracteolatis. 5 in Pyrenæis et Alpibus. — Fl. fr. 4. p. 515. — Lob. ic. 2. p. 73. f. 1. (v. v.)

β. *aristata* (DC.) pedunculis trifloris aristatis.

10. *O. FRUTICOSA* (L. sp. 1010.) fruticosa, foliis trifoliolatis, foliolis sessilibus lanceolatis serratis, stipulis in unicam vaginantem 4-aristatam connatis, summis aphyllis, pedunculis trifloris in racemum dispositis. 5 in apricis montosis Delphini. et Galloprovinciæ. — Fl. fr. 4. p. 514. — Duh. ed. nov. 1. t. 58. (v. v.)

11. *O. CERNISA* (L. mant. 267.) basi suffruticosa multicaulis cespitoso-prostrata, glabra, foliis palmatim trifoliolatis, foliolis cuneatis stipulisque serratis, pedunculis 1-floris muticis folio longioribus. 3 ½ in petrosis Alpium Delphini. et Galloprovinciæ. — Fl. fr. 4. p. 512. All. ped. t. 10. f. 2. — Barr. ic. t. 1104. (v. v. DC.)

β. *subaristata* (DC.) pedunculis subaristatis. In Pyrenæis orientalibus, et centralibus. (v. s.)

12. *O. RECLINATA* (L. sp. 1011.) herbacea diffusa, foliis 3-foliolatis, foliolis obovatis serratis viscoso-pubescentibus, stipulis late ovatis acutis serratis, pedunculis 1-floris longitudine folii subapice subbracteolatis, calyce corollâ longiore. ① in Gallia australis maritima. — Fl. fr. 4. p. 512. — Barr. ic. t. 761. (v. s. sine fructu.)

13. *O. MOLLE* (Lag. nov. sp. p. 22.) herbacea erectiuscula, foliis trifoliolatis, foliolis oblongo-obovatis apice serratis, stipulis integrisculis, pedunculis unifloris muticis folii longitudine, calyce corollæ et demum legumini æquali. ① in Corsicâ collibus circâ Bastia reperiit Ph. Thomas. (v. s.)

14. *O. CNELEURI* (L. sp. 1007.) herbacea hirsuta, foliis 3-foliolatis, foliolis oblongo-cuneatis apice serratis, stipulis subintegris, pedunculis 1-floris muticis folio brevioribus in racemum aggregatis, calyce corollâ longiore legumini æquali. ① in petrosis aridis Corsicæ, Nicæ, Massiliæ, Monspelii. — Fl. fr. 4. p. 512. — J. Bauh. hist. 2. p. 394. f. 2. (v. s.)

\* \* \* *Floribus sessilibus aut brevissimè pedunculatis, purpureis rariis albis.*

15. *O. ALTISSIMA* (Lam. dict. 1. p. 506.) erecta, ramulis inermibus undiquè villosis subviscidis, foliis trifoliolatis, foliolis oblongo-lanceolatis acutis serratis, floribus subgeminatis, lobis calycinis villosis leguminis longitudine. 3 ½ in arvis et arenosis Delphinatûs? — Fl. fr. 4. p. 509. *O. hircina* Jacq. hort. Vind. t. 93. — Clus. hist. 1. p. 99. f. 1. (v. s. ex Valesii.)

16. *O. PROCURRENS* (Wallr. sch. crit. p. 38.) caulibus basi radicante procurentibus diffusis, ramulis floriferis ascendentibus undiquè pubescentibus, foliis trifoliolatis, foliolis ovato-subrotundis utrinquè glandulosis subserratis, lobis calycinis legum superantibus. 3 ½ in campis arenosis Gallie. — *O. arvensis* Lam. dict. 1. p. 505. Fl. fr. 4. p. 509. — Fuchs. hist. 60. ic.

β. *repens* (DC.) ramis foliisque hirsutioribus, foliolis obtusissimis. — *O. repens* L. — Dill. elth. t. 25. f. 28. In arenosis maritimis. (v. v.)

- γ? *caduca* (DC.) ramis villosis-viscosis, foliolis oblongis pubescentibus. In montibus Delphinatūs. O. *caduca* Vill. (v. s.)
17. O. *spinosa* (Wall. sch. crit. p. 379.) caulibus erectisculis ramulisque spinosis bifariam unifariamve pubescentibus, foliis trifoliolatis, foliolis oblongis basi cuneatis subintegris, floribus solitariis, calycinis lobis legumine brevioribus. ♂ ♀ ad vias et fossas. — O. *spinosa* L. L. sp. 1006. — Tabern. hist. 913. ic. An. O. *senescens* Lapeyr. abr. p. 405? (v. v.)
- β. *glabra* (DC.) foliolis oblongis, ramis glabris. ♂ ♀ in Gallis præsertim australis siccis. — O. *antiquorum* L. Fl. fr. 3. p. 509. (v. v.)
18. O. *mitissima* (L. sp. 1007.) caulibus erectis subpilosis albidis, foliis trifoliolatis, foliolis ovalibus serratis, floribus in spicam elongatam bracteis stipularibus scarioso-albidis variegatam dispositis. (1) in Corsicâ et agro Ruscitonensi? ex Lapeyr. — Dill. elth. t. 24. f. 27. (v. s.)
19. O. *scabrata* (Forak. desc. 131? Vahl. symb. 1. p. 52.) tota pubescenti-viscosa multicaulis decumbens, foliis trifoliolatis, foliolis obovatis oblongisve serratis, stipulis subdenticulatis, floribus spiesto-racemosis, calycibus corollæ et legumini æqualibus, laciniis calycinis subtrinerviis. (1) in Corsiam arenosis. Flores albidî et purpurei. (v. s.)
20. O. *alopercuoides* (L. sp. 1008.) caule solitario erecto subpiloso, foliis 1-foliolatis, foliolo elliptico obtuso stipulisque maximis subserratis, floribus dense spicatis, calycinis lobis hispidis corollâ et legumine duplò ferè longioribus. (1) in Corsicâ circâ Bonifacio ex cl. Viviani (cors. 13.) sed ait calycis laciniis corollâ tantum paulò longiores. — Schkuhr. handb. t. 194. Corolla rosea. (v. s. c.)

\*\*\* *Floribus sensilibus aut subsensilibus flavis.*

21. O. *arragonensis* (Asso. syn. arr. 96. t. 6. f. 2.) fruticosa, foliis trifoliolatis glabris, foliolis subrotundis serratis, floribus subsensilibus geminatis in racemum apyllum dispositis, calycibus villosis corollâ dimidio brevioribus. ♂ in Pyrenæis Gallis conterminis circâ Benasque. — Fl. fr. 5. p. 552. O. *dumosa* Lapeyr. — Magn. hort. Monsp. 17. t. 21. (v. s.)
22. O. *striata* (Gouan. ill. 47.) caulibus diffusis-prostratis ramosis, foliis trifoliolatis subscabris, foliolis obovato-cuneatis striatis stipulisque serratis, floribus capitatis, corollis calyce hirsuto-viscoso longioribus. ♂ ♀ in montanis Pyr., Cebenn., Galloprov. superioris, Delphinatūs. — Fl. fr. 4. p. 511. Variat ferè glabra et glanduloso-sabra subhirsuta. (v. v.)
23. O. *columnæ* (All. ped. 1. t. 20. f. 3.) pubescens, caulibus subcespitosus, foliis trifoliolatis, foliolis obovato-oblongis stipulisque serratis, summis simplicibus, floribus in apicam foliosam aggregatis, corollis calyce brevioribus. ♀ in collibus et sabulosis Gallis occidentalis, mediæ, australisque. — O. *parviflora* Lam. DC. fl. fr. 4. p. 510 non Thunb. O. *minutissima* Jacq. austr. t. 240. non L. (v. v.)
24. O. *minutissima* (L. sp. 1007.) glabra, caulibus subcespitosus, foliis trifoliolatis, foliolis oblongo-obovato-cuneatis serratis, stipulis subulatis integris, floribus in apicam foliosam aggregatis; corollis calyce brevioribus. ♀ in saxosis apricis provinc. australium. — Fl. fr. 4. p. 510. O. *barbata* Cav. ic. t. 153. Radix interdum frutescit. Calycis lobi subulati elongati. Flores vernalis plerumque apetalî ex cl. Benth. (v. v.)
25. O. *variegata* (L. sp. 1008.) glabra diffusa, foliis unifoliolatis, foliolo sessili obovato striato serrato, stipulis amplis subvaginantibus serratis, floribus subsensilibus axillaribus, corollis calyce pubescente longioribus. (1) in arenosis maritimis Corsicæ, insularum Stæchadum ex Ger. — Desf. atl. t. 185. (v. s.)

VIII. ANTHYLLIS. L. gen. n. 864.

Cal. tubulosus 5-dentatus post anthesin persistens, plus minus vesticarius inflatus; cor. alæ carinæ et vexillo subæquales; stam. omnia

connexa; legumen ovatum 1-2-spermum rarius oblongo-lineare polyspermum semper calyce indusiato tectum.

Sect. I. DORYCNOIDES. DC. *prod.* 2. p. 168.

Calyx vix inflatus, legumen 1-loculare 1-spermum. — Herbæ perennes.

1. A. GERARDI (L. mant. 100.) herbacea diffusa, foliis pinnatis glabresculis, foliolis 5-9 oblongo-linearibus mucronatis, capitalis 15-20-floris. 2/ in apricis saxosis sylvaticisque subherbosis Galloprovinciæ, Corsicæ et agri Ruscinonensis. — Fl. fr. 4. p. 517. — Ger. gallop. 490. t. 18. (v. v. DC.)

Sect. II. ASPALATHOIDES. DC. *l. c.*

Calyx vix inflatus, legumen 1-locul. 1-2-spermum. — Frutices humiles.

2. A. CYTHOIDES (L. sp. 1013.) fruticosa inermis ramosissima virgata, foliis simplicibus aut trifoliolatis, foliolo impari lateralibus multò majore, ramis canis adpressè velutinis, floribus paucis ad axillas superiores sessilibus in apicem interruptam digestis. 3/ ad rupes et colles propè Perpignanum et Telonem, in Corsicâ ex Viviani. — Fl. fr. 4. p. 518. — Barr. ic. t. 118a. fl. flavi. (v. v. DC.)
3. A. HERMANNIÆ (L. sp. 1014.) fruticosa ramosissima ramis spinoscentibus glabresculis, foliis subsessilibus simplicibus 3-foliolatisve, foliolis oblongo-cuneatis glabris aut adpressè pubescentibus, capitalis paucifloris ad axillas superiores subsessilibus. 3/ in Corsicâ. — Fl. fr. 4. p. 518. — Alp. exot. t. 26. Flores flavi parvi. (v. s.)

Sect. III. ERINACEA. *Clus. hist.* 1. p. 107. *ic.*

Calyces post anthesin inflato-vesicarii; leg. 1-loc. 1-2-ovul. (ovulo inferiore solo superstitè). — Frutex ramosissimus spinosus ferè aphyllus.

4. A. ERINACEA (L. sp. 1014.) fruticosa, ramis spinoscentibus, foliis paucissimis ovalibus oblongisve, floribus subcapitatis. 3/ in Corsicâ ex Viv. et Pyren. Catalonensibus Galliæ conterminis ex Benth. — Bot. mag. t. 676. Flores cæruleo-purpurei. (v. s. c.)

Sect. IV. VULNERARIA. DC. *l. c.*

Calyces post anthesin inflato-vesicarii, legum. 1-locul. 1-2-sperma. — Frutices aut herbæ perennes, foliis impari-pinnatis.

5. A. BARBA-JOVIS (L. sp. 1013.) fruticosa, foliis pinnatis ramulisque sericeo-tomentosis, foliolis 9-13 oblongo-linearibus æqualibus, bracteis capitula globosa multiflora æquantibus. 3/ in rupibus maritimis Corsicæ, Nicæ, Galloprovinciæ orientalis. — Fl. fr. 4. p. 517. — Barr. ic. t. 378. Flores pallidè flavi. (v. v. D.)
6. A. MONTANA (L. sp. 1012.) herbacea caespitose-ascendens, foliis pinnatis ramisque villosis, foliolis 15-19 ovali-oblongis, capitalis solitariis longè pedunculatis. 2/ in montibus præsertim provinc. australium. — Fl. fr. 4. p. 516. — Barr. ic. t. 722. Flores purpurascetes. (v. v.)
7. A. VULNERARIA (L. sp. 1012.) herbacea erectiuscula, foliis pinnatis, foliolis 5-13 inæqualibus infimis minoribus, terminalibus multò majoribus, capitalis subduplicatis. 2/ in pratis et pascuis frequens. — Fl. fr. 4. p. 516.
  - a. *vulgaris*, floribus flavis, foliis caulibusque parè villosis aut subglabris. — Lam. ill. t. 615. f. 1. (v. v.)
  - β. *rubriflora* (DC.) floribus coccineis rubrisve, foliis caulibusque parè villosis aut subglabris. — In montosis. — Dill. elth. t. 3æ. f. 413. (v. v.)
  - γ. *hirsutissima* (DC.) floribus coccineis rubrisve, foliis caulibus capitulisque hirsutissimis. — In Corsicâ et Pyrenæis. (v. v.)

Sect. V. CORNICINA. DC. *l. c.* p. 170.

Calyces inflati, legumina glabra intus septulis transversis semina sejungentibus 2-aut multilocularia. — Herbæ annuæ.

8. *A. TETRAPHYLLA* (L. sp. 1012.) herbacea procumbens, foliis pinnatis, foliolo terminali ovato maximo, cæteris 3 minimis acutis, capitulis axillaribus sessilibus paucifloris, legumine recto 2-spermo 2-locul. intrâ calycem incluso. (1) in agris et cultis apricis Galliæ australis. — Fl. fr. 4. p. 515. — Barr. ic. t. 554. (v. s.)

Subtribus II. *TRIFOLIÆ*. Bronn. diss. DC. prod. 2. p. 171.

Legumen 1-loculare. Stamina diadelpha. Caules herbacei rariùs frutescentes. Folia sæpius palmata 3-5-foliolata, rarissimè impari-pinnata.

IX. *MEDICAGO*. L. gen. n. 1214. Ser. in DC. prod. 2. p. 171.

Calyx subcylindricus 5-fidus; carina à vexillo subremota; stamina diadelpha; legumen polyspermum semper falcatum vel in spiram contortum.

Seqt. I. *HYMENOCARPOS*. Ser. I. c.

Legumina submembranacea compressa arcuata; folia 3-5-foliolata.

1. *M. CIRCINNATA* (L. sp. 1096.) foliis pinnatis 5-foliolatis foliolis integris ovatis terminali maximo oblongo, pedunculis 2-4-floris subcorymbosis, leguminibus reniformibus pilosis, margine dentato non alato. (1) in maritimis Niceæ, Galloprov., Corsicæ. — Fl. fr. 4. p. 548. — Barr. ic. t. 576. (v. s.)
2. *M. RADICATA* (L. sp. 1096.) foliis 3-foliolatis, foliolis obovatis argutè denticulatis, stipulis dentatis, pedicellis 1-floris, leguminibus latis compressis reniformibus margine lato alato dentato, alâ reticulato-venosâ. (1) circa Niceam ex All., Narbonam ex Benth. — Moris. hist. 2. t. 15. n. 3. (v. s. c.)

Sect. II. *LUPULARIA*. Ser. I. c.

Legumina reniformia, falcata vel subcochleata, margine integro. — Folia 3-foliolata.

3. *M. LUPULINA* (L. sp. 1097.) caule procumbente, foliolis obovato-cuneatis apice denticulatis, stipulis lanceolatis acutis integrisculis, pedunculis racemoso-spicatis, floribus sessilibus, leguminibus reniformibus 1-spermiis arcuato-reticulatis, seminibus ovato-subreniformibus. (2) in arvis. — Fl. fr. 4. p. 541. Melilotus Willdenowii Mer. par. t. 2. p. 411. — Fuchs. hist. 819. ic. Variat caule foliisque plus minusve pilosis. (v. v.)
4. *M. FALCATA* (L. sp. 1096.) caulibus prostratis, foliolis oblongis apice dentatis, stipulis integerrimis, pedunculis racemosis, leguminibus falcato-cochleatis glabrisculis, seminibus oblongis, radicula vix prominulâ. ♀ in pratis siccis montanis. — Fl. fr. 4. p. 540. eng. bot. t. 1016. (v. v.)
5. *M. SUFFRUTICOSA* (Ram. in DC. fl. fr. 4. p. 541.) caulibus prostratis suffruticosis, foliolis obovato-rotundis subintegris vel obcordatis, stipulis lanceolatis latis dentatis, pedunculis racemosis petiolo duplò longioribus, leguminibus pubescentibus subspiralibus arcuato-reticulatis, seminibus irregulariter cordatis, radicula prominulâ, ♀ in Pyrenæis et montosis agri Ruscinonensis. (cl. Arnott.) (v. v. DC.)
6. *B. Benthamii*, caulibus foliolis pedunculisque pubescentibus, leguminibus villosis subreticulatis cochleato-contortis. — In Pyrenæis orientalibus editoribusque ex cl. Benth. cat. 100.
6. *M. LEOCARPA* (Benth. l. c.) glaberrima, caulibus prostratis suffruticosis, foliolis obovato-rotundis subintegris obcordatisve, stipulis latis integris vel subdentatis, pedunculis racemosis petiolo duplò longioribus, leguminibus cochleato-contortis, cyclois 2-5 arcuato-reticulatis, seminibus subreniformibus. ♀ in collibus aridis Galliæ Narbonensis detexit cl. Benth. (v. s.)
7. *M. SATIVA* (L. sp. 1096.) caule erecto glabro, foliolis obovato-oblongis dentatis mucronatis, stipulis lanceolatis subdentatis, pedunculis racemosis, leguminibus lævibus cochleato-contortis tennissimè reticulatis, seminibus

irregularibus ovatis cordatisve.  $\mathcal{L}$  colitur in pratia et inde nunc forè in omni Galliâ spontanea. — Fl. fr. 4. p. 539. — Lob. ic. 2. p. 36. f. 2. (v. v.)

Sect. III. SPIROCARPOS. Ser. I. c.

Legumina cochleato-orbiculata, margine integro rugoso vel spinoso. Folia 3-foliolata.

\* *Leguminibus inermibus.*

8. *M. obscura* (Retz. obs. fasc. 1. p. 24. t. 1.) caule decumbente, foliis rhombeo-obovatis denticulatis, nervis numerosis prominentibus, stipulis lanceolatis profundè dentatis, pedunculis multifloris, leguminibus reniformibus vel orbiculariter cochleatis integerrimis glabris venosis 2-spermis, venis marginem versùs anastomosantibus, seminibus reniformibus. ① in Galliâ australi ex cl. Loiseleur. — Fl. fr. 4. p. 566. (v. s. ex Italiâ.)
9. *M. Soleirolii* (Duby mss.) caule decumbente angulato piloso, foliis trifoliolatis, foliolis latè rhombeo-obovatis denticulatis subtùs pubescentibus nervosis, stipulis dentato-pinnatifidis laciniis setiformibus, pedunculis racemosis multifloris, floribus magnis, laciniis calycinis villosis setaceis, leguminibus congestis inermibus cochleatis plano-convexis longitudinaliter reticulatis. ① in Corsicâ, propè Calvi (cl. Soleirol.). (v. s. D.)
10. *M. orbicularis* (All. ped. n. 1150.) caulibus diffusis, foliolis obcordatis apice dentatis, stipulis laciniatis, laciniis angustissimis divergentibus, pedunculis 1-2-floris, leguminibus cochleatis compressis glabris reticulato-venosis polyspermis, seminibus subtriangularibus. ① in Galliâ australi et circâ Parisios. — Fl. fr. 4. p. 542. — Moris. hist. s. 2. t. 15. n. 2. (v. v.)
11. *M. scutellata* (All. ped. n. 1155.) pilosa, caulibus diffusis, foliolis ovatis ambitu dentatis, stipulis lanceolatis dentatis, pedunculis 1-2-floris, leguminibus cochleatis polyspermis suprà convexo-hemisphæricis subtùs planis venoso-reticulatis, venis margine tenui anastomosantibus, seminibus maximis reniformibus. ① in arvis et messibus Nicæ, Galloprov., Delphinatùs ex Vill., Mompellii, Meloduni ex Merat. — Fl. fr. 4. p. 543. — Moris. hist. s. 2. t. 15. n. 3. (v. v.)
12. *M. rugosa* (Lam. dict. 3. p. 632.) caulibus procumbentibus, foliolis rhombeo-obovatis versùs apicem denticulatis, stipulis lanceolatis dentatis, pedunculis 2-4-floris, leguminibus 2-3-cyclis reticulato-venosis, seminibus reniformibus compressis apice subtruncatis. ① in Galliâ meridionali ex cl. Loiseleur. — Fl. fr. 5. p. 566. — Moris. hist. s. 2. t. 15. n. 4. (v. s. c.)
13. *M. tornata* (W. sp. 3. p. 1409.) caulibus diffusis, foliolis obovatis dentatis, stipulis ciliato-dentatis, pedunculis multifloris, leguminibus cochleato-cylindricis utrinquè truncatis glabris lævissimis, cyclis subconcretis, seminibus reniformibus apice truncatis. ① in arvis Galloprovinciæ ex Ger., circâ Bayonam ex Lois. — Fl. fr. 4. p. 543. (v. s. c.)
14. *M. articulata* (Benth. cat. 101.) subglabra, stipulis laciniatis, legum. adpressè cochleatis 4-6-cyclis cylindricis utrinquè planis reticulatis, margine crasso bisulcato tuberculoso. ① in segetibus demissiorum Corbariensium propè Castet detexit cl. Benth.
15. *M. turbinata* (W. sp. 3. p. 1409.) caulibus diffusis, foliolis rhombeo-obovatis stipulisque dentatis, pedunculis subbifloris, leguminibus cochleato-ovatis utrinquè convexis, margine crassissimo subconcreto, seminibus reniformibus apice truncatis. ① in messibus prov. austral. Nicæ, Galloprovinciæ ex Ger., Occitanie ex Gouan. — Fl. fr. 4. p. 544. — Moris. hist. s. 2. t. 15. n. 5. (v. s.)
16. *M. turriculata* (W. sp. 3. p. 1410.) caulibus procumbentibus, foliolis rhombeo-obovatis dentatis, stipulis lanceolatis dentatis, pedunculis 1-2-floris, leguminibus cochleato-ovatis, margine crassissimo subconcreto tuberculis duplici serie dispositis ætate leprosis, seminibus reniformibus apice truncatis. ① in provinciis australibus ex cl. Desf. — Fl. fr. 4. p. 544. J. Bauh. hist. 2. p. 385. f. 1. — Moris. hist. s. 2. t. 15. n. 6. (v. s. c.)

17. *M. STRIATA* (East. Journ. bot. 1814. 3. p. 19.) caulibus procumbentibus, foliolis obovatis denticulatis, stipulis lanceolatis dentatis, pedunculis 5-6-floris leguminibus 3-5-cyclis cochleato-cylindricis, margine crassiusculo supernè punctis distantibus submamillaribus subtùs lineolis notato, seminibus reniformibus apice truncatis parvis. ① in arenosis maritimis Pictaviæ. — Fl. fr. 5. p. 567. Hanc cum *M. littoralis* conjungit cl. Benth. (v. s.)

\* \* *Leguminibus spinosis.*

18. *M. APICULATA* (W. sp. 3. p. 1414.) caulibus prostratis, foliolis obovatis apice vix denticulatis, stipulis ciliato-dentatis, pedunculis 3-7-floris, leguminibus cochleatis utrinquè planis 3-cyclis reticulato-sublacunosus margine muricatis glabris, spinis divergentibus rectis, seminibus oblongo-reniformibus. ① in pratis et messibus Gallie occidentalis australisque. — Fl. fr. 4. p. 548. (v. v.)

19. *M. DENTICULATA* (W. sp. 3. p. 1414.) caulibus prostratis, foliolis obovatis obcordatisve denticulatis, stipulis ciliato-dentatis, leguminibus utrinquè planis 2-cyclis reticulatis margine aculeatis glabris, aculeis divergentibus hamatis, seminibus oblongo-reniformibus apice subtruncatis. ① circà Niceam, Monspelium, Aginnum, Burdigalam (Des Moulins). — Fl. fr. 4. p. 548. An species distincta? (v. v.)

β. *ciliaris* (Ser.) aculeis elongatis. In prov. mediterraneis. (v. v.)

20. *M. PUBESCENS* (DC. cat. monsp. p. 124.) molliter villosa, caulibus prostratis elongatis, foliolis latis obovatis denticulatis, stipulis lanceolatis acuminatis denticulatis, pedunculis 2-8-floris, leguminibus cochleatis 2-4-cyclis, glabris infra marginem aculeatis, aculeis crassis convergentibus adpressis margine longioribus rectis, sem. apice subtruncatis. ① circà Niceam, Monspelium, et in agro Andegaviensi. — Fl. fr. 5. p. 570. — J. Bauh. hist. 2. p. 385. ic? excl. spec. inf. *M. spinulosa*. Fl. fr. 5. p. 569. Ser. in DC. prod. 2. p. 176. Hanc *M. tuberculata* refert cl. Benth. (v. v. DC.)

21. *M. TUBERELLUM* (W. sp. 3. p. 1416.) caulibus prostratis, foliolis obovatis retusis dentatis, stipulis ciliato-dentatis, pedunculis multifloris, leguminibus cochleato-cylindricis utrinquè planis, cyclis 5 reticulatis glabris, aculeis brevissimis distichis subulatis patentissimis, seminibus reniformibus. ① in Corsicâ? et provinciis australibus. — Fl. fr. 4. p. 549. Var. *M. pentacyclæ* ex Benth. (v. s.)

22. *M. MARINA* (L. sp. 1097.) densissime tomentosa, caulibus erectis, foliolis cuneiformibus integerrimis, stipulis lanceolatis integris, pedunculis multifloris, leguminibus cochleatis nervosis subspinosis, seminibus reniformibus. ② in arenosis secùs mare Mediterraneum, secùs Oceanum rariùs. — Fl. fr. 4. p. 546. — Clus. hist. 2. p. 243. f. 2. (v. v.)

23. *M. CORONATA* (Lam. dict. 3. p. 636. non Gærtn.) villosa, caulibus prostratis, foliolis obovato vel obcordato-subrotundis parvis denticulatis, stipulis lanceolatis integris, pedunculis 4-7-floris folio multò longioribus, leguminibus cochleatis 1-2-cyclis villosis reticulatis, aculeis divaricatis rectis coroniformibus, seminibus reniformi-lunulatis parvis. ① in Galloprovincia, Occitania. — Fl. fr. 4. p. 549. — Moris. hist. s. 2. t. 15. n. 16. (v. s.)

24. *M. LITTORALIS* (Rohde in Lois not. 118.) villosa, caulibus procumbentibus, foliolis obcordatis vel obovato-cuneiformibus apice dentatis, stipulis lanceolatis dentatis, pedunculis 2-4-floris, leguminibus cochleatis subcylindricis glabris tetracyclis reticulatis margine aculeatis, cyclis crassis coriaceis, aculeis rigidis subhamosis, seminibus reniformibus. ②? in arenosis maritimis Niceæ, Galloprovinciæ, Occitaniæ. — Fl. fr. 5. p. 568. — J. Bauh. hist. 2. p. 385. ic? (v. v.)

β. *breviseta* (DC.) aculeis brevibus rectis. (v. v.)

25. *M. LAPPACEA* (Lam. dict. 3. p. 637.) glabriuscula, caulibus procumbentibus foliolis obcordatis dentatis, stipulis ciliato-dentatis, pedunculis 3-5-floris, leguminibus cochleatis glabris 3-cyclis nervosis margine aculeatis, aculeis longis hamosis, seminibus reniformibus subtruncatis. ① in Corsicâ, circà

*Monspeliu* et Beziers. — Fl. fr. 5. p. 569. Var. *M. denticulata* ex cl. Benth. (v. s.)

26. *M. PENTACTYLA* (DC. cat. monsp. p. 124.) pilosiuscula, caulibus prostratis, foliolis obovatis denticulatis, stipulis ciliato-denticulatis, pedunculis multifloris, leguminibus cochleatis glabris utrinque convexis, cyclis 5 supernè reticulatis, spinis longis subdivergentibus apice uncinatis, seminibus ovato-subreniformibus compressis. (1) circà Narbonam, Perpignanum, ipse in insulis Stachadam legi. — Fl. fr. 5. p. 569. An spec. satis distincta? (v. v.)
27. *M. DISCIFORMIS* (DC. cat. monsp. p. 124.) pilosa, caulibus procumbentibus, foliolis obcordatis dentatis, pedunculis 3-4-floris, leguminibus cochleatis glabris, cyclis 5 approximatis infrà venosis, supernè planis, 4 inferioribus margine spinulas longas setaceas apice uncinatas gerentibus, quino abortu nudo enervio, seminibus subreniformibus. (1) in sterilibus Monspeli, et Olette Pyren. orient. (Ph. Thomas.) — Fl. fr. 5. p. 567. (v. v. DC.)
28. *M. TRIBULOIDES* (Lam. dict. 3. p. 635.) pilosa, caulibus prostratis, foliolis obovatis dentatis, stipulis profunde dentatis, pedunculis 2-floris, leguminibus cochleatis cylindricis utrinque planis 5-cyclis rugosis, spinis lateralibus non marginalibus crassis divergentibus apice subuncinatis, margine crassiusculo elevato non canaliculato, seminibus reniformibus apice truncatis. (1) in Galloprovincia propè Arelatem. — Fl. fr. 5. p. 568. (v. s.)
- β. Narbonensis* (DC.) spinis adpressis brevibus, cyclisque turgioribus, leguminibus tumido-verrucosis. Circà Carcassonam et Narbonam. (v. v. DC.)
29. *M. PÆRICOX* (DC. cat. monsp. p. 123.) caulibus prostratis, foliolis obcordatis denticulatis, stipulis ciliato-dentatis, pedunculis 1-2-floris brevibus, leguminibus cochleatis glabris utrinque planis, cyclis tennibus, margine crasso lateraliter spinoso, spinis subdivergentibus apice subuncinatis, seminibus ovato-reniformibus. (1) in sterilibus Corsicæ repariit Ph. Thomas, Nizzae Stire, circà Forum-Julianum cl. Balbis, *Monspeliu* cl. Benth. — Fl. fr. 5. p. 570. (v. s.)
30. *M. MINIMA* (Lam. dict. 3. p. 636.) pilosa, caulibus prostratis, foliolis obovatis vel obcordatis integris apice 3-dentatis, pedunculis 1-3-floris brevibus, leguminibus cochleatis villosulis globosis, cyclis 3-5 angustissimis planiusculis, spinis compressis uncinatis basi canaliculatis, seminibus reniformibus. (1) in campestribus. — Fl. fr. 4. p. 545. — Moris. hist. 5. 2. t. 15. n. 15. (v. v.)
- β. canescens* (Ser.) caulibus foliis calycibus leguminibusque canescentibus, spinis longiusculis. — In aridis.
- γ. longiseta* (DC.) spinis longissimis, caule elongato, pedunculis multifloris elongatis. — In provinciis australibus. (v. v.)
31. *M. MACULATA* (W. sp. 3. p. 1412.) caulibus prostratis, foliolis obcordatis obovatisve dentatis maculatis, stipulis dentatis, pedunculis 3-5-floris, leguminibus cochleatis utrinque compressis albidis, cyclis 3-5 circulariter nervulosis subquadriariam spinosis, spinis anastomosantibus reflexis, seminibus reniformibus. (1) in sabulosis Gallie occidentalis, mediæ et australis passim. — Fl. fr. 4. p. 548. — Moris s. 2. t. 15. n. 12. (v. v.)
32. *M. GERARDI* (W. sp. 3. p. 1415.) caulibus prostratis, foliolis obovatis denticulatis, stipulis setaceo-dentatis, pedunculis subbifloris, leguminibus cochleatis villosotomentosis utrinque planiusculis, cyclis 5 enerviis crassis, spinis apice uncinatis vix divergentibus, seminibus reniformibus. (1) in provinciis australibus usque ad Aginnum. — *M. villosa* Fl. fr. 4. p. 545. — Moris. hist. 2. s. 2. t. 15. n. 18. (v. v.)
- β. minor* (Ser.) leguminibus cylindricis, aculeis brevioribus.
33. *M. RIGIDULA* (Lam. dic. 3. p. 634.) caulibus prostratis, foliolis obovatis apice denticulatis, stipulis basi dentatis, pedunculis 2-3-floris, leguminibus cylindraceis utrinque appressatis dorso muricato-spinosis, spinis vix diver-



gentibus. († In Gallia australi ex Lam. — DC. fl. fr. 4. p. 544. Cam priori conjungit cl. Benth. (v. s. c.)

34. *M. MURICATA* (All. ped. n. 1158.) villosissima, caulibus prostratis, foliolis rhombeo ovatis dentatis, stipulis ciliato-dentatis, pedunculis 1-3-floris, leguminibus cochleato-ovatis glabris, cyclis 5 crassissubnervosis, spinis rigidis conicis arcuatis, seminibus irregulariter reniformibus. († in pratis Monspelii, Niceæ, Galloprov. ex Ger., Delphin. ex Vill., Parisiorum ex Vaill., agri Mimatenis (Prost.) — Fl. fr. 4. p. 547. — Vaill. bot. par. t. 33. f. 7. (v. s.)
35. *M. SPÆNOCARPUS* (Bertol. ital. dec. 3. p. 60.) glabra, foliolis rhombeco-ovatis argutè serrulatis, stipulis laciniatis, pedunculis multifloris, leguminibus cochleato-globosis, cyclis arctissime adpressis subconcretis, margine crasso longitudinaliter bisulcato, aculeis brevibus crassiusculis. († in Corsica (Ph. Thomas), circa Olbia ex Benth. — Moris. hist. s. 2. t. 15. n. 12? (v. s.)
36. *M. LACINIATA* (All. ped. n. 1159.) caule erecto, foliolis linearibus inciso-dentatis truncatis, stipulis ciliato-dentatis, pedunculis 1-bifloris, leguminibus cochleato-subglobosis spinosissimis, aculeis rectis subulatis uncinatis compresso-canaliculatis, margine crassiusculo planiusculo, seminibus oblongo-reniformibus. († circa Monspelium. — Fl. fr. 4. p. 547. — Breyn. cent. 81. t. 34. — Magn. Monsp. 270. ic. (v. v. c.)
37. *M. INTERTEXTA* (W. sp. 3. p. 1411.) caule procumbente, foliolis obovatis dentatis, stipulis ciliato-dentatis, pedunculis subbifloris, leguminibus pilosis cochleatis ovato-disciformibus reticulatis, aculeis rectis crassiusculis acutis adpressis. († in provinciis australibus ex Lam. — Fl. fr. 4. p. 546. — Moris. hist. s. 2. t. 15. n. 8. (v. s. c.)
38. *M. ECHINUS* (DC. fl. fr. 4. p. 546.) caulibus prostratis, foliolis obovatis obcordatisve obsolete dentatis, pedunculis 5-6-floris petiolis longioribus, leguminibus cochleato-ovalibus grossè reticulatis, spinis compressis divaricatis adpressis longissimis acutissimis, cyclis 6-7, seminibus reniformibus. († circa Niceam ex Allioni. — Moris. hist. s. 2. t. 15. n. 9. (v. s.)

X. TRIGONELLA. *Lin. gen. n. 1213.*

Calyx campanulatus 5-fidus; carina minima alæ vexillumque subpetentes corollam 3-petalam mentientes; legumen oblongum compressum vel cylindricum acuminatum erectiusculum polyspermum.

Sect. I. *GRAMMOCARPUS. Ser. in DC. prod. 2. p. 181.*

Flores capitati umbellative; legumina ovata longitudinaliter nervosa longè rostrata.

1. *T. UNCINATA* (Ser. l. c.) caulibus ascendentibus, foliolis oblongo-ovatis denticulatis, stipulis linearis-setaceis, floribus dense spicatis, calycibus 2-labiatis, labio superiore 1-dentato brevi, inferiore 4-fido, leguminibus pilosis 1-spermis, rostro longo curvo, seminibus ovato-compressis punctato-rugosis. († in Gallia Burdigalensi circa Libonno ex Ser. l. c. — *Mentha uncinata* Poir. supp. 3. p. 648. (v. s. ex Iberiâ.)

Sect. II. *FÆNUMGRÆCUM. Tourn. inst. 1. t. 236.*

Flores sessiles solitarii geminique, legumina elongata compressa longè rostrata longitudinaliter reticulata.

2. *T. PROSTRATA* (DC. fl. fr. 5. p. 571.) caulibus diffuso-procumbentibus, foliolis obovato-oblongis cuneiformibus argutè serratis, calycibus pilosis, dentibus subulatis longitudine tubi, leguminibus falcatis rostro longioribus 6-spermis, seminibus parvis ovato-subreniformibus punctato-rugosis. († in sterilibus Galloprovinciæ et Occitanie inferioris. — *T. Fænum-Græcum* fl. fr. 4. p. 551. — J. Bauh. hist. 2. p. 365. f. 2. (v. v. DC.)
3. *T. FÆNUM-GRÆCUM* (L. sp. 1022.) caule erecto simplici, foliolis obovatis obsolete dentatis, stipulis lanceolato-falcatis integris, calycibus pilosis, dentibus subulatis longitudine tubi, leguminibus falcatis rostro duplè longiori-

bus polyspermis, seminibus magnis ovatis punctato-rugosis. ① circa Mons-pelium spontanea, in Alsatia culta. — Fl. fr. 4. p. 551 et 6. p. 571. Gærtn. fruct. t. 152. f. 3. (v. s.)

Sect. III. BUGERAS. Ser. l. c.

Flores racemoso-umbellati, legumina cylindraceo-compressa subarcuata reticulata mucronata.

4. T. MONSPELIACA (L. sp. 1095.) pilosa, caulibus prostratis, foliolis obovato-cuneatis serrulatis, stipulis lineari-subulatis, umbellis sessilibus, lacinis calycinis subulatis longitudine tubi, leguminibus subsenis sessilibus congestis arcuatis divaricatis, seminibus oblongis. ① in provinciis australibus et circa Parisios. — Fl. fr. 4. p. 552. — J. Bauh. hist. 2. p. 273. f. 1. (v. v.)
5. T. POLYCARPA (L. sp. 1093.) caulibus diffusis, foliolis cuneiformibus apice dentatis, stipulis lanceolatis basi subdentatis, umbellis sessilibus, leguminibus subsessilibus subquaternis erectis subrectis longis linearibus, seminibus oblongis subtruncatis. ① in aridis et incultis Nices ex All., Galloprov. ex Ger., Occitanis. — Fl. fr. 4. p. 552. (v. s. c.)

Sect. IV. FALCATULA. Brot. phyt. 160.

Flores umbellati pedunculati, legumina compressa falcata reticulata mucronata.

6. T. HYBRIDA (Pourr. act. toul. 3. p. 33.) caulibus ascendentibus, foliolis obovato-cuneatis integris glabris, stipulis inæqualiter lanceolatis denticulatis, floribus racemosis 2-5, leguminibus pedicellatis glabris, seminibus irregulariter cordatis. ② in montibus Corbariensibus, et propè Bsyonam ex Lois. — Fl. fr. 4. p. 350. ic. rar. t. 29. (v. s.)
7. T. ORNITHOPOLIOIDES (DC. fl. fr. 4. p. 550.) caulibus prostratis, foliolis obcordatis denticulatis, stipulis lanceolatis integris submembranaceis acutissimis, floribus subumbellatis 2-4, leguminibus subfalcatis calyce duplò longioribus. ① in herbosis sterilibus Andegaviæ, Monspelii, Pyren. ex Benth. — Pluck. t. 68. f. 1. (v. s.)
8. T. CORNICULATA (L. sp. 1094.) caule erecto, foliolis ovatis apice dentatis, stipulis lanceolatis integris, floribus racemosis, leguminibus declinatis subfalcatis, seminibus inæqualiter cordatis. ① in Delphin. australi ex Vill., Galloprovincia ex Tourn., Nices ex All. — Fl. fr. 4. p. 550. — J. Bauh. hist. 2. p. 372. f. 2? (v. v. c.)

XI. MELILOTUS. Tourn. inst. 406. t. 229.

Calyx tubulosus 5-dentatus; carina simplex, alæ vexillo breviores, legumen calyce longius, coriaceum mono-aut oligospermum vix dehiscens. — Folia pinnatim 3-foliolata.

Sect. I. CELORUTIS. Ser. in DC. prod. 2. p. 186.

Legumina lacunos-rugosa.

1. M. KOCHIANA (Wild. enum. 790.) caule ascendente, foliolis lineari-oblongis denticulatis, stipulis dentatis, dentibus calycinis tubum æquantibus, vexillo carinâ longiore, leguminibus 1-2-spermis ovatis acutis compressis longitudine styli, seminibus ovoideis subcompressis. ② in agro Monspelienſi et Pyrenæis orientalibus. — Fl. fr. 5. p. 564. Flores flavi. (v. s.)
2. M. OFFICINALIS (Wild. enum. 790.) caule erecto ramoso, ramis patentissimis, foliolis lanceolato-oblongis obtusis remotè serratis, stipulis setaceis, dentibus calycinis inæqualibus longitudine tubi, vexillo atque carinam æquantibus, leguminibus 2-spermis obovatis pilosis oculis demum nigricantibus subgibbosis longitudine styli, seminibus inæqualiter cordatis. ① in campestribus. — Fl. fr. 4. p. 537. Oed. fl. dan. t. 934. (v. v.)
3. M. LEUCANTHA (Koch. ex DC. fl. fr. 5. p. 554.) caule erecto ramoso, foliis ovato-oblongis apice truncatis mucronatisque remotè serratis, stipulis setaceis, dentibus calycinis inæqualibus longitudine tubi, vexillo carinâ atque longiore, leguminibus 1-spermis ovatis viridibus, seminibus exactè

ovatis. ② in agris. — Trifolium officinale. L. sp. 1078. Sturm. dent. 1. fasc. 15. f. k. Flores albi. (v. v.)

4. *M. PARVIFLORA* (Desf. atl. 2. p. 192.) caule ascendente, foliolis inferioribus obovato-subrotundis integris, superioribus oblongis obsolete dentatis, stipulis lineari-setaceis, floribus minutissimis dense spicatis, dentibus calycinis subæqualibus latis, alis carinæ vexilloque subæqualibus, leguminibus ovatis 1-spermis, seminibus ovoideis legumini adherentibus. ① in pratis siccis collibusque apricis Galloprov., Occit., agri Ruscinonensis et Santoniz (Des Moulins). — Fl. fr. 4. p. 538. Flores flavi. (v. s.)
5. *M. ITALICA* (Lam. dict. 4. p. 65.) caule ramisque ascendentibus crassis, foliolis obovato-subrotundis obsolete denticulatis, stipulis lanceolatis acutis basi sublaciniatis, dentibus calycinis inæqualibus vix longitudine tubi, alis carinæ æqualibus vexillo brevioribus, leguminibus globosis 1-2-spermis. seminibus orbiculato-compressis. ① in Corsicâ reperit Ph. Thomas. Fl. flavi. Variet foliolis subintegris. (v. s.)
6. *M. GRACILIS* (DC. fl. fr. 5. p. 565.) caule ramisque ascendentibus gracilibus, foliolis obovatis angustis apice denticulatis, stipulis subsetaceis, dentibus calycinis subæqualibus tubo vix brevioribus, alis carinæ vexilloque æqualibus, leguminibus globosis 2-spermis, seminibus orbiculato-compressis. ① in arenosis Corsicæ, Niceæ, Fori-Julii, Nemausi, Perpignani. — Flores flavi. (v. s.)

Sect. II. CAMPYLORUTIS. Ser. I. c.

Legumina arcuatim dense venosa.

7. *M. MISSANENSIS* (Desf. atl. 2. p. 192.) caule erecto, foliolis obovato-cuneatis denticulatis, stipulis basi latis dentatis apice linearibus, racemis paucifloris, dentibus calycinis subæqualibus tubo vix brevioribus, leguminibus inæqualiter lanceolatis acutis 1-spermis, seminibus ovato-compressis magnis. ① in agris circâ Telonem ex cl. Loisel. — Fl. fr. 4. p. 539. — Moris. a. 2. t. 16. n. 9. Flores lutei. (v. s. ex Pedem.)
8. *M. SULCATA* (Desf. atl. 2. p. 193.) caule erecto, foliolis obovatis obsolete dentatis, stipulis lineari-subulatis basi dilatatis integris dentatisve, dentibus calycinis latiusculis tubo brevioribus, leguminibus obovato-subglobosis arcuatis 2-spermis, seminibus suborbiculatis. ① circâ Niceam, Mompelium, Burdigalam ex Des Moulins. — Fl. fr. 4. p. 538. — J. Bauh. 2. p. 371. f. 1. Fl. flavi. (v. s.)

XII. TRIFOLIUM. Tourn. inst. 404. t. 228. Ser. in DC. prod. 2. p. 189.

Calyx tubulosus persistens, 5-fidus; carina alis et vexillo brevior; stamina diadelphea; legumen parvum vix dehiscens sæpius ovatum 1-2-spermum calyce brevius, rarius oblongum 3-4-spermum calycem paululum superans. — Folia palmatim 3-foliolata.

Sect. I. LAGOPUS. Ser. I. c.

Flores spicati; spicæ oblongæ basi ebracteatae, calyx post anthesin non inflatus villosus.

1. *T. ANGUSTIFOLIUM* (L. sp. 1083.) caule erecto, foliolis lineari-lanceolatis acutissimis ciliatis, stipulis longissimis angustis apice subulatis, spicis solitariis terminalibus conico-oblongis, calycibus costatis, laciniis setaceis apice spinulentibus corollam gamopetalam vix superantibus, laciniâ infimâ longiore. ① in arvis Galliz mediterraneæ et austro-occidentalis. — Fl. fr. 4. p. 529. J. Bauh. 2. p. 376. f. 3. (v. v.)
2. *T. PARVUM* (Lois. gall. t. 14.) caule erecto, foliolis lineari-lanceolatis acutis integerrimis ciliatis, stipulis angustis apice subsetaceis; apicis ovato-oblongis terminalibus solitariis, calycibus costatis, laciniis setaceis corollâ

- multò brevioribus, lacinia infimà longiore. ① circà Monspelium. — Fl. fr. 5. p. 557. (v. s.)
3. *T. RUBENS* (L. sp. 1081.) caule erecto stricto, foliolis oblongis obtusissimis glaberrimis, valde denticulatis, stipulis longissimis latissimisque apice lanceolatis, spicis terminalibus subgeminis, calycibus valde striatis, laciniiis setaceis brevissimis, infimà corollam gamopetalam vix æquante. ② in pratis et ad margines sylvarum montium. — Fl. fr. 4. p. 525. — J. Bauh. a. p. 375. f. 1. (v. v.)
4. *T. INCARNATUM* (L. sp. 1033.) caule erecto, foliolis subrotundo-obcordatis crenatis villosis, stipulis latis brevissimis apice obtuso sphacelato, spicis terminalibus solitariis longè pedunculatis, calycibus costatis pilosissimis, laciniiis lanceolato-setaceis æqualibus subatellatis corollam gamopetalam æquantibus. ① in pratis subhumidis Gallie australis et occidentalis, sæpè cultum. — Fl. fr. 4. p. 528. — Mill. ic. t. 267. f. 1. (v. v. c.)
- $\beta$ . *Molinerii* (DC.) floribus albedo-carneis, stipulis vix sphacelatis. T. Molinerii Balb. — In Corsicà et circà Nicæam sæpè cum var.  $\alpha$  colitur.
5. *T. LACINIOSUM* (Poirr. ex W. sp. 3. p. 1365.) hirsutissimum, caule ramosissimo, foliolis obovato-cuneatis denticulatis, stipulis lanceolatis latissimis brevibus nervosis, spicis terminalibus oblongis solitariis sessilibus, calycibus costatis pilosissimis, laciniiis setaceis æqualibus corollà brevioribus. ① circà Prades in agro Ruscinonensi reperiit Ph. Thomas. (v. s.)
6. *T. ARVENSE* (L. sp. 1083.) caule erecto, foliolis spathulatis linearibus apice subtridentatis, stipulis angustis membranaceis longissimè subulatis pilosis, spicis oblongis villosissimis mollibus, calycibus pilosissimis, laciniiis æqualibus corollà polypetalà longioribus. ① in arvis frequens. — Fl. fr. 4. p. 530. — Fuchs. hist. 494. ic. (v. v.)
- $\beta$ . *gracile* (DC.) caule foliisque glabresculis, laciniiis calycinis apice villosis coloratis. T. gracile Thail. — Barr. ic. 901. (v. v.)
- $\gamma$ . *capitatum* (Ser.) capitulis globosis sessilibus. — Barr. ic. t. 902.
7. *T. LIGUSTICUM* (Balb. in act. acad. ital. 1. p. 192. f. 2.) caule ramoso diffuso, foliolis obovatis obcordatisve latis apice subdenticulatis, stipulis latiusculis lanceolatis nervosis parvis acuminatis, spicis oblongis geminis divaricatis pedunculatis pilosis, laciniiis calycinis subulatis æqualibus tubo duplò longioribus corollamque superantibus. ① in Corsicà, insulis Stæchadum, et agro Ruscinonensi propè Bagnols ex cl. Benth. — Fl. fr. 5. p. 557. L. arrectisetum. Brot. phyt. t. 63. f. 1. (v. v.)
8. *T. LAPPACEUM* (L. sp. 1082.) caule ramoso diffuso, foliolis obovatis obcordatisve denticulatis, stipulis nervosis angustis longè subulatis, spicis subglobosis hispidis terminalibus subsolitariis, laciniiis calycinis æqualibus subulatis tubo duplò triplòve longioribus et corollam æquantibus. ① in provinciis anstraliis. — Fl. fr. 4. p. 725. — Barr. ic. 871. (v. v. DC.)

#### SECT. II. PHLEASTRUM. Ser. L. c.

Flores capitati; capitula ovato-conica; calyx post anthesin non inflatus.

9. *T. BOCCONI* (Savi. obs. trif. p. 37. n. 14.) caule erecto subramoso, foliolis oblongo-obovatis apice denticulatis, stipulis longissimè angustissimè subulatis, capitulis ovatis densifloris geminis sessilibus terminalibus basi bracteatis, laciniiis calycinis subæqualibus angustis corollam æquantibus. ① in rupestribus Corsicæ, Occitanie, agri Ruscinon., Andegavie. — Fl. fr. 5. p. 560 et T. collinum. fl. fr. 5. p. 560. T. gemellum Lapeyr. abr. T. semiglabrum Brot. phyt. t. 63. f. 2. (v. s.)
10. *T. STRIATUM* (L. sp. 1085.) caule ramoso, foliolis obovato-oblongis apice denticulatis, stipulis membranaceis latis breviter apiculatis, capitulis ovato-conicis densis terminalibus lateralibusque solitariis sessilibus basi bracteatis, tubo calycis ventricoso, laciniiis parvis inæqualibus divaricatis aristatis corollà brevioribus. ① in rupestribus Gallie occident., mediæ australisque

passim, circa Nancy (Moumier). — Fl. fr. 4. p. 53a. — Vaill. bot. t. 33. f. 2. (v. v.)

11. *T. SCABRUM* (L. sp. 1084.) caule procumbente, foliolis obovatis denticulatis, stipulis parvis submembranaceis, capitulis ovatis sessilibus terminalibus lateralibusque, lacinii calycinis lineari-lanceolatis rigidis recurvis subtrinnerviis inæqualibus corollam polypetalam vix æquantibus. ① in siccis et arenosis provinc. occident., mediarum et australium passim. — Fl. fr. 4. p. 53a. — Vaill. bot. t. 33. f. 1. (v. v.)

### Sect. III. EUTRIPHYLLUM. Ser. I. c.

Flores capitati; capitula ovata sæpè bracteata; calyx non inflatus villosus.

12. *T. MARITIMUM* (Huds. ang. ed. 1. p. 284.) caule erecto patulo, foliolis oblongo-obovatis obtusis emarginatisve subdenticulatis, stipulis angustis longè subulatis, capitula ovato-globois subsessilibus, lacinii calycinis tubo corollæ polypetalæ brevioribus rigidis inæqualibus, inferiore 3-nervi longiore, reliquis 1-nerviis. ① in Gallia australi et occidentali. — T. irregulare Pourr. et Fl. fr. 4. p. 531. — Pluckn. t. 113. f. 4. (v. v. DC.)

β. *Bastardianum* (Ser.) spicis pedunculatis, lacinii calycinis majoribus tenuiter nervosis. (v. s.)

13. *T. SUPINUM* (Savi trif. p. 46. f. 2.) caulibus patulis ramosis, foliolis obovatis latis ciliatis, stipulis angustis nervosis parvis subulatis, capitulis subovatis pedunculatis post anthesin ovoideis, lacinii calycinis inæqualibus linearibus rigidis patentibus corollæ brevioribus. ① circa Nemausum, Monspelium. — Fl. fr. 5. p. 558. (v. s.)

14. *T. XATARDII* (DC. fl. fr. 5. p. 568.) caule erecto villosa ramoso, foliolis obovato-oblongis subciliatis, stipulis angustis nervosis, capitulis ovatis pedunculatis, calycibus striatis glandulosis pilosisque, lacinii vix inæqualibus corollæ minoribus. ① in Pyrenæis orient. (v. s.)

15. *T. CINCYRUM* (DC. cat. monsp. p. 152.) caulibus erectis pilosis subramosis, foliolis ovali-oblongis villosis, stipulis latis nervosis duabus superioribus subpalmatis, capitulis ovatis pedunculatis, calycibus striatis, lacinii inæqualibus, inferiore maximâ corollæ minore. ① circa Monspelium (Saltmann). — Fl. fr. 5. p. 559. Fl. ochroleuci. (v. s.)

16. *T. OCHROLEUCUM* (L. syst. nat. 3. p. 233.) pilosum, caulibus ascendentibus gracilibus, foliis distantibus, foliolis ovato-ellipticis obtusis retusisve ciliatis, superioribus angustioribus, stipulis angustis nervosis petiolo multò brevioribus, capitulis terminalibus ovato-oblongis subsessilibus, calycibus costatis glabrisculis, lacinii lineari-setaceis inæqualibus rigidiusculis semipatentibus corollâ multò brevioribus, inferiore longissimâ. ② in pratis siccis passim. — Fl. fr. 4. p. 428. — Fuohs. hist. 818. ic. (v. v.)

β. *saxatile* (DC.) minus villosum, foliolis latis, inferioribus emarginatis.

*T. ochroleucum humile* Ser. (v. v. DC.)

γ. *dichotomum* (Ser.) caule supernè furcato, spicis geminis approximatis. (v. s.)

17. *T. BARBATUM* (DC. cat. monsp. 152.) pilosissimum, caule subramoso crasso, foliolis lanceolatis acutis integris ciliatis, stipulis basi latis nervosis lanceolato-acuminatis petiolo longioribus, capitulis terminalibus pedunculatis ovato-oblongis crassis, calycibus hirsutis sublævibus, lacinii setaceis, infimâ cæteris duplò longiore corollam æquante. ② circa Monspelium ex Saltzm. — Fl. fr. 5. p. 557. *T. pannonicum* β. Ser. in DC. prod. 2. p. 194. (v. s.)

18. *T. SQUARROSUM* (L. sp. 1082.) caule ramoso, foliis supremis oppositis alternisve subpilosis, foliolis lanceolatis aut ovalibus subretusis, stipulis angustis glabrisculis submembranaceis longè acuminatis, capitulis ovatis, calycibus costatis lacinii inæqualibus 3-nerviis ciliatis, inferiore corollæ longitudine de-

- flexa. ① in Corsicâ, circâ Parisios, in Galloprovinciâ ex Ger. — Fl. fr. 4. p. 53r. — Moris. hist. 2. s. 2. t. 13. n. 1. (v. s.)
19. *T. ALPESINUM* (L. sp. 1082.) caule erecto simplici, foliolis lanceolatis coriaceis integris valdè nervosis, stipulis angustis subsessilibus apice longè lineari instructis, capitulo ovato, calycibus leviter multistriatis, lacinii inæqualibus spinulosis, infimâ corollâ monopetalâ brevior, reliquis brevissimis denticulatis. ② in montosis Delph., agri Ligericini (Prost.), Pyren. — Fl. fr. 4. p. 527. Jacq. obs. 3. t. 64. (v. v.)
20. *T. MEDIUM* (L. faun. suec. ed. 2. p. 558.) glabriusculum, caule flexuoso ramoso, foliolis oblongis coriaceis ciliatis multinerviis, stipulis angustis apice lineari-lanceolatis, capitalis subglobosis subpedunculatis, lacinii calycinis inæqualibus pilosiusculis, inferiore reliquis duplò longiore et corollâ monopetalâ brevior. ② in sylvis. — Fl. fr. 4. p. 526. eng. bot. t. 190. (v. v.)
21. *T. PRATENSE* (L. sp. 1082.) caulibus ascendentibus, foliolis ovatis obtusis retusive integris, stipulis latis nervosis glabris breviter acuminatis inflexis, capitalis ovatis obtusis subsessilibus, lacinii calycinis subpilosis subæqualibus corollâ gamopetalâ duplò triplòve brevioribus. ② in Alpibus pratisque frequens. — Fuchs. 817. ic. Foliola sæpè maculâ centrali notata. (v. v.)
- f. multifidum* (Ser.) lacinii calycinis 6-7 piloso-hispidis. In Corsicâ. (v. s.)
22. *T. MICROPHYLLUM* (Desv. journ. bot. 2. p. 316.) glabriusculum, caulibus ascendentibus, foliolis ovatis obtusis retusive denticulatis, superioribus mucronulatis, stipulis latiusculis nervosis breviter acuminatis inflexis, capitalis ovatis obtusis sessilibus, lacinii calycinis pilosis inæqualibus corollâ gamopetalâ brevioribus. ② in nemoribus siccis Pictaviæ et Parisiorum. — Fl. fr. 5. p. 656. An *T. pratensis* var? (v. v.)
23. *T. DIVERGNUM* (Ehrh. beytr. 7. p. 165.) molliter villosum glaucum, caule diffusum, foliolis ovato-lanceolatis obtusis retusive, stipulis angustis apice lineari-lanceolatis, capitulis ovatis subhispidis, lacinii calycinis setaceis villosissimis rectis æqualibus corollâ gamopetalâ brevioribus, legumine circumscisso dispermo. ① in arenosis circâ Parisios. — Wald. et Kit. hung. 1. t. 50. *T. ciliosum* Thuill. par. ed. 2. p. 380. Fl. fr. 4. p. 525. An sequentis var? (v. v.)
24. *T. HIRTUM* (All. auct. p. 20.) caule erecto, foliolis obovato-cuneatis subintegris, stipulis infimis angustis longè acuminatis, superioribus brevibus latissimis, capitalis subglobosis hispidis, lacinii calycinis longissimis corollam subæquantibus vix inæqualibus. ① in Corsicâ, agro Monspelienâ et Ruscionensi. — T. hispidum Desf. atl. 2. t. 209. f. 1. Fl. fr. 4. p. 524. (v. v. DC.)
- f. pictum* (Ser.) foliolis denticulatis rubro-maculatis. *T. pictum* Roth. (v. s.)
25. *T. CHELLERI* (L. amæn. 4. p. 286.) caule procumbente, foliolis obcordatis integrinerviis, stipulis latis apice linearibus incurvis, capitulis globosis hispidis sessilibus, stipulis a bractæformibus rotundatis, lacinii calycinis hispidis corollâ longioribus. ① in maritimis Corsicæ et Gallie mediterraneæ. — Fl. fr. 4. p. 524. — Barr. ic. 859. (v. v.)
26. *T. STELLATUM* (L. sp. 1083.) villosum, caule diffuso, foliolis obcordato-cuneatis subtriangularibus margine superiore denticulato, stipulis obovatis latissimis serrulatis, capitulis globosis, lacinii calycinis foliaceis lineari-lanceolatis patentibus æqualibus longitudine corollæ, leguminibus 1-spermis. ① in sterilibus, ad vias provinc. mediterr., circâ Dax ex Thore, Burdigalam ex Laterrade. — Fl. fr. 4. p. 530. — Barr. ic. p. 860. (v. v.)
27. *T. LEUCANTHUM* (Bieb. fl. taur. 2. p. 214.) caulibus erectiusculis villosis, foliolis obovato-oblongis apice subserratis, stipulis lanceolato-subulatis integerrimis, capitalis subglobosis pedunculatis, lacinii calycinis patentibus subæqualibus corollâ brevioribus, leguminibus 1-spermis. ① in Corsicâ reperit Ph. Thomas. (v. s.)

28. *T. saxatile* (All ped. t. 59. f. 3.) caule erectiusculo, foliis villosis, foliolis cuneatis apice emarginatis parvulis integris, stipulis ovatis mucronulatis nervosis, capitulis globoso-depressis sessilibus bracteatis, calycibus tomentosis, laciniiis minimis æqualibus longitudine corollæ. (1)(2) in editioribus Alpibus Delphinatûs. — Fl. fr. 4. p. 523. (v. s.)

#### Sect. IV. TRIFOLIASTRUM. Ser. I. c.

Flores capitati sæpè post anthesin deflexi, capitula globosa, calyces non inflati.

29. *T. suffocatum* (L. mant. 276.) glabrum, caulibus brevissimis procumbentibus, petiolis longis, foliolis obcordato-cuneiformibus denticulatis, stipulis scariosis subnervosis breviter acuminatis, capitulis axillaribus sessilibus subglobosis glabris approximatis, floribus densis sessilibus, calycis laciniiis lanceolatis acutis subæqualibus patentibus corollâ multo longioribus, leguminibus 2-spermis. (1) in siccis Corsicæ, Nicæ, Telouis, Monspelii, Aginui ex St.-Am., Andegaviæ ex Bast. — Fl. fr. 4. p. 521. Brot. phyt. t. 64. (v. v. DC.)
30. *T. elomeratum* (L. sp. 1084.) glabrum, caule patulo, foliolis obovatis argutè dentatis, stipulis scariosis nervosis longè acuminatis, capitulis axillaribus sessilibus distantibus globosis, floribus densis sessilibus, laciniiis calycinis dentiformibus ovatis brevibus acuminatis corollâ polypetalâ brevioribus, leguminibus dispermis. (1) in pratis siccis et petrosis Galliæ mediterraneæ et austro-occidentalis, propè Falaise (de Brébisson). — Fl. fr. 4. p. 522. — Barr. ic. 882. (v. v.)
31. *T. parviflorum* (Ehrh. beit. 7. p. 165.) caule diffuso, foliolis obovatis nervosis argutè serratis, stipulis scariosis angustè acuminatis, capitulis axillaribus pedunculatis sessilibusque globosis, floribus densis sessilibus, laciniiis calycinis nervosis lanceolatis acutis inæqualibus, 2 superioribus corollâ longioribus, leguminibus dispermis. (1) in Pyrenæis orient. propè Saillagouze ex cl. Arnott. (v. s. ex Hispaniâ.)
32. *T. stractum* (L. sp. 1079.) glabrum, foliolis inferioribus obovatis, superioribus oblongo-ellipticis serrulatis, stipulis subscariosis latissimis obtusiusculis, capitulis axillaribus longè pedunculatis globosis, bracteis membranaceis sublanceolatis, floribus densis sessilibus, laciniiis calycinis subulatis inæqualibus corollâ brevioribus, leguminibus subexsertis dispermis. (1) in pratis siccis Fontis-Blesudi, Aginui ex St.-Am., Galloprov., Corsicæ. — Fl. fr. 4. p. 520. — Michel. gen. t. 25. f. 7. (v. s.)
33. *T. repens* (L. sp. 1080.) caulibus repentibus subradicantibus diffusis basi ramosis, foliolis obovato-subrotundis subretusis denticulatis, stipulis scariosis angustis lanceolatis longè mucronatis, capitulis axillaribus longissimè pedunculatis, floribus pedicellatis post anthesin deflexis, laciniiis calycinis inæqualibus dentiformibus corollâ brevioribus, leguminibus 4-spermis. (1) in pratis. — Fl. fr. 4. p. 520. — J. Bauh. hist. 2. p. 380. f. 3. (v. v.)
34. *T. cespitosum* (Reyn. mem. 1. p. 162.) glaberrimum, caulibus cespitosis brevissimis, foliolis obovatis subretusis denticulatis, stipulis scariosis angustis lanceolatis acutissimis 1-nerviis, capitulis axillaribus longè pedunculatis, floribus subsessilibus post anthesin patentibus, laciniiis calycinis subæqualibus lanceolatis corollâ brevioribus, leguminibus 3-4-spermis. (1) in pratis Alpium et Pyrenæorum. — Fl. fr. 4. p. 521. T. Thalii Vill. dauph. 3. t. 41. (v. v.)
35. *T. angulatum* (Wald. et Kit. hung. 1. t. 27.) glabriusculum, caule decumbente angulato, foliolis obovatis obcordatisve argutè serratis, stipulis scariosis angustis longè acutis, capitulis axillaribus pedunculatis, floribus pedicellatis post anthesin deflexis, laciniiis calycinis angustis acutis corollâ brevioribus, leguminibus exsertis 4-spermis subarticulatis. (1) circà Mons-peliom. — Fl. fr. 5. p. 555. (v. s.)
36. *T. pyramum* (Savi pis. 2. p. 90.) glabrum, caule ascendente solido, foliolis obcordato-cuneiformibus denticulatis, stipulis latis submembranaceis angus-

- tissimis acutis, capitulis subumbellatis axillaribus paucifloris, floribus pedicellatis post anthesin deflexis, laciniis calycinis inæqualibus tubo brevioribus latiusculis corollâ multò brevioribus, leguminibus exsertis 4-spermis. ① in provinciis australibus. — Fl. fr. 5. p. 555. Michel. nov. gen. t. 25. f. 6. T. pallescens Schreb? T. intermedium Lapeyr. ab. 437. (v. s.)
37. T. MICHELIANUM (Savi fl. pis. 2. p. 139.) caule ascendente fistuloso, foliis obovatis obcordatisque serrulatis, stipulis foliaceis lanceolatis acutis, capitulis subumbellatis axillaribus, floribus longè pedicellatis post anthesin deflexis, laciniis calycinis setaceis subpatentibus tubo duplò triplòve longioribus et corollâ brevioribus, leguminibus exsertis 2-spermis. ① in Corsicâ et Galliâ occident. — Fl. fr. 5. p. 554. — Mich. gen. t. 25. f. 2. et 5. — Vaill. bot. t. 22. f. 5. (v. s.)
38. T. ELEGANS (Savi pis. p. 161. t. 1. f. 2.) caule ascendente solido, foliolis obovatis denticulatis, stipulis foliaceis longè angustèque acuminatis, capitulis globosis axillaribus, floribus breviter pedicellatis post anthesin deflexis, laciniis calycinis subæqualibus triquetris tubo longioribus corollâque brevioribus, leguminibus dispersis. ½ in pratis Galliæ mediæ, orientalis et occidentalis. — Fl. fr. 5. p. 554. Vaill. bot. par. t. 22. f. 1. Flores pallidè rubri. (v. s.)
39. T. MONTANUM (L. sp. 1087.) pubescens, caule erecto subsimplici, foliolis lanceolato-oblongis obtusis denticulatis valdè nervosis, stipulis lanceolatis acutissimis, capitulis subglobosis axillaribus pedunculatis dein oblongis, floribus subsessilibus congestis patentibus dein deflexis, laciniis calycinis inæqualibus angustis tubi longitudine et carinâ brevioribus, leguminibus monospermis. ½ in pratis montosis. — Fl. fr. 4. p. 529. — J. Bauh. 2. p. 380. f. 2. Fl. albi. (v. v.)
40. T. BALBINIANUM (Ser. in DC. prod. 2. p. 201.) villosum, caulibus cespitosis simplicibus, foliolis ellipticis obtusis denticulatis valdè nervosis, stipulis lanceolatis acutis, capitulis hemisphæricis terminalibus solitariis, pedunculis caule multò longioribus, laciniis calycinis æqualibus patulis angustis tubo cylindrico multò longioribus et carinâ brevioribus. ½ in monte Lachen Galloprovinciæ ex Ser. — Flores purpurei.
- Sect. v. VESICASTRUM. Ser. l. c. p. 202.
- Flores capitati densi, calycis labium superius post anthesin accrescens inflatum.
41. T. SUBTERRANEUM (L. sp. 1080.) villosum, caulibus procumbentibus, foliolis obcordatis denticulatis, stipulis lanceolatis latis acutis, capitulis paucifloris subglobosis post anthesin subterraneis, florum inferiorum fertiliū calycibus inflatis, superiorum steriliū vacnis elongatis deflexis, leguminibus 1-spermis. ① in graminosis, collibus et ad oras sylvarum Galliæ australis, occidentalis et circâ Parisios. — Fl. fr. 4. p. 522. — Barr. ic. t. 881. (v. v.)
42. T. VESICULOSUM (Savi pis. 2. p. 165.) caule erecto firmo striato, foliolis lanceolatis argutè serrulatis, stipulis angustis submembranaceis longè acuminatis, capitulis ovatis crassis longè pedunculatis, calycibus scariosis inflatis, laciniis subulatis acutissimis æqualibus corollâ multò brevioribus, leguminibus 2-spermis. ① in Corsicâ. — Fl. fr. 5. p. 561. Lois. gall. 2. t. 15. T. recurvum. Walde. et Kit. 2. t. 165. Fl. rubelli. (v. s.)
43. T. SPUMOSUM (L. sp. 1085.) caule patulo ramoso, foliolis obovatis obcordatisve integræsculis, stipulis latis acutissimis, capitulis ovatis crassis subsessilibus, calycibus scariosis inflatis striatis transversè venosis, laciniis setaceis divergentibus corollam subsequantibus, bracteis lanceolatis calyce brevioribus, leguminibus 4-spermis. ① in Corsicâ, Galloprovinciâ ex Ger. Occitan. ex Gouan. — Fl. fr. 4. p. 533. — Cupan. panphyt. t. 69. f. 1. (v. s.)
44. T. RESUPINATUM (L. sp. 1086.) caule prostrato, foliolis obovatis argutè ser-



rubatis basi maculatis, stipulis brevibus lineari-lanceolatis acutis, capitulis hemisphericis dein globosis, pedunculis petiolo brevioribus, floribus sessilibus, calycibus post anthesin inflatis scariosis longitudinaliter transversèque venosis, lacinii lanceolatis acutissimis corollâ brevioribus, leguminibus subspermis. ① in herbosis siccis provinc. austral. et rariis occidentaliis. — Fl. fr. 4. p. 534. — Clns. hist. 2. p. 247. ic. (v. v.)

45. *T. FRAGILE* (L. sp. 1086.) caule prostrato, foliolis ovatis obovatisve, stipulis angustis linearibus elongatis, capitulis ovato-globosis longè pedunculatis, floribus sessilibus, calycibus post anthesin inflatis membranaceis reticulatis pilosis, lacinii inæqualibus corollâ brevioribus, leguminibus 2-spermis. ② ad vias et in graminosis passim. — Fl. fr. 4. p. 534. — Vaill. bot. t. 22. f. 3. (v. v.)

46. *T. TOMENTOSUM* (L. sp. 1086.) caule prostrato, foliolis obcordato-cuneiformibus argutè serrulatis, stipulis lanceolatis acutis scariosis, capitulis globosis brevissimè pedunculatis, floribus sessilibus, calycibus post anthesin inflatis membranaceis tomentosis, lacinii corollâ brevioribus, leguminibus 1-spermis. ② (f) in herbosis maritimis Gallie mediterraneæ. — Fl. fr. 4. p. 534. — Barr. ic. 864. (v. v.)

#### Sect. VI. LUPINASTER. *Mæsch. meth. supp. p. 30.*

Flores maximè petala persistentia; lacinie calycium subulata erectæ.

47. *T. UNIFLORUM* (L. sp. 1085.) caule brevissimo, foliolis ovatis acuminatis dentatis nervosis, stipulis vaginantibus longè acuminatis, floribus strobilatis breviter pedunculatis, calycibus cylindricis striatis, lacinii brevibus subulatis aequalibus, corollâ calyce multò longiore, leguminibus 2-spermis. ② in siccis maritimis circa Masiliam. — Fl. fr. 4. p. 556. — Roxb. cent. 3. t. 51. f. 1. (v. s.)

48. *T. ALBUM* (L. sp. 1080.) glaberrimum, caulibus erectissimis crassis subterraneis, petiolis longis, foliolis ternis lanceolato-linearibus obtusiusculis denticulatis, stipulis longissimis angustissimis linearibus acutis, capitulis umbellatis longè pedunculatis, pedicellis minimis subverticillatis, calycibus campanulatis, lacinii æqualibus corollâ multò brevioribus, leguminibus 2-spermis. ② in pascuis editiorum montium. — Fl. fr. 4. p. 519. — J. Bauh. hist. 2. p. 376. f. 1. Corollâ purpurea. (v. v.)

#### Sect. VII. CHAEROSIUM. *Ser. in dc. prod. 2. p. 204.*

Flores in capitulum ovatum pedunculatum dispositi; petala scariosa flava, post anthesin badia deflexa persistentia.

49. *T. RADIUM* (Schreb. in Sturm. deutsch. fl. 1. fasc. 16. ic.) caulibus ascendentibus, foliis petiolatis, foliolis sessilibus obcordatis denticulatis, stipulis lanceolatis acutis, capitulis pedunculatis subglobosis laxiusculis, vexillo obovato, calycibus brevissimis campanulatis, lacinii inæqualibus superiore minimâ, leguminibus subglobosis 1-spermis. ② in pratis editiorum montium. — Fl. fr. 5. p. 561. *T. spadiceum* Vill. et Fl. fr. 4. p. 535. non L. — Barr. ic. 1024. (v. v.)

50. *T. AGRARIUM* (L. sp. 1087.) caule ascendente ramoso firmo, foliis breviter petiolatis, foliolis oblongo-ovatis sessilibus denticulatis, stipulis foliaceis lanceolatis acutis petiolo longioribus, capitulis longè pedunculatis ovoideis, vexillo obcordato, calycis brevis campanulati lacinii inæqualibus glabris elongatis superiore minore, leguminibus 1-spermis. ① in pratis subhumidis. — Fl. fr. 4. p. 585. — Vaill. bot. t. 22. f. 4. (v. v.)

51. *T. SPADICEUM* (L. sp. 1087.) caule erecto subsimplici gracili, foliis petiolatis, foliolis oblongo-ovatis sessilibus denticulatis, stipulis foliaceis angustis acuminatis, capitulis longè pedunculatis ovoideis, vexillo obcordato, lacinii calycis inæqualibus exterioribus longis pilosis, 2 interioribus minimis glabris, leguminibus ovoideis compressis 1-spermis. ② in pratis montosis. — Fl. fr. 5. p. 561. Sturm. deutsch. 1. fasc. 16. ic. (v. v.)

52. *T. racemosum* (L. sp. 1088.) caulibus procumbentibus, foliis breviter petiolatis, foliolis obovatis obcordatisve denticulatis terminali petiolulato, stipulis ovatis ciliatis petiolo duplò brevioribus, capitulis axillaribus ovatis, pedunculis folia superantibus æquantibusve, floribus numerosis densis, laciniiis calycinis inæqualibus, 2 superioribus brevissimis, leguminibus 1-spermis. ① in arvis et pratis siccis. — Fl. fr. 4. p. 436. — Vaill. bot. t. 22. f. 3. (v. v.)
- β. *campestre* (Ser.) caule erecto ramoso, pedunculis longitudine foliorum. — T. campestre Schreb. in Sturm. deutsch. 1. fasc. 16. ic. Fl. fr. 5. p. 562. (v. v.)
- γ. *nanum* (Ser.) caule minimo, capitulis paucis, pedunculo terminali longitudine foliorum, foliolis ovato-subcordatis.
53. *T. Parisiense* (DC. fl. fr. 5. p. 562.) caulibus procumbentibus, foliis breviter petiolatis, foliolis obovatis obcordatisve denticulatis, terminali sæpè petiolulato, stipulis obovatis acuminatis ciliatis petiolo plerumque subæquantibus, pedunculis filiformibus longissimis, capitulis subumbellatis paucifloris, laciniiis calycinis inæqualibus, 2 superioribus brevissimis tubo longioribus, leguminibus pedicellatis 1-spermis. ① circa Parisios; Burdigalam (Des Moulins), Tolosam et in Pyrenæis ex Benth., in Galloprovincia. (v. s.)
54. *T. filiforme* (L. sp. 1088.) caule diffuso, foliolis obovatis obcordatisve subdenticulatis terminali petiolulato, stipulis latis ovatis petioli longitudine, pedunculis filiformibus longis, floribus subumbellatis sessilibus, laciniiis calycinis inæqualibus, 2 superioribus brevissimis tubo brevioribus, leguminibus 1-2-spermis substipitatis. ① in pratis. — Fl. fr. 4. p. 556. Eng. bot. t. 2257. — Ray. syn. t. 14. f. 4. (v. v.)

### XIII. DORYCNIUM. Tourn. inst. 391. t. 211. f. 3. Ser. in DC. prod. 2. p. 208.

Calyx subcampanulatus bilabiatus sursum subgibbosus; alæ vexillo breviores; carina submutica; stylus rectus; stigma oculo nudo capitatum. — Herbæ vel suffrutices.

1. *D. rectum* (Ser. l. c.) villosum, caulibus erectis suffruticosis, foliis petiolatis, foliolis obovatis mucronatis, stipulis ovato-cordatis, pedunculis bracteatis multifloris folio vix duplò longioribus, pedicellis lanatis calyce longitudine, laciniiis angustissimis tubo longioribus corollæque brevioribus, leguminibus calyce duplò longioribus. 3 in provinciis australibus. — Lotus rectus L. sp. 1092. Fl. fr. 4. p. 557. — Barr. ic. t. 544. Fl. rosei. (v. v.)
2. *D. hirsutum* (Ser. l. c.) incano-tomentosum, caule erecto suffruticoso, foliis sessilibus, stipulis lanceolatis, pedicellis fructiferis folio duplò longioribus, bracteis lanceolatis calycem æquantibus, pedicellis calyce multò brevioribus, laciniiis subulatis tubo multò longioribus et corollà brevioribus, leguminibus calyce vix longioribus. 3 in provinciis australibus. — Lotus hirsutus L. sp. 1091. Fl. fr. 4. p. 556. — J. Bauh. hist. 2. p. 360. ic. Fl. magni rosei. (v. v.)
- β. *incanum* (Ser.) foliis sericeo-incanis, capitulis breviter pedunculatis densioribus. — In locis apricis Nicæ et Stæchadam. — L. sericeus. Fl. fr. 5. p. 573. (v. v.)
3. *D. parviflorum* (Ser. l. c.) hispidum, caulibus subprostratis, foliolis lanceolatis, stipulis ovatis, capitulis 4-5-floris, bracteis lanceolato-linearibus, pedunculis folio longioribus, laciniiis calycinis corollam æquantibus, leguminibus oblongis calyce vix longioribus. ① in Corsicâ et insulis Stæchadam. — Lotus parviflorus Desf. atl. 2. t. 211. DC. ic. rar. t. 30. L. hispidus. Fl. fr. 4. p. 556. non Desf. — Fl. parvi flavi. (v. s.)
4. *D. merbaorum* (Vill. dauph. 3. p. 417. t. 41.) caulibus herbaceis, foliolis stipulisque obovatis obtusis, calycibus pilosis, dentibus calycinis tubo multò brevioribus, capitulis longè pedunculatis, bracteis monophyllis à capitulo distantibus, leguminibus ovatis calyce duplò triplòve longioribus po-

*lypermis*. *ℒ* circa Nicæam, in Delphinatâ propè Gratianopolim ex Vill. — *Fl. fr.* 4. p. 558. (v. v. DC.)

5. *D. suffruticosum* (Vill. l. c. p. 416.) caulibus suffruticosis, foliolis stipulisque lineari-lanceolatis acutis, calycibus pilosis, dentibus tubo multò brevioribus, capitulis longè pedunculatis, bracteis 1-3-phyllis, leguminibus globosis calyce duplò longioribus monospermis. *♂* in provinciis australibus et austro-occidentalibus usque ad Santoniam (Des Moulins.) — *Fl. fr.* 4. p. 557. — *Lob. ic.* 2. p. 51. f. 1. 2. (v. v.)

XIV. LOTUS. *Seringe in DC. prod.* 2. p. 209. — *Loti sp. Lin.*

Calyx tubulosus 5-fidus; alæ vexillum subæquantes; carina rostrata; legumen cylindraceum vel compressum apterum; stylus rectus; stigma oculo nudo subulatum. — *Herbæ.*

#### SECT. I. KROKZIA. *Mœnch. meth.* t. 43.

Legumen turgidum curvatum, flores 1-3.

1. *L. axialis* (L. sp. 1090.) pilosus, caulibus erectis, foliolis obovatis, floribus 1-3, bracteis ovatis calycem æquantibus, leguminibus glabris. (1) in incultis et sterilibus maritimis Corsicæ, Nicæ, Galloprovinciæ. — *Fl. fr.* 4. p. 554. — *J. Bauh.* 2. p. 365. f. 2. (v. v.)

SECT. II. LOTEA. *Medic. phil. bot. fasc.* 1. p. 204.

Legumen compressum longum, flores subumbellati.

2. *L. oanthorhomboides* (L. sp. 1091.) glabriusculæ, caulibus diffusis, foliolis obovato-rhomboidis integris, stipulis ovatis, floribus 3-5, bracteis calycem multò longioribus, leguminibus compressis submentaceis glabris curvulis flexis. (1) in maritimis Corsicæ, Nicæ, Galloprovinciæ, Occitanizæ Gousn, agri Syrtici? ex Thore. — *Fl. fr.* 4. p. 554. *Cav. ic.* 2. p. 48. t. 163. — *J. Bauh.* 2. p. 369. f. 1. Flores flavi. (v. v. DC.)

SECT. III. EULOTUS. *Ser. l. c.* p. 210.

Legumen cylindraceum longum, flores corymbosi.

3. *L. caeticus* (L. sp. 1091.) sericeus, caulibus suffruticosis suberectis, foliolis obovatis, stipulis ovatis pedunculo fructifero triplò brevioribus, bracteis lanceolato-linearibus calyce minoribus, capitulis sub-4-floris, laciniiis calycinis lanceolatis longitudine tubi corollæque multò brevioribus, stylo exserto, leguminibus teretibus glabriusculis nutantibus subtorulosis. *♂* circa Bonifacio reperierunt Requien et Soleirol. — *Cav. ic.* p. 44. t. 156. Flores flavi. (v. v.)
4. *L. cymoides* (L. sp. 1092.) caulibus basi prostratis subramosis, pedicellis calycibusque incanis, foliolis obovatis subcarnosis pilis adpressis tectis, laciniiis calycinis tubo brevioribus, leguminibus subtorulosis glabris. (1) in maritimis Corsicæ, Nicæ, Galloprovinciæ, agri Ruscimonensis ex Lapeyr. — *Fl. fr.* 4. p. 555. *All. ped.* t. 20. f. 1. — *Barr. ic.* t. 1031. Fl. flavi. (v. v.)
5. *L. xilosissimus* (Poir. supp. 3. p. 504) caulibus ramosis subprostratis, foliolis ovato-lanceolatis, stipulis ovatis subacutis, capitulis paucifloris longè pedunculatis, bracteis angustissimis lanceolatis, laciniiis calycinis angustis acutis tubo longioribus, leguminibus subteretibus glabris. (1) in Galliâ australi ex Poiret. — *Cor. latea* exsiccatione viridis. An var. *L. hispidus*? †
6. *L. hispidus* (Desf. cat. par. 190.) caulibus numerosis prostratis, foliolis hispidis oblongo-lanceolatis, stipulis ovatis, capitulis paucifloris, pedunculis folio multò longioribus, bracteis lanceolatis longitudine calycis, laciniiis angustis tubo longioribus corollæque brevioribus, leguminibus compressis punctatis. (1) in Corsicâ et agro Syrtico. — *Loisel. gall.* 490. t. 16. *Fl. fr.* 5. p. 572. (v. v.)
7. *L. angustissimus* (L. sp. 1090.) caulibus prostratis folisque hispidis,

foliolis stipulisque oblongo-linearibus, capitulis 1-3-floris, pedunculis folio longioribus, lacinii calycinis tubo longioribus corollâque brevioribus, leguminibus compressis angustissimis subsolitariis calyce sextuplò longioribus. (I) in incultis et siccis provinc. austral. occidentaliumque passim. — Fl. fr. 5. p. 573. — J. Bauh. hist. 2. p. 356. f. 2. (v. v.)

β. *diffusus*, minus hispidus, caulibus elongatis, pedunculis folio plerumquè triplò longioribus. — L. *diffusus* Soland. Sm. eng. bot. t. 925. Fl. fr. 5. p. 573. Ser. in DC. prod. 2. p. 213. (v. v.)

8. L. *COMPLANATUS* (Brot. phyt. p. 127. t. 53.) glabriusculus; caulibus subramosis procumbentibus, foliis glaucescentibus, foliolo intermedio obovato, lateralibus lanceolatis stipulisque ovatis apice ciliatis, pedunculis brevissimis 1-floris, lacinii calycinis angustis longitudine tubi corollâque brevioribus apice ciliato-aristatis, leguminibus cylindricis longissimis arcuatis glabris. (I) in Corsicâ et Galloprovinciâ. — L. *aristatus* DC. fl. fr. 5. p. 572. Fl. albi cum carinâ purpureâ. (v. s.)

9. L. *COARCTULATUS* (L. sp. 1092.) caulibus prostratis, foliolis obovatis vel linearibus glabris pilosisve, stipulis ovatis, bracteis lanceolatis linearibusve, pedunculis longissimis, capitulis depressis 8-10-floris, calycibus campanulatis, lacinii acutis longitudine tubi et corollâ multò brevioribus, leguminibus teretibus. 2/ in pratis frequens. — Fl. fr. 4. p. 555. Fl. flavi sæpè extus rubri exsiccatione virides.

α. *arvensis* (Ser.) glabriusculus, caulibus subprostratis, foliolis obovatis, floribus luteis. — Oed. fl. dan. t. 99r. (v. v.)

β. *major* (Ser.) caulibus erectis plus minusve pilosis majoribus fistulosis. In sepiis humidis. — Sow. eng. bot. t. 209r. (v. v.)

γ. *villosus* (Ser.) caulibus foliisque villosis erectis. L. *villosus* Thuil. (v. v.)

δ. *crassifolius* (Pers.) pilosus, foliolis ovatis subcarnosis, caulibus foliosis prostratis. — In sterilibus maritimis. (v. v.)

ε. *Alpinus* (Schleich.) caulibus foliisque minimis, floribus sæpè extus ruberrimis. In Alpibus. (v. v.)

ζ. *tenuifolius* (Poll.) caulibus suberectis filiformibus, foliolis stipulisque lineari-lanceolatis. — In rudertis. L. *tenuis* Kit.

#### XV. TETRAGONOLOBUS. Scop. carn. 2. p. 87. Ser. in DC. prod. 2. p. 215.

Calyx tubulosus 5-fidus; alæ vexillo breviores; carina rostrata; stylus flexuosus, stigma infundibuliforme demum obliquè rostratum; legumen cylindraceum alis foliaceis 4 marginatum. — Herbæ.

1. T. *RUPESTRIS* (Mœnch. meth. p. 164.) pilosus, caulibus subdecumbentibus, foliolis obovatis, floribus solitariis geminatisve, bracteis calyce longioribus, leguminibus glabris latè alatis. (I) circâ Niceam ex All. — Lotus tetragonolobus L. et Fl. fr. 4. p. 553. — J. Bauh. hist. 2. p. 258. f. 2. Fl. atro-purpurei. (v. v. c.)

2. T. *SILIQVOSUS* (Roth. germ. 1. p. 323.) pilosus, caulibus subdecumbentibus, foliolis obovatis, stipulis obtusiusculis, floribus solitariis longè pedunculatis, bracteis obovato-linearibus calyce brevioribus, leguminibus glabris angustissimè alatis. 2/ in pratis humidis. — L. *siliquosus* L. sp. 1089. Fl. fr. 4. p. 553. — J. Bauh. 2. p. 359. f. 2. Flores flavi. (v. v.)

β. *maritimus* (DC.) foliis carnosis glabriusculis. — In maritimis. L. *maritimus* L. (v. v.)

3. T. *CONJUGATUS* (Ser. l. c.) pilosus, caulibus subdecumbentibus, foliolis obovatis, stipulis parvis acuminatis, floribus conjugatis, bracteis obovatis calyce longioribus, leguminibus glabris subteretibus angustissimè alatis. (I) circâ Monapeliū? ex L. et Gouan. Valletrium? ex Lapeyr. — L. *conjugatus* L. et Fl. fr. 4. p. 554. Fl. purpurei. Dubia civis. (v. s. c.)

Subtrib. III. *CLITORIÆ*. DC. prod. 2. p. 216. leg. mem. vi.

Legumen 1-loculare. Stam. sæpius diadelpha. Caules herbacei aut suffrutescentes.

XVI. *PSORALEA*. L. gen. n. 894.

Sepala 5 ad medium concreta in calycem 5-fidum persistentem, lobis acuminatis, infimo paulò productiore; legumen cnlycis longitudine evalvè 1-spermum interdum in rostrum desinens.

1. *P. PALMETINA* (Gou. ill. 51.) foliis pinnatim 3-foliolatis, foliolis inferiorum ovatis, superiorum lanceolatis, petiolis pubescentibus sulcatis, pedunculis axillaribus folio triplò quadruplòve longioribus, spicis capitatis, calycibus pubescentibus demum subvesicariis. 2/ circà Monspelium propè Pont-Juvénal ex seminibus in lanis extraneis allatis. — Fl. fr. 5. p. 554. Jacq. vind. 2. t. 184. Sequenti similis sed subinodora. (v. v.)
2. *P. BITUMINOSA* (L. sp. 1075.) foliis pinnatim 3-foliolatis, foliolis ovato-lanceolatis, petiolis pubescentibus lævibus, pedunculis axillaribus folio triplò quadruplòve longioribus, spicis capitatis, calycibus pubescentibus. 2/ 3/ in provinciis australibus usque ad Lugdunum ex cl. Balbis. — Fl. fr. 4. p. 518. Lam. ill. t. 614. f. 1. Bitumen tedolens. (v. v.)

Subtrib. IV. *GALEGÆ*. Bronn. diss. DC. prod. 2. p. 243.

Legumen 1-loculare. Stamina diadelpha, rariùs monadelpha. Caules herbacei fruticosi aut arborei.

XVII. *GLYCYRHIZA*. Tourn. inst. 210. L. gen. n. 882.

Calyx nudus tubulosus 5-fidus bilabiatus; cor. vexillum ovato-lanceolatum rectum, carina biceps aut dipetala recta acuta; stam. diadelpha; stylus filiformis; legumen compressum 1-loculare 1-4-spermum.

1. *G. GLABRA* (L. sp. 1046.) foliolis ovatis subrétusis subtùs subglutinosiis, stipulis nullis, spicis pedunculatis folio brevioribus, floribus distantibus, leguminibus glabris 3-4-spermis. 2/ in provinciis australioribus. — Fl. fr. 4. p. 560. Lam. ill. t. 625. f. 2. (v. v. c.)

XVIII. *GALEGA*. Tourn. inst. t. 222. Lam. ill. t. 625.

Calyx dentibus 5 subulatis subæqualibus; corolla vexillo obovato oblongo; carina obtusa; stam. monadelpha, decimo nempe ad medium concreto; stylus filiformis glaber; stigma terminale punctiforme; legumen teretiusculum torulosum obliquè striatum.

1. *G. OFFICINALIS* (L. sp. 1063.) foliolis lanceolatis mucronatis glabris, stipulis lato-lanceolatis, racemis folio longioribus. 2/ in glareosis Gallie passim. — Fl. fr. 4. p. 560. Mill. ic. t. 137. (v. s. ex Pedem.)

XIX. *ROBINIA*. DC. prod. 2. p. 261. — Robinie spec. Lin.

Calycis dentes 5 lanceolati, 2 superiores breviores approximati; cor. papilionacea vexillo amplo, carina obtusa; stam. diadelpha decidua; ovarium 16-20-ovulatum; stylus anticè barbatus; legumen compressum subsessile polyspermum, valvis planis tenuibus, suturà seminiferà marginatà.

1. *R. PSEUDACACIA* (L. sp. 1043.) spinis stipularibus, ramis virgatis, racemis laxis pendulis leguminibusque glabris, foliolis ovatis. 3/ ex Americà boreali in Galliam ubi nunc subspontanea introducta. — Fl. fr. 4. p. 561. Lam. ill. t. 606. f. 1. (v. v.)

XX. COLUTEA. *R. Brown. hort. kew. ed. 2. vol. 4. p. 325. — Colutea sp. L.*

Calyx 5-dentatus; vexillum explanatum bicalcosum carinâ obtusâ majus; stamina diadelpa (9 et 1); stigma laterale sub apice styli uncinato; stylus posticè longitudinaliter barbatus; legumen stipitatum cymbiformi-ovatum inflatum scariosum. — Frutices.

1. *C. ARBORÆSCENS* (L. sp. 1045.) foliolis ellipticis retusis, pedunculis subseffloris, vexilli gibbis abbreviatis, leguminibus clausis. § ad sepes in Galliâ australi. — Fl. fr. 4. p. 561. Duham. arb. ed. nov. 1. t. 22. (v. v.)

Subtrib. V. *ASTRAGALÆ*. Adans. fam. DC. prod. 2. p. 273.

Legumen suturâ alterâ introflexâ longitudinaliter 2-loculare aut semibiloculare. Stamina diadelpa (9 et 1). Caules herbacei aut suffrutescentes. Folia pinnata.

XXI. PHACA. *L. gen. n. 891.*

Cal. 5-dentatus, dentibus 2 superioribus remotioribus; carinâ obtusa; stylus imberbis; stigma capitatum; legumen subturgidum uniloculare suturâ superiore seminiferâ tumidâ.

\* *Floribus albidis aut ochroleucis.*

1. *P. ALPINA* (Jacq. ic. rar. 1. t. 151.) caule erecto ramoso pubescente, stipulis lineari-lanceolatis, foliolis 9-12-jugis oblongis obtusis pubescentibus, leguminibus compressis semi-ovatis acutis glabrisculis. ¶ in apricis asperis Alpium et Pyrenæorum. — Fl. fr. 4. p. 562. — Till. pis. t. 914. f. 2. (v. v.)

\*\* *Floribus purpurascens aut albo et violaceo pictis.*

2. *P. GLABRA* (Clar. bull. philom. n. 61. DC. astr. n. 4.) caule ramoso prostrato glabro, stipulis latis ovatis, foliolis 6-7-jugis ovali-oblongis acutis glabris, pedunculis folio longioribus, alis integris, leguminibus stipitatis turgidis glabris. ¶ in Alpibus demissioribus Galloprovinciæ. Corolla alba apice et margine violacea. — Fl. fr. 4. p. 563. (v. s.)
3. *P. AUSTRALIS* (Linn. mant. 103.) caule ramoso cespitoso ascendente glabro, stipulis ovatis, foliolis 6-8-jugis lineari-lanceolatis glabris, impari sessili, pedunculis folio longioribus, alis apice bifidis carinâ longioribus, leguminibus stipitatis ovoideis demum glabris. ¶ in asperis Alpium et Pyrenæorum. — Fl. fr. 4. p. 563. Jacq. misc. 2. t. 3. Corolla pallida apice purpurascens. (v. v.)
4. *P. ASTRAGALINA* (DC. astr. n. 9.) caule brevi procumbente adnurgente glabresculo, stipulis ovatis acutis, foliolis 10-11-jugis ovatis adpressè utrinquè pubescentibus, pedunculis folio longioribus, alis integris carinâ brevioribus, leguminibus stipitatis pendulis, junioribus nigro-hirsutis. ¶ in editioribus Alpibus et Pyrenæis. — Fl. fr. 4. p. 564. — Schenck. itin. alp. 509. f. 7. Fl. violacei. (v. v.)

XXII. OXYTROPIS. *DC. astr. n. xv. ed. maj. p. 19. et 53. — Astragali spec. Linn.*

Calyx 5-dentatus; corollæ carina in mucronem exsertum summo dorso desinens; legumen biloculare aut semibiloculare, suturâ superiore introflexâ. — Herbæ.

- §. 1. Acaules, *nempè caulibus subnullis, stipulis petiolo adnatis, foliolis conjugatis.*

\* *Floribus purpurascens aut albis.*

1. *O. MONTANA* (DC. l. c. n. 1.) subacaulis villosa, pilis petiolorum et scapi

petalis, foliolis elliptico-lanceolatis, scapis vix folio longioribus, racemis abbreviatis, bracteis calyce dimidiò brevioribus, leguminibus erectis tereti-oblongis villosis stylo acuminatis semibilocularibus.  $\frac{1}{2}$  in Alpibus, Pyrenæis, montibus editioribus Arvernæ. — Fl. fr. 4. p. 565. Astragalus montanus L. Scop. carn. t. 45. Fl. cæruleo-purpurei. (v. v.)

2. *O. URALENSIS* (DC. astr. n. 3.) acaulis villosa-sericea, foliolis ovato-lanceolatis, scapis folio longioribus calycibusque hirsuto-lanatis, capitulis multifloris ovalis, floribus erectis, bracteis etiam infimis calyce paulò brevioribus, leguminibus confertis erectis ovatis acuminatis pubescentibus bilocularibus.  $\frac{1}{2}$  in herbosis fertilibus Alpium Galloprovinciæ et Pyrenæorum. — Fl. fr. 4. p. 565. Astr. Uralensis. Jacq. ic. rar. 1. t. 155. eng. bot. t. 466. Fl. cæruleo-purpurei. — Differre videtur ab *A. Uralensi* L. sp. 1071. Conf. DC. prod. 2. p. 276. (v. v. DC.)

\*\* *Floribus ochroleucis.*

3. *O. CAMPESTRIS* (DC. astr. n. 10.) acaulis, foliolis multijugis lanceolatis sericeis, scapo aspiùs decumbente foliis subæquali, spicis capitatis, bracteis calyce paulò brevioribus, floribus erectis, leguminibus erectis ovatis inflatis pubescentibus semibilocularibus.  $\frac{1}{2}$  in declivibus herbosis Alpium et Pyrenæorum. — Fl. fr. 4. p. 566. — Hall. helv. t. 13. (v. v.)

$\beta$ . *viscosa* (DC.) foliolis pubescentibus subviscosis. — Astr. viscosus. Will. danph. 3. t. 43. (v. v.)

4. *O. FORTIDA* (DC. astr. n. 11.) acaulis, foliolis multijugis lanceolato-linearibus glabris viscosis, scapo foliis sublongiore superne lanato, spicis capitatis paucifloris, bracteis calyce brevioribus, leguminibus erectis tereti-inflatis subincurvis pubescentibus.  $\frac{1}{2}$  in editioribus Alpibus Delphinatûs ex Villars. — Fl. fr. 4. p. 566. Astr. fortidus. Will. danph. 3. t. 43. A. Halleri. All. (v. v. in Helv.)

§. 2. *Caulescentes, nempe caulibus elongatis, stipulis caulinis nec petiolo adnatis, foliolis conjugatis.*

5. *O. PILOSA* (DC. astr. n. 27.) caulescens erecta molliter pilosa, foliolis lanceolatis acutis, pedunculis axillaribus folio longioribus, spicis ovato-oblongis, leguminibus erectis tereti-subulatis villosis.  $\frac{1}{2}$  in rupestribus montosis Galliæ austro-orientalis. — Fl. fr. 4. p. 567. — Astr. pilosus. L. Jacq. austr. t. 51. Fl. ochroleuci. (v. v.)

XXIII. *ASTRAGALUS*. DC. astr. n. v. ed. maj. p. 22 et 79. — Astragali spec. Linn.

Calyx 5-dentatus; corolla carinâ obtusâ; stamina diadelpa; legumen biloculare aut semibiloculare, suturâ inferiore introflexâ. — Herbæ aut suffrutices.

#### Series I. PURPURASCENTES.

*Stipulis à petiolo liberis, floribus purpurascentibus.*

§. 1. *HYPOGLOTTIDEI*. DC. prod. 2. p. 281.

Floribus purpureis aut albis, stipulis à petiolo liberis inter se coalitis et ideò quasi bifidis et oppositifoliis.

1. *A. HYPOGLOTTIS* (Linn. mant. 274.) diffuso-procumbens piloso-subvillosus, stipulis (saltem junioribus) concretis oppositifoliis, foliolis obovatis oblongis sæpè emarginatis 8-10-jugis, spicis capitatis, pedunculis folio longioribus, bracteis calycis tubo dimidiò brevioribus, calycibus nigro-pilosis, leguminibus erectis capitatis hirsutis ovato-triquetris, localis monospermis.  $\frac{1}{2}$  in collibus arenosis passim. — Fl. fr. 4. p. 570. DC. astr. n. 18. t. 14. (v. v.)
2. *A. PURPUREUS* (Law. diet. 1. p. 314.) diffuso-procumbens subvillosus, stipulis (saltem junioribus) concretis oppositifoliis, foliolis obovatis apice

bidentatis, apicis capitatis, pedunculis folio longioribus, leguminibus erectis capitatis hirsutis ovato-triquetris, localis trispermis.  $\mathcal{V}$  in montanis Delphinatûs, Galloprov., Niocœ, agri Mimatensis (Prost.). — DC. fl. fr. 4. p. 569. astr. n. 17. t. 12. (v. s.)

3. A. *BAYONENSIS* (Lois. fl. gall. 474.) diffuso-procumbens pube adpressâ albâ subtomentosa, stipulis concretis oppositifoliis, pedunculis 4-6-floris folio subæqualibus, leguminibus subsessilibus tomidiusculis tomentosis latitudine duplò longioribus.  $\mathcal{V}$  in arenosis maritimis Gallis occid. à Bayonâ ad Ruppellam passim. — Fl. fr. 5. p. 573. A. *Austriacus* Thore non L. (v. v. D.)

### §. II. DISSITIFLORI. (DC. l. c.)

Stipulis nec petiolo nec inter se adnatis, floribus purpurascenscentibus aut albis distantibus nec densè spicato-racemosis, leguminibus rectis.

4. A. *AUSTRIACUS* (L. spec. 1070.) diffuso-procumbens, foliolis glabris linearibus truncato-emarginatis, racemis pedunculatis folio longioribus, alia corollâ bifidis, leguminibus subtriquetris pendulis.  $\mathcal{V}$  in rupestribus apricis Delphinatûs ad arces Brigantii. — Fl. fr. 4. p. 567. Jacq. Aust. t. 175. (v. v. DC.)

### §. III. ONOBRÛCHOIDEI. (DC. l. c.)

Stipulis à petiolo et inter se distinctis, floribus purpurascenscentibus densè spicatis capitativè, vexillis linearibus elongatis, leguminibus rectis rariùs falcatis, radicibus perennibus.

5. A. *ONONCHIS* (Linn. spec. 1070.) diffusus pubescens, foliolis oblongis 7-9-jugis, apicis oblongo-ovatis folio longioribus, vexillo linearis alis duplò longiore, leguminibus ovato-triquetris subvillosis erectis calyce duplò longioribus utrinquè 4-spermis, acumine brevi erecto.  $\mathcal{V}$  in pratis siccis montanis Delphinatûs, Galloprovinciis. — Fl. fr. 4. p. 571. — Clus. hist. 2. p. 238. f. 2. (v. v. in Helv. D.)

### §. IV. SESAMEI. (DC. l. c.)

Stipulis à petiolo et inter se liberis, floribus purpurascenscentibus densè spicatis capitativè, leguminibus rectis, radicibus annuis.

6. A. *PENTAOGOTTIS* (L. mant. 274.) diffuso-procumbens subvillosus, stipulis ovatis, foliolis obovatis retuso-emarginatis 4-6-jugis, pedunculis folio sublongioribus, leguminibus capitatis æmiovatis subtriquetris acutis tuberculato-echinatis, papillis sæpè piliferis, localis monospermis.  $\mathcal{V}$  in Corsicâ ex Valle, propè Cascastel in demissioribus Corbariensibus ex cl. Benth. — Fl. fr. 4. p. 569. Cav. ic. t. 188. (v. s. ex Hisp.)
7. A. *GLAUX* (L. sp. 197.) procumbens villosus subcanescens, stipulis ovato-lanceolatis, foliolis oblongis acutiusculis 8-9-jugis, pedunculis folio longioribus, vexillo lineari, leguminibus capitatis erectis ovato-triquetris mucronatis villosis, localis monospermis.  $\textcircled{1}$  in Castello dicto Bombax inter Avenionem et Cabellionem et propè Béziers. — Fl. fr. 4. p. 924. — Clus. hist. 2. p. 241. f. 1. (v. v. DC.)
8. A. *STELLA* (Gouan. ill. 50.) diffuso-procumbens cano-villosus, stipulis lanceolatis, foliolis elliptico-oblongis 8-10-jugis, pedunculis folii longitudine aut brevioribus, leguminibus capitatis demùm stellatis subteretibus dorso sulcatis mucronatis hirsutis, localis 5-10-spermis.  $\textcircled{1}$  Monspelii. — Fl. fr. 4. p. 568. — Pluck. t. 79. f. 4. Lœb. ic. 2. t. 83. (v. v. DC.)
9. A. *SESAMEUS* (L. sp. 1068.) diffusus villosus subcanescens, stipulis lanceolatis, foliolis elliptico-oblongis 9-10-jugis, capitulis axillaribus sessilibus, leguminibus capitatis erectis subteretibus acuminatis dorso sulcatis villosis, localis 7-8-spermis.  $\textcircled{1}$  circâ Telonem, Aquas-Sextias, Monspelium, Cascastel. — Fl. fr. 4. p. 568. — Garid. aix. t. 12. (v. m.)



§. v. VESICARIÆ. (DC. l. c.)

*Stipulis nec petiolo nec inter se adnatis, floribus purpurascensibus aut albidis, calycibus vesicariis.*

10. *A. VESICARIUS* (L. sp. 1071.) diffusoprocumbens pube adpressâ sericeo-canus, foliolis 5-7-jugis ellipticis, pedunculis folio longioribus, calycibus vesicariis pube nigrâ adpressâ et pilis albis patulis vestitis, leguminibus hirsutis calyce paulò longioribus.  $\mathcal{L}$  in Galloprovinciâ super. et Delphinatûs apricis asperis. — Fl. fr. 4. p. 569. Vill. dauph. t. 42. f. 1. (v. v.)

Series 11. OCHROLEUCI. (DC.)

*Stipulis à petiolo liberis, floribus ochroleucis.*

§. VI. BUCERATES. (DC. l. c.)

*Stipulis nec petiolo nec inter se adnatis, floribus ochroleucis, leguminibus sæpius arcuatis, radicibus annuis.*

11. *A. RAMOSUS* (L. sp. 1067.) diffusus pubescens, folioli cuneatis emarginatis supernè glabris, pedunculis sub-6-floris folio brevioribus, leguminibus hamatis dorso sulcatis teretibus apice subulatis, junioribus pubescentibus, adultis glabris. ① in siccis et lapidosis provinc. australiam et in Arvernâ propè Claromontem. — Fl. fr. 4. p. 572. — Moris. hist. s. 2. t. 9. n. 10. (v. v.)
12. *A. EPIGLOTTIS* (L. mant. 274.) procumbens villosio-incanus, stipulis lanceolatis calycibusque nigro-villosis, foliolis oblongo-linearibus 6-7-jugis, apicis capitatis brevissimè pedunculatis, leguminibus depressis subcoarctatis acuminatis margine subreplicatis patulis pubescentibus. ① in nemoribus et montibus Galloprovinciâ. — Fl. fr. 4. p. 673. — Riv. tetr. t. 109. (v. v. c.)
13. *A. BATICUS* (L. sp. 1068.) procumbens pubescens, stipulis membranaceis ovatis acuminatis, foliolis obovatis retosis 10-15-jugis, apicis brevissimè pedunculatis, leguminibus erectis triquetrop-rismaticis glabris rectis apice mucronato-uncinatis. ① in Corsicâ propè Bonifacio. — Riv. tetr. irr. t. 105. (v. s.)

§. VII. CICEAÏDEI. (DC. l. c.)

*Stipulis nec petiolo nec inter se adnatis, apicis pedunculatis, floribus ochroleucis, radicibus perennibus, leguminibus sessilibus.*

14. *A. GLYCYPHYLLOS* (L. sp. 1067.) procumbens glabrescens, stipulis ovatis mucronatis acuminatisve, foliolis 5-6-jugis ovatis, pedunculis folio brevioribus, apicis ovato-oblongis, leguminibus oblongo-subulatis subtriquetris subincurvis sessilibus glabris erectis.  $\mathcal{L}$  in pratibus, sepibus, dametis. — Fl. fr. 4. p. 572. — Riv. tetr. t. 103. (v. v.)
15. *A. DEPRESSUS* (L. sp. 1073.) subcaulis aut diffusus pubescenti-canescens, stipulis ovatis membranaceis, foliolis 9-11-jugis ovatis, pedunculis folio brevioribus, leguminibus tereti-depressis rectis patulo-deflexis glabris.  $\mathcal{L}$  in Pyrenæis et editioribus Alpibus Delphinatûs. — Fl. fr. 4. p. 571. All. ped. t. 19. f. 3. (v. v. DC.)
16. *A. CICKA* (L. sp. 1067.) diffusoprocumbens subpubescens, stipulis lanceolatis, foliolis 10-13-jugis elliptico-oblongis mucronatis, pedunculis folio brevioribus, floribus spicato-capitatis, leguminibus inflatis hirsutis mucronatis.  $\mathcal{L}$  in siccis montosis Galliæ orientalis passim. — Fl. fr. 4. p. 573. — Moris. hist. s. 2. t. 9. n. 9. (v. v.)

§. VIII. ALOPECUROÏDEI. (DC. l. c.)

*Stipulis petiolo non adnatis, caulibus erectis strictis, floribus flavis dense congestis in spicas ad foliorum axillas sessiles aut brevissimè pedunculatas.*

17. *A. NARBONENSIS* (Gouan. ill. 49. Pall. astr. t. 10.) caulescens erectus, caule molliter hirsuto, stipulis lanceolatis, foliolis oblongo-linearibus villosis, apicis subglobosis, dentibus calycinis setaceis corollâ brevioribus tubi

longitudine.  $\mathcal{L}$  in asperis circa Narbonam, Biterras (*Béziers*. ) — Fl. fr. 4. p. 574. (v. v. DC.)

Series III. TRAGACANTHACEI. (DC. l. c.)

Stipulis petiolo adnatis, petiolis indurato-persistentibus.

§. IX. TRAGACANTHÆ. (DC. l. c.)

Petiolis persistentibus spinescentibus, stipulis petiolo adnatis.

18. A. MASSILIENSIS (Lam. dict. 1. p. 320. DC. astr. n. 96) pedunculis sub-4-floris folio subæqualibus, calycibus cylindraceis obtusè et breviter 5-dentatis, foliolis 9-11-jugis ellipticis incanis.  $\mathfrak{z}$  in arenosis maritimis Massiliæ, Narbonæ. — Fl. fr. 4. p. 574. — Lob. ic. 2. t. 27. Fl. albi. (v. v.)
19. A. ARISTATUS (L'Her. stirp. 170.) pedunculis sub-6-floris brevissimis, calycis dentibus longis setaceis, foliolis 6-9-jugis oblongis macronatis pilosis, legumine vix semibiloculari.  $\mathfrak{z}$  in sterilibus rupetribus Alpium et Pyrenæorum. — Fl. fr. 4. p. 575. — Garid. aix. t. 104. Fl. albi aut purpurascens. (v. v.)

Series IV. PODOCHREATI.

Stipulis petiolo adnatis, petiolis non indurato-spinosis.

§. X. INCANI. (DC. l. c.)

Calycibus leguminibusque non vesicariis, vexillis valdè elongatis, leguminibus perfectè bilocularibus.

20. A. MONASPULANUS (L. sp. 1072.) subacaulis incano-pubescentis aut glabrescentis, foliolis 21-41 ovatis lanceolatisve, extimis subminoribus, scapis folia superantibus, calycis dentibus longis subulatis, leguminibus teretibus subulatis subarcanatis, adultis glabris.  $\mathcal{L}$  in provinciis australibus, Pyrenæis et propè Medhntam (Mantes), Rothomagus ex cl. Le Prévost. — Fl. fr. 4. p. 576. Curt. bot. mag. t. 219. Cam. epit. 929. ic. Variet floribus purpureis pellidis aut albis. (v. v.)
21. A. INCANUS (L. sp. 1072.) acaulis incanus, foliolis 13-17 ovatis, scapis folia vix superantibus, spicis capitatis, dentibus calycinis subulatis brevibus, leguminibus subcylindraceis pube brevissimâ canis apice incurvis subulato-macronatis calyce vix triplò longioribus, loculis 5-6-spermis.  $\mathcal{L}$  in campis apricis et lapidosis Gallie australis. — Fl. fr. 4. p. 376. — Magn. bot. p. 32. ic. (v. v. D.)

XXIV. BISERRULA. L. gen. n. 893.

Calyx 5-fidus; carina obtusa; stamina diadelpha; legumen depresso-planum biloculare valvis carinato-compressis dorso sinuato-dentatis. — Herba.

1. B. PELACIUS (L. sp. 1073.) diffusa pubescens, foliis impari-pinnatis, foliolis multijugis. ① in Galloprovincia orientali, et Mospelii ex Gouan. — Fl. fr. 4. p. 577. — Barr. ic. t. 1137. (v. s.)

Trib. III. HEDYSARÆ. DC. prod. 2. p. 307. leg. mem. VII.

Embryo homotropus. Corolla papilionacea, stamina rarò libera, sæpiùs monadelpha aut diadelpha. Legumen in loculos articulosve monospermos transversim divisum. Cotyledones planiusculæ per germinationem in folia stomatibus donata conversæ.

Subtrib. I. CORONILLÆ. DC. prod. 2. p. 308.

Flores umbellati. Legumina teretia aut compressa.

XXV. SCORPIURUS. *L. gen. n. 876. Gartn. fruct. 2. t. 155.*

Calyx 5-fidus, lobis æqualibus acutis; corolla carinā bicipiti; stamina diadelphæ (9 et 1) 5 longiora sub antherā paululum dilatata, ovarium sulcatum; stylus filiformis acutus; legumen revolutum subspirale, articulis constans 3-6 monospermis longitudinaliter sulcatis; nervis sæpè aculeatis muricatisve; embryo replicatus; cotyledones lineares. — Herbæ annuæ.

1. *S. MURICATA* (L. sp. 1050.) leguminis glabri costis interioribus lævibus, externis tuberculis minimis subobtusis muricatis. (1) ad vias et margines agrorum Monspeli, Nicæ, Delphinatūs ex Vill. — Fl. fr. 4. p. 601. — Moris. hist. 2. t. 11. n. 4. Valdè similis sequenti, sed leguminis costæ exteriores muricatæ nec aculeatæ. (v. s.)
2. *S. SULCATA* (L. sp. 1050.) leguminis glabri costis interioribus lævibus, 4 externis aculeos rigidulos distantes apice subuncinatos gerentibus. (1) in arvis et ad vias Galliæ australis. — Fl. fr. 4. p. 601. Gartn. fruct. 2. t. 155 (v. v.)
3. *S. SUBVILLOSA* (L. sp. 1050.) leguminis glabri costis interioribus lævibus, externis 6-8 aculeos rigidulos apice subuncinatos subconfertos gerentibus. (1) Hab. cum priore. — Fl. fr. 4. p. 602. — Moris. hist. 2. t. 11. n. 2. Legumen crassius et magis revolutum quàm in *S. sulcata*. (v. v.)
4. *S. ACUTIFOLIA* (Viv. fl. lyb. 43. t. 19. f. 4.) leguminis brevissimè hirti costis intimis lævibus, externis 8-10 aculeos rigidulos brevissimos confertos gerentibus. (1) in Corsicâ circa Bonifacio. — An satis à præcedente distincta? (v. s.)
5. *S. VERMICULATA* (L. sp. 1050.) leguminis glabri costis intimis subnullis, externis 10 tuberculos stipitatos confertos apice obtuse dilatatos gerentibus. (1) in provinc. australium arvis. — Fl. fr. 4. p. 601. — Moris. s. 2. t. 12. n. 3. Legumina crassa. (v. s.)

XXVI. CORONILLA. *Neck. elem. n. 1319. Lam. ill. t. 630. — Coronilla sp. L.*

Calyx campanulatus brevis 5-dentatus, dentibus 2 super. approximatis subcoarctatis; corolla petalorum unguibus calyce sæpius longioribus, carinā acutâ; stamina diadelphæ (9 et 1); legumen teretiusculum, gracile, demum in articulos oblongos 1-spermos secedens; semina ovata aut cylindracea. — Frutices aut Herbæ.

\*Sect. I. *EMERUS. Tourn. inst. t. 418. Desv. journ. bot. 3. p. 121. t. 4. f. 9.*

Legumen teretiusculum tardiùs et obscuriùs secedens, petalorum ungues calyce triplò longiores.

1. *C. EMERUS* (L. sp. 1146.) fruticosa, glabra, stipulis minimis, foliolis 5-7 obovatis, pedunculis 3-floris. ♂ in sepibus et dumetis Galliæ australis orientalisque. — Fl. fr. 4. p. 606. — Mill. ic. t. 132. f. 1 et 2. (v. v.)

Sect. II. *CORONILLA. Tourn. inst. t. 419. Desv. journ. bot. 3. p. 119. t. 4. f. 3.*

Legumen compressiusculum evidenter secedens; petalorum ungues vix calyce longiores.

\* *Floribus flavis.*

2. *C. JUNCEA* (L. sp. 1047.) fruticosa glabra, stipulis minimis, foliolis 3-7 lineari-oblongis obtusis subcarnosis, infimis à caule remotis, umbellis 5-7-floris. ♂ in collibus et dumetis Nicæ, Gailoprovinciæ. — Fl. fr. 4. p. 606. — Barr. ic. t. 133. J. Bauh. hist. 1. p. 2. p. 383. f. 2. Rami juncei teretes subnudi. (v. v. DC.)
3. *C. STIPULARIS* (Lam. dict. 2. p. 120.) fruticosa glabra, stipulis subrotundis maximis deciduis, foliolis 7-9 obovatis mucronulatis glaucis, infimis à caule

remotis, umbellis 6-8-floris.  $\beta$  propè Niceam (D. Stire). — C. Valentina bot. mag. t. 185. (v. v.)

4. C. GLAUCA (L. sp. 1047.) fruticosa glabra, stipulis parvis lanceolatis, foliolis 5-7 obovatis obtusissimis glaucis, infimis à caule remotis, umbellis 7-8-floris.  $\beta$  in maritimis prov. austral. præsertim in Galliâ Narbonensi et Occitanâ. — Fl. fr. 4. p. 607. Curt. bot. mag. t. 13. (v. s.)
5. C. MINIMA (L. sp. 1048.) suffruticosa diffusa glabra, stipulis concretis oppositifoliis bidentatis, summis majusculis membranaceis deciduis, foliolis 7-11 ovatis obtusis retusisve, infimis à caule remotis, umbellis 7-8-floris.  $\gamma$   $\beta$  in arenosis aridis submontanis Galliæ austro-orientalis, propè Falais ex cl. de Bréhisson. — Fl. fr. 4. p. 607. — J. Bauh. hist. 2. p. 351. f. 2. (v. v.)
6. C. OORONATA (L. sp. 1048.) suffruticosa suberecta aut ascendens glabra, stipulis concretis oppositifoliis bidentatis parvis, foliolis 5-9 obovatis submacronatis glaucis, infimis cauli approximatis, umbellis 8-10-floris.  $\gamma$   $\beta$  in aridis prov. austral., agri Ligericini, Arvernæ. — Fl. fr. 4. p. 608. — Clus. hist. 1. p. 98. f. 2. (v. v.)
7. C. MONTANA (Scop. carn. ed. 2. n. 912. t. 44.) herbacea erecta glabra, stipulis concretis oppositifoliis oblongis emarginatis deciduis, foliolis 7 ovatis macronatis subglaucis, inferioribus cauli approximatis, umbellis 15-20-floris.  $\gamma$  in Juraaso. — Fl. fr. 5. p. 533. — Riv. tetrap. t. 207. (v. s.)

\*\* *Floribus purpurascentibus aut albis.*

8. C. VARIA (L. sp. 1048.) herbacea diffusa glabra, stipulis minimis acutis, foliolis 9-13 oblongis macronatis, infimis cauli approximatis, umbellis 16-20-floris, leguminibus erectis.  $\gamma$  in pratibus et sterilibus. — Fl. fr. 4. p. 608. — Clus. hist. 2. p. 237. f. 2. (v. v.)

XXVII. ASTROLOBIUM. Desv. journ. bot. 3. p. 121. t. 4. f. 10. — Ornithopi sp. L.

Calyx ebracteatus tubulosus subæqualiter 5-dentatus; corollæ carina minima compressa; stamina diadelphe (9 et 1); legumen teretiusculum articulis plurimis 1-spermis indehiscentibus cylindraceis utrinquè truncatis constans. — Herbhæ.

1. A. EBRACTEATUM (DC. prod. 2. p. 311.) pedunculis folio subæqualibus apice ebracteatis, stipulis subnullis, foliis omnibus pinnatis, foliolis elliptico-oblongis multijugis æqualibus, infimis à caule distantibus, leguminibus teretibus arcuatis. (1) in Galliæ occidentalis australisque sabulosis passim. — Ornithopis ebracteatus Brot. lus. 2. p. 159. Fl. fr. 5. p. 582. — O. durus. Fl. fr. 4. p. 603, non Cav. — Dalech. hist. 1. p. 487. f. 1. (v. v.)
2. A. SCORPIOIDES (DC. l. c.) pedunculis folio longioribus ebracteatis, stipulis concretis vaginantibus, foliis 3-foliolatis, foliolis infimis minimis subrotundis cauli approximatis, terminali maximo ovali, leguminibus subincurvis subnodosis. (1) in arvis et segetibus Galliæ australis ex Nicæâ ad Aginnum. — Ornith. scorpioides L. sp. 1049. Cav. ic. t. 37. Fl. fr. 4. p. 603. (v. v.)

XXVIII. ORNITHOPUS. Desv. journ. 3. p. 121. t. 5. f. 14. — Ornithopi sp. L. gen. n. 884.

Calyx bracteatus tubulosus subæqualiter 5-dentatus; corollæ carina minima compressa; stamina diadelphe (9 et 1); legumen compressum articulis plurimis monospermis indehiscentibus utrinquè æqualiter truncatis ad margines parallelis constans. — Herbhæ.

1. O. COMPRESSUS (L. sp. 1049.) foliis villosis pedunculos subsuperantibus, foliolis lanceolatis acuminate subsessilibus, leguminibus compressis rugosis pubescentibus apice rostro adnato articulo ultimo longiore instructis, articulis ovalibus, calyce hirsuto 5-fido corollâ vix duplè brevior. (1) in arenosis provinciarum australium. — Fl. fr. 4. p. 603. Flores flavi. (v. v.)
2. O. ROSÆUS (Dufour in ann. sc. nat. t. 5. p. 84.) foliis pedunculo brevior-

ribus villosis, foliolis lanceolatis acuminatis subsessilibus, leguminibus subcompressis rugosis glabris apice rostro recto articulo ultimo breviori instructis, articulis rotundis, calyce hirsuto quinquefido corollâ duplò breviori. (1) in sabulosis agri Syrtici propè Mont-de-Marsan. — *O. perpusillus*  $\beta$ . intermedius. DC. prod. 2. p. 312. non Roth. An *O. sativus* Brot. lueit. 2. p. 160? Fl. rosei. (v. s.)

3. *O. perpusillus* (L. sp. 1049.) pedunculis folia superantibus subæquantibusve, foliis villosis, foliolis subsessilibus ovatis obtusis aut vix mucronatis, leguminibus vix compressis hirsutis rostro recto brevissimo terminatis, articulis rotundis, calyce hirsuto 5-dentato, dentibus brevibus tubo quintuplò brevioribus. (1) in sabulosis umbrosis passim. — Fl. fr. 4. p. 602. eng. bot. t. 369. Flores ex rubro albo et paulinùm flavo mixti. (v. v.)

$\beta$ . *intermedius* (Roth. germ. 2. p. 2. p. 215. non DC. prod.) major, caulibus vix pubescentibus, foliis subpetiolatis, bracteis leguminibus majoribus. (Roth. l. c.)

#### XXIX. HIPPOCREPIS. L. gen. n. 885.

Calyx 5-fidus, lobis æqualibus acutis; corolla carinâ bicipiti; stamina diadelphea (9 et 1); stylus filiformis acutus; legumen constans articulis plurimis 1-spermis in ferri equini formam curvis; undè legumen ad latus superius sinubus plurimis rotundatis quasi incisum videtur; semina cylindracea aut compressa oblonga ad partem articuli cuiusque mediam adfixa curva, umbilico in medio curvatura.

##### §. 1. *Radice perenni, leguminibus sinuatis nempe articulis tantum curvis et sinus latè apertos efficientibus.*

1. *H. comosa* (L. sp. 1050.) glanduloso-pubescent, caulibus herbaceis diffusis, pedunculis folio longioribus apice umbelliferis, leguminibus subarcuatis, articulis 2-3 ad semina puberulis. (1) in cretaceis et calcareis vulgaris. — Fl. fr. 4. p. 605. — Garid. six. t. 34. (v. v.)

$\beta$ . *scorpioides*, legum. angustioribus, articulis ad semina glanduloso-hispidis. — In Galloprovinc. et Occitan. *H. scorpioides* Req. in h. DC. (v. s.)

##### §. 2. *Radice annua, leguminibus orbiculariter incisis, nempe articulis sinus subrotundos angustè apertos efficientibus.*

2. *H. multisiliquosa* (L. sp. 1030.) pedunculis folio paulò brevioribus 2-4-floris, leguminibus glaberrimis circulariter incurvis. (1) in provinciis australibus indicata sed forsàn per confusionem cum *H. ciliatâ*. — Fl. fr. 4. p. 605. — Ferrum equinum polyceratum. Col. ecphr. p. 300. ic. (v. s. c.)

3. *H. velutina* (Req. in h. DC.) pedunculis folio paulò brevioribus 1-2-floris, leguminibus pube adpressâ velutinis ad semina glanduloso-hirtis circulariter incurvis, sinuum extremitatibus in cornu productis. (1) circâ Monspelium propè Pont-Juvenal reperit èl. Requier. (v. s.)

4. *H. unisiliquosa* (L. sp. 1050.) floribus axillaribus subsessilibus solitariis, leguminibus subincurvis glabris aut in parte mediâ articulorum puberulo-hirtis. (1) in apricis lapideis provinc. australium. — Fl. fr. 4. p. 604. Lam. ill. t. 630. f. 3. (v. v.)

5. *H. ciliatâ* (Wild. mag. nat. ges. Berol. 1808. p. 173.) pedunculis 4-2-floris folio subbrevioribus, leguminibus arcuatis ad semina glanduloso-hispidis ad alterum latus subciliatis. (1) in Galloprovincia circâ Avenionem et probabiliter in omnibus provinciis australibus. (v. v. D.)

#### XXX. SECURIGERA. DC. fl. fr. 4. p. 609.

Calyx brevis bilabiatus, labio inf. 3-partito, super. bidentato; petalorum ungues calyce sublongiores; carina acuta; stamina diadelphea (9 et 1); legumen compresso-planum, suturis prominulis, inter se-

mina isthmis notatum, sed ob suturas prominulas non articulatam, lineare in rostrum longum apice desinens; semina 8-10 compressa parallelogramma.

1. *S. CORONILLA* (DC. l. c.) ① in arvis Niceæ, Corsicæ. — *Coronilla Securidaca* L. sp. 1048. — Gærtn. fruct. 2. t. 153. f. 3. (v. v.)

Subtrib. II. *EUHEDYSARÆ*. DC. prod. 2. p. 313.

Flores racemosi. Legumina compressa.

- XXXI. *HEDYSARUM*. *Jaum. journ. bot.* 3. p. 61. excl. spec. DC. prod. 2. p. 340.

Calyx 5-fidus laciniis lineari-subulatis subæqualibus; corolla vexillo amplo, carinâ obliquè truncatâ, alis carinâ multò brevioribus; stamina diadelphâ (9 et 1); genitalium fasciculus abruptè infractus; legumen constans articulis plurimis compressis monospermis orbiculatis lenticularibusve regularibus medio inter se adnexis et ideò ad utramque suturam convexis.

Sect. I. *ECHINOLOBIUM*. DC. l. c.

Leguminum articuli in disco villosi costati muricati aut glochidiati.

1. *H. CORONARIUM* (L. sp. 1058.) caule diffuso, foliolis 3-5-jugis ellipticis aut subrotundis subtus et margine pubescentibus, apicis ovatis confertis, alis calyce duplò longioribus, leguminis articulis 2-5 orbiculatis aculeatis, glabris. ② ex Italiæ pratis indigenum, in hortis sæpè cultum. — Fl. fr. 4. p. 610. — Dod. pempt. 549. (v. v. c.)
2. *H. HUMILE* (L. sp. 1058.) caule erectiusculo, foliolis 7-9-jugis lineari-cuneiformibus obtusis, floribus racemosis, alis carinâ dimidiò brevioribus, vexillo carinâ brevior, leguminis articulis 2-3 orbiculatis junioribus tomentosis tuberculato-muricatis et villosis. ② in Cebennis, circà Monspelium ex Gouan, *Aqua-Sextias* ex Ger. — Fl. fr. 4. p. 610. — J. Bauh. hist. 2. p. 336. f. 1. (v. v. DC.)
3. *H. SPINOSISSIMUM* (L. sp. 1058. excl. syn.) caule diffuso, foliolis 5-7-jugis obovatis aut oblongis emarginatis glabriusculis, apicis capitatis paucifloris, alis carinâ  $\frac{1}{2}$  brevioribus, vexillo carinam æquante, leguminis articulis 2-3 orbiculatis pubescentibus aculeatis, aculeis apice hamulosis. ① in arenosis maritimis Galloprovinciæ, Niceæ. — Fl. fr. 5. p. 588. (v. v. D.)
4. *H. CAPITATUM* (Desf. atl. 2. p. 177. non Burm.) caule decumbente, foliolis 6-7-jugis oblongis acutis, apicis ovatis laxis, alis calyce duplò longioribus, leguminis articulis orbiculatis aculeatis villosis. ① ex Corsicâ acceptumq. misit el. Solier. — Sims. bot. mag. t. 1251. *H. Corsicum* Balb. (v. s.)

Sect. II. *LEIOLOBIUM*. DC. l. c.

Leguminum articuli læves nec echinati nec pilosi; calycis laciniae corollâ breviores.

5. *H. OSCURUM* (L. sp. 1057.) caule erecto, foliolis ovatis glabris 5-9-jugis, stipulis etiam summis concretis oppositifoliis vaginantibus, bracteis pedicello longioribus, leguminibus pendulis glaberrimis. ② in glareosis ad torrentes editiorum Alpium Delphin. — Fl. fr. 4. p. 610. — Hall. enum. 1. t. 14. hist. 2. t. 12. *H. controversum* Grants aust. 425. t. 2. f. 3. (v. v. in Pedem.)

- XXXII. *ONOBRYCHIS*. *Tourn. inst.* 2. 211. *Gærtn. fruct.* 2. t. 148. DC. fl. fr. 4. p. 511.

Calyx 5-fidus, laciniis subulatis subæqualibus; corolla papilionacea; carina quasi obliquè truncata; alae breves; stamina diadelphâ (9 et 1); legumen sessile 1-articulatum, compressum 1-spermum indehiscens subcoriaceum echinatum cristatum aut alatum, latere superiore crassiore recto, inferiore convexo tenuiore.

Sect. I. EUBRYCHIS, DC. prod. 2. p. 344.

Legumen obliquum in disco rugosum aculeatumve, dorso dentatum aut interruptè cristatum.

1. O. SATIVA (Lam. fl. fr. 2. p. 652.) caule erectiusculo, stipulis sæpius distinctis, foliis cuneato-lanceolatis mucronatis glabris, spicis elongatis, carinâ vexillo brevior, alia calyce brevioribus, leguminis pubescentis dorso denticulato, lateribus rugosis subaculeatis. ½ in cretaceis, siccis, collibus. — Fl. fr. 4. p. 611. — Jacq. aust. t. 352. (v. v.)
2. O. MONTANA (DC. fl. fr. 4. p. 611.) caule subdecumbente, stipulis coalitis oppositifoliis, foliis cuneato-lanceolatis mucronatis glabris, spicis abbreviatis, carinâ vexillo longiore, alia calyce brevioribus, leguminis dorso denticulato, lateribus rugosis pubescentibus. ½ in Alpium et Pyrenæorum pratis elatis. — An prioris varietas? (v. v.)
3. O. SUPINA (DC. fl. fr. 5. p. 612.) caule diffuso subhirsuto, foliis oblongis vix mucronulatis, spicis ovato-oblongis pedunculatis, alia calyce brevioribus, carinâ calyce brevior, leguminis villosi dorso cristato dentato, discis rugosis subaculeatis. ½ in pratis et ruderalis Delph., Galloprov. super., Occitanis, Cebenn., Pyren. orient. et Santonis (Des Moulins). (v. v.)
4. O. SAXATILIS (All. ped. n. 1191. t. 19. f. 1.) caule ascendente, foliis linearibus mucronatis, spicis cylindricis pedunculatis, alia calyce longioribus, carinâ vexillo paulò brevior, leguminis glabri dorso cristato integro, lateribus rugosis edentulis. ½ in apricis, collibus, petrosis Galloprovinciæ, Delphinatûs australis, Nicææ. — Fl. fr. 4. p. 612. Variat foliis glabris et villosis, leguminibus glabris et tenuissimè pubescentibus. (v. v.)
5. O. CAPUT-GALLI (Lam. fl. fr. 2. p. 651.) caule erecto aut diffuso, foliis oblongis aut cuneato-obovatis mucronatis pubescentibus, spicis paucifloris, alia calyce sublongioribus, carinâ vexillo paulò brevior, leguminibus subpubescentibus undiquè aculeatissimis. ½ in Galliæ australis apricis asperis. — Fl. fr. 4. p. 613. — Lob. ic. 2. p. 81. f. 1. (v. v.)
6. O. CRISTA-GALLI (Lam. fl. fr. 2. p. 652.) caule supino, foliis cuneatis oblongis aut obovatis obtusis retusive pubescentibus, spicis paucifloris, calycibus corollæ subæqualibus, alia et carinâ vexillum subæquantibus, leguminis glabri dorso cristato, cristâ in lacinias planas oblongas dentatas partitâ, disco rugoso subaculeato. ① in sterilibus Nicææ ex All., Galloprovinc. austral. ex Gerard. — Fl. fr. 4. p. 613. Gartn. fruct. 2. t. 148. (v. v. c.)

Trib. IV. VICIÆ. Bronn. diss. p. 331. DC. prod. 2. p. 353. leg. mem. VII.

Corolla papilionacea. Stam. diadelphe (9 et 1). Legumen continuum. Cotyledones crassæ farinaceæ per germinationem immutata subterrâ intra spermodermium inclusæ remanentes. Radicula inflexo-curva. — Folia abruptè-pinnata (excepto Cicer), petiolo communi apice in setulam cirrhumve desinente basi super caulem non articulato.

XXXIII. CICER. Tourn. inst. 389. t. 210. f. 2. Linn. gen. n. 1189. Gartn. fruct. 2. p. 326. t. 151.

Calyx 5-lobus, tubo ad latus superius plus minùs basi gibbo, lobis acuminatis 2-4 superioribus vexillo incumbentibus; legumen turgidum dispermium, semina gibbosa mucronata.

1. C. ARIETINUM (L. sp. 1040.) foliis impari-pinnatis, foliis ovatis serratis æqualibus, stipulis lanceolatis subdenticulatis, calycis vix gibbosi lacinia alarum longitudine. ① Cultum præcipuè in Galliâ australi. — Fl. fr. 4. p. 600. — Cam. epit. 204. ic. (v. v.)

XXXIV. FABA. *Tourn. inst. t. 222. DC. fl. fr. 4. p. 598.* — *Viciæ sp. L.*

Omnia Viciæ; sed legumen grande coriaceum subtumidum; seminibus oblongis; hilo terminali.

1. *F. VULGARIS* (Mœnch. meth. p. 130.) foliis crassis, foliolis 2-5 ovalibus mucronatis, stipulis semisagittatis ovalibus, cirrhis subnullis, dentibus calycinis sublinearibus. ① è finitimis Caspii maris orta et in hortis et agris culta. — *Fl. fr. 4. p. 598. Vicia Faba. L. sp. 1039. Potamogeton bifolium Lapeyr. (v. v.)*

XXXV. VICIA. *Tourn. inst. t. 221. Juss. gén. 360. DC. fl. fr. 4. p. 589.*

Calyx tubulosus, 5-fidus aut 5-dentatus, dentibus 2 superioribus brevioribus; corolla papilionacea; stylus filiformis angulum ferè rectum cum ovario conficiens, supernè et infrà apicem subtus villosus; legumen oblongum, 1-loculare, polyspermum; semina hilo laterali ovali aut lineari.

#### §. 1. Floribus pedunculatis.

1. *V. FISIFORMIS* (L. sp. 1034.) glaberrima, foliis 4-jugis, foliolis ovato-cordatis obtusis reticulato-venosis distantibus, infimis sessilibus, stipulis semisagittatis dentatis parvis, pedunculis multifloris longitudine folii, dentibus calycinis subæqualibus tubo brevioribus, leguminibus oblongis compressis, seminibus globosis. ② in sylvaticis Alsatiæ, Burgundiæ ex Dur., Galloprovinciæ ex Ger. — *Fl. fr. 4. p. 589.* — *Clus. hist. 2. p. 229. f. 1. Fl. ochroleuci. (v. s.)*
2. *V. DUMETORUM* (L. sp. 1035.) glabriuscula, foliolis reflexis ovato-lanceolatis mucronulatis reticulato-venosis alternis, stipulis semisagittatis dentatis, pedunculis multifloris folii longitudine, laciniis calycinis brevissimis subæqualibus, stylis apice barbatis, leguminibus oblongis compressis, seminibus subrotundis. ② in sylvis et dometis Galliæ orientalis passim, Monspelii ex Gou. — *Fl. fr. 4. p. 590.* — *J. Bauh. 2. p. 316. f. 1. Corollæ violaceæ. (v. s.)*
3. *V. SYLVATICA* (L. sp. 1035.) glabra, foliolis numerosis alternis vel oppositis elliptico-oblongis mucronulatis tenuissimè reticulato-venosis, stipulis semisagittato-reniformibus setaceo-dentatis, pedunculis folio longioribus multifloris, laciniis calycinis tubi vix longitudine, stylis apice villosulis, leguminibus oblongo-linearibus compressis apice incurvis, seminibus globosis. ② in sylvis montosis Corsicæ, Galloprovinciæ, Occitanicæ. — *Fl. fr. 4. p. 590.* — *Hall. helv. t. 12. f. 2. Fl. albido-violacei. (v. s.)*
4. *V. CASSUBICA* (L. sp. 1035.) foliolis oppositis alternisve numerosis ellipticis mucronatis subglabris, nervis numerosissimis firmis, stipulis inferioribus semisagittato-lanceolatis, superioribus linearibus subintegris, pedunculis multifloris folium æquantibus, racemis confertis secundis, dentibus calycinis inæqualibus tubo brevioribus, stylis apice villosis, leguminibus coriaceis vix reticulatis oblongis brevibus compressis glabriusculis, seminibus globosis. ② in collibus herbosis.
  - a. *glabriuscula* (Ser. in DC. prod. 2. p. 356.) caule piloso, petiolis cirrhosis villosis, foliolis glabris. — *V. multiflora* Pollich? DC. fl. fr. 5. p. 577.
  - V. *cassubica* Sturm. deutsch. 1. fasc. 31. — in Andegaviâ. (v. s.)
  - β. *Orobis* (Ser. l. c.) tota pilosa, foliolis pilosis vel ciliatis, cirrhis nullis vel subfoliaceis. — *V. Orobis*. *Fl. fr. 5. p. 577. Orobis sylvaticus. Fl. fr. 4. p. 586. O. aristatus* Lapeyr. supp. — In pratis montosis Arverniciæ, Cebenn., Pyren. (v. v. DC.)
5. *V. CRACCA* (L. sp. 1095.) caulibus ramosis, foliis apice cirrhosis, foliolis numerosis oblongis alternis oppositisque puberulis mucronatis, nervis subparallelis simpliciusculis, stipulis semisagittato-linearibus, pedunculis multifloris folium æquantibus vel superantibus, racemis confertis secundis, dentibus calycinis inæqualibus, superioribus brevissimis inferioribus tubo brevioribus, stylis apice pilosis, leguminibus oblongis coriaceis compressis



- glabris, seminibus globosis.  $\mathcal{L}$  in arvis. — Fl. fr. 4. p. 591. — Riv. tetrap. t. 50. Fl. violacei. (v. v.)
6. *V. GERARDI* (Jacq. fl. austr. t. 229<sup>2</sup> DC. fl. fr. 4. p. 591. excl. syn.) caulibus simpliciusculis, foliis apice cirrhosis, foliolis numerosis oblongo-lanceolatis alternis oppositive sublanatis mucronatis, nervis subparallelis simpliciusculis, stipulis semisagittato-linearibus, pedunculis multifloris folio sublongioribus, racemis confertis secundis, dentibus calycinis inæqualibus, inferioribus tubo sublongioribus pilosis, stylis apice pilosis, leguminibus oblongis coriaceis compressis glabris.  $\mathcal{L}$  in pascuis Galloprovinciæ, Monsp. ex Benth. agri Ruscinonensis, Aurelianensis ex St. Hil., Audegaviæ ex Bast. — Fl. violacei. An var. *V. craccæ*? (v. s.)
7. *V. PSEUDOCRACCA* (Bert. pl. rar. 58.) pubescens, caule basi ramoso diffuso, foliolis 8-12 oblongis obtusis mucronatis, stipulis semisagittato-linearibus utrinque acuminatis, pedunculis subseffloris flexuosis folio longioribus, floribus secundis laxis deflexis, dentibus calycinis inæqualibus linearibus acutis tubo brevioribus, stylis apice villosis, leguminibus ellipticis brevibus glabris. (1) in Corsicâ reperit Ph. Thomas. — Fl. cærulei. (v. s.)
8. *V. TENUIFOLIA* (Roth. germ. 2. p. 183? Ser. in DC. prod. 2. p. 358.) caule erecto flexuoso, foliis cirrhosis, foliolis numerosis linearibus alternis oppositive mucronatis subtus adpressè villosis, stipulis inferioribus semisagittato-linearibus, superioribus lineari-setaceis, pedunculis rigidis folio longioribus, racemis confertis, dentibus calycinis inæqualibus, inferioribus tubi longitudine, stylis apice pilosis, legum. lanceolatis.  $\mathcal{L}$  in Corsicâ, Pyr. orient., Cebennis. — Fl. fr. 5. p. 577. Sturm. deutsch. 1. fasc. 31. (v. s.)
9. *V. STRICTA* (Duby mss.) caule flexuoso striato, foliis cirrhosis, foliolis 8-12 distantibus glabriusculis linearibus apice retusis mucronatis, stipulis semisagittatis sublaciniatis, pedicellis in axillis 2, uno brevi 1-floro, altero 3-4-floro folio vix dimidiò brevior, dentibus calycinis linearibus æqualibus longitudine tubi, stylis apice barbatis. (1) inter segetes circa Aquas Tarbellicas (Dax). — Fl. flavi. (v. v. D.)
10. *V. ONORATYCHOIDES* (L. sp. 1036.) caulibus striatis, cirrhis simpliciusculis, foliolis numerosis linearibus alternis oppositive obtusis mucronatisque, nervis subparallelis, stipulis semisagittato-linearibus dentatis, pedunculis multifloris longissimis, floribus distantibus, dentibus calycinis lanceolatis tubi longitudine, stylis clavatis apice barbatis, leguminibus lanceolatis. (1) in arvis montium provinc. australium. — Fl. fr. 4. p. 591. All. ped. t. 42. f. 1. Fl. purpurei. (v. v.)
- β. angustissima* (Ser. in DC. prod. 2. p. 358.) foliis setaceis brevioribus. Circâ Monspelium. (v. s.)
11. *V. ATROPURPUREA* (Desf. fl. atl. 2. p. 164.) villosa, caulibus tetragonis, cirrhis simpliciusculis, foliolis oblongis mucronatis numerosis oppositis alternisque, stipulis semisagittato-lanceolatis latis basi sæpè dentatis, pedunculis multifloris folii longitudine, floribus secundis approximatis, dentibus calycinis setaceis pilosis tubo longioribus, stylis elongatis subclavatis apice barbatis, leguminibus oblongis compressis hirsutissimis, seminibus globosis subvelutinis. (1) in Corsicâ, insulis Stæchadum, Galloprovinciâ propè Telsonem et St. Chinian. — Fl. fr. 4. p. 592. Vent. hort. cels. t. 84. Fl. purpurei. (v. s.)
12. *V. PERENNIS* (DC. cat. hort. monsp. 155. et fl. fr. 5. p. 578.) villosa, caulibus tetragonis, cirrhis simpliciusculis, foliolis oblongo-linearibus mucronatis villosis oppositis alternisque, stipulis semisagittato-lanceolatis basi sæpè dentatis, pedunculis paucifloris folii longitudine, floribus secundis laxiusculis, dentibus calycis setaceis pilosis tubi longitudine, stylis elongatis subclavatis apice barbatis, leguminibus oblongis compressis pubescentibus, seminibus orbiculato-compressis.  $\mathcal{L}$  in messibus agri Ruscinonensis. — Fl. purpurei. An satis à præcedente distincta? (v. v. DC.)

13. *V. ARGENTEA* (Lapeyr. abr. 417.) canescens, caulibus tetragonis, foliis cinereo-argenteis cirrho destitutis, foliolis oblongo-linearibus mucronatis, stipulis semisagittato-lanceolatis, pedunculis multifloris folio sublongioribus, floribus secundis laxiusculis, laciniis calycinis subæqualibus longitudine tubi, stylis elongatis subclavatis apice barbatis, leguminibus oblongis compressis tomentosis. ♀ in Pyrenæis editioribus Hispaniæ conterminis. — Fl. fr. 5. p. 576. Fl. subrosei, carina apice nigro maculata. (v. s.)
14. *V. DISPERMA* (DC. cat. monsp. 154.) subpilosa, caulibus tetragonis, cirrhis simpliciusculis, foliis 8-10-jugis, foliolis lineari-oblongis mucronatis, stipulis semisagittatis integris, pedunculis 2-3-floris folio brevioribus, floribus minimis, dentibus calycinis subæqualibus lanceolato-subulatis tubi longitudine et corollam subæquantibus, stylis apice villosis, leguminibus oblongis compressis glabris dispermis, seminibus subglobosis. ① in petrosis sterilibus Corsicæ, Telonis ex Lois., Mouspelii, agri Ruscinonensis ex Lapeyr. — Fl. fr. 5. p. 578. Fl. parvi cærulescentes. (v. s.)

### §. 2. *Floribus subsessilibus.*

15. *V. SATIVA* (L. sp. 1037.) foliis cirrhosis, foliolis 10-12 obovato-retusis vel oblongo-retusis mucronulatis pilosis vel glabriusculis, stipulis semisagittato-dentatis, floribus subgeminis sessilibus, calycibus cylindricis, laciniis lanceolato-linearibus subæqualibus parallelis tubi longitudine, stylis apice barbatis, leguminibus compressis subtorulosis oblongis erectiusculis, seminibus subglobosis subvelutinis. ② inter segetes Europæ. Species valde polymorpha. — Fl. fr. 4. p. 593. Fl. purpurei.
- a. obovata* (Ser.) foliolis obovato-cordatis pilosis, leguminibus pubescentibus. — *V. sativa* Hoppe. in Sturm. deutschl. fl. 1. fasc. 31. icon. (v. v.)
- β. segetalis* (Ser.) foliolis oblongis truncato-acuminatis pilosis, leguminibus pubescentibus. — *V. segetalis* Thuill. fl. par. ed. 2. p. 367. et Hoppe in Sturm. l. c. icon. Smith engl. bot. t. 334. Fl. fr. 5. p. 599. (v. v.)
- γ. angustifolia* (Ser.) foliolis linearibus longissimis subretusis mucronulatis, seminibus subglobosis. — *V. angustifolia* Roth. germ. 310. Fl. fr. 5. p. 599. (v. v.)
16. *V. PARGURIA* (L. sp. 1038.) foliis cirrhosis, foliolis 10-12 linearibus truncatis mucronatisque, stipulis semisagittato-linearibus integris, floribus solitariis pedunculatis, calycibus campanulatis, laciniis lanceolato-linearibus subæqualibus divergentibus tubi longitudine, stylis apice barbatis, leguminibus compressis lanceolatis reticulatis inclinatissimis puberulis, seminibus subglobosis. ① in siccis provinciarum australium. — Fl. fr. 5. p. 580. — Pluckn. t. 233. f. 6. Fl. purpureo-violacei. (v. v.)
17. *V. AMPHICARPA* (Dorth. journ. phys. 35. p. 131.) foliis subcirrhosis, foliolis inferioribus obcordatis mucronatis, superioribus linearibus retusis mucronatisque, stipulis semi-sagittatis integris vel dentatis, floribus solitariis subsessilibus, calycibus cylindricis, laciniis æqualibus linearibus parallelis tubo brevioribus, stylis apice barbatis, leguminibus puberulis, superioribus lineari-oblongis inclinatissimis, inferioribus subterraneis ovatis compressis brevibus. ① in aridis provinc. australiorum rariis. — Fl. fr. 4. p. 594. — J. Bauh. hist. 2. p. 323. f. 1 et 2. Fl. purpurei. (v. v. DC.)
18. *V. PYRENAICA* (Pourr. act. toul. 3. p. 333.) foliis subcirrhosis, foliolis obcordatis mucronatis, stipulis semisagittatis integris vel denticulatis, floribus solitariis sessilibus, calycibus subcampanulatis, laciniis inæqualibus lanceolatis subpatentibus tubo brevioribus, stylis apice barbatis, leguminibus oblongo-linearibus glabriusculis inclinatissimis. ♀ in pratis Pyrenæorum editiorum. — Fl. fr. 4. p. 595 excl. syn. J. Bauh. Ic. rar. p. 10. t. 33. *V. Fagonii* Lapeyr. abr. 419. Fl. purpurei. (v. v.)
19. *V. LATHYROIDES* (L. sp. 1037.) subvillosa, caulibus ramosis, foliis subcirrhosis, foliolis 4-6 obovatis retusis mucronatis, superioribus oblongis linearibus truncatis mucronulatis, stipulis semisagittatis integris, floribus so-

- latis sessilibus, calycibus cylindricis, laciniis linearibus subæqualibus parallelis tubo brevioribus, stylis apice barbatis, leguminibus compressis glabrisculis rugosis inclinatiss, seminibus globosis verrucoso-punctatis.  $\mathcal{L}$  (I in arenosis Gallie occidentalis, mediz, australisque passim. — Fl. fr. 4. p. 595. *Ervum soloniense* L. sp. 1040. Lam. ill. t. 634. f. 2. Fl. violacei. (v. v.)
20. *V. LUTEA* (L. sp. 1037.) pilosa, caulibus ramosis, foliis cirrhosis, foliolis oblongo-ovatis, stipulis brevissimis subsemisagittatis linearibusque basi denticulatis, floribus solitariis subsessilibus, laciniis calycinis inæqualibus divergentibus, superioribus brevibus, inferioribus tubo brevioribus, vexillo emarginato glabro, stylis apice barbatis, leguminibus compressis lanceolato-oblongis pilosis, seminibus subglobosis. (I ad vias et inter segetes Gallie occidentalis, mediz, et australis. — Fl. fr. 4. p. 596. — Moris. s. 2. t. 21. n. 5. Fl. lutei. (v. v.)
- $\beta$ . *pallidiflora* (Ser.) magis hirsuta, foliolis angustioribus, floribus albidis. — V. hirta Balb. Fl. fr. 5. p. 581. (v. v.)
21. *V. HYBRIDA* (L. sp. 1037.) pilosa, caulibus ramosis, foliis cirrhosis, foliolis obovato-cordatis mucronulatis, stipulis semisagittatis subdentatis, floribus solitariis subsessilibus subdeflexis, laciniis calycinis angustis subæqualibus longitudine tubi, vexillo emarginato piloso, stylis apice barbatis, leguminibus lanceolato-oblongis compressis pilosis. (I ad vias in Gallia occidentali, mediâ et australi. — Fl. fr. 4. p. 596. erg. bot. t. 482. Fl. lutei rubro striati. (v. v.)
22. *V. SARPIUM* (L. sp. 1038.) caule subramoso, foliis cirrhosis, foliolis 10-12 ovato-retosis submucronulatis ciliatis, stipulis semisagittatis, inferioribus dentatis, floribus 3-4 inclinatiss, calyce obliquè campanulato, dentibus inæqualibus supernè deflexis tubo brevioribus, stylis apice barbatis, leguminibus lanceolato-oblongis inclinatiss subciliatis, seminibus globosis.  $\mathcal{L}$  in umbrosis. — Fl. fr. 4. p. 596. — J. Bauh. 2. p. 313. f. 2. Fl. sordidè violacei. (v. v.)
- $\beta$ . *ochroleuca* (Bast.) floribus ochroleucis. — in Pictaviâ. — Fl. fr. 5. p. 581. (v. s.)
23. *V. PANNONICA* (Jacq. fl. sust. t. 34.) pilosa, caule simplici, foliis cirrhosis, foliolis oblongis numerosis, stipulis lanceolatis basi maculatis, floribus 2-4 inclinatiss subsessilibus, laciniis calycinis setaceis subæqualibus tubum æquantibus superantibusve, vexillo emarginato piloso, stylis apice barbatis, leguminibus lanceolato-oblongis compressis pilosis deflexis. (I  $\mathcal{L}$  in montibus Arverniz, Occitaniz, Bayonz ex Lois. — V. purpurascens DC. cat. monsp. 155. Fl. fr. 5. p. 580. Fl. purpurei. (v. v. DC.)
24. *V. NARBOYENSIS* (L. sp. 1038.) caule ascendente tetragono striato quadrifariam piloso, foliis cirrhosis, foliolis ovatis serratis integrisve, stipulis semisagittato-dentatis, inferioribus integris, floribus 3-4 breviter pedicellatis, calycibus campanulatis, laciniis ovatis triuerviis lateralibus supernè inflexis, stylis apice barbatis, leguminibus oblongis compressis glabris, suturis serrato-ciliatis, seminibus subglobosis. (I in Cebennis, agro Ruscinonensi, Narbonensi, Monspeliensti, Niceensi; circâ Dax ex Thore, in Arvernâ ex Delarbre. — Fl. fr. 4. p. 597. — Jacq. aust. app. t. 8. Fl. atropurpurei. Folia crassa. (v. s.)

XXXVI. *ERVUM*. L. gen. n. 874. DC. fl. fr. 4. p. 598.

Calyx 5-fidus, laciniis linearibus acutis corollam subæquantibus; stigma glabrum; legumen oblongum 2-4-spermum.

Sect. I. *LENS*. Ser. in DC. prod. 2. p. 366.

Legumen subinflatum lato-oblongum, 2-spermum.

1. *E. LENS* (L. sp. 1039.) caule ramoso, foliolis oblongis suboctonis glabrisculis, stipulis lanceolatis ciliatis, cirrhis subsimplicibus, pedunculis 2-3-floris folium æquantibus, leguminibus latis brevibus subtruncatis glabris,

13. *V. ARGENTEA* (Lapeyr. abr. 417.) canescens, caulibus tetragonis, foliis cinereo-argenteis cirrho destitutis, foliolis oblongo-linearibus mucronatis, stipulis semisagittato-lanceolatis, pedunculis multifloris folio sublongioribus, floribus secundis laxiusculis, laciniis calycinis subæqualibus longitudine tubi, stylis elongatis subclavatis apice barbatis, leguminibus oblongis compressis tomentosis.  $\mathcal{V}$  in Pyrenæis. editoribus Hispaniæ conterminis.  $\nearrow$  Fl. fr. 5. p. 576. Fl. subrosei, carina apice nigro maculata. (v. s.)
14. *V. DISPERMA* (DC. cat. monsp. 154.) subpilosa, caulibus tetragonis, cirrho simpliciusculis, foliis 8-10-jugis, foliolis lineari-oblongis mucronatis, stipulis semisagittatis integris, pedunculis 2-3-floris folio brevioribus, floribus minimis, dentibus calycinis subæqualibus lanceolato-subulatis tubi longitudine et corollam subæquantibus, stylis apice villosis, leguminibus oblongis compressis glabris dispermis, seminibus subglobosis. ① in petrosis sterilibus Corsicæ, Telonis ex Lois., Monspeli, agri Rusciniensis ex Lapeyr. — Fl. fr. 5. p. 578. Fl. parvi cærulescentes. (v. s.)

### §. 2. *Floribus subsessilibus.*

15. *V. SATIVA* (L. sp. 1037.) foliis cirrhosis, foliolis 10-12 obovato-retusis vel oblongo-retusis mucronulatis pilosis vel glabriusculis, stipulis semisagittato-dentatis, floribus subgeminis sessilibus, calycibus cylindricis, laciniis lanceolato-linearibus subæqualibus parallelis tubi longitudine, stylis apice barbatis, leguminibus compressis subtorulosis oblongis erectiusculis, seminibus subglobosis subvelutinis. ② inter segetes Europæ. Species valde polymorpha. — Fl. fr. 4. p. 593. Fl. purpurei.

a. *obovata* (Ser.) foliolis obovato-cordatis pilosis, leguminibus pubescentibus. — *V. sativa* Hoppe. in Sturm. deutschl. fl. t. fasc. 31. icon. (v. v.)

ß. *segetalis* (Ser.) foliolis oblongis truncato-acuminatis pilosis, leguminibus pubescentibus. — *V. segetalis* Thuill. fl. par. ed. 2. p. 367. et Hoppe in Sturm. l. c. icon. Smith engl. bot. t. 334. Fl. fr. 5. p. 579. (v. v.)

γ. *angustifolia* (Ser.) foliolis linearibus longissimis subretusis mucronulatis, seminibus subglobosis. — *V. angustifolia* Roth. germ. 310. Fl. fr. 5. p. 579. (v. v.)

16. *V. PÆRAGRINA* (L. sp. 1038.) foliis cirrhosis, foliolis 10-12 linearibus truncatis mucronatisque, stipulis semisagittato-linearibus integris, floribus solitariis pedunculatis, calycibus campanulatis, laciniis lanceolato-linearibus subæqualibus divergentibus tubi longitudine, stylis apice barbatis, leguminibus compressis lanceolatis reticulatis inclinatissimis puberulis, seminibus subglobosis. ① in siccis provinciarum australium. — Fl. fr. 5. p. 580. — Pluckn. t. 233. f. 6. Fl. purpureo-violacei. (v. v.)

17. *V. AMPHICARPA* (Dorth. journ. phys. 35. p. 131.) foliis subcirrhosis, foliolis inferioribus obcordatis mucronatis, superioribus linearibus retusis mucronatisque, stipulis semi-sagittatis integris vel dentatis, floribus solitariis subsessilibus, calycibus cylindricis, laciniis æqualibus linearibus parallelis tubo brevioribus, stylis apice barbatis, leguminibus puberulis, superioribus lineari-oblongis inclinatissimis, inferioribus subterraneis ovatis compressis brevibus. ① in aridis provinc. australiorum rariis. — Fl. fr. 4. p. 594. — J. Bauh. hist. 2. p. 323. f. 1 et 2. Fl. purpurei. (v. v. DC.)

18. *V. PYRÆNAICA* (Pourr. act. toul. 3. p. 333.) foliis subcirrhosis, foliolis obcordatis mucronatis, stipulis semisagittatis integris vel denticulatis, floribus solitariis sessilibus, calycibus subcampanulatis, laciniis inæqualibus lanceolatis subpatentibus tubo brevioribus, stylis apice barbatis, leguminibus oblongo-linearibus glabriusculis inclinatissimis.  $\mathcal{V}$  in pratis Pyrenæorum editorum. — Fl. fr. 4. p. 595 excl. syn. J. Bauh. Ic. rar. p. 10. t. 33. *V. Fagonii* Lapeyr. abr. 419. Fl. purpurei. (v. v.)

19. *V. LATHYROIDES* (L. sp. 1037.) subvillosa, caulibus ramosis, foliis subcirrhosis, foliolis 4-6 obovatis retusis mucronatis, superioribus oblongis linearibusve truncatis mucronulatis, stipulis semisagittatis integris, floribus so-

- litaris sessilibus, calycibus cylindricis, laciniis linearibus subæqualibus parallelis tubo brevioribus, stylis apice barbatis, leguminibus compressis glabrisculis rugosive inclinatiss, seminibus globosis verrucoso-punctatis.  $\mathcal{L}$  (1) in arenosis Galliæ occidentalis, mediæ, australisque passim. — Fl. fr. 4. p. 594. Ervum soloniense L. sp. 1040. Lam. ill. t. 634. f. 2. Fl. violacei. (v. v.)
20. *V. LUTEA* (L. sp. 1037.) pilosa, caulibus ramosis, foliis cirrhosis, foliolis oblongo-ovatis, stipulis brevissimis subæmisagittatis linearibusque basi denticulatis, floribus solitariis subsessilibus, laciniis calycinis inæqualibus divergentibus, superioribus brevibus, inferioribus tubo brevioribus, vexillo emarginato glabro, stylis apice barbatis, leguminibus compressis lanceolato-oblongis pilosis, seminibus subglobosis. (1) ad vias et inter segetes Galliæ occidentalis, mediæ, et australis. — Fl. fr. 4. p. 596. — Moris. s. 2. t. 21. n. 5. Fl. lutei. (v. v.)
- β. pallidiflora* (Ser.) magis hirsuta, foliolis angustioribus, floribus albidis. — V. hirta Balb. Fl. fr. 5. p. 581. (v. v.)
21. *V. HYBRIDA* (L. sp. 1037.) pilosa, caulibus ramosis, foliis cirrhosis, foliolis obovato-cordatis mucronulatis, stipulis semisagittatis subdentatis, floribus solitariis subsessilibus subdeflexis, laciniis calycinis angustis subæqualibus longitudine tubi, vexillo emarginato piloso, stylis apice barbatis, leguminibus lanceolato-oblongis compressis pilosis. (1) ad vias in Galliâ occidentali, mediâ et australi. — Fl. fr. 4. p. 596. eng. bot. t. 482. Fl. lutei rubro striati. (v. v.)
22. *V. SERIUM* (L. sp. 1038.) caule subramoso, foliis cirrhosis, foliolis 10-12 ovato-retusis submucronulatis ciliatis, stipulis semisagittatis, inferioribus dentatis, floribus 3-4 inclinatiss, calyce obliquè campanulato, dentibus inæqualibus supernè deflexis tubo brevioribus, stylis apice barbatis, leguminibus lanceolato-oblongis inclinatiss subciliatis, seminibus globosis.  $\mathcal{L}$  in umbrosis. — Fl. fr. 4. p. 596. — J. Bauh. 2. p. 313. f. 2. Fl. sordide violacei. (v. v.)
- β. ochroleuca* (Bast.) floribus ochroleucis. — in Pictaviâ. — Fl. fr. 5. p. 581. (v. s.)
23. *V. PANNONICA* (Jacq. fl. aust. t. 34.) pilosa, caule simplici, foliis cirrhosis, foliolis oblongis numerosis, stipulis lanceolatis basi maculatis, floribus 2-4 inclinatiss subsessilibus, laciniis calycinis setaceis subæqualibus tubum æquantibus superantibusve, vexillo emarginato piloso, stylis apice barbatis, leguminibus lanceolato-oblongis compressis pilosis deflexis. (1)  $\mathcal{L}$  in messibus Arvernæ, Occitanæ, Bayonæ ex Lois. — V. purpurascens DC. cat. monsp. 155. Fl. fr. 5. p. 580. Fl. purpurei. (v. v. DC.)
24. *V. NARBONENSIS* (L. sp. 1038.) caule ascendente tetragono striato quadrifariam piloso, foliis cirrhosis, foliolis ovatis serratis integrisve, stipulis semisagittato-dentatis, inferioribus integris, floribus 3-4 breviter pedicellatis, calycibus campanulatis, laciniis ovatis trinerviis lateralibus supernè inflexis, stylis apice barbatis, leguminibus oblongis compressis glabris, suturis serrato-ciliatis, seminibus subglobosis. (1) in Cebennis, agro Ruacionensi, Narbonensi, Monspelienſi, Niceensi; circa Dax ex Thore, in Arvernâ ex Delarbre. — Fl. fr. 4. p. 597. — Jacq. aust. app. t. 8. Fl. atropurpurei. Folia crassa. (v. s.)

XXXVI. ERVUM. L. gen. n. 874. DC. fl. fr. 4. p. 598.

Calyx 5-fidus, laciniis linearibus acutis corollam subæquantibus; stigma glabrum; legumen oblongum 2-4-spermum.

Sect. I. LENS. Ser. in DC. prod. 2. p. 366.

Legumen subinflatum lato-oblongum, 2-spermum.

1. *E. LENS* (L. sp. 1039.) caule ramoso, foliolis oblongis suboculis glabrisculis, stipulis lanceolatis ciliatis, cirrhis subsimplicibus, pedunculis 2-3-floris folium æquantibus, leguminibus latis brevibus subtruncatis glabris,

- seminibus compressis. ① in arvis inter segetes. — Fl. fr. 4. p. 599. — Fuchs. hist. 859. ic. (v. v.)
2. *E. KIASUTUM* (L. sp. 1039.) foliis cirrhosis, foliolis linearibus obtusis mucronulatis, stipulis semisagittatis angustis, pedunculis 3-6-floris folio brevioribus, lacinii calycinis lanceolato-linearibus aequalibus tubo longioribus, leguminibus oblongis compressis subtruncatis hirsutis nutantibus, seminibus globosis variegatis. ① in arvis Galliae mediæ et australis. — Fl. fr. 4. p. 599. — J. Bauh. 2. p. 315. f. 1. (v. v.)

Sect. II. *ERVILIA*. Ser. I. c.

Legumen subinflatum oblongo-lineare 4-6-spernum.

3. *E. ERVILIA* (L. sp. 1040.) glabrum, foliis polyphyllis subcirrhosis, foliolis oblongis mucronulatis, floribus subgeminis pedunculatis, stipulis sublanceolatis dentatis, lacinii calycinis aequalibus angustissimis tubo multò longioribus, leguminibus torulosis tetraspermis glabris, seminibus angulato-subrotundis. ① in arvis passim. — *Vicia Ervilia* Wild. Fl. fr. 4. p. 593. — Cam. epit. 215. ic. (v. v.)
4. *E. MONANTHOS* (L. sp. 1040.) caulibus cespitosis simplicibus, foliis numero-  
sis linearibus truncatis mucronatisque, cirrhis subimPLICIBUS, stipulis inæqualibus, illà lineari-lanceolatà integrâ, alterâ angustissimè fimbriatâ, pedunculis unifloris folium subæquantibus, lacinii calycinis linearibus aequalibus tubo longioribus, leguminibus ovalibus compressis glabris 3-4-sper-  
mis torulosis. ① in segetibus agri Ruscitonensis, Mimatenis (Prost), Arvernæ (Bast.). — *Vicia monantha* Desf. Fl. fr. 4. p. 592. (v. v.)
5. *E. TETRASPERMUM* (L. sp. 1039.) caulibus cespitosis ramosis, foliis cirrho-  
sis, foliolis 4-6 oblongis mucronulatis, stipulis lanceolato-semisagittatis, pedunculis 1-4-floris filiformibus, lacinii calycinis inæqualibus latiusculis tubo brevioribus, leguminibus oblongis compressis glabris subtorulosis, seminibus subglobosis. ① inter segetes, ad vias et colles passim. — Fl. fr. 4. p. 599. *Vicia parviflora*. Lois gall. 460 in Lapeyr. abr. 418 ex cl. Arnott. — J. Bauh. 2. p. 315. f. 2. (v. v.)

β. *gracile* (Ser.) foliolis angustissimis, pedunculis longioribus sæpissimè unifloris. — *E. gracile* DC. monsp. 190. fl. fr. 4. p. 581. *Vicia gracilis* Lois. gall. t. 12. (v. v.)

6. *E. RUBESCENS* (DC. cat. monsp. p. 109. fl. fr. 5. p. 582.) caulibus cespito-  
sis ramosis, foliis cirrhosis, cirrhis furcatis, foliolis 4-6 ellipticis mucro-  
nulatis, stipulis linearibus, pedunculis 1-4-floris filiformibus, lacinii cal-  
ycinis inæqualibus latiusculis tubo brevioribus, leguminibus oblongis com-  
pressis villosulis subtorulosis. ① in Galloprovincia propè Olbias. (v. s.)

XXXVII. *PISUM*. *Tourn. inst.* 394. t. 215. — *Pisi sp. L.*

Calycis laciniae foliaceae, 2 superioribus brevioribus; vexillum am-  
plum reflexum; stylus compressus carinatus supra villosus; legumen  
oblongum compressum non alatum; semina plura subglobosa, hilo  
subrotundo. — Stipulae amplae.

1. *P. SATIVUM* (L. sp. 1026.) petiolis teretibus trijugis, foliolis ovatis integris  
margine undulatis sæpè oppositis mucronulatis, stipulis ovato-subsemicor-  
datis crenatis, pedunculis bi-vel multifloris, leguminibus subcarnosis. ①  
ubique colitur, patria ignoratur. — Fl. fr. 4. p. 584. Lam. ill. t. 633. (v. v.)
- β. *umbellatum* (L. sp. 1027.) stipulis quadrifidis acutis, pedunculis  
multifloris terminalibus subumbellatis. — Tabern. ic. t. 495. An sp.  
propria? (v. v.)
2. *P. ARVENSE* (L. sp. 1027.) petiolis teretibus? 2-3-jugis, foliolis ovato-rotun-  
datis crenulatis mucronulatis, stipulis ovato-subsemicordatis denticulatis,  
pedunculis subunifloris brevissimis. ① patria ign. — Fl. fr. 4. p. 585. —  
Moris. s. 2. t. 1. n. 4. (v. v.)

- β. versicolor*, caule erecto flexuoso, petioli teretibus hexaphyllis, foliolis lanceolato-oblongis acuminatis, stipulis infernè rotundatis crenatis, pedunculis 2-3-floris folio sublongioribus. ① inter segetes circa Aquas Tarbellicas repertum misit cl. Grateloup. — An idem ac *P. elatum* Bieb. taur. 2. p. 151. aut distincta species? (v. s.)
3. *P. MARITIMUM* (L. sp. 1027.) caule angulato, petioli supra planiusculis, foliolis 5-8 ovatis vel rotundatis sæpè alternis puberulis, stipulis ovato-semisagittatis, pedunculis multifloris folio brevioribus, leguminibus oblongis parvis, seminibus parvis numerosis approximatis rotundatis. ① ad ostium Somonæ, in maritimis Nicæ? ex All. — Fl. fr. 4. p. 585. — Moris. s. 2. t. 1. n. 5. (v. s.)

XXXVIII. LATHYRUS. L. gen. n. 1186. Ser. in DC. prod. 2. p. 369.

Cal. campanulatus 5-fidus, lobis 2 superioribus brevioribus; cor. papilionacea; stam. diadelpa; stylus complanatus, apice dilatatus, anticè villosus aut pubescens; legumen oblongum, polyspermum, bivalve, 1-locul.; semina globosa aut angulata. — Stipulæ semisagittatæ.

Sect. 1. EULATHYRUS. Ser. l. c.

Vexillum basi edentatum; foliola opposita vel abortu nulla; petioli angustè-alatus.

§. 1. *Perennes, pedunculis multifloris.*

\* *Folii unijugis.*

1. *L. SYLVESTRIS* (L. sp. 1033.) glaberrimus, caulibus prostratis alatis, foliis unijugis, foliolis lineari-lanceolatis attenuatis coriaceis, stipulis semisagittatis folio brevioribus, pedunculis 6-12-floris folii longitudine, leguminibus compressis, seminibus subrotundis verrucoso-punctatis. 2℥ in sylvis et sepibus montanis. — Fl. fr. 4. p. 583. — Lob. ic. 2. p. 68. f. 2. Fl. rosei aut purpurei.
- α. ensifolius* (Ser.) foliolis valdè elongatis angustissimis. — In provinciis australibus. (v. s.)
- β. oblongus* (Ser.) foliolis brevioribus elliptico-oblongis, stipulis multo minoribus. — OEd. fl. dan. t. 325. An sp. distincta?
2. *L. LATIFOLIUS* (L. sp. 1083.) glaberrimus, caulibus prostratis alatis, foliis unijugis, foliolis ovato-ellipticis subglaucescentibus 3-5-nerviis obtusis mucronatisque, stipulis semisagittatis latis, pedunculis multifloris folio longioribus, leguminibus compressis. 2℥ in vineis, sylvis, pratisque umbrosis provinc. austral. passim. — Fl. fr. 4. p. 583. Eng. bot. t. 1108. Fl. purpurei. (v. v. DC.)
3. *L. PRATENSIS* (L. sp. 1033.) glabriusculus, caulibus erectiusculis tetragonis, foliis unijugis, foliolis oblongis vel lineari-lanceolatis trinerviis, stipulis sagittatis ovatis foliolis brevioribus, pedunculis multifloris folio duplò longioribus, calycibus nervosis, dentibus subæqualibus longitudine tubi, leguminibus compressis, seminibus globosis lævissimis. 2℥ in pratis humidis. — Fl. fr. 4. p. 583. — J. Bauh. 2. p. 304. f. 2. Fl. flavi. (v. v.)
- β. velutinus* (DC.) caule foliolisque velutinis, leguminibus subpilosis. — In locis aridis. — Fl. fr. 5. p. 575. (v. v.)
4. *L. TUMEROSUS* (L. sp. 1033.) caule debili tetragono, foliis unijugis, foliolis oblongo-ellipticis submucronulatis, stipulis semi-sagittatis angustis acutis petiolum æquantibus, pedunculis 3-6-floris folio duplò triplòve longioribus, calycibus aveniis, dentibus subæqualibus latis tubum subæquantibus, stylis arcuatis, leguminibus compressis, seminibus subrotundis lævibus. 2℥ ad margines agrorum passim. — Fl. fr. 4. p. 582. — Lob. ic. 2. p. 70. f. 2. Fl. rosei. (v. v.)

\*\* *Folii multijugis.*

5. *L. PALUSTRIS* (L. sp. 1034.) glaberrimus, caulibus alatis erectiusculis, fo-

liolis 3-jugis, petiolo subalato, foliis oblongis mucronulatis, stipulis semisagittatis acutis parvis, pedunculis 3-5-floris folio vix longioribus, lacinii calycinis inæqualibus sublinearibus longitudine tubi, leguminibus compressis.  $\mathcal{V}$  in pascuis paludosis Gallix mediæ, borealis, orientalisque, Pyren. centr. ex cl. Benth. — Fl. fr. 4. p. 584. fl. dan. t. 399. Fl. pallide purpurei. (v. v.)

*f. subexstipulatus* (Ser.) foliis lineari-oblongis 2-jugis, stipulis minimis setaceis. — Circà Argentoratun. (v. s.)

6. *L. HYKROPHYLLOS* (L. sp. 1034.) caule erecto rigido alato, foliis uni et 2-jugis, foliolis lanceolatis mucronulatis, petiolis basi alatis, stipulis lanceolato-semisagittatis, pedunculis 6-8-floris, leguminibus compressis glabris.  $\mathcal{V}$  in petrosis passim præsertim in provinc. australibus. — Fl. fr. 4. p. 583. — J. Bauh. hist. 2. p. 304. f. 1. Fl. purpurei. (v. s. c.)

§. 2. *Anhui, pedunculis 1-3-floris.*

\* *Petioli aphyllis.*

7. *L. AFRAGA* (L. sp. 1029.) caule erecto, petiolis cylindricis filiformibus aphyllis, stipulis sagittato-ovatis maximis, pedunculis 1-floris apice articulatis, bractea minimâ angustissimâ, lacinii calycinis tubo duplò longioribus, leguminibus valdè compressis oligospermis latis, seminibus compressis. ① inter segetes. — Fl. fr. 4. p. 577. — Lob. ic. 2. p. 70. f. 1. Petioli rarissimè 1-jugi, pedunculi rarissimè 2-flori. Fl. latei. (v. v.)

8. *L. NISSOLIA* (L. sp. 1029.) caule erecto, petiolis dilatatis foliaceo-gramineis 3-5-nerviis aphyllis, stipulis minimis subulatis sæpè nullis, floribus solitariis longè pedunculatis, pedunculis apice articulatis ebracteatis, leguminibus compressis angustis nervosis reflexis. ① in arvis lapidosis, dumetis et ad margines pratorum Gallix occidentalis et australis passim. — Fl. fr. 4. p. 578. — Lob. ic. 2. p. 71. f. 1. Fl. purpurei. (v. v.)

\*\* *Foliis unijugis.*

9. *L. INCONSPICUUS* (Linn. sp. 1030.) caulibus obsoletè triquetris, foliolis lanceolatis subtus striatis acuminatis, stipulis semisagittato-lanceolatis, cirrhis inferioribus ferè nullis, superioribus filiformibus elongatis, pedunculis brevissimis unifloris. ① in agro Occitano et Ruscinonensi ex cl. Benth. — Corolla pallida tam parva ut vix calyce longior.
10. *L. SPHERICUS* (Retz. obs. 3. p. 39.) glabriusculus, caulibus erectis tetragonis apice subulatis, foliolis ensiformibus mucronatis nervosis, stipulis semisagittato-linearibus longitudine pedunculi et petioli, pedunculis unifloris crassiusculis, dentibus calycinis angustis tubo longioribus, leguminibus subtorulosis nervosis, nervis valdè numerosis crassis, seminibus sphericis. ① in siccis et lapidosis provinc. austr. — Fl. fr. 4. p. 580. ic. rar. t. 32. L. axillaris. Fl. fr. 5. p. 574. Fl. rubri. (v. v.)
11. *L. MICRANTHUS* (Gér. in Lois. not. 106.) caulibus tetragonis, cirrhis simplicibus brevissimis, foliolis lanceolato-linearibus, pedunculis brevissimis 1-floris, lacinii calycinis corollam subæquantibus, leguminibus subsessilibus strictis patentibus cylindricis subvillosis. ①? in arvis Galloprov. ex Ger. — Fl. fr. 5. p. 575. An var. præcedentis. †
12. *L. ANGULATUS* (L. sp. 1031.) glaber, caulibus erectis tetragonis, foliolis linearibus acuminatis, cirrhis 3-fidis, stipulis semisagittato-linearibus acutis petiolo vix longioribus, pedunculis unifloris filiformibus stipulis duplò triplòve longioribus foliumque vix æquantibus longè aristatis, dentibus calycinis acutis tubi longitudine, leguminibus angustis compressis. ① inter segetes, in sterilibus provinc. australium et aliquandò occidentalium et mediarum. — Fl. fr. 4. p. 580. — Buxb. cent. 3. t. 42. f. 2. Fl. purpurei minores. (v. v.)
13. *L. SETIFOLIUS* (L. sp. 1031.) glaber, caulibus erectis tetragonis, foliolis linearibus angustissimis, stipulis semisagittatis angustissimis longitudine



petioli, cirrhis trifidis, pedunculis unifloris filiformibus folio duplò brevioribus apice articulatis bracteolâ minimâ instructis, dentibus calycinis acutis tubi longitudine, leguminibus ovato-oblongis brevibus, seminibus globosis verrucoso-rugosis. (1) in provinciis australibus. — Fl. fr. 4. p. 581. — J. Bauh. 2. p. 308. ic. Fl. rosei. (v. v.)

14. *L. SATIVUS* (L. sp. 1030.) glabrescens, caulibus diffusis alatis, foliolis lineari-oblongis, cirrhis trifidis, stipulis semisagittato-ovatis ciliatis vix petioli longitudine, pedunculis unifloris petiolo longioribus apice articulatis bracteolatis, lacinii calycinis lanceolatis foliaceis tubo ferè triplò longioribus, leguminibus ovatis latis brevibus dorso alatis, seminibus trigonis subtruncatis lævibus. (1) ex Hispaniâ ortus, in agris reperitur. — Fl. fr. 4. p. 579. — J. Bauh. 2. p. 306. f. 2. Flores albi, rosei aut violacei. (v. v.)

15. *L. CACERA* (L. sp. 1030.) glabrescens, caulibus diffusis alatis, foliolis lineari-oblongis, cirrhis 3-4-fidis, stipulis semisagittato-lanceolatis subdentatis ciliatis petioli longitudine, pedunculis uniloricis stipulâ longioribus, bracteolis minimis, lacinii calycinis lanceolatis foliaceis tubo ferè triplò longioribus, leguminibus oblongis dorso canaliculatis non alatis, seminibus trigonis subtruncatis lævibus. (1) ex Hispaniâ orta, in arvis Gallie australis colitur. — Fl. fr. 4. p. 579. Jacq. f. ecl. t. 115. Fl. purpurei. (v. v.)

16. *L. ANTHUS* (L. sp. 1032.) glaber, caulibus diffusis alatis, foliolis linearibus elongatis, cirrhis 3-fidis, stipulis semisagittatis angustissimis petiolo multò brevioribus, pedunculis 1-2-floris folioli longitudine, bracteolis minimis, lacinii calycinis lanceolatis tubo vix longioribus, leguminibus oblongo-lanceolatis non alatis. (1) in arvis provinc. australium. — Fl. fr. 4. p. 581. — Bauh. cent. 3. t. 42. f. 1. Fl. lutei parvi. (v. s.)

17. *L. HIRAUTUS* (L. sp. 1032.) caulibus diffusis alatis, foliolis lineari-oblongis, stipulis semisagittato-linearibus petiolum subæquantibus, pedunculis 1-3-floris folio vix longioribus, lacinii calycinis ovatis tubi longitudine, leguminibus oblongis hirsutis, seminibus globosis verrucoso-punctatis. (1) inter ægetes Gallie australis orientalisque. — Fl. fr. 4. p. 582. — J. Bauh. hist. 2. p. 305. ic. (v. s.)

18. *L. QUORATUS* (L. sp. 1032.) hirsutus, caulibus diffusis alatis, foliolis ovatis mucronulatis, stipulis semisagittato-lanceolatis petiolo multò brevioribus, pedunculis 2-3-floris folio multò longioribus, dentibus calycinis latis tubo longioribus, leguminibus oblongo-linearibus compressis hirsutis. (1) in hortis colitur. — Fl. fr. 4. p. 582. Fl. odori. (v. v.)

\*\*\* *Folii 2-3-jugis.*

19. *L. CIRRANOSUS* (Ser. in DC. prod. 2. p. 374.) glaberrimus, caulibus tetragonis alatis, petioliis 5-6-phyllis alatis, cirrhis ramosissimis, foliolis alternis ellipticis mucronulatis reticulato-venosis, stipulis semisagittato-linearibus acutis, pedunculis multifloris folio longioribus, lacinii calycinis inæqualibus angustis tubo brevioribus, leguminibus oblongis, stylis arcuatis. — In Pyrenæis orientalibus circa Olette reperiit Ph. Thomas. (v. s.)

20. *L. BITHYNICUS* (Lam. dict. 2. p. 706.) pilosus, caulibus diffusis tetragonis, foliis inferioribus unijugis ovato-oblongis mucronatis, superioribus bijugis lineari-lanceolatis, stipulis semisagittatis dentatis, pedunculis brevibus elongatisve 1-2-floris, lacinii calycinis linearibus tubum æquantibus supernantibusve, stylis barbatis, leguminibus oblongis compressis villosis turgidis, seminibus subrotundis. (1) in provinciis mediterraneis et austro-occidentibus usque ad Bardigalam. — Vicia Bithynica L. sp. 1038. All. ped. 1. 26. f. 2. Fl. fr. 4. p. 597. (v. v.)

#### Sect. II. CLYMENUM. DC. prod. 2. p. 375.

Vexillum basi utrinque gibbositate conicâ notatum; folia inferiora è foliolata, superiora 2-6 foliolata, foliolis sæpè alternis; petioli sæpè alatus.

21. *L. ALATUS* (Tenor. prod. 42. non Sibth.) caule tetragono alato, foliolis 6-8 lanceolato-oblongis mucronulatis alternis, petioliis alatis, stipulis lanceolatis

subagittatis inæqualibus, pedunculis 2-3-floris folio longioribus, dentibus calycinis inæqualibus tubo brevioribus, leguminibus compressis planis polyspermis. ① in Galloprovincia orientali reperiit Stire. — Fl. purpurei magni. Valde affinis sequenti. (v. s.)

22. L. CLYMENUM (Linn. sp. 1032.) caule tetragono alato, petiolis infimis dilatatis aphyllis lineari-lanceolatis, superioribus 5-6-phyllis linearibus, stipulis semisagittato-linearibus, pedunculis 1-6-floris folium æquantibus, dentibus calycinis inæqualibus tubo longioribus, leguminibus oblongis compressis, suturis seminiferis tumidis, seminibus compressis variegatis. ① in Corsicâ, insulis Stæchadum et collibus circâ Telonem, Perpignanum. — Fl. fr. 5. p. 574. Variat vexillo rubro, alis cæruleis et flore cæruleo. (v. v. DC.)
23. L. ARTICULATUS (Linn. spec. 1031.) caule tetragono alato, petiolis inferioribus aphyllis linearium-acuminatis, superioribus 5-6-phyllis linearibus, stipulis semisagittato-lanceolatis, pedunculis 1-3-floris folium æquantibus, dentibus calycinis subæqualibus tubo brevioribus, leguminibus turgescentiâ seminum nodosis, suturis seminiferis tumidis, seminibus compressis subvelutinis. ① Niceæ, in Corsicâ, Occitaniâ, agro Ruscitonensi et circâ Aquas Tarbellicas ex Thore. — Fl. fr. 4. p. 579. Gærtn. fruct. 2. t. 152. f. 2. — Mill. ic. t. 96. Fl. purpurei. (v. s.)
24. L. OCHRUS (DC. fl. fr. 4. p. 578.) caule prostrato tetragono alato, petiolis latis plerisque aphyllis, supremis 2-3-phyllis cirrhosis, stipulis inferioribus nullis, superioribus ovatis, foliolis ovatis mucronatis, pedunculis unifloris brevibus, dentibus calycinis subinæqualibus calycis longitudine, leguminum suturis seminiferis alato-membranaceis polyspermis. ① In arvis et inter segetes Corsicæ, Niceæ ex All., MonsPELLI ex Gouan, Aq. Tarbellicarum ex Thore. — Pisum Ochrus L. sp. 1027. — Moris. s. 2. t. 3. n. 8. (v. s.)

XXXIX. OROBUS. Tourn. inst. t. 214. L. gen. n. 871.

Calyx campanulatus 5-fidus, lobis 2 superioribus brevibus; cor. papilionacea; stam. diadelphe; stylus gracilis linearis apice villosus; legumen cylindraceum oblongum 2-valve polyspermum; semina hylo lineari. — Stipulæ semi-sagittatæ.

§. 1. *Foliis multijugis, foliolis latis, stipulis folio multò minoribus.*

1. O. VERNUS (L. sp. 1028.) subpubescens, caule simplicei, foliis trijugis, foliolis lanceolatis acuminatis, stipulis semisagittatis integris, pedunculis axillaribus multifloris folio brevioribus, floribus subsecundis nutantibus, lacinii calycinis latis vix longitudine tubi, leguminibus subcompressis, stylis geniculatis. 2. in nemorosis. — Fl. fr. 4. p. 587. — Clus. hist. 2. p. 230. f. 1. (v. v.)
2. O. VARIEGATUS (Tenor. prod. suppl. 1. p. 62. et fl. napol. t. 68. non Lapeyr.) pilosus, caule simplici flexuoso, foliis 2-3-jugis, foliolis lanceolatis acuminatis, stipulis ovato-semisagittatis integerrimis acutis foliolis multò minoribus, pedunculis axillaribus multifloris folium æquantibus, lacinii calycinis angustis acutis longitudine tubi, leguminibus compressis punctatis; stylis rectis. 2. in Corsicâ propè Bastia reperiit Ph. Thomas. — Columna phytob. p. 59. ic. (v. s.)
3. O. LUTAEUS (L. sp. 1028.) glabriusculus, caule simplici anguloso, foliis 3-5-jugis, foliolis elliptico-lanceolatis antrorsum, glaucis mucronulatis, stipulis semisagittatis basi dentatis foliolis multò minoribus, pedunculis elongatis multifloris folium vix æquantibus ascendentes, calycibus glabriusculis, dentibus inæqualibus brevibus, leguminibus compressis sessilibus, seminibus globosis variegatis. 2. in montibus. — Fl. fr. 4. p. 587. O. Tournefortii Lapeyr. mem. mus. 2. p. 298. t. 10. — J. Bauh. 2. p. 343. f. 1. (v. v.)
4. O. NIGER (L. sp. 1028.) glabriusculus, caule ramoso angulato flexuoso, foliis 3-6 jugis, foliolis ellipticis mucronulatis, stipulis lineari-lanceolatis acutis, pedunculis multifloris folio longioribus, dentibus calycinis inæquali-

bus tubo brevioribus, leguminibus compressis, stylis geniculatis, seminibus globosis.  $\mathcal{L}$  in nemorosis montanis. — Fl. fr. 4. p. 586. — Clus. hist. 2. p. 230. f. 2. Fl. purpurei. (v. v.)

5. *O. rupestris* (L. sp. 1028.) glaberrimus, caulibus prostratis internè tuberculatis, foliis 2-4-jugis, foliolis ellipticis mucronulatis punctato-rugosis, stipulis semisagittatis, pedunculis paucifloris folium vix superantibus, dentibus calycinis inæqualibus ovatis obtusiusculis tubo brevioribus, leguminibus compressis, stylis geniculatis, seminibus globosis.  $\mathcal{L}$  in sylvis. — Fl. fr. 4. p. 587. Fl. purpurei rarò albi. (v. v.)

$\beta$ . *Pyrenaicus* (Ser.) foliolis ovato-ellipticis, stipulis lanceolato-linearibus ex ic. Pluk. phyt. t. 210. — *O. Pyrenaicus* L. sp. 1029. *O. Plucknetii* Lapeyr.

$\gamma$ . *divaricatus* (Ser.) foliolis ellipticis oblongis, dentibus calycinis angustioribus, stipulis plus minùsve dentatis. — *O. divaricatus* Lapeyr. mem. mus. 2. p. 302. t. 2.

$\delta$ . *tenuifolius* (Ser.) foliolis lineari-lanceolatis.

§. 2. *Foliis multijugis, foliolis angustissimis.*

6. *O. canescens* (L. f. suppl. p. 327.) caule tetragono, foliis subtrijugis, foliolis linearibus obtusiusculis pubescentibus vel punctato-rugosis, stipulis semisagittatis linearibus acutis petiolum alatum subæquantibus, pedunculis 3-5-floris folio duplò longioribus, dentibus calycinis latis tubo brevioribus, leguminibus rectis compressis glabris, stylo geniculato rhomboides.  $\mathcal{L}$  in montosis Galloprov., Delphin. ex Vill., Pyrenæorum. — *O. filiformis* Lam. DC. fl. fr. 4. p. 588. excl. syn. Bauh. Cor. albo et cæruleo variegata. (v. v. DC.)

$\beta$ . *ensifolius* (Ser.) foliis latioribus acutis, pedunculis multifloris. — *O. ensifolius* Lapeyr. mem. mus. 2. p. 303. t. 12. *O. atropurpureus* Lapeyr. abr. 413 non Desf. — In Pyrenæis centralibus. (v. v. DC.)

$\gamma$ . *tenuis* (DC.) foliis angustissimis acutis, pedunculis multifloris. — In Pyrenæis orient. (v. s.)

7. *O. albus* (L. f. suppl. 327.) caule simplici, foliis subtrijugis, foliolis linearibus mucronatis, stipulis semisagittatis latiusculis petiolo alato brevioribus, pedunculis multifloris folio longioribus, dentibus calycinis lanceolatis valde inæqualibus inferioribus multò longioribus, leguminibus compressis glabris subflexuosis, stylo lineari.  $\mathcal{L}$  in Andegaviâ (Bast.) et Delphinatùs inferioris montosis. — Fl. fr. 4. p. 588. — J. Bauh. hist. 2. p. 326. f. 2. (v. v. D.)

$\beta$ . *asphodeloides* (DC.) foliolis brevioribus latioribusque, tuberibus radicalibus oblongis. — In Pictaviâ et Cebennis. (v. s.)

8. *O. saxatilis* (Vent. hort. cels. t. 94.) caulibus simplicibus debilibus, foliis 2-jugis, foliolis linearibus, stipulis semisagittatis minimis, pedunculis unifloris folio multò brevioribus articulatis bracteolatis, lacinii calycinis lanceolatis subæqualibus tubo brevioribus, leguminibus subcylindricis cinereis. (1) in collibus aridis Galloprovinciæ Mompelii, Narbonæ et agri Ruscinonensis ex cl. Benth. — Fl. fr. 4. p. 589. (v. s.)

Trib. V. *PHASEOLEÆ*. Bronn. diss. p. 133. DC. prod. 2. p. 381. leg. mem. ix.

Corolla papilionacea. Stam. monadelphæ aut sæpè diadelphæ (9 et 1). Legumen polyspermum dehiscens continuum membranæ cellulosis transversis inter semina sitis sæpè subdivisum non vèrè articulatum. Radicula super loborum commissuram inflexa. Cotyledones crassæ per germinationem immutatæ aut in folia crassa vix stomatibus donata conversæ, extra terram exsertæ. Folia primordialia opposita.

XL. PHASEOLUS. *L. gen. n. 866. Savi diss. 1824.*

**Calyx** campanulatus 2-labiatus, labio superiore 2-dentato, inferiore tripartito; corolla papilionacea, carinā cum staminibus diadelphis styloque spiraliter tortā vel rariū incurvā; legumen compressum aut cylindricum bivalve intus isthmis cellulosis subdistinctum polyspermum; semina hilo ovali-oblongo.

1. *P. MULTIFLORUS* (Wild. sp. 3. p. 1030.) volubilis glabriusculus, foliolis ovatis acuminatis, racemis pedunculatis folio longioribus, pedicellis geminatis, bracteolis calyce subbrevioribus adpressis, leguminibus pendulis subfalcatis torulosis scabris. ① ex Americā calidiore orta, in hortis ornamenti causā colitur. — Fl. fr. 4. p. 559. — Moris. hist. s. 2. t. 5. n. 4. Variat floribus coccineis et albis. (v. v. c.)
2. *P. VULGARIS* (Savi mem. 3. p. 14.) volubilis glabriusculus, foliolis ovatis acuminatis, racemis pedunculatis folio brevioribus, pedicellis geminis, leguminibus pendulis rectiusculis subtorulosis longē mucronatis, semine ovato subcompresso. ① ex Indiā orient. ortus, in hortis frequentissimus, valdē variabilis, longē volubilis aut subnanus, floribus albis lilacinisve. seminibus colore valdē variis, Ph. vulgaris et Ph. nani L. sp. 1016. varietates diversæ ad hunc aut sequentem pertinent. — Fl. fr. 4. p. 558. — Lob. ic. 2. t. 59. f. 2. Moris. s. 2. t. 5. n. 1. (v. v. c.)
3. *P. COMPRESSUS* (DC. prod. 2. p. 392.) subvolubilis glabriusculus, foliolis ovatis acuminatis, racemis pedunculatis folio brevioribus, pedicellis geminis, legumine compresso subtoruloso mucronato, seminibus compressis. ① Patr. ign. Fl. albi. — *P. Romanus* Savi mem. 3. p. 17. t. 10. f. 20. (v. v. c.)
  - a. *humilis* (DC.) caule humili recto, seminibus minoribus. — *P. nanus* Fl. fr. 4. p. 559.
  - β. *major* (DC.) caule procero, leguminibus subtortis.
4. *P. TUMIDUS* (Savi mem. 3. p. 19. f. 16.) humilis subvolubilis glabriusculus, foliolis ovatis acuminatis, racemis folio brevioribus, legumine rectiusculo mucronato plus minus toruloso, semine sphaerico vel ovato-turgido albo immaculato, ventre tumido. ① Patr. ign. Fl. albi. (v. v. c.)
5. *P. SPHERICUS* (Savi mem. 3. p. 20 f. 17.) procerus volubilis glabriusculus, foliolis ovatis acuminatis, racemis folio brevioribus, legumine rectiusculo toruloso-mucronato, semine subrotundo nunquā albo. ① Patr. ignota. Flores pallidē violacei. (v. v. c.)

XLI. LUPINUS. *Tourn. inst. 392. t. 213. L. gen. n. 1176.*

**Calyx** profundē bilabiatus; cor. papilionacea, vexilli lateribus reflexis, carinā acuminatā: stam. monadelpha vaginā integrā, antheris 5 parvis subrotundioribus præcioribus, 5 oblongis serioribus; stylus filiformis; stigma terminale subrotundum barbatum; legumen coriaceum oblongum compressum obliquē torulosum. — Stipulæ petiolo adnatæ; folia sæpius foliolis 5-9 digitatis constantia.

1. *L. ALBUS* (L. sp. 1015.) floribus alternis pedicellatis ebracteolatis, calycis labio superiore integro, infer. tridentato, foliolis obovato-oblongis subtus villosis. ① in provinciis australibus cultus. — Fl. fr. 4. p. 506. — Clus. hist. 2. p. 228. f. 1. Fl. albi. (v. v.)
2. *L. VARIUS* (L. sp. 1015.) floribus semiverticillatis aut subalternis pedicellatis bracteolatis, calycis labio superiore bidentato, inferiore vix tridentato, foliolis oblongis subtus villosis. ① inter segetes agri Syrtici propē St.-Sever ex Thore, Narbonæ, Mompelii, Galloprovinciæ ex Ger, Nicææ ex All. — Fl. fr. 4. p. 507. Fl. cærulei et purpurascens. (v. s.)
3. *L. HIRAUTUS* (L. sp. 1015.) floribus alternis bracteolatis, calycis labio superiore bipartito, inferiore trifido, foliolis oblongis utrinquē hirsutis, leguminibus hirsutissimis. ① in cultis Corsicæ, Mompelii. — Fl. fr. 4. p. 508. — J. Bauh. hist. 2. p. 289. ic. Flores minores quam prioris, cærulei. (v. s.)

4. *L. ANOUSTROPIUS* (L. sp. 1016.) floribus alternis breviter pedicellatis ebracteolatis, calycis labio superiore bifido, inferiore integro, foliolis linearibus adpresse pubescentibus, leguminibus villosis 6-spermis. ① inter segetes Corsicæ, Galloprovinciæ, agri Ruscinonensis ex cl. Benth., Syrtici propè St.-Sever (cl. Dufour). — Fl. fr. 4. p. 507. — J. Bauh. hist. 2. p. 291. ic. excl. descr. Flores cærulei. (v. v.)
5. *L. LUTRUS* (L. sp. 1015.) floribus verticillatis sessilibus bracteolatis, calycis labio superiore bipartito, inferiore 3-dentato, foliolis oblongis, infimis obovatis. ① in arvis circà Mûnsipium? ex Gouan. — Fl. fr. 4. p. 508. — J. Bauh. hist. 2. p. 290. ic. Fl. lutei odorati. (v. s. ex Hispania.)

Subordo II. *CÆSALPINIÆ*. R. Brown. gen. rem. p. 19. DC. prod. 2. p. 473. leg. mem. XIII.

Flores plus minùs irregulares, nunc papilionacèi nunc subrosacei. Petala per æstivationem irregulariter imbricata nec valvata, interdum nulla, nunquam inter se coalita. Stamina plus minùs inæqualia, semper perigyna, sæpius libera. Embryonis recti plumula sæpius conspicua magna.

Trib. VI. *CASSIÆ*. DC. prod. 2. p. 478.

Calycis lobi ante explicationem imbricati. Petala perigyna subæqualia rarò subpapilionacea, per æstivationem imbricata. Legumen sæpius siccum bivalve. Cotyledones foliaceæ aut rarius carnosæ.

XLII. *CERATONIA*. L. gen. n. 1167. Lam. ill. t. 859.

Fl. polygami aut dioici; sepala 5 in calycem 5-partitum basi coalita; pet. 0; stam. 5; stigma sessile orbiculatum; legumen lineare coriaceum indehiscens polyspermum, isthmis subcarnosis multiloculare, valvis crassis intus pulposis. — Arbores trunco crasso.

1. *C. SILIQUA* (L. sp. 1513.) inermis, foliolis ovalibus obtusis planis. 5 in rupestribus maritimis Nicæ, Galloprovinciæ, Corsicæ. — Fl. fr. 4. p. 490. Cav. ic. t. 113. (v. v.)

XLIII. *CERCIS*. L. gen. n. 510. Gærtn. t. 144.

Cal. sepala 5 concreta in urceolum obtusè 5-dentatum infernè gibbum; pet. 5 unguiculata subpapilionacea, omnia distincta, alis majoribus; stam. 10, libera inæqualia; legumen oblongum tenuè compressum 1-loc. polyspermum, suturâ superâ seminiferâ subulatâ, alterâ dehiscente; semina obovata, endoplevrâ tumidâ albumen mentiente; embryo rectus, radiculâ brevi, cotyl. planis, plumulâ inconspicuâ. — Arbores.

1. *C. SILIQUASTRUM* (L. sp. 534.) foliis simplicibus basi cordatis integris obtusissimis glaberrimis. 5 in rupestribus et sylvis Narbonæ ex Lin., Mûnsipii ex Gouan, Nicæ ex All., Delph. australis ex Vill. spontanea, in sylvis sæpè colitur. — Fl. fr. 4. p. 490. (v. v. c.)

Ordo XXXIV. ROSACEÆ. *Juss. gen. 334. DC. prod. 2. p. 525.*

Calyx constans sepalis sæpius 5, inter se basi in tubum cohærentibus et ideo 5-lobus, plerumque persistens, sæpius liber, interdum ovario adhærens. Petala tot quot sepala, rarius abortu nulla, calyci inserta, æstivatione quincunciali, ferè semper regularia. Stamina cum petalis inserta, sæpissimè infinita, filamentis per æstivationem incurvis, antheris bilocularibus rimâ duplici dehiscentibus. Carpella plurima, nunc abortu solitaria, nunc coalitione inter se aut cum calycis tubo in ovarium primo adpectu unicum conversa. Ovaria 1-locul. Styli simplices, ex ovario sæpius lateraliter orti, nunc distincti, rarius inter se coaliti. Semina in carpellis sæpius 1-2, rarius plurima, erecta aut inversa, exalbuminosa. Embryo rectus, cotyledonibus nunc foliaceis, nunc carnosus. — Herbar frutices aut arbores, foliis alternis basi bistipulatis.

Trib. I. *AMYGDALÆÆ. Juss. gen. 340. excl. gen. DC. prod. 2. p. 529. — Drupacæ. Fl. fr. 4. p. 479.*

Carpella sæpissimè abortu solitaria, rarius 2 aut plurima, stylum filiformem ex apice ferè gerentia, desinentia in drupam nucleo solitario 2-ovulato 1-2-spermo. Funiculus umbilicalis è basi nuclei ortus, in canali laterali ad apicem ferè loculi locatus et sic semen pendulum gerens. Endoplevra subtumida. Cotyledones 2 crassæ. Calyx ovario non adhærens deciduus 5-fidus, gerens petala 5 et stam. 20-30 libera subæqualia. — Arbores fruticesve, foliorum serraturis infimis petiolisve glandulosi, stipulis liberis.

I. *AMYGDALUS. Tourn. inst. t. 402. DC. fl. fr. 4. p. 486.*

Drupa pubescenti-velutina, cortice fibrosâ exsuccâ irregulariter secedente, putamine foraminuloso aut lævi. — Folia juniora conduplicata; flores subsessiles foliis præcociore; fructus lanati.

1. *A. communis* (L. sp. 677.) foliis oblongo-lanceolatis serrulatis, floribus solitariis, calycibus campanulatis, fructibus ovoideo-compressis tomentosis.  $\beta$  è Mauritaniâ orta, nunc in provinciis australibus subsponsanea. — Fl. fr. 4. p. 486.

*a. amara* (DC.) stylis ferè longitudine staminum infra tomentosis, seminibus amaris. (v. v. c.)

*β. dulcis* (DC.) stylis stamina multò superantibus, fructibus acuminatis, seminibus dulcibus. — Lam. ill. t. 430. f. 2. (v. v. c.)

II. *PERSICA. Tourn. inst. t. 400. DC. fl. fr. 4. p. 487.*

Omnia Amygdali, sed drupa valdè carnosâ, epicarpio nunc velutino nunc glaberrimo, putamine sulcis insigniter rugoso.

1. *P. vulgaris* (Mill. dict. n. 1.) fructibus tomentosis.  $\beta$  ex Persiâ orta, in hortis colitur. — Fl. fr. 4. p. 487. *Amygdalus Persica* L. sp. 677. (v. v. c.)

2. *P. LÆVIS* (DC. fl. fr. 4. p. 487.) fructibus lævibus. ♂ Patr. ign. in hortis colitur. (v. v. c.)

III. ARMENIACA. *Tourn. inst. t. 399. Juss. gen. 341. DC. fl. fr. 4. p. 485.*

Drupa ovato-globosa carnosa extus velutina, putamine hinc obtuso illinc acuto, utrinquè sulcato, cæterum lævi nec rugoso. — Folia juniora convolutiva; flores subsessiles.

1. *A. VULGARIS* (Lam. dict. t. p. 2.) floribus sessilibus, foliis cordatis vel ovatis. ♂ ex Armeniâ orta, in hortis colitur. — Fl. fr. 4. p. 486. *Prunus Armeniaca* L. sp. 679. (v. v. c.)
2. *A. BRIGANTIAICA* (Pers. ench. 2. p. 36.) foliis subcordatis acuminatis argutè dentatis, dentibus numerosis subimbricatis, floribus agglomeratis subsessilibus subcoctaneis. ♂ in Alpibus Galloprovinciæ et Delphinatûs propè Brigantium (*Briançon*). — *Prunus Brigantiaca* Vill. dauph. 3. p. 535. Fl. fr. 4. p. 484. Lois. in Duham. ed. nov. 5. t. 59. (v. v. DC.)

IV. PRUNUS. *Tourn. inst. t. 398. Juss. gen. 341. DC. fl. fr. 4. p. 483.*

Drupa ovata aut oblonga, carnosa, glaberrima, polline cæσιο tecta; putamine compresso utrinquè acuto, ad margines subsulcato; cæterum lævi. — Folia juniora convolutiva; florum pedicelli sæpiùs umbellato-fasciculati 1-flori.

1. *P. SPINOSA* (L. sp. 681.) ramis aculeatis, pedunculis solitariis, calycibus campanulatis, lobis obtusis tubo longioribus, foliis ovato-ellipticis vel ovatis subtus pubescentibus argutè duplicatoque dentatis, fructibus globosis. ♂ in collibus apricis. — Fl. fr. 4. p. 484. Fl. dan. t. 926. Drupa acerba.
  - a. *vulgaris* (Ser.) foliis ovato-ellipticis, fructibus atropurpureis. — *P. spinosa* Lois. in Duham. ed. nov. 5. t. 54. f. 1. (v. v.)
  - β. *microcarpa* (Wallr.) foliis ellipticis angustis obtusiusculis, fructibus minoribus. — In sylvis agri Syrtici propè La Teste detexit cl. Des Moulins. (v. s.)
2. *P. INSITITA* (L. sp. 680.) ramis spinescentibus, pedunculis geminis, foliis ovatis subtus villosis convolutis, fructibus subrotundis. — In Galliâ australi? *P. domestica* β. Fl. fr. 4. p. 484. An var. *P. spinosa*? Planta valdè incerta. †
3. *P. DOMESTICA* (L. sp. 680.) floribus subsolitariis, foliis lanceolato-ovatis convolutis, ramis muticis, ♂ in pomariis colitur. — Fl. fr. 4. p. 484. excl. var. β. — Quoad var. confer. Ser. in DC. prod. 2. p. 533. (v. v.)
  - β. *pyramidalis*, ramis pyramidalis, fructibus ovatis plus minùs obtusis elongatis violaceis, umbilico exserto, nucleo valdè compresso. — *P. pyramidalis* Fl. fr. 4. p. 485. *P. domestica prunaeuliensis* Ser. l. c.

V. CERASUS. *Juss. gen. 340. DC. fl. fr. 4. p. 479.*

Drupa globosa aut basi umbilicata carnosa, glaberrima, polline cæσιο destituta, nucleo subgloboso lævi. — Folia juniora conduplicata; pedicelli nunc 1-flori, nunc ramosi.

SECT. 1. CERASOPHORA. *DC. prod. 2. p. 535.*

Flores umbellati, pedicelli uniflori è gemmis orti.

1. *C. AVIUM* (Mœnch meth. 672. DC. fl. fr. 4. p. 482.) floribus subcoctaneis, fructibus rotundo-ovatis depressis, pedunculis tenuibus, carne valdè succulentâ, epicarpio carni adhærente, foliis subtus albidis subpubescentibus, ramis validis divaricatis. ♂ in sylvis præsertim montosis et exindè in pomariis culta. Varietates coluntur subnomine vernaculo *Merisier*. (v. v.)
2. *C. DURACINA* (DC. fl. fr. 4. p. 483.) ramis junioribus ascendentibus, adul-

tis vix patulis, floribus subcœtaneis, fructibus cordatis, pedunculis longis tenuibus, suturâ valde depressâ, carne dariusculâ, epicarpio carni adhærente, putamine ovato. § Patr. ign. varietates coluntur subnomine *Bigarreau*, quoad earum distributionem cf. Ser. in DC. prod. 2. p. 535. (v. v.)

3. C. JULIANA (DC. fl. fr. 4. p. 482.) ramis junioribus ascendentibus, adultis vix patulis, floribus subcœtaneis, fructibus ovatis depressis cordiformibus, carne molliusculâ non acidâ, epicarpio carni valdè adhærente, foliis glabris. § Patr. ign. Nomen vernaculum *Guignier* aut *Cerisier*. (v. v.)
4. C. CAPRONIANA (DC. fl. fr. 4. p. 482.) ramis patulis, floribus subcœtaneis, calycibus campanulatis amplis, pedunculis sæpissimè crassis rigidiusculis, fructibus globoso-depressis, suturâ vix depressâ, carne molli plus minus acidâ et stipticâ, epicarpio carni non adhærente, putamine rotundo. § Patr. ign. Vulgò *Cerisiers*, *Griottiers*. (v. v.)
5. C. SEMPERFLORENS (DC. fl. fr. 4. p. 481.) ramis nutantibus, foliis ovatis serratis, floribus serotinis axillaribus solitariis, calycibus serratis, fructibus globosis rubris. § Patr. ign. — Lois. in Duham. nov. ed. t. 5. f. A. (v. v.)

Sect. II. PADUS. DC. prod. 2. p. 539.

Flores racemosi è ramis orti.

6. C. MAHALEB (Mill. dict. n. 4.) racemis subcorymbosis foliosis, foliis cordato-subrotundis denticulatis glandulosis curvatis, fructibus ovato-subrotundis nigris. § in incultis, sylvaticis, et sepibus passim. — Fl. fr. 4. p. 480. Lois. in Duham. ed. nov. 5. t. 2. Prunus Mahaleb L. Jacq. austr. t. 227. (v. v.)
7. C. PADUS (DC. fl. fr. 4. p. 580.) racemis elongatis foliosis, foliis ovato-lanceolatis subacuminatis tenuibus patentim serrulatis, fructibus rotundis amaris. § in collibus et sepibus. — Lois. in Duh. l. c. 5. t. 1. Pr. Padus L. sp. 667. Fl. dan. t. 205. (v. v.)
8. C. LAUROCERASUS (Lois. in Duham. ed. nov. 5. p. 6.) racemis folio brevioribus, foliis ovato-lanceolatis remote serratis subtus 2-4-glandulosis, fructibus ovatis acutis. § ex Trapezunte orta, frequentissimè in hortis colitur. — Prunus Laurocerasus L. sp. 678. — Fl. fr. 4. p. 480. — Blackw. herb. t. 512. (v. v.)

Trib. II. SPIREACEÆ. DC. prod. 2. p. 541.

Carpella plurima a calyce libera, inter se distincta aut rarissimè subconnata, circâ axim floris idealem verticillata, sæpius 5, abortu interdum pauciora, stylo apiculata, demum capsularia suturâ interiore dehiscencia, subbivalvia. Semina 2-4, rariùs abortu 1-3 è suturæ interioris margine ad medium aut propè basin orta, exarillata, exalbuminosa. Embryo rectus, cotyledonibus planis crassiusculis. — Frutices aut Herbæ.

VI. SPIRÆA. L. gen. n. 630.

Calyx 5-fidus persistens; stamina 10-50 cum petalis toro calyci adhærenti inserta; carpella 1-∞ distincta, rarò basi coalita, sessilia rarò stipitata; semina 2-6; embryo inversus.

Sect. I. CHAMÆDRYON. Ser. in DC. prod. 2. p. 542.

Torus apice liber. — Frutices floribus hermaphroditis umbellatis vel subcorymbosis, foliis integris vel dentatis exstipulatis.

1. S. HYPERICIFOLIA (DC. fl. fr. 5. p. 645.) folis obovato-oblongis 3-4-nerviis integris dentatisve glabris puberulisve, corymbis pedunculatis aut umbellis sessilibus, pedicellis glabris puberulisve, sepalis ascendentibus. § in hortis frequenter colitur.



- a. Plukenetiana* (Ser.) foliis integerrimis glabris, corymbis sessilibus. — In sylvis Biturigum (Berri) reperit cl. Gay ex DC. l. c. — *S. hypericifolia* L. sp. 701. — Pluck. phyt. t. 218. f. 5. (v. v. c.)  
*β. crenata* (Ser.) foliis obovatis apice crenulatis subpubescentibus, corymbis sessilibus, pedunculis puberulis. — In Cebennis ex Gouan. — *S. hypericifolia* γ. DC. fl. fr. 5. p. 645. *S. crenata* β. fl. fr. 4. p. 477. — Barr. ic. t. 564. (v. v. c.)

Sect. II. *ARUNCUS*. Ser. l. c.

Carpella 5 libera pendula, torus apice liber crassissimus. — Herba floribus abortu dioicis, foliis tripinnatisectis, stipulis nullis.

2. *S. ARUNCUS* (L. sp. 702.) foliis tripinnatisectis, lobis acuminate terminali ovato, lateralibus lanceolatis, floribus numerosissimis, carpellis glaberrimis. ♀ in sylvis montosis. — Fl. fr. 4. p. 479. — Cam. hort. 26. t. 9. (v. v.)

Sect. III. *ULMARIA*. Cambess. spir. in ann. sc. nat. 1. p. 378.

Torus obsoletus, stylus clavatus retroflexus; ovula 2 circa mediam ovarii suturam affixa. — Herba floribus hermaphroditis cymoso-umbellatis, foliis pinnatisectis stipulatis.

3. *S. ULMARIA* (L. sp. 702.) foliis interruptè pinnatisectis subtus tomentoso-niveis (raro glabris), lobo terminali majore trilobato, sepalis reflexis, stylis elongatis, carpellis glabris contortis. ♀ in pratis humidis. — Fl. fr. 4. p. 478. Eng. bot. t. 960. (v. v.)  
 4. *S. FILIPENDULA* (L. sp. 702.) radice tuberosa, foliis interruptè pinnatisectis, lobis oblongo-linearibus acutè dentatis, stipulis subreniformibus amplexicaulibus dentatis, corymbis laxis, sepalis reflexis, carpellis villosis. ♀ in sylvis et pascuis umbrosis.  
*a. vulgaris* (Camb.) caule elato foliisque glabris. (v. v.)  
*β. pubescens* (Camb.) foliis subincano-pubescentibus. — In Galloprovincia propè Fouchateau. — *S. pubescens* DC. fl. fr. 5. p. 546. (v. a.)

Trib. III. *DRYADEÆ*. Vent. tabl. 3. p. 349. DC. prod. 2. p. 549.

Calyx 5-fidus (rarius 4-fidus aut pluri-divisus), per æstivationem valvatus, sæpè extus bracteolas lobulose exteriores lobis alternos gerens. Pet. tot quot calycis lobi veri et iis alterna. Stam. ∞, rarius 5 et tunc lobis cal. opposita, apice tubi calycini expansi inserta. Carpella ∞, rarius pauca congesta, toro inserta, inter se et à calyce libera, et latere propè apicem stylifera. Akenia uni-ovulata omnino libera, exsucca aut baccata. Semen solitarium erectum aut inversum, exalbuminosum. Embryo rectus, cotyled. planiusculis. — Herbae aut frutices, stipulis 2 lateraliter petiolo adnatis.

VII. *DRYAS*. L. gen. n. 637.

Cal. 8-g-partitus extus nudus, tubo subconcavo; pet. 8-g; stam. ∞; carpella ∞ stylo terminali demum in caudam barbato-plumosam desinente, semen ascendens. — Frutices humillimi.

1. *D. OCTOPETALA* (L. sp. 717.) foliis crenato-serratis obtusis ovatis aut subcordatis. ♀ in excelsis montibus. — Fl. fr. 4. p. 473. Eng. bot. t. 451. (v. v.)

VIII. *GEUM*. L. gen. n. 867.

Cal. tubus concavus, limbus 5-fidus extus 5-bracteolatus; pet. 5;

stam.  $\infty$ ; carpella exsucca caudata in capitulum disposita, stylo post anthesin articulado aut barbato; semen ascendens. — Herbas, foliis pinnatisectis.

Sect. I. CARIOPHYLLASTRUM. *Ser. in DC. prod. 2. p. 550.*

Flores ascendentes; calyces reflexi; styli deflexi geniculati, appendices plerumque stylo breviores.

1. *G. URBANUM* (L. sp. 716.) caule erecto ramoso piloso, foliis radicalibus quinato-pinnatisectis, caulinis ternato-palmatisectis, lobis ovatis latis dentato-crenatis, supremis unilobatis ovatis, stipulis suborbiculatis magnis, petalis obovatis longitudine calycis, carpellis pilosis, stylis glabris, appendicibus pilosisculis.  $\frac{1}{2}$  in sepibus. — Fl. fr. 4. p. 270. Eng. bot. t. 1400. (v. v.)

Sect. II. CARYOPHYLLATA. *Tourn. inst. t. 151. Ser. L. c.*

Flores erecti vel cernui; calyces erecti; styli deflexi geniculati, appendice stylum subæquante.

2. *G. RIVALE* (L. sp. 717.) pilosum, caulibus erectis simpliciusculis 1-4-floris, foliis interruptè pinnatisectis lobis obovatis biserratis, foliis caulinis trifloris lobis acutis, stipulis ovatis dentatis, pedunculis pilosis elongatis, floribus nantantibus, petalis obcordatis basi angustatis longitudine calycis, ovaria valde pilosis, stylis elongatis flexilibus pilosis.  $\frac{1}{2}$  in pratis humidis montosis. — Fl. fr. 4. p. 471. — Lob. ic. t. 694. Fl. rosei. (v. v.)
3. *G. PYRÆNICUM* (Ram. bull. phil. n. 42. t. 10. f. 3.) pilosum, caulibus erectis simpliciusculis 1-4-floris, foliis interruptè pinnatisectis, lobulis inferioribus ovatis dentatis minimis, terminali oordato-reniformi maximo biserrato, stipulis ovatis obtusè dentatis, floribus nantantibus, petalis obcordato-rotundatis brevissimè unguiculatis calyce longioribus, stylis deflexis rigidis longitudine carpellorum, carpellis valde pilosis, appendicibus pilosis.  $\frac{1}{2}$  in editioribus Pyrenæis. — Fl. fr. 4. p. 472. *G. Tournefortii* Lapeyr. (v. v. DC.)
4. *G. SYLVATICUM* (Pourr. act. toul. ex DC. fl. fr. 5. p. 544.) pilosum, caulibus erectis simplicibus, foliis interruptè pinnatisectis, lobis inferioribus ovatis dentatis, terminali subcordato biserrato maximo, foliis caulinis simplicibus profundè dentatis, stipulis lanceolatis subdentatis, floribus ascendentibus, petalis obcordatis calyce longioribus, stylis deflexis rigidis longitudine ovarii pilosi, appendicibus glabris.  $\frac{1}{2}$  in sylvaticis circa Narbonam, Monspelium, Nemausum. — *G. atlanticum* Desf. atl. 1. p. 401. (v. v. DC.)
5. *G. THOMASIANUM* (Ser. in mem. soc. phys. gen. 2. p. 140.) pilosum, caulibus erectis 1-3-floris, foliis radicalibus subinterruptè pinnatisectis, lobis subæqualibus obovatis subduplicato-serratis, caulinis ternatisectis, lobo terminali 3-lobo vel lanceolato, stipulis ovatis subduplicato-serratis, floribus ascendentibus, lobis calycinis ovatis brevibus, petalis obovatis calyce vix longioribus.  $\frac{1}{2}$  in Pyrenæis orientalibus circa Mont-Louis reperiit Ph. Thomas. (v. s.)

Sect. III. SIEVERSIA. *Wild. berol. mag. 1811. p. 397. — Oreogeeum. Ser. L. c.*

Flores calycisque erecti; styli patentes non geniculati pilosi.

6. *G. REPTANS* (L. sp. 717.) caulibus sterilibus reptantibus, floriferis erectis indivisis, foliis radicalibus interruptè pinnatifidis, lobis majoribus obovato-cuneiformibus apice profundè dentatis, minoribus ovatis integris tridentatisve, foliis caulinis trilobis, flore maximo unico, laciniis calycinis elongatis sæpè apice trifidis, petalis obcordatis calyce longioribus, stylis patentibus valde pilosis.  $\frac{1}{2}$  in editioribus Alpibus. — Fl. fr. 4. p. 472. — Barr. ic. t. 400. Fl. flavi. (v. v.)
7. *G. MONTANUM* (L. sp. 717.) caulibus erectis unifloris, stolonibus nullis, foliolis radicalibus interruptè pinnatifidis, lobis lateralibus minoribus den-

tatis, terminali ovato-oblongo obtuso maxime obtusè biserrato, caulinis unilobis stipellisque profunde dentatis, flore magno, laciniis calycinis indivisis, petalis obo cordatis calyce longioribus, stylis patentibus valdè pilosis. ½ in editoribus montibus. — Fl. fr. 4. p. 472. — Cam. epit. 727. ic. Fl. flavi. (v. v.)

IX. RUBUS. L. gen. n. 864.

Calyx fundo planiusculus 5-fidus nudus; pet. 5 et stam. oo calyci inserta; carpella plurima tero manifeste protuberanti conico non carnoso inserta, stylo sublaterali superata, in drupellas carnosas conversa; semen inversum. — Frutices rariùs herbæ perennes, foliis petiolatis.

§. 1. *Foliis pinnatis ternatisve.*

1. R. INZUS (L. sp. 706.) villosus, caulibus teretibus, aculeis tenuibus recurvis, foliis ramorum sterilibus pinnatis, fertiliùm trifoliolatis, pinnis ovatis inciso-serratis steriliùm subpalmato-quinatis subtus niveo-tomentosis, stipulis angustissimis setaceis, pedunculis subcorymbosis, laciniis calycinis ovato-lanceolatis niveo-tomentosis submucronulatis, petalis obovato-cuneatis integris conniventibus calyce brevioribus, carpellis numerosis tomentosis. ½ in petrosis montium. — Fl. fr. 4. p. 476. Engl. bot. t. 2442. (v. v.)

§. 2. *Foliis palmatis 3-5-foliolatis.*

\* *Fruticosi, stipulis linearibus petiolaribus.*

2. R. CÆSUS (L. sp. 706.) caule repente tereti sæpè cæsioprofusosus, aculeis tenuibus subrecurvis, foliis ternato-quinatoque-palmatis, foliolis ovatis duplicato-serratis vel crenatis glabris subciliatisve, paniculâ simpliciusculâ, laciniis calycinis longè ovato-acuminatis, carpellis crassis paucis subcæcis. ½ ad marginas agrorum. — Fl. fr. 4. p. 474. — J. Bauh. hist. 2. p. 59. f. 1. (v. v.)
3. R. CORYMBOSUS (Smith. fl. brit. p. 542; eng. bot. t. 827.) caule fruticoso angulato, aculeis rectiusculis, foliis ternato-quinatoque-palmatis, foliolis crenato-ovatis firmis duplicato-serratis subtus pilosis, paniculâ simpliciusculâ, laciniis calycinis ovato-acuminatis demùm reflexis. ½ in sepibus passim. — Fl. fr. 4. p. 475. — Schmid. ic. t. 2. Carpella purpureo-cærules magus. (v. v.)  
*β. villosus* (DC.) foliis utrinque villosis. (v. v.)
4. R. FRUTICOSUS (L. sp. 707.) caule erecto 5-angulari subtomentoso, aculeis recurvis, foliis 3-5-palmatis, foliolis petiolulatis ovato-oblongis acutis glabris subtus adpressè tomentoso-niveis, paniculâ decompositâ angustâ strictâ, laciniis calycinis reflexis subinermibus. ½ in sepibus et dumetis. — Fl. fr. 4. p. 475. Lam. ill. t. 441. f. 2. Fructus purpureo-atric. (v. v.)  
*β. inermis* (DC.) caule inermi. — Barr. ic. t. 395. (v. a.)
5. R. TOMENTOSUS (Wild. sp. 2. p. 1083.) caule erecto angulato aculeato, foliis 3-5-palmatis, foliolis breviter petiolulatis obovato-cuneatis suprè pubescentibus subtus lanuginoso-tomentosis, paniculâ decompositâ angustâ patente, laciniis calycinis subinermibus reflexis. ½ in collibus siccis passim. — Fl. fr. 4. p. 476. (v. v. DC.)
6. R. COLLINUS (DC. cat. monsp. 139.) caule fruticoso, petiolis aculeatis, foliis quinato rariùs ternato-palmatis, foliolis subrotundis dentatis, laterali-bus subsessilibus supernè villosis, subtus tomento vix adpresso candicantibus, floribus laxè paniculatis, paniculâ ramosâ, laciniis calycinis ovato-acuminatis. ½ in collibus sterilibus Monspeli. — Fl. fr. 5. p. 545. — Neoca et Balb. fl. ticin. t. 9. (v. v. DC.)
7. R. GLANDULOSUS (Bell. act. taur. 3. p. 230.) caule debili teretiusculo piloso-hispido vel villosus, aculeis subrecurvis tenuibus acutissimis, foliis ternatis rariter quinato-palmatis villosis, foliolis ovatis duplicato-serratis, pedunculis villosis glanduloso-hispidis, floribus laxè paniculatis, calycis laciniis lan-

ceolatis acuminatis.  $\zeta$  in sylvis umbrosis montium. — Fl. fr. 4. p. 474. R. hirtus Walde. et Kit. hung. 2. t. 141. R. villosus a glandulosus Ser. in DC. prod. 2. p. 564. (v. v.)

\*\* *Herbacei, stipulis ovatis.*

8. R. *PARATILIS* (L. sp. 708.) caule herbaceo obtusangulo, foliis ternato-palmatis glabriusculis, foliolis ovato-rhomboides crenato-serratis, lateralibus sessilibus, stipulis ovatis latiusculis, floribus subcorymbosis subsolitariisque breviter pedunculatis, laciniis calycinis submentosis deflexis corollam æquantibus, carpellis paucis rubris maximis.  $\zeta$  in rupestribus montosis. — Fl. fr. 4. p. 473. Eng. bot. t. 2233. (v. v.)

X. *FRAGARIA*. Tourn. inst. t. 152. Ser. in DC. prod. 2. p. 569.

Calycis tubus concavus 5-fidus extus 6-bracteolatus; pet. 5; stam.  $\infty$ ; carpella  $\infty$  receptaculo demum deciduo carnosio-succulento sparse inserta exsucca, stylus lateralis; semen appensum. — *Herbæ foliis 3-foliolatis.*

\* *Carpellis numerosissimis superficialibus, staminibus numerosis, floribus hermaphroditis.*

1. F. *vesca* (L. sp. 705.) foliis ternatis plicatis supra pubescentibus subtus pilosis, foliolis sessilibus cuneato-subrotundis grosse dentatis dentibus acuminatis, pedunculis lanuginosis, fructibus pendulis, sepalis post anthesin reflexis petalis brevioribus.  $\mathcal{Z}$  in sylvis collibusque et in hortis culta. — Fl. fr. 4. p. 468. — Blackw. herb. 77. — Quoad varietates conf. quæ scripsit cl. Ser. in DC. prod. 2. p. 569. (v. v.)
2. F. *CALYCEA* (Lois. gall. 1. p. 299.) foliis ternatis pilosiusculis, foliolis sessilibus cuneato-subrotundis grosse dentatis, floribus corymbosis, pedunculis scapo longioribus, calycibus corollam subæquantibus.  $\mathcal{Z}$  in sylvis circa Lutetiam ex Lois. — F. vesca grandiflora. Fl. fr. 4. p. 468.

\*\* *Carpellis paucis in foveolis receptaculi immersis, staminibus paucis, floribus abortu sæpe dioicis.*

3. F. *MAJAUFRA* (Duch. in Lam. dict. 2. p. 533.) stolonifera, lobis foliorum plicatis tennibus subtus pilosis, floribus hermaphroditis, staminibus longioribus, carpellis numerosis, calycibus post anthesin erecto-adpressis.  $\mathcal{Z}$  in Galliâ, et in hortis colitur.
4. F. *BRESLINGEA* (Duch. in Lam. dict. 2. p. 533.) lobis foliorum plicatis subcoriaceis viridioribus, pilis longioribus firmioribus, petalis angustioribus albo-flavicantibus, sepalis longioribus post anthesin erecto adpressis, receptaculis adhærentibus, carpellis turgidiaribus.  $\mathcal{Z}$  in Galliâ, et in hortis culta. Quoad varietates conf. quæ scripsit cl. Ser. l. c.
5. F. *CHULENSIS* (Ehr. beytr. 7. p. 26.) scrotina, semper abortu dioica, foliis glaucis coriaceis latè crenatis subtus et supra pilosis, pedunculis sepalis receptaculisque erectis pendulisve crassioribus.  $\mathcal{Z}$  in Americâ meridionali orta, in hortis culta.

XI. *POTENTILLA*. Nestl. potent. diss. Ser. in DC. prod. 2. p. 571. — Tormentilla, Potentilla et Comarum L.

Cal. tubus concavus, limbus 4-5-fidus extus 4-5-bracteolatus; pet. 4-5; stam.  $\infty$ ; carpella  $\infty$  stylo laterali, receptaculum persistens exsuccum; semen appensum. — *Herbæ aut suffrutices, foliis compo-*  
*sitis.*

Sect. I. *POTENTILLASTRUM*. Ser. l. c.

Petala obtusa obcordatave lutea; folia palmata vel pinnatisecta.

§. 1. *Foliis palmatisectis.*

1. *P. MINIMA* (Hall. fl. in Ser. mus. helv. t. p. 51. t. 8.) caulibus diffusis filiformibus unifloris, foliis ternato-palmatisectis supra glabresculis subtus pilosis, lobis obovatis brevibus subretusis crenatis, laciniis calycinis inæqualibus obtusis corollâ brevioribus, carpellis subovoideis lævibus.  $\frac{1}{2}$  in editioribus Alpibus Delphin. et Galloprovinciæ. — *P. frigida*  $\beta$  Fl. fr. 4. p. 462. *P. verna*  $\gamma$ . Fl. fr. 5. p. 542. (v. v. DC.)
2. *P. FRIGIDA* (Vill. dauph. 3. p. 563.) pilosissima, radice crassâ, caulibus cespitosius subdiffusis, foliis ternato-palmatisectis, lobis ovatis crenatis summo prominulo, stipulis ovato-lanceolatis, laciniis calycinis lanceolatis acutiusculis petala obcordato-rotunda superantibus æquantibusve, carpellis lævibus.  $\frac{1}{2}$  in summis Alpibus Delphinatûs ex Vill. — Nestl. pot. 70. t. 10. f. 3. non Fl. fr. 4. p. 462 ex Ser. in DC. prod. p. 462.
3. *P. GRANDIFLORA* (L. sp. 715.) caule ascendente paucifloro, foliis pilosis ternato-palmatisectis, lobis obovato-cuneatis lanceolatis profunde simpliciter vel subduplicato-dentatis dentibus obtusis, stipulis lanceolatis, laciniis calycinis lanceolatis acutis, bracteis calycinis ellipticis pilosis brevioribus, petalis obcordatis calyce longioribus, carpellis ovoideis rugosis.  $\frac{1}{2}$  in pascuis saxosis Alpium. — Fl. fr. 4. p. 463. — Hall. helv. t. 20. f. 1. (v. v.)
4. *P. SUBCAULIS* (L. sp. 715. non Lehm.) incano-tomentosa, caule brevi, foliis ternato-palmatisectis, lobis obovato-cuneiformibus apice crenatis venosis, stipulis linearibus, floribus subpaniculatis paucis, laciniis calycinis ovatis obtusiusculis stellato-tomentosis, petalis obovatis calyce brevioribus, receptaculo villoso, carpellis rugosis.  $\frac{1}{2}$  in montosis rupestribus apricis Galloprov. et agri Ruscinonensis. — Fl. fr. 4. p. 465. *P. grandiflora* Scop. carn. 1. t. 22. non L. (v. s.)
5. *P. TORMENTILLA* (Nestl. pot. 65.) pilosa, radice tuberosâ, caule ascendente dichotomo, foliis ternato-palmatisectis, caulinis sessilibus, lobis obovato-cuneiformibus plus minusve profunde dentatis, stipulis nullis vel 3-dentatis, floribus axillaribus solitariis longè pedunculatis, bracteis palmato-incisis, laciniis calycinis lanceolato-linearibus corollam æquantibus, receptaculo villoso, carpellis rugosis.  $\frac{1}{2}$  in graminosis ad vias et oras sylvarum. — *Tormentilla erecta* L. sp. 716. — Fl. fr. 4. p. 454. — Cam. epit. 685. ic. Calyx 4-5-fidus; pet. 4-5. (v. v.)  
*ℓ. nemoralis* (Ser.) caule debili prostrato, stipulis frequentissimè lanceolatis indivisis rarè 2-3-dentatis, foliis petiolatis sæpè oppositis. — *Torment. reptans* L. Fl. fr. 4. p. 454. (v. v.)
6. *P. REPTANS* (L. sp. 714.) caule flagelliformi reptante subindiviso, foliis plus minusve petiolatis 3-5-7-palmatisectis, lobis obovatis dentatis lateralibus basi manifestè coëctis, stipulis linearibus parvis vel nullis, bracteis ovatis 2-3-lobis, floribus axillaribus solitariis folio longioribus, laciniis calycinis ovatis latis corollâ brevioribus, petalis obcordatis, receptaculo piloso.  $\frac{1}{2}$  in argillosis et glareosis subhumidis. — Fl. fr. 4. p. 461. — Fuchs. hist. 624. ic. (v. v.)
7. *P. VERNA* (L. sp. 712.) hirsuta, caulibus declinatis, foliis infimis quinato-palmatisectis, lobis obovato-cuneiformibus dentatis dente ultimo minore, stipulis inferioribus linearibus integris acutis, floribus subpaniculatis, laciniis calycinis lanceolatis acutis, petalis obovatis calycem superantibus, carpellis lævibus.  $\frac{1}{2}$  in siccis et ad vias. — Fl. fr. 4. p. 459.  
*α. vulgaris* (Ser.) foliis pilosis hirsutius profunde dentatis, dentibus acutis, bracteis calycis laciniisque subæqualibus conformibusque. — *P. verna*  $\alpha$  et  $\gamma$ . Ser. in DC. prod. 2. p. 575. — Clus. hist. 2. p. 106. f. 2. (v. v.)  
*β. montana* (Wallr.) hirsuta, caulibus prostratis filiformibus elongatis, lobis foliorum obovato-cuneatis profunde dentatis. — In pratis et rupestribus Delphin., Jurassi, Alsatie, Pyrenæorum. — *P. filiformis* Vill. dauph. 3. p. 564. Fl. fr. 5. p. 542. *P. heterophylla* Lapeyr. (v. v.)

- γ. *cinerea* (Ser.) caule foliisque pube stellatâ canescentibus, foliis dentatis dentibus obtusiusculis, bracteis calycis laciniis angustioribus obtusis. In Gallia boreali et Delphinatâ ex Vill. — *P. cinerea* Chaix et DC. fl. fr. 4. p. 461. — Barr. ic. 709. An merito cum *P. vernâ* conjuncta? (v. s. ex Mogantiâ.)
8. *P. OFACA* (L. sp. 713.) hirsutissima, caulibus declinatis filiformibus, foliis infimis 5-7-palmatisectis, lobis obovato-oblongis profunde dentatis, dente terminali minore, stipulis lanceolatis raro integris, floribus laxè subcorymbosis, pedunculis filiformibus, laciniis calycinis lanceolatis oblongis, petalis obovatis calycem superantibus, carpellis lævibus. ♀ in Galloprovinciâ ex Ger., Pyren. ex Lois. — Fl. fr. 4. p. 460. — Clus. hist. 2. p. 106. f. 3. (v. s. ex Pedemontio et Palatinatâ.)
9. *P. SABAUDA* (DC. fl. fr. 4. p. 459.) caulibus subdeclinatis, foliis radicalibus quinato-palmatisectis pilosis, lobis latè obovatis obtusè dentatis cuneatis, dente ultimo subprominulo, stipulis lanceolatis subacutis, floribus laxè subcorymbosis, pedunculis filiformibus, calycis laciniis lanceolatis acutis bracteis angustiores obtusiusculas superantibus, petalis obovatis calycem superantibus. ♀ in Vogesis, Alpihus, Pyrenæis. — *P. aurea* Ser. in mus. helv. 1. t. 8. f. a. b. et in DC. prod. 2. p. 576. *P. Salisburgensis*? Hænke in Jacq. rar. 3. t. 490. An characteres qui hanc a sequente distinguunt satis constantes sunt? (v. v.)
10. *P. AUREA* (L. sp. 712.) caulibus declinatis, foliis radicalibus quinato-palmatisectis nitidis, lobis obovatis acutè dentatis sericeo-ciliatis, stipulis lanceolatis subacutis, floribus laxè subcorymbosis, laciniis calycinis lanceolatis acutis, bracteis angustioribus subobtusis laciniis subæquantibus corollâ dimidio brevioribus, petalis obcordatis. ♀ in Alpibus, editioribus montibus Arvernæ, Pyrenæis. — Fl. fr. 4. p. 459. *P. Halleri* Ser. in DC. prod. 2. p. 576. — Hall. helv. t. 21. f. 4. (v. v.)
11. *P. PYRÆNAICA* (Ram. in DC. fl. fr. 4. p. 459. 5. p. 542.) caule ascendente, foliis palmatisectis inferioribus quinatis, superioribus ternatis, lobis obovato-cuneatis acutè profundeque dentatis pilosis, stipulis ovatis acutis, laciniis calycinis lanceolatis acutis bracteis linearilanceolatis æquantibus, petalis obcordatis calyce longioribus, carpellis ovoideis lævibus. ♀ in editioribus Pyrenæis. — *P. grandiflora* γ. Ser. in DC. prod. (v. v.)
12. *P. ARGENTRA* (L. sp. 712.) caule ascendente erectove tomentoso, foliis quinato-palmatis, lobis obovato-cuneatis subtus incano-tomentosis supernè plerumque glabris plus minusve incisim pinnatifidiave, stipulis ovato-acuminatis integris dentatisque, floribus corymbosis numerosis, laciniis calycinis lanceolatis corollâ brevioribus, carpellis ovatis rugulosis. ♀ in siccis et incultis. — Fl. fr. 4. p. 461. — Camer. epit. 760. ic. (v. v.)
13. *P. INTERMEDIA* (L. mant. 76. non Poir.) hirsuta, caulibus, cespitosus subprostratis dichotomis, foliis radicalibus longè petiolatis 5-7-palmatisectis pallidis, lobis obovato-lanceolatis retusis incisim dente summo breviorè, stipulis inferioribus angustissimis acutis, floribus numerosis longè pedunculatis, petalis obcordatis laciniis calycinas lanceolatas acutas æquantibus superantibusve, carpellis rugosis. ♀ in rupestribus Pyren. et montium Delphin. — Fl. fr. 4. p. 458. Nestl. pot. t. 8. Eng. bot. t. 2449. (v. s. ex Valesiâ.)
14. *P. CANESCENS* (Besser galic. 1. p. 330.) tomentoso-canescens, caule erecto firmo, foliis quinato-rarius 7-palmatisectis, lobis obovato-oblongis dentato-pectinatis, stipulis lanceolatis integris dentatisve, floribus paniculatis numerosis, petalis obcordatis laciniis calycinas lanceolato-lineares incanas æquantibus, carpellis rugosis. ♀ in Alsatiâ propè Argentoratam. — Fl. fr. 5. p. 541. (v. s.)
- β. *inclinata* (Ser.) caule inclinato debili, foliis suprâ glabriusculis. — In Delphinatâ ex Vill. — *P. inclinata* Vill. dauph. 3. t. 45. — Fl. fr. 4. p. 461. (v. v. c.)
15. *P. BERTA* (L. sp. 712.) pilosa, caule rubello panicifero erecto ascendente,

foliis 5-7-palmatisectis, lobis oblanceolatis angustis facialis dentibus subobtusis, stipulis lanceolatis acutis integris basi latis, caulinis petiolo brevioribus, floribus corymbosis lacinii calycinis lanceolato-ovatis acutis bracteis lineari-lanceolatis acutas superantibus, petalis obcordatis calycem superantibus subæquantibusve, carpellis rugosis marginulatis. 2/ in siccis et petrosis Galliæ australis et Alsatiz. — Fl. fr. 4. p. 457. All. ped. t. 71. P. pilosa Wild. Fl. fr. 5. p. 540. (v. v.)

β. *angustifolia* (Ser.) foliorum lobis cuneatis subintegris apice 3-4-dentatis, stipulis subintegris. — In Pyren. orientalibus. — P. angustifolia Fl. fr. 5. p. 540. (v. s.)

16. P. RECTA (L. sp. 711.) pilosa, caule multifloro erecto, foliis quinato-ternatoque-palmatisectis adpressè pilosis, lobis lanceolatis latis incisive dentibus acutis, stipulis lanceolato-linearibus elongatis acutis basi dentatis incisive, caulinis petiolo longioribus, calycis laciniiis bracteisque lanceolatis acutis æqualibus nervosis, carpellis rugosis marginulatis. 2/ in Galloprovinciâ ex Ger., Occitaniâ, Delph. ex Vill., Bressiâ ex Latour, Burgundiâ ex Dur. — Fl. fr. 4. p. 457. All. ped. t. 71. f. 1. P. hirta s. Ser. in DC. prod. — Lob. ic. 689. f. 2. An ut vult cl. Seringe præced. var? (v. v. c.)

§. 3. *Foliis pinnatisectis.*

17. P. FRUTICOSA (L. sp. 709.) caule fruticoso, foliis pinnatisectis hirsutis, lobis oblongo-lanceolatis integerrimis approximatis, stipulis lanceolatis membranaceis acutis, floribus subcorymbosis, lacinii calycinis pilosis lanceolatis acutis basi latis, bracteis lineari-lanceolatis basi attenuatis, corollâ calyce longiore, receptaculo pilosissimo. 3/ in Pyrenæorum valle Eynes et sepibus Alsatiz superioris. — Fl. fr. 4. p. 455. Eng. bot. t. 88. P. prostrata Lap. excl. Benth. (v. s.)
18. P. SUPINA (L. sp. 711.) caule humifuso herbaceo, foliis pinnatisectis, lobis obovatis vel oblongis glabriusculis plus minusve dentatis, stipulis lanceolatis integris, pedunculis axillaribus solitariis, lacinii calycinis triangulari-lanceolatis, bracteis lanceolatis, petalis obovatis longitudinè calycis, carpellis ovatis subcompressis. ① in glareosis vel lapidosis inundatis Galliz boreali-occidentalis. — Fl. fr. 4. p. 456. — Clus. hist. 2. p. 107. f. 2. (v. s.)
19. P. PENNSYLVANICA. (L. mant. 76.) incano-tomentosa, caule erecto herbaceo, foliis subinterruptè pinnatisectis, caulinis trilobis, lobis oblongo-obovatis profundè dentatis, stipulis lanceolatis sublaciniatis, floribus corymboso-paniculatis, lacinii calycinis triangulari-lanceolatis, bracteis oblongo-lanceolatis petala obcordata æquantibus, receptaculo piloso, carpellis rugulosis. 2/ copiosè nunc spontanea circâ Parisios in sylvâ Bononiensi ubi seminibus ex Americâ boreali certò introducta. — Fl. fr. 4. p. 456. in not. — Jacq. vind. 2. t. 189. — Gmel. sib. 3. t. 34. f. 1. (v. s.)
20. P. ANSERINA (L. sp. 710.) caulibus filiformibus radicanthibus, foliis interruptè pinnatisectis, lobis obovato-oblongis plus minusve profundè argute serratis, suprâ glabriusculis subtùs sericeis, stipulis caulinis multifidis, floribus solitariis geminisve, pedunculis longitudine foliorum, lacinii calycinis lanceolatis integris, bracteis sapius 3-5-lobis, petalis obovatis calyce longioribus, receptaculo piloso. 2/ frequens ad vias et in subhumidis. — Fl. fr. 4. p. 455. — Camer. epit. 758. ic. — Moris. hist. s. 2. t. 20. n. 4. (v. v.)

Sect. II. COMARUM. Ser. in DC. prod. 2. p. 583.

Petala acuminata rubra; folia pinnatisecta, bractes minores deflexæ.

21. P. COMARUM (Scop. carn. ed. 2. t. p. 359.) lobis foliorum approximatis, stipulis latis coriaceis, petalis calyce minoribus, carpellis ovato-compressis lævibus. 2/ in paludosis Galliz occid., mediæ, borealis orientalisque. — Comarum palustre. L. sp. 718. Fl. fr. 4. p. 469. Lam. ill. t. 444. (v. v.)

Sect. III. FRAGARIASTRUM. Ser. l. c.

Petala obtusa obcordatave alba vel rubra; folia pinnata vel palmatisecta.

§. 1. *Foliis pinnatisectis.*

22. *P. AUFRESTRIS* (L. sp. 711.) caule erecto dichotomo virgato, foliis pilosis radicalibus pinnatisectis, caulinis trilobis, lobis ovato-subrotundis dentatis, stipulis latis brevibus obtusis, laciniis calycinis latis lanceolatis, bracteis minimis sæpè 2-lobis, petalis obovatis calyce multò longioribus, carpellis lævibus. ♀ in saxosis montosis. — Fl. fr. 4. p. 464. Jacq. austr. t. 114. — J. Bauh. hist. 2. p. 398. d. f. 2. (v. v.)

β. *pygmaea*, procumbens, vix 2-pollicaris, caule simplici 1-2-floro. — In pascuis humidis montium Corsicæ. (cl. Soleirol.) (v. s.)

§. 2. *Foliis palmatisectis.*

23. *P. ALBA* (L. sp. 713.) caulibus debilibus procumbentibus paucifloris, foliis radicalibus quinato-palmatis, caulinis ternato-palmatis, lobis tenuibus ovato-oblongis apice serratis serraturis subconniventibus, suprà glabriusculis subtùs sericeis, stipulis linearibus acutis, laciniis calycinis lineari-lanceolatis acutis sericeis petala obcordata æquantibus, receptaculo piloso, carpelliis rugulosis glabris. ♀ in sabulosis et graminosis montosis Gallis orientalis. — Fl. fr. 4. p. 466. — Clus. hist. 2. p. 105. f. 1. (v. s.)

β. *splendens* (Ser.) lobis foliorum ternis quinque latioribus suprà sericeo-pilosis, sepalis latioribus. — In arenosis et graminosis Gallis occident., Pyren., et circà Parisios. — *P. splendens* Ram. in DC. fl. fr. 4. p. 467. (v. v.)

24. *P. ALCHIMILLOIDES* (Lapeyr. act. toul. 1. t. 17.) radice crassà, caulibus firmis crassiusculis suberectis, foliis radicalibus 5-7-palmatis, caulinis 3-lobatis, lobis subcoriaceis oblongis apice vix denticulatis suprà glabris subtùs incano-tomentosis, stipulis linearibus acutis, laciniis calycinis lanceolatis tomentosis, exterioribus linearibus, petalis obcordato-subrotundis vix calyce longioribus, receptaculo piloso, carpellis valdè pilosis. ♀ in-Pyrenæis. — Fl. fr. 4. p. 466. (v. v.)

25. *P. CAULESCENS* (L. sp. 713.) caulibus tomentosis suberectis multifloris, foliis radicalibus quinato-palmatisectis, superioribus ternatisectis, lobis obovato-cuneiformibus apice dentatis vix retusis ciliatis, stipulis inferioribus longissimis angustissimisque, superioribus ovato-lanceolatis, floribus subcorymbosis laxis, sepalis lanceolato-linearibus acutis petala obovato-cuneata sub-æquantibus, receptaculo piloso, carpellis non rugosis pilosis. ♀ in fissuris rupium montium. — Fl. fr. 4. p. 464. Jacq. austr. t. 220. (v. v.)

β. *nivalis* (Ser.) caule foliis pedunculisque villosissimis, stipulis superioribus dilatatis, floribus corymboso-glomeratis. — In Corsicà, Pyrenæis et editioribus Alpibus Galloprovinciæ, Delphin. — *P. nivalis* Lapeyr. act. toul. 1. p. 210. t. 16. Fl. fr. 4. p. 465. (v. v.)

26. *P. NITIDA* (L. sp. 714.) sericea, caulibus cespitosis subbifloris, foliis radicalibus subquinato-palmatisectis, superioribus ternato-palmatis, lobis obovato-cuneiformibus sericeis integris apiceque tridentatis, stipulis angustis, laciniis calycinis lanceolatis acutis, bracteis angustissimis, petalis obcordatis latissimis calyce longioribus, receptaculo lauto, carpellis non rugosis valdè pilosis. ♀ in montibus Delphinatùs. — Fl. fr. 4. p. 467. Jacq. aust. app. 3. 25. — Bocc. mus. t. 9. f. 4. Fl. magni. (v. s.)

27. *P. MICRANTHRA* (Ram. in DC. fl. fr. 4. p. 468.) pubescens, non stolonifera, foliis radicalibus ternato-palmatisectis, lobis ovato-rhomboidæis dentatis subsessilibus subtùs glaucescentibus, terminali subretuso, foliis caulinis 1-lobis, stipulis latis brevibus, laciniis calycinis ovatis, bracteis subdentatis, petalis obcordato-cuneatis subacuminatisque calyce brevioribus, receptaculo villosq. carpellis rugosis. ♀ in Pyrenæis. — Ser. mus. helv. 1. p. 60. t. 5. Fl. albi vix aperti. (v. v.)

28. *P. FRAGARIA* (Poir. dict. 5. p. 599.) pubescens stolonifera, caulibus decumbentibus 2-floris, foliis radicalibus caulinisque ternato-palmatisectis,



lobis obovato-subrotundis, intermedio retuso dente summo minore, stipulis latis membranaceis obtusiusculis, laciniis calycinis lanceolatis, bracteis oblongis, petalis obcordatis calycem æquantibus superantibusve, receptaculo subvillosa, carpellis non rugosis.  $\mathcal{L}$  in sylvis et aridis. — Fl. fr. 4. p. 468. — *Fragaria sterilis* L. sp. 709. — Lob. ic. t. 698. f. 1. (v. v.)

XII. SIBBALDIA. L. gen. n. 393.

Omnia Potentillæ sed petala minuta; carpella et stam. sæpius 5; calyx bracteoliferus; semen in akenio inversum, receptaculum exsuccum. — Herbæ humiles.

1. *S. procumbens* (L. sp. 406.) caulibus procumbentibus, foliis 3-foliolatis, foliolis obovatis apice tridentatis superne glabris subtus pilosis, floribus corymbosis, petalis lanceolatis acutis vix longitudine calycis.  $\mathcal{L}$  in editioribus Alpibus et Pyrenæis. — Fl. fr. 4. p. 453. Lam. ill. t. 221. f. 1. Fl. dan. t. 32. (v. v.)

XIII. AGRIMONIA. Tourn. inst. t. 155. Nestl. pot. 16. — Agrimonie sp. L.

Cal. tubus turbinatus foveolatus suprâ lappaceus, extus setis involucriformibus onustus, limbo 5-fido; pet. 5; stam. 15; carpella 2 stylo terminali; akenia tubo cal. indurato inclusa membranacea, semine appenso. — Herbæ foliis pinnatisectis.

1. *A. EUPATORIA* (L. sp. 643.) hirsuta, foliis interruptè pinnatisectis, lobis oblongo-ovatis crenato-dentatis, petalis calyce duplò longioribus, fructibus distantibus, tubo calycis campanulato piloso setoso, setis patulis, foveolis usque ad basin productis.  $\mathcal{L}$  ad margines viarum sylvarumque. — Fl. fr. 4. p. 450. Lam. ill. t. 409. f. 1. Fl. dan. t. 588. (v. v.)
2. *A. ODORATA* (Camer. hort. 7.) hirsuta, foliis interruptè pinnatisectis, lobis oblongo-lanceolatis profundè crenato-dentatis, petalis calyce duplò longioribus, fructibus distantibus, tubo calycis campanulato piloso, foveolis obovatis usque ad basin productis, setis ascendentibus brevibus.  $\mathcal{L}$  circà Parisios, in Andegaviâ et agro Mimatensi (Mende) (Prost). — Fl. fr. 4. p. 451. (v. s.)

Trib. IV. SANGUISORBEÆ. Juss. gen. p. 336. excl. gen. DC. prod. 2. p. 588.

Flores sæpè polygamo-dioici. Calyx 3-5-fidus, lobis per æstivationem valvatis, tubo apice contracto carpella foverit et cum iis demùm sæpè subaccreto. Pet. 0 aut 4 basi in corollam rotatam coalita. Stam. tot quot lobi calycini; nonnulla sæpè abortiva. Carpella pauca 1-2, stylo sublaterali, stigmate rariùs capitato sæpius penicilliformi aut barbato. Akenia uni-ovulata sicca indehiscentia. Semen inversum. — Herbæ aut suffrutices, foliis sæpius compositis, floribus parvis.

XIV. ALCHIMILLA. Tourn. inst. t. 289. DC. prod. l. c. — Alchemilla et Aphanes. L. gen. n. 165 et 166.

Calyx tubulosus tubo apice subcontracto, limbo 8-partito, bracteis alternis minoribus interdum minimis dentiformibus; pet. 0; stam. 1-4; carpella 1-2 lateraliter stylum filiformem apice capitatum gerentia, demùm sicca indurata 1-sperma; semen inversum. — Herbæ, foliis palmatim lobatis aut sectis.

Sect. I. *ALCHEMILLA*. *L. l. c.*

Cal. 8-fidus, lobis alterdis parum minoribus; stam. 2-4. — Species perennes.

\* *Foliis 7-9-nerviis 7-9-fidis.*

1. *A. VULGARIS* (L. sp. 178.) foliis reniformibus plicato-concavis 9-lobis serratis, caule petiolisque glabriusculis, floribus dichotomis corymbosis.  $\frac{1}{2}$  in sylvis et pascuis frequens. — Fl. dan. t. 963. Fl. fr. 4. p. 451. excl. var.  $\beta$ . *Stipulæ incisæ*. (v. v.)
2. *A. HYBRIDA* (Hoffm. germ. 1. p. 79.) foliis reniformibus 9-lobis serratis subtus caule petiolisque sericeis, floribus fastigiatis glomeratis sessilibus.  $\frac{1}{2}$  in pascuis siccis montanis. — *A. vulgaris hybrida* Wild. Fl. fr. 4. p. 451. — Pluck. t. 240. f. 2. *Stipulæ serratæ*. (v. v.)

\*\* *Foliis digitato-5-7-sectis, foliolis apice serratis.*

3. *A. ALPINA* (L. sp. 179. var.  $\alpha$ .) foliis digitato-5-7-sectis, foliolis lanceolato-cuneatis obtusis apice adpresso-serratis subtus sericeis.  $\frac{1}{2}$  in pascuis montium. — Fl. fr. 4. p. 452. Eng. bot. t. 244. — Cam. epit. 909. ic. (v. v.)

\*\*\* *Foliis digitato-3-5-sectis, foliolis multifidis aut per totum ambitum serratis.*

4. *A. PENTAPHYLLA* (L. sp. 179.) foliis 3-5-sectis, segmentis multifidis glabris subciliatis, stipulis foliaceis apice 2-3-dentatis.  $\frac{1}{2}$  in pascuis excoelsis Alpium Delphinatūs. — Fl. fr. 4. p. 452. — Bocc. mus. p. 18. t. 1. (v. v. in Helv.)

Sect. II. *APHANES*. *L. l. c.*

Cal. 4-fidus, rariùs 5-fidus, dentibus minimis inter lobos interjectis; stam. 1-2 fertilia, cætera abortiva. — Species annuæ.

5. *A. ARVENSIS* (Scop. carn. 1. p. 115.) foliis pubescentibus petiolatis 3-partitis, lacinii 2-3-fidis, floribus axillaribus glomeratis. ① in arvis et agris. — Fl. fr. 4. p. 453. — *Aphanes arvensis* L. sp. 179. OEd. fl. dan. t. 973. (v. v.)

XV. *SANGUISORBA*. *L. gen. n. 146.*

Flores hermaphroditi; cal. 4-fidus extus basi bisquamosus; pet. 0; stam. 4; carpella 2 intrâ calycis tubum inclusa, stylo apice penicilliformi superata, in akenia sicca indehiscencia monosperma conversa; semen inversum. — Herbæ perennes, floribus in spicas densissimas aggregatis.

1. *S. OFFICINALIS* (L. sp. 169.) spicis ovatis, staminibus calyci æqualibus, calycibus bracteis folisque glabris, foliolis ovato-subcordatis.  $\frac{1}{2}$  in pratibus. — Fl. fr. 4. p. 449. Eng. bot. 1312. (v. v.)

XVI. *POTERIUM*. *L. gen. n. 106g.*

Flores monoici seu polygami; calyx basi 3-squamosus, tubo apice coarctato, limbo 4-partito; pet. 0; stam. 20-30; ovaria 2 stylo filiformi et stigmate penicilliformi superata; akenia intrâ calycem induratum sicca 1-sperma; semen inversum. — Flores in spicas dense aggregati.

1. *P. SANGUISORBA* (L. sp. 1411.) herbaceum, caulibus subangulosis folisque glabris, foliolis ovato-subretundis, spicis globosis, floribus inferioribus masculis superioribus feminis.  $\frac{1}{2}$  in pratibus asperis. — Fl. fr. 4. p. 449. Cam. epit. 777. (v. v.)

$\beta$ . *puberulum* (DC.) caulibus folisque hinc inde puberulis. (v. v.)

Trib. V. *ROSEÆ*. DC. *prod.* 2. p. 596.

Calycis tubus apice contractus, limbo 5-partito, lobis per æstivationem apice subspiraliter imbricatis sæpè pinnatim sectis. Pet. 5. Stam. ∞. Carpella plurima calycis tubo demùm baccato inserta et in eo inclusa sicca indehiscencia subcrustacea, è latere interiore stylum gerentia, stylis è calycis tubo coarctato exsertis, nunc omnino liberis, nunc in stylum columnarem accretis. Semen in akenio solitarium exalbuminosum inversum. Embryo rectus, cotyledonibus planiusculis. — Frutices aut arbusculæ.

XVII. ROSA. *Tourn. inst. t.* 408. *L. gen. n.* 631. *Ser. in DC. prod.* 2. p. 597.

Character isdem ac tribûs.

Sect. 1. *SYNSTYLÆ*. DC. *cat. monsp.* p. 137.

Styli in columnam cohærentes; fructus ovati vel subglobosi; folia sæpè sempervirentia; stipulæ adnatæ.

1. *R. ARVENSIS* (L. mant. 245.) surculis flagelliformibus, aculeis inæqualibus falcatis, foliis 5-7-nis subtus glaucescentibus glabris vel subciliatis deciduis, floribus solitariis vel corymbosis, sepalis subintegris brevibus, stylis in columnam elongatam glabram cohærentibus, fructibus ovatis vel ovato-globosis coriaceis pedunculisque glabris vel hispidulis. ♀ in sylvis et sepius. Stipulæ apice divergentes, fructus carnosini.

α. *vulgaris* (Ser.) foliolis ovatis acutis. — *R. arvensis* α. Fl. fr. 4. p. 438. Eng. bot. t. 188. — J. Bauh. 2. p. 44. f. 1. (v. v.)

β. *bibracteata* (Red. et Thor.) pedunculis bi-vel-multibracteatis, floribus numerosioribus, foliis latiusculis. — *R. bibracteata* Bast. in DC. fl. fr. 5. p. 537. (v. s.)

γ. *prostrata* (Desv.) foliis crassiusculis sempervirentibus. — In dumetis Tolosæ. — *R. prostrata* Fl. fr. 5. p. 536. (v. v. DC.)

2. *R. SEMPERVIRENS* (L. sp. 704.) surculis scandentibus, aculeis subæqualibus falcatis, foliis persistentibus 5-7-nis coriaceis, floribus subsolitariis corymbosisque, sepalis subintegris longiusculis, stylis in columnam elongatam pilosam cohærentibus, fructibus ovatis vel ovato-globosis pedunculisque sæpissimè glanduloso-hispidis. ♀ in sepius ad vias et sylvas. — Fl. fr. 5. p. 533, 4. p. 446. Stipulæ apice subfalcatæ acutiores, fructus aurantiaci.

α. *scandens* (DC.) pedunculis fructibusque hispidulis. — In provinciis mediterraneis et austro-occidentalibus. — *R. atrovirens* Viv. fragm. 1. p. 4. t. 6. (v. v.)

β. *microphylla* (DC.) foliolis minoribus, pedunculis fructibusque hispidulis. — in aridis apricis circà Monspelium. (v. v.)

3. *R. MOSCHATA* (Mill. dict. n. 13.) surculis ascendentibus, aculeis caulinis tenuibus recurvis, foliolis 5-7 lanceolatis acuminatis glabriusculis, stipulis angustissimis acutis, floribus sæpè numerosissimis, pedunculis lateralibus articulatim calycibusque pilosis subhispidis, sepalis subpinnatisectis elongato-appendiculatis, fructibus ovatis rubris. ♀ ex agro Tunetano orta? in hortis colitur. — Fl. fr. 4. p. 447. Red. et Thor. ros. 1. p. 33 et 99 ic. — J. Bauh. 2. p. 46 ic. Fl. albi odoratissimi. (v. v. c.)

4. *R. STYLOSÀ* (Desv. journ. bot. 1809, v. 2. p. 317.) surculis assurgentibus, aculeis validis aduncis, foliolis 5 ovato-acuminatis argutè serratis pubescentibus, stipulis amplis breviter acuminatis, petiolis tomentosis, floribus subsolitariis, sepalis pinnatisectis subappendiculatis, stylis in columnam glabram clavatam cohærentibus. ♀ Fl. pallide rosei.

α. *Desvauxiana* (Ser.) foliis pubescentibus non glandulosis, pedunculis

calycibusque glabris. — In Pictaviâ superiore. — Desv. journ. bot. 1813. v. 2. t. 14. Fl. fr. 5. p. 536. (v. s.)

- β. *leucochroa* (Ser.) foliis glabriusculis non glandulosis, pedunculis hispidulis, calycibus glabris. — in Pictaviâ superiore et in Andegaviâ. — R. leucochroa Desv. journ. bot. 1819, t. 15. R. brevistyla a. et γ. Fl. fr. 5. p. 537. (v. s.)

Sect. II. CHINENSIS. DC. in Ser. mus. helv. 1. p. 2.

Styli liberi subinclusi; sepala integrinscula deflexa; fructus ovati vel globosi; folia persistentia nitida coriacea; stipulæ subliberæ subulatæ vel angustissimæ sæpius deciduæ.

5. R. *INDICA* (L. sp. 705.) caulibus erectis subcanescentibus viridibus vel purpureis, aculeis validis falcatis distantibus, foliolis 3-5 ovato-acuminatis coriaceis glabris serrulatis, stipulis valdè angustis cum petiolo coalitis integrinsculis vel serratis, staminibus inflexis, pedunculis subarticulatis sæpè incrassatis calycibusque lævibus vel rugoso-setosis. ♂ Ex Chinâ orta, nunc in hortis frequentissima. — Red. et Thor. ros. 1. p. 49 et 51 ic. (v. v. c.)

Sect. III. CINNAMOMÆ Ser. mus. helv. 1. p. 2. et DC. prod. 2. p. 602.

Styli liberi inclusi rarè exserti; sepala integerrima rarè subpinnatisecta post anthesin sæpè conniventia; stipulæ nullæ aut adnatæ; aculei stipulares gemini; fructus globosi aut globoso-depressi.

6. R. *TURKNATA* (Ait. kew. ed. 1. v. 2. p. 206.) caule subinermi, ramis lævibus, foliolis 5-7 ovato-cordatis bullato-rugosis amplis simpliciter serratis subtus villosulis, stipulis maximis amplexicaulibus, floribus subcorymbosis amplis, pedunculis hispidulis, calycibus turbinate sublævibus, sepalis indivisis subspathulatis, fructibus globosis. ♂ Patr. ign. — Fl. fr. 4. p. 441. Red. et Thor. 1. p. 127. ic. Fl. rubro-violaceæ. (v. v. c.)

7. R. *GALLICA* (L. sp. 704.) aculeis inæqualibus, stipulis angustis apicè divaricatis, foliolis 5-7 coriaceis rigidis ovatis vel lanceolatis deflexis, sepalis per anthesin patentibus, fructibus subglobosis valdè coriaceis calycibus pedunculisque plus minus glanduloso-hispidis subviscosis. ♂ in dumetis apricis.

α. *pumila* (DC.) foliolis ovato-subrotundis rariter lanceolatis, stipulis angustissimis apicè acutis, ramis plus minus aculeatis, pedunculis calycibusque hispidulo-glandulosis. — In dumetis imprimis provinc. austr. passim. — R. pumila Jacq. austr. 2. t. 198. (v. v.)

β. *officinalis* (Red. et Thor. 1. p. 73. ic.) foliolis ovatis lanceolatisve, pedunculis calycibusque glandulosis, sepalis vix pinnatisectis. — In hortis culta. (v. v. c.)

γ. *parvifolia* (Ser.) parvula subinermis, foliolis minimis subtus petiolisque puberulis, tubo calycis apicè dilatato pedunculisque sublævibus. — In montibus Divionensibus ex Dur. et in hortis culta. — R. Remensis fl. fr. 4. p. 443. (v. v. c.)

8. R. *CINNAMOMÆ* (L. sp. 703.) cinerea, ramis strictis, aculeis geminis falcatis stipularibus, foliolis 5-7 ovato-ellipticis serrulatis subtus pubescenti-cinereis, stipulis dilatatis undulatis jugo foliolorum inferiorum valdè proximis, sepalis apicè spathulatis patulis, pedunculis brevibus, fructibus globosis vel sphaerico-ovatis sepalis conniventibus coronatis. ♂ in Galliâ mediâ et austro-orientali passim. — Fl. fr. 4. p. 439. — J. Bauh. hist. 2. p. 39. f. 1. (v. v.)

9. R. *EGLANTERIA* (L. sp. 703.) caulibus junioribus aculeatissimis, adultis armatis aculeis sparsis rarè stipularibus, foliolis ovatis obovatisve argutè serratis suprâ lævibus subtus glandulosis, stipulis angustis integrinsculis apicè divergentibus, pedunculis calycibusque lævibus, sepalis patentibus pinnatifidis, fructibus globosis. ♂ Patria ign.

α. *lutea* (Red. et Thor.) petalis concoloribus, stigmatibus purpureis. — In

- sepibus propè Avenionem (Requien) et in Galloprovinciâ ex Vill. — Fl. fr. 4. p. 437. (v. s. c.)
- β. *punicea* (Red. et Thor.) petalis suprâ puniceis infrâ luteis, stigmatibus purpureis. — R. bicolor Jacq. vind. t. 1. (v. v. c.)
10. R. *SULFURÆA* (Ait. kew. 2. p. 201.) caule juniore aculeatissimo, aculeis ramorum sparsis tenuibus falcatis, foliolis 5-7 glaucis obovatis denticulatis glabris, stipulis planiusculis angustis subfimbriatis apice dilatatis, calyce globoso aut hemisphærico, sepalis subindivisis. ζ ex Oriente verosimiliter indigena, in hortis culta. — Fl. fr. 4. p. 438. Red. et Thor. 1. p. 29. ic. (v. v. c.)
11. R. *PIMPINELLIFOLIA* (L. sp. 307.) caule juniore aculeatissimo, aculeis inæqualibus horizontalibus, foliolis 5-9 ovato-rotundis coriaceis tenuibus dentatis, stipulis angustis apice dilatatis, sepalis indivisis patentibus, pedunculis sæpissime incrassatis, fructibus coriaceis sepalis convergentibus coronatis. ζ in dumetis passim. Fructus nigri. — Fl. fr. 4. p. 438.
- α. *vulgaris* (Ser.) pedunculis fructibusque lævibus, floribus albis. — R. spinosissima Jacq. fragm. 79. t. 124. — Clus. hist. 1. p. 116. f. 1. 2. (v. v.)
- β. *myriacantha* (Ser.) foliolis minimis, aculeis valdè numerosis, inferioribus sæpè reversis, pedunculis calycibusque hispidis. — Propè Lugdunum et Mompelium. — R. myriacantha Fl. fr. 4. p. 439 et 5. p. 533. Red. et Thor. 3. p. 11. ic. Fl. albi. (v. v. DC.)
- γ. *inermis* (DC.) caule ramisque inermibus. — In montosis rariùs. (v. v.)
12. R. *RUBRIFOLIA* (Vill. dauph. 3. p. 549.) glauca pruinosa, aculeis paucis falcatis, foliolis ovatis dentatis 5-7, stipulis latiusculis acuminatis, pedunculis subcorymbosis brevibus hispidulis glabrisve, sepalis subspathulatis rarò pinnatisectis corollam subsuperantibus, fructibus ovato-globosis nitido-subpelucidis. ζ in Alpibus et montanis. Fructus rubri.
- α. *lævis* (Ser.) foliolis ovatis, floribus rubris, fructibus corymbosis pedunculisque lævibus, sepalis integris. — In pratis Vogesorum, Alpium, montium Arvernæ, Cebenn., Pyren. — Fl. fr. 4. p. 445. Red. et Thor. 1. p. 31. (v. v.)
- β. *pinnatifida* (Ser. mus. helv. 1. p. 11. t. 2. f. 11.) foliolis ovatis, floribus solitariis terminalibus, sepalis pinnatifidis, fructibus globosis pedunculisque lævibus. — In Cebennis et Pyrenæis.
13. R. *GLANDULOSA* (Bell. act. tur. 1790. p. 230.) glauca, junior rubicunda, aculeis geminis stipularibus rectis validis, foliolis 5-7 ovato-rotundis duplicato-serratis stipulisque latiusculis acutis margine glandulosis, floribus subsolitariis, tubo calycis subgloboso pedunculisque glanduloso-hispidis, sepalis integris longè subspathulatis corollam superantibus. ζ in Alpibus Delph. propè Brigantium. — Fl. fr. 5. p. 539. R. rubrifolia var. ζ. Ser. (v. v. DC.)

Sect. IV. CANINÆ Ser. in DC. prod. 2. p. 611.

Styli liberi inclusi vel exserti; sepala pinnatifida post anthesin deflexa sæpissime decidua; fructus sæpiùs ovatus; stipulæ adnatæ, aculei sparsi non stipulares.

§. 1. Folii non glandulosi.

14. R. *ALPINA* (L. sp. 703.) caulibus junioribus aculeis debilibus numerosis echinatis, adultis lævibus, foliolis 5-11 ovatis obovatisve argutè serratis vel duplicato-serratis, stipulis angustis apice divergentibus, pedunculis post anthesin deflexis, sepalis indivisis patulis, fructibus ovatis rarò subglobosis pendulis plus minusve elongatis calyce connivente coronatis. 4 in montibus.
- α. *vulgaris* (Desv.) pedunculis hispidis elongatis, calycibus lævibus. — R. Alpina DC. fl. fr. 4. p. 446. — J. Bauh. 2. p. 39. f. 2. (v. v.)
- β. *Pyrenaica* (DC.) pedunculis elongatis calycibusque hispidis. — R. Alpina δ. Fl. fr. 5. p. 536. R. Pyrenaica Fl. fr. 4. p. 446. (v. v.)
15. R. *CANINA* (L. sp. 704.) aculeis distantibus validis compressis falcatisque,

foliolis subcoriaceis acutè et sæpè duplicato-serratis, stipulis latiusculis acutis tenuissimè serrulatis, sepalis pinnatifidis post anthesin deflexis et dein deciduis, fructibus ovatis rarè subglobosis coriaceis.  $\S$  in sepibus dumetis sterilibusque frequens. — Fructus punicei.

*a. glabra* (Desv.) petiolis glabris vix puberulis, foliolis ovatis acutis glabris, fructibus ovatis pedunculisque lævibus. Fl. fr. 4. p. 447. — R. nitens Desv. in Mer. fl. par. 192. (v. v.)

*ß. leucantha*, petiolis puberulis sæpè glandulosis, foliolis ovato-subrotundis acuminatis subtus pilosulis, fructibus ovatis pedunculisque lævibus. — In Neustriâ, Andegaviâ, Pictaviâ, Occitaniâ. — R. leucantha Lois. not. 82. DC. fl. fr. 5. p. 535. (v. s.)

*γ. fastigiata* (Desv.) aculeis validis, petiolis puberulis, foliolis ovatis acutis subtus puberulis, fructibus ovatis lævibus, pedunculis hispidis. — In Andegaviâ et Delphinatâ. — R. fastigiata Bast. supp. 30. Fl. fr. 5. p. 535. Styli approximati non coacti brevissimè exserti. (v. v. DC.)

*δ. hispida* (Desv.) ramis aculeatis, petiolis glabresculis, foliolis ovatis acutis glabris, fructibus ovatis pedunculisque hispidulis. — In agro Aurelianensi, Andegaviensi, Mimatensi. — R. Andegavensis Bast. ess. 189. Fl. fr. 5. p. 539. (v. s.)

*ε. dumetorum* (Desv.) ramis aculeatis, petiolis tomentosis lanatisque, foliolis ovatis acutiusculis subtus puberulis tomentosive, fructibus ovatis pedunculisque sublævibus. — R. dumetorum Thail. fl. par. ed. 2. p. 250. Fl. fr. 5. 534. R. collina Jacq. aust. 2 t. 197. Fl. fr. 4. p. 441.

### $\S$ . 2. *Foliis subtus plus minùsve glandulosis.*

16. R. *aveternosa* (L. mant. 564.) aculeis validis compressis uncinatis rarè rectis, foliolis 5-7 ovatis vel subrotundis serratis subtus glanduloso-rubiginosis, ovaris ellipticis brevibus pedunculisque hispidis duris.  $\S$  in sepibus et dumetis. Fl. roseo-purpurei. Fructus punicei.

*a. vulgaris* (Wild.) germinibus subsolitaris pedunculisque plus minùsve glanduloso-hispidis, foliolis ovatis obtusis glabris glandulosisque, aculeis aduncis. — R. rubiginosa L. mant. 564. Fl. fr. 4. p. 445 et 5. p. 525. var.  $\beta$ . Eng. bot. 991. (v. v.)

*ß. umbellata* (Lindl.) floribus 3-6 fasciculatis, fructibus ovato-globosis sublævibus, pedunculis hispidis, foliolis ovato-rotundis glabris subtus glandulosis, aculeis aduncis. — R. umbellata Fl. fr. 5. p. 532. — Red. et Thor. 1. p. 93. ic. (v. v.)

*γ. parvifolia* (Wild.) pumila, aculeis numerosis, floribus subsolitaris, fructibus ovato-globosis pedunculisque sublævibus, foliolis minimis subtus rubiginoso-glandulosis glabris. — Monapellii in aridis montis Lapi. — R. micrantha Fl. fr. 5. p. 539. non Sm. Flores alborosei. (v. s.)

*δ. sepium* (Ser.) aculeis validis, foliolis obovatis ellipticis lanceolatisve supra glabris subtus glandulosis, petiolo glabro, floribus solitariis vel corymbosis, fructibus ovatis pedunculisque lævibus. — In collibus siccis. — R. sepium Thail. Fl. fr. 5. p. 538. — Red. et Thor. ros. 2. p. 61. ic. et 3. p. 51. ic. (v. v.)

17. R. *tomentosa* (Sm. brit. 2. p. 539.) aculeis tenuibus basi vix compressis subaduncis, foliolis ovatis subduplicato-serratis plus minùsve tomentosis, serratis rectis, fructibus ovatis pedunculisque plus minùsve hispidis.  $\S$  in montanis. — Fr. punicei.\*

*a. Smithiana* (Ser.) foliolis ovatis vel ovato-lanceolatis molliter tomentosis, pedunculis fructibusque plus minùsve hispidulis. — R. tomentosa Fl. fr. 4. p. 440. eng. bot. t. 990. (v. v.)

*ß. scabriuscula* (Ser.) foliolis ovatis supra scabriusculis subtus tomentosis, fructibus ovatis pedunculisque plus minùsve hispidis. — In Juraso. — R. scabriuscula Sm. eng. bot. t. 1896. — R. montana Fl. fr. 5. p. 532. non Vill. (v. s.)

*γ. fistida* (Ser.) foliolis ovatis infra pubescentibus supra glabresculis ru-

gulosis, pedunculis terminalibus subsolitariis hispidis, fructibus ovatis foetidis. — Ad ripas Ligeris. — R. foetida Bast. supp. 29. Fl. fr. 5. p. 634. Red. et Thor. 1. p. 131. ic. (v. s.)

18. *R. villosa* (L. sp. 704.) aculeis tenuibus basi vix compressis rectis, foliolis ovatis subduplicato-serratis plus minus tomentosis, serraturis rectis, ovarii globosis sæpius valde hispido-aculeatis subpendulis. § in calidis. — Fructus purpurei maximi. — Fl. fr. 4. p. 440.

a. *sylvestris* (Desv.) foliolis lanceolatis ellipticisve molliter villosis dense glandulosis, calycibus ovato-globosis valde aculeatis. — In collibus siccis. — J. Bauh. hist. 2. p. 38. f. 1. (v. v.)

β. *nuda* (Desv.) foliolis ovatis obtusis villosis glutinosis, fructibus obovato-globosis lævibus, pedunculis hispidulis. — In sterilibus. (v. v.)

19. *R. centifolia* (L. sp. 704.) aculeis subrectis basi vix dilatatis, foliolis 5-7 ovatis margine glandulosis flaccidiusculis subtus pilosis, stipulis latiusculis acuminatis, sepalis per anthesin patentibus, fructibus conico-ovatis calycibus pedunculisque glanduloso-hispidis viscosis fragrantibus. § Patr. ign., in hortis colitur.

a. *vulgaris* (Ser.) aculeis aparsis, foliolis ovatis planis serratis, sepalis ciliatis tomentosis. — Fl. fr. 4. p. 442. Red. et Thor. 1. p. 77. ic. et p. 25. ic. (v. v. c.)

β. *muscosa* (Ser.) aculeis inæqualibus parvis numerosis, foliolis ovatis planis serratis, pedunculis calycibusque glanduloso-mucosis. — R. muscosa Fl. fr. 4. p. 442. — Red. et Thor. 1. p. 39. et 41. ic. (v. v. c.)

γ. *Pomponia* (Lindl.) parvula, foliolis floribusque minimis. — R. Pomponia Fl. fr. 4. p. 443. Red. et Thor. 1. p. 65. ic. (v. v. c.)

20. *R. DAMASCENA* (Mill. dict. n. 15.) aculeis numerosis inæqualibus validis basi dilatatis, foliolis 5-7 ovatis rigidiusculis, sepalis per anthesin deflexis, ovario elliptico elongato apice sæpe dilatato, calycibus pedunculisque glanduloso-hispidis viscosis, fructibus ovatis. § ex Syria orta, in hortis colitur. — Red. et Thor. 1. p. 63. 53. 37. ic. R. semperflorens Desf. cat. 175. Fl. fr. 4. p. 443. non Curt. nec Willd. Fl. plus minus corymbosi suaveolentes.

21. *R. ALBA* (L. sp. 705.) glaucescens, aculeis tenuibus recurvis, foliolis sæpissime ovato-rotundis breviter acuminatis, petiolis nervisque foliorum subtomentosis glandulosisque, sepalis pinnatifidis, petalis patentibus, fructibus ovatis. § in hortis frequentissimè culta. — Fl. fr. 4. p. 448. — Fl. albi vel pallide rosei. (v. v. c.)

Trib. VI. *POMACEÆ* Juss. gen. p. 384. DC. prod. 2. p. 626.

Calycis tubus campanulatus urceolatusque ad maturitatem carnosus carpella fovens et cum iis adhærens; limbus 5-lobus. Pet. 5. fauci calycis inserta decidua, æstivatione quinquunciali. Stam. ∞ æstivatione inflexa. Discus sæpius carnosus. Ovaria sæpius 5 unilocularia in stylos totidem simplices coalitosve desinentia. Pomum calyce baccato carpellisque constans. Carpella cartilaginea aut ossea bivalvia aut indehiscientia. Semina sæpius in quoque carpello 1-2; plurima in Cydonia, erecta, spermodermio cartilagineo aut osseo. Cotyledones ovales carnosæ. — Frutices aut Arbores, foliis simplicibus rariùs pinnatis stipulatis.

XVIII. *CRATÆGUS* Lindl. trans. lin. 13. p. 105. DC. prod. 2. p. 626. — *Cratægi* et *Mespili* sp. auct.

Cal. tubus urceolatus, limbus 5-fidus; pet. patentia orbiculata; ovarium 2-5-loculare: styli totidem glabri; pomum carnosum ovatum

dentibus calycinis vel disco incrassato clausum, putamine osseo. — Frutices spinosi, foliis angulatis aut dentatis.

§. 1. *Foliis dentatis subintegrise nec angulato-lobatis.*

1. C. PYRACANTHA (Pers. ench. 2. p. 37.) foliis glabris persistentibus ovato-lanceolatis crenatis, calycis lobis obtusis, stylis 5. ♂ in asperis et sepibus provinc. australiorum et austro-occident. usque ad Burdigalam ex Laterrade. — *Mespilus Pyracantha* L. sp. 685. Fl. fr. 4. p. 434. — Lob. ic. 2. p. 182. f. 1. (v. v.)

§. 2. *Foliis sæpius variè lobatis incisive.*

2. C. OXYACANTHA (L. sp. 683.) foliis obovato-cuneiformibus subintegrise trifidis laciniatisve glabris subnitidis, floribus corymbosis 1-3-gynis, calycibus eglandulosis acutis. ♂ in sepibus vulgaris, valde varians.
  - a. *obtusata* (DC.) foliis subrhombeis basi obovatis indivisis aut obtusè trifidis crenatis utrinque concoloribus, floribus 1-3-gynis. — *Mespilus oxyacanthoides* Fl. fr. 4. p. 433. — Fl. dan. t. 334. (v. v.)
  - β. *vulgaris* (DC.) foliis ovatis basi cuneatis profundè trifidis pinnatifidisve utrinque concoloribus, lobis acutis divergentibus subserratis, floribus 1-3-gynis. (v. v.)
  - γ. *laciniata* (Wallr.) *Mespilus Oxyacantha* Fl. fr. 4. p. 433. (v. v.)
3. C. AZAROLUS (L. sp. 683.) foliis pubescentibus basi cuneatis trifidis, lobis obtusis grossè paniculatis, ramulis corymbis calycibusque pubescentibus, cal. lobis obtusis, floribus 1-3-gynis. ♂ in sylvis asperis Gallie australis. — *Mesp. Azarolus* Fl. fr. 4. p. 434. — J. Bauh. 1. p. 67. ic. (v. v.)

XIX. COTONEASTER *Medik. gesch. bot. 1793. DC. prod. 2. p. 632.*

Flores abortu polygami. Cal. turbinatus obtusè 5-dentatus; pet. brevia erecta; stam. dentium calycis longitudine; styli glabri; staminibus breviores; carpella 2-3 parietalia calyce inclusa biovulata. — Frutices, foliis simplicibus integerrimis.

1. C. VULGARIS (Lindl. trans. linn. 13. p. 101.) foliis ovatis basi rotundatis, calycibus pedunculisque glabris. ♂ in rupestribus montosis Gallie australis orientalisque. — *Mespilus Cotoneaster* L. sp. 686. Fl. fr. 4. p. 435. — Clus. hist. 1. p. 60. f. 2. (v. v.)
2. C. TOMENTOSA (Lindl. l. c.) foliis ellipticis utrinque obtusis, calycibus pedunculisque lanatis. ♂ in rupestribus Jurassi Alpium, et Pyr. or. loco dicto Font de Comps ex cl. Benth. — *Mespilus eriocarpa* DC. Fl. gall. synops. et Fl. fr. 5. p. 532. (v. v.)

XX. AMELANCHIER *Medik. gesch. 1793. Lindl. trans. linn. 13. p. 100.*

Cal. 5-fidus; petala lanceolata; stam. calyce subbreviora; ovarii loculi 10 seu potius 5 bipartiti; ovula 10 in loculorum partitionibus solitaria; styli 5 basi subcoaliti; pomum maturum 3-5-loculare; semina 3-5 endocarpio cartilagineo. — Arbusculæ, foliis simplicibus serratis.

1. A. VULGARIS (Mœnch meth. 682.) foliis subrotundo-ovalibus obtusiusculis subtus pubescentibus demum glabris. ♂ in asperis montosis. — *Cratægus Amelanchier* Fl. fr. 4. p. 432. — Clus. hist. 1. p. 62. ic. (v. v.)

XXI. MESPILUS *Lindl. l. c. p. 99. DC. l. c. — Mespili sp. L.*

Cal. 5-fidus, laciniis foliaceis; petala suborbiculata; discus magnus mellifluus; styli 2-5 glabri; pomum turbinatum apertum 5-loculare, endocarpio osseo. — Arbusculæ, foliis lanceolatis serrulatis.

1. M. GERMANICA (L. sp. 684.) foliis lanceolatis subtus tomentosis indivisis, floribus solitariis. ♂ in dumetis et sylvaticis; colitur. — Fl. fr. 4. p. 434. Pall. fl. ross. t. 13. f. 1. (v. v.)



XXII. PYRUS *Lindl. l. c. p. 97. DC. prod. 2. p. 633.* — *Pyrus* et *Sorbus* *L.*

Cal. tubus urceolatus, limbus 5-lobus; petala subrotunda; styli sæpius 5, rarius 2-3; pomum clausum 5-loculare putaminibus cartilagineis. Semina in quoque loculo 2, testâ cartilagineâ. — Arbores fruticesve.

Sect. I. PYROPHORUM *DC. l. c.*

Petala patentia plana, styli 5 liberi; pomum plus minus turbinatum aut subglobosum nec basi umbilicatum; pedicelli simplices umbellati, folia simplicia eglandulosa.

1. *P. COMMUNE* (*L. sp. 686.*) foliis ovatis serratis utrinque gemmis ramulisque glabris, pedunculis corymbosis. § in nemoribus sylvestris et in hortis culta. — *Fl. fr. 4. p. 430.*

*a. Achras* (*Wallr.*) spinosa, foliis ovatis acuminatis integerrimis longè petiolatis junioribus calycisque tubo lanatis demùm glabris, pomis basi productis. § sylvestris. (v. v.)

*ß. Pyrausta* (*Wallr.*) spinosa, foliis subrotundis acutis argutè serratis junioribus calycisque tubo glabris, pomis basi rotundatis. § sylvestris. — *Gærtn. fr. t. 87. f. 2. (v. v.)*

*γ. sativa* (*DC.*) inermis. — Quoad innumeras varietates in hortis cultas vide *Mill. dict. Duham. arb. fruit.*

2. *P. BOLLWYLLERIANA* (*DC. fl. fr. 5. p. 530.*) foliis ovatis grossè serratis subtus gemmisque tomentosis, corymbis multifloris. § in sylvis Alsatiæ circa Bollwyller. — *P. Pollveria* *L.* — *J. Bauh. hist. 1. p. 59. ic.* Fructus turbinati parvi intus flavidi. (v. v. c.)

3. *P. SALICIFOLIA* (*DC. prod. 2. p. 634.*) foliis lanceolatis integerrimis subtus gemmisque tomentosis, superne junioribus velutinis adultis glabris. § in Galliâ circa Aureliam et Cænomanum sylvestris et culta. — *Fl. fr. 5. p. 531. in adnot. (v. v. DC. sine fl.)*

4. *P. AMYGDALIFORMIS* (*Vill. cat. strab. 322.*) spinosa, foliis oblongis acutis integerrimis subtus gemmisque tomentosis, superne junioribus velutinis, adultis glabris petiolo sextuplo longioribus, floribus corymbosis. § sylvestris in asperis Galloprovinciæ, Delphinatûs, Occitanîæ. — *Fl. fr. 5. p. 531. (v. v. DC.)*

Sect. II. MALUS *Tourn. inst. t. 406.*

Petala patentia plana; styli 5 basi subcoaliti; pomum sæpius globoso-depressum semper basi umbilicatum; pedicelli simplices umbellati; folia simplicia eglandulosa.

5. *P. ACERBA* (*DC. prod. 2. p. 635.*) foliis ovatis acutis crenatis, junioribus calycinisque tubis glaberrimis, floribus corymbosis. § in sylvis et versuriis. — *Malus acerba* *Mer. fl. par. 187. Fl. fr. 5. p. 530.* — Fructus acerbi. (v. v.)

6. *P. MALUS* (*L. sp. 686.*) foliis ovatis acutis crenatis subtus calycisque tubis lanatis, floribus corymbosis, stylis glabris. § in nemoribus et versuriis sylvestris. In hortis culta innumeras varietates profert, de quibus videndum *Duham. arb. fruit. (v. v.)*

Sect. III. ARIA *DC. prod. 2. p. 635.*

Petala patentia plana; styli sæpius 2-3; pomum globosum. — Flores racemoso-corymbosi, pedunculi ramosi; folia simplicia eglandulosa subtus albotomentosa.

7. *P. ARIA* (*Ehr. Beitr. 4. p. 20.*) foliis ovatis duplicato-serratis subtus adpresse albotomentosis, corymbis planis. § 5 in nemoribus. — *Cratægus Aria* *Fl. fr. 4. p. 432. Sorbus Aria* *Crantz austr. t. 2. f. 2. (v. v.)*

8. *P. INTERMEDIA* (Ehr. Beitr. 4. p. 20.) foliis ovatis inciso-lobatis dentatis subtus adpressè albo-tomentosis, corymbis planis.  $\frac{5}{5}$  in sylvis Delphinatùs et propè Fontembellaqueum. — *Crat. latifolia* Fl. fr. 4. p. 431. Duham. arb. 1. t. 80. An satia distincta species? (v. s.)

Sect. IV. *TORMINARIA* DC. l. c.

Petala patentia plana subunguiculata; styli 2-5 inter se connexi glabri; pomum basi turbinatum subexsaccum calycinis lobis deciduis truncatum. — Folia lobato-angulata, adulta glabra, flores corymbosi, pedunculi ramosi.

9. *P. TORMINALIS* (Ehr. Beitr. 6. p. 90.) foliis cordato-ovatis penninerviis pinnatifido-lobatis, junioribus subtus puberulis, adultis glabris, lobis serratis acuminatis, infimis divaricatis.  $\frac{5}{5}$  in sylvis asperis passim. — *Cratægus torminalis* L. sp. 681. Fl. fr. 4. p. 431. — Cam. epit. 162. ic. (v. v.)

Sect. V. *SORBUS* L. gen. n. 623.

Petala patentia plana; styli 2-5; pomum globosum turbinatumve. — Folia impari-pinnata aut pinnatisecta; flores corymbosi, pedunculi ramosi.

10. *P. AUCUPARIA* (Gærtn. fruct. 2. p. 45. t. 87.) foliis pinnatis glabrinerviis, foliolis serratis, gemmis molliter tomentosis, pomis globosis.  $\frac{5}{5}$  in nemoribus montosis. — *Sorbus Aucuparia* L. sp. 663. Fl. fr. 4. p. 436. Crantz. austr. t. 1. f. 4. (v. v.)
11. *P. SORBUS* (Gærtn. fruct. 2. p. 45. f. 87.) foliis pinnatis, foliolis serratis subtus villosis demùm glabris, gemmis glabris glutinosis acuminatis, pomis pyriformi-obovatis.  $\frac{5}{5}$  in montosis sylvaticis versaurisque, sylvestris et culta. — *Sorbus domestica* L. sp. 684. Fl. fr. 4. p. 436. — Cam. epit. 160. ic. (v. v.)

Sect. VI. *CHAMÆMESPILUS* DC. l. c.

Petala erecta conniventia concava; styli 2; pomum ovatum. — Folia simplicia eglandulosa; flores capitato-corymbosi.

12. *P. CHAMÆMESPILUS* (Lindl. tr. lin. 13. p. 98.) foliis ovatis serratis glabris.  $\frac{5}{5}$  in montibus. — *Crat. Chamæmespilus* Jacq. austr. t. 231. Fl. fr. 4. p. 432. — Clus. hist. 1. p. 63. f. 1. Folia juniora subtus pube decidua instructa. (v. v.)

XXIII. *CYDONIA* Tourn. 632. inst. 2. 405. DC. prod. 2. p. 638.

Cal. 5-fidus; pet. suborbiculata; stem. erecta; styli 5; pomum clausum 5-locul. loculis polyspermis cartilagineis; semina pulpâ mucilagineâ obducta. — Arbusculæ, foliis indivisis.

1. *C. VULGARIS* (Pers. ench. 2. p. 40.) foliis ovatis basi obtusis integerrimis subtus calycibusque tomentosis.  $\frac{5}{5}$  in Occitanis et Galloprovincia sepibus, in hortis frequenter culta. — *Pyrus Cydonia* L. sp. 667. Fl. fr. 4. p. 430. (v. v.)

Ordo XXXV. GRANATEÆ *Don DC. prod. t. 3. ined.*

Calycis tubus turbinatus, limbus 5-7-fidus coriaceus, lobis per æstivationem valvatis. Petala 5-7. Stam.  $\infty$  filamentis liberis, antheris anticis bilocularibus rimâ duplici dehiscentibus. Stylus filiformis. Stigma capitatum. Fructus magnus sphaericus calycis limbo subtubuloso coronatus ejusdem tubo corticatus, indehiscens, diaphragmate horizontali inæqualiter bicameratus, camerâ superiore 5-7-locul., inferiore minore 3-loculari, septis utriusque membranaceis loculos separantibus. Placentæ cameræ super. carnosæ à parietibus ad centrum tendentes, in inferiori processus irregulares ab ipso fundo. Semina  $\infty$  pulpâ pellucidâ baccata, exalbuminosa. Embryo oblongus, radiculâ brevi rectâ, cotyledonibus foliaceis spiralliter convolutis. — Arbusculæ, ramulis subspinescentibus, foliis deciduis sæpius oppositis oblongis integris impunctatis.

I. PUNICA *Tourn. inst. t. 407. L. gen. n. 618.*

Car. isdem ac ordinis.

1. P. GRANATUM (L. sp. 676.) foliis lanceolatis, caule arboreo. § 5 in Galliâ australiori. — Fl. fr. 4. p. 427. Sims. bot. mag. t. 1832. (v. v.)

Ordo XXXVI. MYRTACEÆ *Brown gen. rem. p. 14. DC. prod. t. 3. ined. — Myrti. Juss. gen. 323.*

Calycis sepala 4-6 sæpius 5 concreta in tubum ovario adnatum, apice nunc usque ad ovarii marginem libera, nunc paulò ultrâ in faucem concreta. Petala calyci inserta tot quot sepala, iis alterna, æstivatione quinquunciali. Stamina cum petalis inserta, sæpè multiseriaria, numero petalorum dupla aut multipla, filamentis nunc liberis nunc variè connexis, antheris 2-locularibus rimâ duplici dehiscentibus. Carpella 4-6 sæpius 5, abortu sæpè pauciora, concreta in ovarium multiloculare calyci adnatum. Stylus sæpius stylis partialibus omnino concretis constans et ideò unicus stigmate simplici dictus, interdum stylis (in Philadelpho) plus minùs ad apicem distinctis et ideò ferè multiplex dicendus. Fructus polyspermus. Embryo sæpius exalbuminosus. Cotyledones non convolutæ. — Arborea aut frutices, foliis sæpius oppositis exstipulatis plerisque pellucido-aut-glanduloso-punctatis.

Trib. I. *MYRTEÆ* DC. l. c.

Sépala 4-5. Petala totidem. Stamina filam. libera. Fructus carnosus plurilocularis. — Pedunculi axillares.

I. *MYRTUS* Lin. gen. 617. DC. l. c.

Calycis tubus subglobosus; pet. 5; bacca 2-3-locul. subglobosa calycis limbo coronata, semina in loculo quoque plurima subossea reniformi-incurva; embryo curvus, cotyledonibus semi-cylindricis brevissimis, radicula cotyledonibus duplo longiore.

1. *M. communis* (L. sp. 673.) pedicellis solitariis 1-floris, bracteolis 2 linearibus deciduis subflore, calycibus 5-fidis, foliis ovatis lanceolatisve acutis. 3 in apricis rupestribus Corsicæ, Nicæ, Galloprovinciæ australis, Narbonæ et agri Ruscinonensis. — Fl. fr. 4. p. 426. Lam. ill. t. 419. (v. v.)

Trib. II. *PHILADELPHEÆ* DC. l. c.

Sépala 4-10. Petala totidem. Stam. libera. Fructus siccus plurilocularis. Semina arillata albuminosa. — Folia impunctata.

I. *PHILADELPHUS* L. gen. n. 614.

Calycis tubus obovato-turbinatus, limbus 4-5-partitus; pet. 4-5; stam. 20-25 libera petalis breviora; styli 4-5; capsula calyci semi-adnata 4-5-locularis polysperma; semina inclusa arillo membranaceo oblongo; albumen carnosum; embryo inversus rectus.

1. *P. coronarius* (L. sp. 671.) foliis ovatis acuminatis dentatis glabresculis, stylis à basi ferè distinctis stamina non superantibus. 3 ad sepes in Delphinatu, frequenter in hortis cultus. — Fl. fr. 4. p. 425. Lam. ill. t. 420. (v. v. c.)

Ordo XXXVII. CUCURBITACEÆ *Juss. gen. p. 393. DC. fl. fr. 3. p. 688. St.-Hil. mem. mus. 9. p. 190. Ser. in mem. soc. Genev. 3. p. 1. p. 5.*

Flores hermaphroditi monoïci vel dioïci. Calyx monosepalus suprâ coarctatus, sepalis 5 plus minus basi inter se et mediante toro cum carpellis coalitis. Corolla marcescens persistens. Petala 5 libera aut coalita à calyce distincta, aut rariùs cum calyce subcontinua ex margine tori nascentia. Stamina 5 libera vel sæpiùs triadelpa; antheræ biloculares longissimæ rarò ovatæ breves. Stylus 1 rarò subnullus; stigmata 3-5 biloba crassa. Carpella 3-5 (abortu 1) carnosa, toro et calyce involuta. Bacca cortice duro 1-5-locularis. Semina angulo sepi-mentorum interno adfixa horizontalia arillata, arillâ aquosâ exsiccatione membranaceâ; funiculus umbilicalis semen versùs tumidus; hylus obliquus ad apicem seminis. Embryo rectus; albumen nullum; cotyledones planæ foliaceæ; radícula hylum versùs directa. — Herbæ scandentes sæpè asperæ, foliis petiolatis alternis, pilis septulatis sæpè munitis, cyrrhis axillaribus, pedunculis axillaribus mediò articulatis.

I. LAGENARIA *Ser. in mem. soc. Genev. 3. p. 1. p. 25. et in DC. prod. t. 3. ined.*

Calyx campanulatus, laciniis subulatis vel latiusculis tubo brevioribus; petala libera et à calyce distincta infra calycis marginem nascentia. *Masc.* Stam. 5 triadelpa. *Fem.* Stylus ferè nullus; stigm. 3; fructus 3-5?-locularis; semina obovata compressa margine tumida apice biloba. — Fl. monoïci albi.

1. *L. vulgaris* (*Ser. l. c.*) molliter pubescens, cyrrhis 3-4-fidis, foliis cordatis integrisculis basi biglandulosis, floribus patentissimis fasciculatis, fructibus pubescentibus maturitate glabris lævissimis. ① ex Indiâ orta, in hortis culta. — *Cucurbita Lagenaria* L. sp. 1434. Fl. fr. 3. p. 69a. — *Moris. hist. s. 1. t. 5. n. 1. 2. 3. (v. v. c.)*

## II. CUCUMIS *L. gen. n. 1479.*

Calyx tubuloso-campanulatus, laciniis subulatis vix longitudine tubi; petala inter se et cum calyce vix coalita. *Masc.* Stam. 5 triadelpa. *Fem.* Stigmata 3 crassa; fructus 3-6-locularis, seminibus ovatis compressis non marginatis. — Fl. monoïci vel hermaphroditi lutei.

1. *C. Melo* (L. sp. 1436.) caule humifuso scabro, foliis rotundatis angulatis petiolatis, florum masculorum calycis tubo basi subventricosso apice dilatato, staminibus inclusis, hermaphroditorum stigmatibus 3-4, fructu ovato vel subgloboso sulcato. ① ex Asiâ ortus, in hortis cultus. — Fl. fr. 3. p. 691. (v. v. c.)
2. *C. sativus* (L. sp. 1437.) caule scabro, foliis cordatis obsolete 5-lobis petiolatis, lobo terminali majore, floribus breviter pedunculatis subternis, masculorum tubo calycis tubuloso-campanulato, petalis acutiusculis, fructibus oblongis sæpè nitidis, ① ex Indiâ orientali ortus, in hortis cultus. — Fl. fr. 3. p. 691. — *Lob. stirp. 363. f. 1. (v. v. c.)*

3. *C. CITRULLUS* (Ser. in DC. prod. t. 3. in ed.) valdè pilosus, caule humifuso, foliis obtusè pinnati-sectis subglaucescentibus, floribus solitariis bracteis oblongè suffaltis, fructibus subglobosis glabris. (1) ex Africâ et Indiâ ortus, in hortis colitur. — Cucurbita Citrullus L. C. Anguria Duch. in Lam. dict. 2. p. 158. Fl. fr. 3. p. 694. — Lob. stirp. 364. f. 2. (v. v. c.)

### III. BRYONIA L. gen. n. 1480.

Flores monoïci vel dioïci; petala basi vix coalita. *Masc.* Calyx 5-dentatus; stam. 3-adelpha. *Fem.* Stylus 3-fidus; fructus ovatus vel globosus oligospermus, semina ovata vix compressa plus minus marginata. — Cyrrhi simplices rarò bifurci.

1. *B. ALBA* (L. sp. 621.) caule scandente, foliis cordatis 5-lobis dentatis calloso-punctatis scabris, lobo terminali vix majore, floribus racemosis monoïcis, fructibus globosis. (2) in Galliâ austro-occidentali et Pyrenæis ex Lapeyr. — Fl. fr. 3. p. 432. Lam. ill. f. 796. (v. s.)
  2. *B. DIOICA* (Jacq. aust. t. 199.) caule scandente, foliis cordato-palmatis 5-lobis dentatis calloso-punctatis, lobo terminali longiore, floribus racemosis dioïcis, filamentis basi pilosis, fructibus globosis. (2) in sepibus. — Fl. fr. 3. p. 689. eng. bot. t. 436. — Mill. ic. t. 71. Fructus rubri. (v. v.)
- β. *lutea* (Ser. l. c.) foliis profundè lobatis, lobis lateralibus sublinearibus terminali lanceolato lateraliter bidentato, fructibus luteis. — in sepibus et sylvaticis Arvernæ. (v. s.)

### IV. MOMORDICA L. gen. n. 1477.

Fl. monoïci. *Masc.* Calyx 5-fidus tubo brevissimo; cor. 5-partita; stamina 3-adelpha, antheris connatis. *Fem.* Filamenta 3 sterilia; stylus trifidus; ovarium 3-loculare; fructus sæpè muricatus, maturus elasticè dissiliens; semina compressa.

1. *M. ELATERIUM* (L. sp. 1434.) hispido-scabra glaucescens, caule nano absque cyrrhis, foliis cordatis sublobatis crenato-dentatis longè petiolatis, fructibus ovatis hispido-scabris. (1) in sterilibus Galliæ Mediterraneæ et austro-occidentalis usque ad Burdigalam ex Laterrade. — Fl. fr. 3. p. 690. bot. mag. t. 1914. — Black. herb. t. 108. (v. v.)

### V. CUCURBITA L. gen. n. 1478.

Flores monoïci; corolla campanulata; petala inter se et cum calyce coalita. *Masc.* Calyx hemisphærico-campanulatus; stam. 5 triadelphæ et syngenesæ. *Fem.* Calyx obovato-clavatus campanulatusve et post anthesin infra limbum semper circumscissus; stigm. 3; fructus 3-5-locularis; semina ovato-compressa margine vix tumidulo.

1. *C. MAXIMA* (Duch. in Lam. dict. 2. p. 151.) foliis cordatis rugosis, petiolis hispidis, tubo calycis obovato in collum breve desinente, fructibus globosis subdepressis. (1) Patria ignoratur, sæpè colitur. — Fl. fr. 3. p. 692. — Lob. ic. 641. f. 2. (v. v. c.)
2. *C. MELOPEPO* (L. sp. 1435.) foliis cordatis obtusis subquinguelobis denticulatis, calyce brevi hemisphærico-campanulato, fauce valdè dilatata, fructibus depressis, carpellis ultra faucem calycis prominentibus 4-5-locularibus. (1) Patria ign., colitur. — C. pepo ? Fl. fr. 3. p. 693. — J. Bauh. hist. 2. p. 224. — Moris. hist. 1. s. 1. t. 8. (v. v. c.)
3. *C. PEPO* (L. sp. 1435.) foliis cordatis obtusis subquinguelobis denticulatis, calycis tubo in collum desinente, fructibus subrotundis oblongisve. (1) ex Oriente indigena, ubique colitur. — Fl. fr. 3. p. 693. excl. var. ζ. (v. v. c.)

Ordo XXXVIII. ONAGRARIÆ *Juss. ann. mus. 3. p. 315.*  
*excl. Haloragcis. DC. prod. t. 3. ined.*

Calycis tubus cum ovario aut totus adnatus aut basi adhærens et ultrà ovarium productus, limbus 2-5 sæpius 4-lobus. Petala tot quot calycis lobi, plerumque regularia, lobis alterna, summo tubi inserta, rarius nulla. Stamina petalis æqualia nunc dupla nunc dimidiò minora, filamentis liberis, antheris oblongis aut ovatis. Ovarium pluriloculare, sæpè glandulâ cupulari coronatum. Stylus filiformis. Stigma capitatum aut lobatum. Fructus capsularis baccatus aut drupaceus 2-aut 4-locularis. Semina plurima (aut rarissimè solitaria) in quoque loculo, angulo centrali adfixa, exalbuminosa (endoplevrâ tumidâ interdum albumen mentiente), embryone recto, radiculâ longâ teretiusculâ, cotyledonibus brevibus. — Herbæ (in nostris), foliis simplicibus integris aut dentatis penninerviis.

Trib. I. *ONAGRÆ DC. l. c.*

Fructus capsularis, loculis polyspermis. Calycis tubus ultrà ovarium productus. Stamina petalorum numero dupla.

I. *EPILOBIUM L. gen. 471. Gertn. fruct. 1. p. 157. t. 31. f. 6.*

Calyx 4-sepalus, sepalis coalitis in tubum longum tetragonum, limbo post anthesin caduco; petala 4; stamina 8 polline non viscoso; capsula linearis obtusè tetragona 4-locularis 4-valvis polysperma; semina papposa.

Sect. I. *CHAMENERION Tourn. inst. p. 302. t. 157. f. A. C. D.*

Flores irregulares, petala ovata; genitalia deflexa.

1. *E. SPICATUM* (Lam. dict. 2. p. 373.) caule erecto stricto, foliis sparsis lanceolatis longissimis integris venosis subpellucidis, bracteis à basi pedicellorum omninò liberis et hinc pedicellis axillaribus, pistillo staminibus longiore.  $\mathcal{L}$  in sylvis montosis, in hortis sæpè cultum. — Fl. fr. 4. p. 421. *E. angustifolium*  $\alpha$  L. sp. 493. — J. Bauh. 2. p. 907. f. 1. (v. v.)
2. *E. ROSMARINIFOLIUM* (Hænicke in Jacq. coll. 2. p. 50.) caule erecto ramoso, foliis linearibus lineari-lanceolatisve obsolete denticulatis aveniis, bractearum parte inferiore cum pedicellis plus minùs coadunatâ, stylo staminibus breviorè.  $\mathcal{L}$  in sylvaticis humidis. — Fl. fr. 4. p. 421. — Moris. 3. t. 11. m. 2. Lob. ic. 343. f. 2. (v. v.)

$\beta$ . *Dodonæi*, caule debili tortuoso, foliis sæpè latioribus. — Ad torrentes Alpium. — *E. Dodonæi* Vill. dauph. 3. p. 507. (v. v.)

Sect. II. *LYSIMACHION Tausch. hort. canal. fasc. 1.*

Flores regulares, petala obcordata, genitalia ascendentia.

3. *E. ALPINUM* (L. sp. 495.) caule basi repente, foliis ovato-oblongis obtusis denticulatis integrisve oppositis glaberrimis in brevem petiolum attenuatis, floribus antè anthesin nutantibus, stigmate indiviso, capsulis lævibus pedi-

- cellatis.  $\mathcal{L}$  in Alpinis et Pyrenæis. — Fl. fr. 4. p. 424. eng. bot. t. 2001. (v. v.)
4. *E. ORIGANIFOLIUM* (Lam. dict. 2. p. 376.) caule ascendente in parte superiori nutante, foliis lanceolato-ovatis acutis denticulatis sessilibus oppositis glabris, floribus antè anthesin erectis, stigmatè indiviso, capsulis lævibus pedicellatis.  $\mathcal{L}$  in Alpinis, Pyrenæis, et editioribus montibus Arverniz. — Fl. fr. 4. p. 424. *E. alsinæfolium* eng. bot. t. 2000. (v. v.)
5. *E. ROSUM* (DC. fl. fr. 4. p. 422. non Smith.) caule erecto simplici, foliis oppositis binis ternisve lanceolato-ovatis acuminatis serrulatis pubescentibus sessilibus, petalis calyce longioribus, stigmatè indiviso, floribus subsessilibus, capsulis breviter pedicellatis.  $\mathcal{L}$  in Alpinis et Pyrenæis. (v. v.)
6. *E. MONTANUM* (L. sp. 494.) caule erecto ramoso aut simplici, foliis alternis oppositisve lanceolatis inæqualiter serratis subsessilibus glabriusculis, petalis calyce longioribus, stigmatè 4-partito, floribus sessilibus, capsulis gracilibus elongatis breviter pedicellatis.  $\mathcal{L}$  in sylvis montosis. — Fl. fr. 4. p. 423. fl. dan. t. 922. (v. v.)
7. *E. FALUSTRE* (L. sp. 495.) caule tereti ramoso aut simplici infernè glabro supernè puberulo, foliis sessilibus lanceolatis subdenticulatis oppositis alternisque glabris, stigmatè indiviso, capsulis pubescentibus.  $\mathcal{L}$  ad marginem fossarum et stagnorum. — Fl. fr. 4. p. 422. eng. bot. t. 346. (v. v.)
8. *E. HIRAUTUM* (L. sp. 494.) caule ramoso hirtio, foliis oppositis alternisque lanceolato-oblongis hirsutis amplexicaulibus irregulariter serratis, serraturis subhamatis, stigmatè 4-partito, sepalis mucronatis.  $\mathcal{L}$  ad aquas. — Fl. fr. 4. p. 421. eng. bot. t. 838.
- a. *vulgare*, caule ramis foliisque pilosis, capsulis pilis brevibus tectis. (v. v.)
- ß. *intermedium* (Ser. in DC. prod.) caule ramis foliisque pilosissimis, capsulis lanatis. — Circà Parisios, Monspelium, Nemausim. — *E. intermedium* Mer. fl. par. t. 2. p. 290. (v. v.)
9. *E. MOLE* (Lam. dict. 2. p. 475.) caule stricto villosa ramoso aut simplici, foliis oblongo-lanceolatis serrulatis acutis molliter pubescentibus oppositis alternisve sessilibus, stigmatè 4-partito, capsulis puberulis subglabrisve.  $\mathcal{L}$  in humidis umbrosis. — Fl. fr. 4. p. 422. *E. parviflorum* Schreb. eng. bot. t. 795. — Moris. s. 3. t. 11. n. 4. (v. v.)
10. *E. TETRAGONUM* (L. sp. 494.) caule erecto tetragono subglabro, foliis oblongo-lanceolatis serratis acutis glabris alternis oppositisve in petiolum attenuatis, stigmatè indiviso, capsulis puberulis.  $\mathcal{L}$  in inundatis et ad marginem stagnorum et fossarum. — Fl. fr. 4. p. 422. eng. bot. t. 1948. (v. v.)

## II. OENOTHERA L. gen. 469.

Calyx 4-sepalus, sepalis coalitis in tubum longum tetragonum aut octocostatum limbo post anthesin caduco; petala 4; stamina 8 poliline viscoso; capsula oblongo-linearis obtusè tetragona aut obovato-clavata 4-locularis 4-valvis polysperma, seminibus nudis.

1. *OE. BIENNIS* (L. sp. 492.) caule muricato-viloso, foliis ovato-lanceolatis planis, staminibus corollà brevioribus, petalis oboordatis, capsulis oblongis subtargidis. ② in humidis passim. — Fl. fr. 4. p. 419. — Mill. ic. t. 189. f. 2. (v. v.)

### Trib. II. JUSSIEÆ DC. prod. t. 3. ined.

Fructus capsularis oculis sæpiùs polyspermis, rariùs 1-spermis. Calycis tubus ultrà ovarium non productus et immediatè partitus.



III. *ISNARDIA* DC. l. c. — *Isnardiæ* et *Ludwigie* sp. L. gen. n. 156 et 153.

Calycis tubus ovatus aut subcylindraceus brevis ovario adhærens, limbus 4-partitus persistens; petala nunc 4 calyci alterna interdum rudimentaria, nunc nulla; stam. antè lobos calycinos; stylus à basi filiformis deciduus; stigma capitatum; capsula obovata aut serè cylindracea tetragona 4-valvis 4-locularis polysperma, loculicido-dehiscens. — *Herbæ aquaticæ aut paludosæ.*

1. *I. PALUSTRIS* (L. sp. 175.) caule procumbente radicante glabro, foliis oppositis in petiolum attenuatis ovatis acutis, floribus axillaribus solitariis sessilibus apetalis. ① in fossis et aquosis passim. — Fl. fr. 4. p. 419. — Bocc. mus. t. 84. f. 2. (v. v.)

IV. *CIRCÆA* Tourn. inst. t. 155. L. gen. p. 24.

Calyx brevis limbo bipartito; petala 2 obcordata; stam. 2 petalis alterna; stigma emarginatum; capsula ovata pilis uncinatis hispida 2-loc. 2-valvis 2-sperma; semina in loculis solitaria erecta.

1. *C. LUTETIANA* (L. sp. 12.) caule erecto pubescente, foliis ovatis acutis denticulatis sæpius subpubescentibus. ② in sepibus et nemoribus umbrosis. — Fl. fr. 4. p. 417. eng. bot. t. 1056. — Lob. ic. 266. f. 2. (v. v.)
2. *C. ALPINA* (L. sp. 12.) caule basi ascendente aut radicante glabro, foliis cordatis acutis dentatis membranaceis levibus. ② in umbrosis et humidis montium. — Fl. fr. 4. p. 417. eng. bot. t. 1057. (v. v.)
2. *intermedia* (Ehr. beit. 4. p. 42.) caule erecto simpliciter glabriusculo, foliis repando-denticulatis acuminatis. — In montosis et umbrosis. — Fl. dan. t. 256. (v. s.)

Trib. III. *HYDROCARYES* Link enum. hort. berol. 1. p. 141. DC. l. c.

Fructus nucamentaceus indehiscens cornutus durus, maturus sæpius 1-loculus. 1-spermus. Semen pendulum. Cotyledones valdè inæquales. — *Herbæ natantes.* — Tribus media inter *Onagrarias* et *Halorageas*.

V. *TRAPA* L. gen. n. 157. Gærtn. fruct. 1. p. 127. t. 26.

Calycis tubus ovario adhærens, limbus 4-partitus; pet. 4; stam. 4; stylus filiformis basi subcrassior; ovarium 2-loculare loculo uno dein abortivo; nux serè cornea 2-4-cornuta, cornubus à basi lorum calycinorum omnibus aut 2 oppositis ortis, semen 1 magnum intrà nucem pendulum exalbuminosum; cotyledones 2 valdè inæquales, 1 maximo farinaceo intus nucem per germinationem remanente, altero minimo exserto.

1. *T. NATANS* (L. sp. 175.) fructibus 4-cornutis, cornubus acutis cruciatim oppositis, 2 superioribus horizontaliter patentibus, 2 inferioribus subascendentibus. ① in stagnis et fossis passim. — Fl. fr. 4. p. 419. — Camer. epit. 716. ic. (v. v. DC.)

Ordo XXXIX. HALORAGÆ *R. Brown. gen. rem. p. 17.*  
*DC. prod. t. 3. ined.*

Calycis tubus per totam longitudinem ovario adhærens, limbus partitus aut subnullus. Petala summo calycis tubo inserta, lobis alterna et iis numero æqualia aut nulla. Stamina ibidem inserta, petalis numero dupla, æqualia rariùs pauciora. Ovarium calyci adhærens, sæpiùs pluriloculare. Stylus o. Stigmata tot quot ovarii loculamenta suprà apicem sessilia. Fructus carpellis plurimis plus minùs intrà calycem concretis constans et ideò plurilocularis indehiscens, membranaceus nucamentaceusve, loculis 1-spermis. Semina in loculis pendula, albumine carnosio, embryone centrali recto, radiculâ suprâ, cotyledonibus 2 brevibus. — (In nostris) Herbæ aquaticæ, floribus axillaribus aut in spicas terminales dispositis, interdum abortu monoïcis aut dioïcis.

Trib. I. *CERCODIANÆ Juss. dict. sc. nat. 7. p. 441. DC. l.c.*

Calycis limbus evidenter partitus. Stamina lobis calycis numero æqualia aut dupla. Petala et fructus loculamenta sæpiùs tot quot calycis lobi.

I. MYRIOPHYLLUM *Vaill. act. acad. Paris. 1719. t. 2. f. 3. L. gen. n. 1066.*

Flores monoïci aut rariùs hermaphroditi. — *Masc.* Calyx 4-partitus; petala 4 lobis calycis alterna valdè caduca; stam. 4, 6, aut sæpiùs 8. — *Fem.* Calyx 4-lobus; petala 0; carpella 4 subconcreta compressa aut subglobosa nucamentacea indehiscencia 1-sperma; semina albumine subnullo, embryone teretiusculo inverso, cotyledonibus obtusis, radiculâ longâ. — Herbæ natantes, apicibus ad anthesin emersis; floribus parvis, superioribus masculis, inferioribus femineis.

1. *M. SPICATUM* (L. sp. 1409.) foliis verticillatis pinnati-partitis lobis capillaceis, spicâ terminali nudiusculâ, foliis floralibus flore brevioribus, infimis subserratis, plerisque integerrimis.  $\mathcal{U}$  in aquis lente fluentibus. — Fl. fr. 4. p. 416. eng. bot. t. 83. — C. Bauh. prod. 73. f. 1. (v. v.)

2. *M. PECTINATUM* (DC. fl. fr. 5. p. 529.) foliis verticillatis pinnati-partitis lobis capillaceis, spicâ terminali nudiusculâ, foliis floralibus flores paulo excedentibus omnibus serrato-pectinatis.  $\mathcal{U}$  in aquis quietis Monspeliî, Arelatis. — M. verticillatum fl. dan. t. 1046. non Lin. (v. v.)

3. *M. ALTERNIFLORUM* (DC. fl. fr. 5. p. 529.) foliis verticillatis pinnati-partitis lobis capillaceis distantibus alternis, spicâ terminali nudiusculâ, floribus alternis, inferioribus subfasciculatis, cæteris solitariis, foliis floralibus inferioribus pinnati-partitis flores excedentibus, super. integris brevibus.  $\mathcal{U}$  in aquis quietis Galliæ occidentalis circâ Naunetes et in Vasconiâ. (v. a.)

4. *M. VERTICILLATUM* (L. sp. 1410.) foliis verticillatis pinnati-partitis lobis capillaceis oppositis, spicâ terminali foliosâ, foliis floralibus omnibus pinnati-partitis flores multò superantibus et à foliis vix distinctis.  $\mathcal{U}$  in inundatis. — Fl. fr. 4. p. 417. eng. bot. t. 218. — Clus. hist. 2. p. 252. f. 1. (v. v.)

$\beta$ . *limosum* (DC.) foliorum lobis linearibus acutis regularibus et ad costam non omninò attingentibus.  $\mathcal{U}$  in inundatis subexsiccatis. (v. a.)

Trib. II. *CALLITRICHINÆ* Link. enum. h. berol. 1. p. 7. DC. l. c.

Calycis limbus non conspicuus. Petala 0. Stam. 1-rarius 2. Fructus 4-locularis 4-spermus. — Herbæ aquaticæ, foliis oppositis.

II. *CALLITRICHE* L. gen. 13. DC. prod. t. 3. ined.

Flores polygami, hermaphroditi, dioici aut sæpius monoici; bracteæ 2 oppositæ petaloideæ ad basin floris (Petala auct.); calyx non conspicuus veròsimiliter ovarii basi adhærens limbo abortivo. Masc. Stam. 1 rarius 2 filamentis filiformibus exsertis, antherâ reniformi 1-loculari suturâ transversâ dehiscente. — Fem. Ovarium bisulcum; styli 2; capsula compressa bisulca constans carpellis 4 concretis per paria approximatis indehiscentibus 1-spermis lenticulari-compressis; semina cum carpello concreta, albuminae carnoso, embryone inverso subincurvo, radicula longâ superâ, cotyledonibus brevissimis. — Herbæ ænuæ.

1. *C. Verna* (L. sp. 6.) foliis trinerviis supernè aggregatis majoribus, fructibus sessilibus, carpellorum dorso obtusiusculo. ① in aquis lentè fluentibus. — *C. sessilis* excl. var. *fl.* fr. 4. p. 414.

a. *vulgaris* (DC.) foliis omnibus elongato-obovatis. — *C. verna* fl. dan. t. 129. (v. v.)

β. *intermedia* (Hoffm.) foliis inferioribus linearibus obtusis aut emarginatis, superioribus ovalibus. — Schkuhr. handb. t. 1. *C. dubia* Thuil. (v. v. DC.)

γ. *stellata* (Hop.) foliis omnibus ovalibus, caulibus abbreviatis. — *C. sessivalis* Thuil. (v. s.)

δ. *cespitosa* (Schult.) foliis omnibus ovalibus rigidulis parvis, caulibus abbreviatis stellato-patentibus. — In humidis vix inundatis. (v. v. DC.)

ε. *tenuifolia* (Pers.) foliis omnibus linearibus, summis subtrinerviis. — In humidis subinundatis. (v. s.)

2. *C. PEDUNCULATA* (DC. fl. fr. 4. p. 415.) foliis inferioribus linearibus, superioribus oblongis trinerviis, fructibus pedunculatis, carpellorum dorso obtusiusculo. ① in aquis cœnosis Galliæ circâ Fontembellequeum et in Andegavia. (v. v. DC.)

3. *C. AUTUMNALIS* (L. sp. 6.) foliis omnibus uninerviis per caulem sparsis aequalibus truncatis, fructibus sessilibus, carpellorum dorso elato-membranaceo. ① ad littora lacuum, fluviorum, rivorum, in aquâ quietâ; primâ fronte distinguenda colore intensius viridioris et fructu majore. — *C. sessilis* fl. fr. 4. p. 414. — Gmel. sib. 3. t. 1. f. 2. (v. v. DC.)

Trib. III. *HIPPURIDÆ* Link. enum. h. berol. 1. p. 5. DC. l. c.

Calycis limbus integer minimus. Pet. 0. Stam. 1. Fructus nucamentaceus 1-locul. 1-spermus.

III. *HIPPURIS* L. gen. n. 11. Juss. ann. mus. 3. t. 30. f. 3.

Calycis tubus ovario adnatus, limbus minimus integer; stamen calycis margini insertum, stylus filiformis antheræ sulco receptus; nux 1-sperma calycis limbo marginato coronata, embryonis radícula cylindrica cotyledonibus longior.

1. *H. VULGARIS* (L. sp. 3.) foliis verticillatis 6-12. ♀ in fossis et aquosis. — Fl. fr. 4. p. 415. eng. bot. t. 763. (v. v.)

β. *fluvialis* (Hoffm.) foliis longioribus membranaceis. ♀ in aquis profundis è quibus planta vix emergit sterilis. (v. s.)

Ordo XI. CERATOPHYLLÆ *Gray. br. pl. arr. 2. p. 554.*  
*DC. prod. t. 3. ined.*

Flores monoïci. Calyx (seu Perigonium) liber multi- (10-12) partitus, lobis æqualibus. Petala 0. — *Masc.* Stam. 2-20, filamentis nullis, antheris ovato-oblongis bilocularibus apice 2-3-cuspidatis sessilibus in centro calycis congestis. — *Fem.* Ovarium liberum ovatum 1-loculare, stylus filiformis incurvus. Stigma simplex. Nux 1-locularis, 1-sperma stylo apiculata, indehiscens. Semen ex apice pendulum. Albumen 0. Embryo rectus, radicula superâ, cotyledonibus 4 verticillatis duobus latioribus oppositis, plumula valdè composita? — *Herbæ* aquaticæ submersæ, foliis verticillatis in lobos filiformes acutos divisus, floribus axillaribus solitariis sessilibus.

I. CERATOPHYLLUM *L. gen. n. 1065. Gartn. fr. 1. p. 212. t. 44.*

Character isdem ac ordinis.

1. *C. DEMERSUM* (*L. sp. 1409.*) nucē trispinosâ, nempe spinâ 1 terminali et 2 lateralibus inæqualibus.  $\frac{1}{2}$  in fossis et aquis quietis. — *Fl. fr. 4. p. 413.* — *Vaill. act. ac. par. 1719. t. 2. f. 2.* Folia juniora dense conferta. (v. v.)
2. *C. SUBMERSUM* (*L. sp. 1409.*) nucē subinermi nempe styli basi tubercula 2 minima obtusa gerente.  $\frac{1}{2}$  in fossis et aquis quietis. — *Fl. fr. 4. p. 413. eng. bot. t. 679.* Folia juniora minus conferta. (v. v.)

Ordo XII. LYTHRARIÆ *Juss. dict. sc. natur. 27. p. 453.*

*DC. in mem. soc. hist. nat. gen. 3. p. 2. p. 65.* — *Salicariæ.*  
*Juss. gen. 330.*

Calyx sepalis numero definitis ultrâ medium concretis constans, liber, tubulosus aut campanulatus, lobis per æstivationem valvatis aut distantibus, sinubus interdum in lobulos conicos dentesve exteriores productis. Petala summo calycis tubo inter lobos inserta, numero varia interdum nulla, sæpius valdè caduca. Stamina tubo calycino infrâ petala inserta, petalis nunc æqualia nunc dupla triplave, nunc pauciora; antheræ ovales biloculares dorso insertæ. Ovarium liberum. Stylus filiformis. Stigma sæpius capitatum. Capsula membranacea, calyce tecta aut cincta, carpellis 2-4 constans, junior sæpius carpellorum marginibus tenuissimis introflexis 2-4-locularis, demum sæpè septis evanidis unilocularis, nunc longitudinaliter dehiscens, nunc rariùs irregulariter circumscissa. Placenta centralis septis si adsunt adnata aut libera. Semina plurima parva exalbuminosa. Embryo rectus. Radicula ad hylum versa; cotyledones planæ foliaceæ. — *Herbæ* (in nostris), foliis oppositis rariùs alternis simplicibus exstipulatis.

I. LYTHRUM *Juss. gen. 33a.*

Calyx cylindricus striatus apice dentatus, dentibus 4-6 latioribus erectis, 4-6 alternis subulatis sæpè corniformibus interdum nullis aut minimis; petala 4-6 (aut rarius nulla) ex apice tubi orta, dentibus erectis alterna; stamina e medio aut basi calycis orta, petalorum numero dupla aut æqualia aut abortu pauciora; capsula oblonga calyce tecta, bilocularis, polysperma.

Sect. I. SALICARIA *DC. prod. t. 3. ined.*

Stamina petalorum numero dupla; flores in axillis superioribus plurimi verticillati et spicato-racemosi.

1. *L. SALICARIA* (L. sp. 642.) foliis lanceolatis basi cordatis, floribus spicatis subsessilibus.  $\frac{1}{2}$  in fossis et aquosis. — Flores purpurei aut albi, petala 6-7, stamina 12-14.

*a. vulgaris* (DC.) spicâ caulina terminali densâ cylindraceâ, lateralibus gracilibus, calycibus hispidulis. — Fl. fr. 4. p. 409. Lam. ill. t. 408. f. 1. (v. v.)

*β. gracilis* (DC.) spicis plurimis gracillimis, foliis caulibus calycibusque velutinis. — In humidis arenosis maritimis propè Monspelium. — Fl. fr. 5. p. 528. (v. v. DC.)

Sect. II. HYSSOPIFOLIA *C. Bauh. pin. 218. DC. l. c.*

Stamina numero petalorum æqualia rarius dupla aut dimidiò minora; flores in axillis foliorum solitarii non spicati.

2. *L. HYSSOPIFOLIA* (L. sp. 642.) foliis alternis oppositisque lineari-lanceolatis obtusiusculis, floribus subsessilibus folio brevioribus, etiam fructiferis erectis, bracteolis acutis minimis, petalis 5-6 oblongis, staminibus 5-10. ① in inundatis humidisque Galliæ occidentalis, mediæ, australisque. — Fl. fr. 4. p. 410. eng. bot. t. 292. (v. v.)
3. *L. THYMIFOLIA* (L. sp. 642.) foliis alternis linearibus acutiusculis, floribus subsessilibus folio brevioribus, etiam fructiferis erectis, bracteolis foliaceis linearibus calyce sæpius longioribus, petalis 4-5, stam. 2. ① in paludosis subaëis Occitanæ, Galloprovinciæ, Nicæ ex All. — Fl. fr. 4. p. 410. L. tribracteatum Saltzm. — Barr. ic. t. 773. f. 2. Flores minimi. (v. v. D.)

Sect. III. AMMANIOIDES *DC. l. c.*

Stamina 4-6; petala nulla; calyx tubuloso-campanulatus. — An hujus generis?

4. *L. NUMMULARIFOLIUM* (Lois. not. 74. non Pers.) glabrum, caule herbaceo ramoso, foliis obovato-subrotundis obtusis oppositis, rameis nonnullis alternis, floribus axillaribus solitariis. ① in Corsicâ. — Calyx ferè Ammanii 8-12-dentatus, dentibus internis apice glandulis purpureis petaloideis instructis. Stam. inclusa. Ex flore Ammanii ex fructu Lythri species; potiùs novum crederem genus. (v. s.)

II. PEPLIS *L. gen. n. 446.*

Calyx campanulatus 12-fidus, lobis 6 latioribus erectis, 6 alternis subulatis patulis e sinibus ortis; petala 6 minima fugacia interdum 0; stamina 6, petalis alterna, lobis calycis latioribus opposita; stylus vix ullus; capsula bilocularis polysperma.

1. *P. PORTULA* (L. sp. 474.) foliis oppositis obovato-subrotundis in petiolum angustatis. ① in inundatis passim. — Fl. fr. 4. p. 412. — Vaill. bot. t. 15. f. 5. Calyx florens subrubens. Petala alba 1-6 aut nulla. (v. v.)

Ordo XLII. TAMARISCINÆ. *Desv. diss. in ann. sc. nat. 4. p. 344. DC. prod. t. 3. ined.*

Calyx sepalis 4-5 basi coalitis constans et ideò 5-partitus, persistens, lobis per æstivationem subimbricatis. Petala tot quot sepala, iis alterna, basi calycis inserta, marcescentia, per æstivationem imbricata. Stam. numero petalorum æqualia aut dupla, filamentis nunc ferè omninò liberis nunc longiùs monadelphis. Ovarium liberum trigonum. Stylus unicus. Stigmata 3. Capsula trigona, trivalvis 1-locularis polysperma. Semina parietalia erecta aut ascendentia apice comosa. Albumen o. Embryo rectus, radicula parvâ inferâ, cotyledonibus plano-convexis oblongis. — Frutices ramis virgatis; foliis alternis parvis persistentibus squamiformibus, floribus spicatis.

I. TAMARIX *Desv. ann. sc. nat. 4. p. 348.* — Tamarices 4-5-andræ Auct.

Calyx 4-5-partitus; petala 4-5; stam. 4-5 petalis alterna, ferè omninò libera; ovarium apice longè attenuatum; stigmata 3 longa divaricata apice glandulosa; semina imæ valvularum basi aut ferè centro capsulæ inserta; seminum coma pilis plurimis simplicibus ex apice ortis constans.

1. T. AFRICANA (Poir. voy. 2. p. 139.) glabra subglauca, foliis lanceolatis subamplexicaulibus imbricatis, spicis sessilibus squamosis densis crassis vix latitudine triplò longioribus. ♂ in maritimis Occitanie et agri Ruscinonensis ex cl. Benth. — Fl. fr. 5. p. 535. (v. s.)
2. T. GALLICA (L. sp. 386.) glabra glauca, foliis minutis amplexicaulibus adpressis acutis, spicis lateralibus subpaniculatis gracilibus latitudine quinatuplò longioribus. ♂ in arenosis Gallie Mediterraneæ frequens et secus Oceanum usque ad Pictaviam et etiam Neustriam ex cl. de Brébisson. — Fl. fr. 4. p. 399. — Lob. ic. 2. t. 218. (v. v.)

II. MYRICARIA *Desv. l. c. p. 349. DC. l. c.* — Tamarices monadelphæ Auct.

Calyx 5-partitus; petala 5; stam. 10, alterna breviora, filamentis ad medium circiter monadelphis; stylus o; stigmata in capitulum congesta; semina secus lineam valvularum mediam inserta, apice in filum barbato-plumosum desinentia.

1. M. GERMANICA (Desv. l. c.) fruticosa glabra, foliis lineari-lanceolatis sessilibus, spicis terminalibus solitariis, bracteis pedicello longioribus. ♂ in arenosis inundatis et fluviorum ripis Gallie australis, medice, orientalis borealisque. — Tamarix Germanica L. — Fl. fr. 4. p. 399. — Lob. ic. 2. t. 218. (v. v.)

Ordo XLIII. PORTULACEÆ. *Juss. gen.* 313. *excl. gen.* DC.  
*fl. fr.* 4. p. 398. *prod. t.* 3. *ined.*

Calyx liber aut imo ovario subadnatus, sæpius disepalus sepalis oppositis, interdum 3-5-sepalus, sepalis basi in calycem 3-5-lobum concretis. Petala rariùs 0, sæpiùs 3, 4 et 6, nunc omninò libera, nunc imò basi in tubum brevissimum subconnexa, sepalis alterna cum numero congruunt. Stamina cum petalis nunc imo calyci, nunc forsan toro inserta, numero in iisdem generibus imò speciebus varia, omnia fertilia; filamenta inter se libera, petalis cum (petala basi coalescunt) concreta et iis opposita (cum numero cum iis congruunt). Antheræ biloculares, rimà duplici dehiscentes, oscillatoriæ. Ovarium 1 sæpiùs subrotundum uniloculare; stylus nunc 1 apice fissus in stigmata plurima; nunc nullus aut subnullus, stigmatibus plurimis liberis ex apice ovarii ortis. Capsula 1-locularis, nunc circumscissa dehiscens, nunc 3-valvis ab apice ad basim dehiscens. Semina ∞ placentæ centrali adfixa. Albumen farinosum centrale. Embryo periphericus; radícula cylindracea longa. — Herbæ (in nostris) pingues. Folia alterna rariùs opposita integra exstipulata. Flores axillares aut terminales.

I. PORTULACA *Juss. gen.* 312.

Calyx persistens compressus 2-valvis; petala 5; stam. 6-12; ovarium calycis basi subadnatum 1-loculare, stylo 1 infra medium 5-fido; capsula circumscissa polysperma, seminibus columnæ centrali 5 placentis formatæ adfixis.

1. *P. OLERACEA* (L. sp. 638.) foliis cuneiformibus confertis lævibus sessilibus, floribus sessilibus. ① in locis cultis. — *Fl. fr.* 4. p. 402. — *Fuchs. hist. p.* 113. ic. (v. v.)

II. MONTIA *Mich. gen. t.* 13. f. 2. *L. gen.* 101.

Calyx persistens 2-3-valvis; corolla 1-petala 5-partita laciniis 3 alternis minoribus; stamina 3-5; stylus 1 brevis caducus apice 3-fidus lobis interne stigmatosis; capsula abortu 3 septorum 1-locularis tri-valvis trisperma, seminibus basi columnæ 3 placentis formatæ adfixis.

1. *M. FONTANA* (L. sp. 129.) foliis oppositis. ① in humidis secùs aquas provinc. austr., occident., mediarum borealiumque. — *Fl. fr.* 4. p. 402. — *Vaill. bot. t.* 3. f. 4. (v. v.)

Ordo XLIV. PARONYCHIEÆ. *St.-Hil. plac. lib. p. 56. Juss. mem. mus. 1. p. 387. DC. prod. t. 3. ined.*

Calyx 5-sepalus rarissimè 3-4-sepalus, sepalis nunc imâ basi (5-partitus) nunc ad medium (5-fidus) et ultrâ (5-dentatus) concretis. Petala parva squamiformia staminum steriliùm æmula, sæpiùs tot quot sepala, inter lobos tubo inserta, rariùs nulla. Stam. calycis tubo antè lobos exactè inserta, etiam in generibus apetalis, sepalis numero æqualia aut abortu pauciora; filamenta distincta; antheræ 2-loculares. Ovarium liberum; styli 2-3 nunc à basi distincti, nunc plus minùs concreti. Fructus siccus parvus sæpiùs membranaceus, nunc evalvis indehiscens, nunc trivalvis. Semina nunc plurima placentæ centrali adfixa, nunc solitaria ex apice funiculi longi à fundo loculi orti subpendula. Albumen farinaceum. Embryo cylindraceus lateralis incurvus aut periphericus; radícula ad hylum versa. — Herbæ aut suffrutices ramosissimi. Folia sæpiùs opposita rariùs alterna, nunc nuda, nunc stipulis scariosis utrinquè instructa, sessilia, parva, integerrima. Flores parvi, nunc in axillis sessiles, nunc in cymas terminales variè dispositi. Bracteæ stipulis analogæ.

Trib. I. *TELEPHIEÆ. DC. l. c.*

Calyx 5-partitus. Petala et Stam. 5. imo calyci inserta. Styli 3 liberi aut basi subconcreti. — Folia alterna stipulacea.

I. *TELEPHIUM Tourn. t. 128. L. gen. 377.*

Calyx ferè quinquepartitus persistens; petala 5 calyci æqualia eidem inserta; stam. 5 pariter perigyna; stylus brevis; stigmata 3; ovarium basi 3-loculare apice 1-loculare septis non ultrâ productis; capsula ovata 3-quetra in mucronem supernè attenuata 1-locularis 3-valvis polysperma; embryo curvus non annularis, albumini implicitus, radiculâ ad receptaculum et seminis hylum spectante.

1. *T. IMPERATI* (L. sp. 388.) procumbens, foliis alternis brevissimè stipulaceis, floribus corymbosis terminalibus. 2/ in siccis et calidioribus Galliæ australis. — Fl. fr. 4. p. 400. — Clus. hist. 2. p. 67. f. 3. (v. v. DC.)

II. *GORRIGIOLA L. gen. 378.*

Calyx ferè 5-partitus persistens; petala 5 calyci subæqualia eidem inserta; stam. 5 pariter perigyna; stylus brevis; stigm. 3; capsula 1-sperma trivalvis sæpè indehiscens triquetra, insertus seminis versùs apicem funiculus ex imâ capsulâ oriens.

1. *C. LITTORALIS* (L. sp. 388.) caulibus diffusis inter flores foliosis, foliis alternis linearibus. ① in arenosis secùs rivos Galliæ austr., occid., mediæ et circâ Argentoratum. — Fl. fr. 4. p. 401. excl. var.  $\beta$ . — Barr. ic. 532. (v. v.)



2. *C. TELEPHIFOLIA* (Pourr. act. toul. 3. p. 316.) caulibus diffusis inter flores aphyllis, foliis alternis, radicalibus linearibus, caulinis ovatis.  $\mathcal{U}$  in agro Ruscinonensi. — Fl. fr. 5. p. 527. An satis à precedente distincta? (v. v.)

Trib. II. *ILLECEBREÆ. DC. l. c.*

Calyx 5-partitus. Pet. 5 aut 0. Stam. 2-5 imo calyci inserta. Styli 2 liberi aut subconcreti. Capsula indehiscens 1-sperma, funiculus e fundo ortus longus apice semen subpendulum gerens. — Herbæ rarò suffrutices. Folia stipulis scariosis aucta opposita.

III. *HERNIARIA Tourn. inst. t. 288. L. gen. n. 308. DC. l. c.*

Calyx ferè 5-partitus intus subcoloratus; petala 5 filiformia integerrima sepalis alterna, interdum 0 aut minima; stam. 5 aut abortu 2-3; styli 2 breves distincti aut basi concreti; caps. 1-sperma indehiscens calyce tecta.

1. *H. POLYGONOIDES* (Cav. ic. 2. t. 137.) caule erecto dichotomo frutescente, foliis ovatis mucronatis, stipulis linearibus acuminatis folio duplò brevioribus, floribus terminalibus glomeratis sessilibus.  $\mathfrak{J}$  in collibus maritimis Galloprovinciæ ex Lam. — Fl. fr. 3. p. 406. — Moris. s. 5. t. 29. u. 5. Illecebrum suffruticosum L. Valdè dubia civis. (v. s. ex Hisp.)
2. *H. GLABRA* (L. sp. 317.) caulibus ramosissimis prostratis glabris, foliis ovatis vix acutis glabris subciliatisve, stipulis minimis ciliatis, florum sessilium glomerulis axillaribus multifloris glabris.  $\textcircled{1}$   $\mathcal{U}$  in arvis et sabulosis. — Fl. fr. 3. p. 405. Fl. dan. t. 529. (v. v.)
3. *H. HIRSA* (L. sp. 317.) caulibus ramosis prostratis hirsutis, foliis ovatis lanceolatisve obtusis hirsutis stipulisque folio quintuplò brevioribus ciliatis, florum sessilium glomerulis axillaribus paucifloris hirsutis.  $\mathcal{U}$  in arvis. — Fl. fr. 3. p. 405. Paronychia pubescens Fl. fr. 3. p. 403. — Zannich. ic. 284. *H. latifolia* Lapeyr. abr. 127. (v. v.)  
 $\beta$ . *incana*, foliis glomerulisque plus minùs incanis. — In sterilibus provinc. australium. — *H. incana* Lam. DC. fl. fr. 5. p. 375. (v. v.)
4. *H. CINEREA* (DC. fl. fr. 5. p. 375.) caulibus ramosis herbaceis ascendentibus tomentosis, foliis lanceolatis acutis canescenti-hispidis, stipulis brevissimis ciliatis, florum sessilium glomerulis axillaribus multifloris canescenti-hispidis.  $\textcircled{1}$  in arvis Occitanicæ. (v. v. D.)
5. *H. ALPINA* (Vill. dauph. 2. p. 556. non Lois.) caulibus prostratis ramosissimis puberulis, foliis obovatis obtusis crassiusculis, stipulis acutis ciliatis folio vix duplò brevioribus, florum subpedicellatorum glomerulis 1-3-floris hirsutis.  $\mathcal{U}$  in Alpibus, Delph. Galloprovinciæ. — Fl. fr. 5. p. 375. *H. Alpina*  $\beta$  Fl. fr. 3. p. 406. (v. v. DC.)

IV. *ILLECEBRUM Gærtn. carp. t. 184. Juss. mem. mus. 1. p. 386. DC. l. c.* — *Illecebrum* sp. L.

Calyx 5-partitus ferè 5-sepalus, sepalis subcucullatis dorso incrassatis ad apicem in cornu aristatum desinentibus; petala 0 aut squamulæ 5 subulatæ sepalis alternæ; stam. 2-5 sepalorum basi inserta; stylus vix ullus; stigmata 2 capitata; capsula calyce tecta 5-valvis aut ad strias 5 scissilis; semen (abortu?) solitarium lateraliter in capsulâ insertum; embryo lateralis vix curvus hinc albumini farinaceo applicitus.

1. *I. VERTICILLATUM* (L. sp. 298.) prostratum, stipulis scariosis, foliis oppositis axillaribus.  $\textcircled{1}$  in humidis arenosis Galliæ mediæ, occidentalis australisque. Fl. fr. 3. p. 403. — Vaill. par. t. 15. f. 1. (v. v.)

V. PARONYCHIA *Juss. l. c. p. 388. DC. prod. t. 3. — Illecebrī sp. L.*

Cal. 5-partitus, sepalis dorso non incrassatis, sæpius apice mucronatis subcucullatis; petala seu squamulæ 5 lineares sepalis alternæ; stam. 5; stylus 1; stigmata 2; capsula 1-sperma indehiscens aut 5-valvis calyce tecta.

Sect. I. CHÆTONYCHIA *DC. l. c.*

Sepala æqualia apice dilatata membranacea, dorso in aristam desinente. — Flores cymosi.

1. *P. cymosa* (Poir. dict. 5. p. 26.) caule ramoso erecto glabriusculo, foliis verticillatis subteretibus glabris apice aristatis, stipulis bracteisque brevissimis, floribus cymosis glaberrimis, sepalis longè aristatis. ① in Cebennis, propè Arausionem ex Vill., Forum Julii (cl. Solier). — Fl. fr. 3. p. 402. — Illec. cymosum L. Schrad. journ. 1801. p. 408. t. 4. (v. s.)

Sect. II. EUNYCHIA *DC. l. c.*

Sepala æqualia apice non dilatata aristata mucronata aut subinermia. — Flores in axillis congesti.

2. *P. echinata* (Lam. fl. fr. 3. p. 232.) caulibus prostratis hirsutis, foliis lanceolato-ovatis acutis glabris, basi dilatatis acutis folio triplo brevioribus, floribus in axillis fasciculatis verticillum mentientibus, bracteis subnullis, sepalis linearibus dorso aristatis. ① in maritimis Corsicæ et Galloprovinciæ. — Fl. fr. 3. p. 402. — Bocc. sic. t. 20. f. 3. (v. s.)
3. *P. polygonifolia* (DC. fl. fr. 3. p. 403.) caulibus prostratis elongatis glabris, foliis lanceolato-ovatis vix acutis glaberrimis, stipulis ovatis acutis folio vix dimidio brevioribus, bracteis lanceolatis acuminatis folio longioribus, floribus sessilibus solitariis lateralibus terminalibusque, sepalis subacutis. ½ in Delphinatū montibus et Pyrenæis editor. ex Benth. — I. polygonifolium. Vill. dauph. 2. p. 557. t. 16. (v. s.)
4. *P. argentea* (Lam. fl. fr. 3. p. 230.) caulibus nodosis ramosis prostratis tomentosius, foliis lanceolatis acutis brevissimè ciliatis, stipulis folio brevioribus ovatis acutis, bracteis argenteis lanceolatis acuminatis flores sessiles hirsutos glomeratos axillares occultantibus, sepalis mucronatis. ½ in siccis Gallis australis. — Fl. fr. 3. p. 404. Illec. Paronychia L. sp. 299. — Barr. ic. 726. (v. v.)
5. *P. capitata* (Lam. fl. fr. 3. p. 229.) caulibus nodosis ramosis prostratis hirsutis, foliis lineari-lanceolatis acuminatis ciliatis, stipulis lanceolato-ovatis acutis folia subsuperantibus, floribus terminalibus congesto-capitatis, bracteis argenteis maximis rotundo-ovatis acuminatis florem sessilem hirsutum occultantibus, sepalis obtusis dorso aristatis. ½ in collibus siccis provinciarum mediterranearum. — Fl. fr. 3. p. 404. Illec. capitatum L. sp. 299. (v. v.)
6. *P. serpyllifolia* (Poir. dict. 5. p. 24.) caulibus nodosis ramosis prostrato-repentibus, foliis ovatis ciliatis subacutis, stipulis linearibus acuminatis folio brevioribus, bracteis argenteis maximis obliquè ovatis acutis flores hirsutos sessiles occultantibus, sepalis obtusiusculis. ½ in Pyrenæis et Alpibus Delphin. — Fl. fr. 3. p. 404. Illecebrum serpyllifolium Vill. Schrad. journ. 1801. p. 413. t. 4. (v. v.)
7. *P. nivea* (DC. prod. t. 3. ined.) caulibus nodosis ramosis prostrato-repentibus, foliis lanceolatis ciliatis acutis, stipulis lanceolatis acuminatis folium superantibus æquantibusve, bracteis argenteis maximis obliquè rotundato-ovatis acutis flores hirsutos sessiles occultantibus, sepalis linearibus obtusiusculis. ½ in rupestribus Narbonnæ loco dicto Pech de l'Agnè et montis Ventosi (cl. Requier). — Illecebrum cephalotes Bieb. (v. s.)

Trib. III. *POLYCARPÆÆ*. DC. l. c.

Calyx 5-partitus. Pet. 5 aut o. Stam. 1-5 imo calyci inserta. Styli 2-3 nunc à basi liberi, nunc concreti. Caps. 1-locularis coosperma. Semina placentæ centrali adfixa. — Herbæ aut suffrutices, folia opposita; stipulæ scariosæ.

VI. *POLYCARPON* L. gen. n. 105. DC. l. c.

Cal. profundè 5-fidus, sepalis basi plus minùs coalitis margine membranaceis concavis carinatis apice mucronatis; petala 5 emarginata; stam. 3-5; styli 2 brevissimi; caps. 1-locul. 3-valvis polysperma.

\* *Floribus plerùmque 3-andris.*

1. *P. TETRAPHYLLUM* (L. sp. 131.) triandrum, petalis emarginatis, foliis caulinis quaternis, rameis oppositis. (I) in cultis et ad sepes umbrosas Galliarum australis et occidentalis. — Fl. fr. 4. p. 767. — Barr. ic. 534. (v. v.)

\* \* *Floribus plerùmque 5-andris.*

2. *P. ALSINÆFOLIUM* (DC. l. c.) pentandrum, petalis subintegerrimis, foliis ovalibus subcarnosis, floribus in cymas terminales congestis. (I) in arenosis maritimis Occitaniarum, Galloprov. — Hægea alsinæfolia Biv. manip. 3. p. 7. — Bocca sic. t. 38. Folia lævia ovalia; flores pauciores magis congesti quàm in *P. tetraphyllo*. (v. v.)

3. *P. PEPLOIDES* (DC. l. c.) pentandrum, petalis integerrimis, foliis oppositis obovatis, floribus in cymas terminales congestis. (I) in cultis et arenosis maritimis agri Ruacinonensis circà Perpignan et Collioure. — Hægea polycarpoides Biv. manip. 2. n. 3. Arenaria peploides Lapeyr. abr. 251. non L. Folia rotundiora, cymæ densiores quàm in *P. tetraphyllo*. (v. v.)

Trib. IV. *SCLERANTHÆÆ*. Linck enum. 417. DC. l. c.

Calyx 4-5-fidus, tubo urceolato. Pet. o. Stam. 1-10 fauci inserta. Styli 2, aut 1 apice emarginatus. Utriculus membranaceus 1-spermus, tubo calycis indurato tectus. Semen funiculo longo è fundo loculiorto apice recurvo suspensum. — Herbæ oppositifoliæ non stipulatæ.

VII. *SCLERANTHUS* L. gen. n. 562.

Calyx tubulosus, collo coarctatus, suprâ 5-fidus; petala nulla; stamina sæpiùs decem minima summo calyci insidentia; styli et stigmata 2; capsula tenuissima monosperma evalvis, tecta calyce suprâ coarctato.

1. *S. FRAXINÆUS* (L. sp. 580.) foliis oppositis linearibus, calycibus fructus margine membranaceis obtusiusculis clausis. (I) in arvis arenosis. — Fl. fr. 4. p. 403. — Vaill. bot. t. 1. f. 5. (v. v.)

2. *S. ANTHUS* (L. sp. 580.) foliis oppositis linearibus, calycibus fructus margine membranaceis acuminatis patulis. (I) in arvis. — Fl. fr. 4. p. 403. Gærtn. fruct. 2. p. 196. t. 126. (v. v.)

Trib. V. *MINUARTIÆÆ*. DC. l. c.

Calyx 5-partitus. Pet. o aut minima. Stam. 3-10 imo calyci inserta. Styli 5. Caps. 1-locul. 3-valvis. Sem. plurima placentæ centrali adfixa. — Herbæ ☉ non stipulatæ oppositifoliæ.

VIII. LOEFLINGIA *L. gen. n. 52.*

Calyx ferè 5-partitus laciniis basi bidentatis; petala 5 minima conniventia, stamina 3; stylus unicus, stigma triplex; capsula unilocularis trivalvis polysperma.

1. *L. HISPANICA* (*L. sp. 50.*) foliis oppositis minimis basi appendiculatis, appendicibus stipulæformibus. ① in maritimis agri Narbonensis et Ruscinnensis. — *Fl. fr. 5. p. 603.* — *Loefl. it. 113. t. 1. f. 2. (v. s.)*

†.

Ordo XLV. CRASSULACEÆ. *DC. bull. phil. 1801. n. 49. p. 1. fl. fr. 4. p. 382. prod. t. 3. ined.* — *Sempervivæ. Juss. gen. 207.*

Calyx constans sepalis plurimis 3-20 basi plus minus concretis et ideò multipartitus. Petala tot quot sepala, cum iis alterna, imò calyci inserta, nunc inter se libera, nunc in corollam gamopetalam concreta. Stam. cum petalis inserta, nunc iis numero æqualia et cum iis alterna, nunc dupla, alia petalis alterna longiora precociora, alia petalis opposita breviora seriora. Filamenta inter se libera subulata; antheræ biloculares rimâ duplici dehiscentes. Squamæ nectariferæ ad basin carpellorum solitariae. Carpella tot quot petala et iis opposita et circâ axim idealem verticillata, inter se libera, unilocularia, rimâ longitudinali ad angulum internum dehiscencia. Semina ad angulum internum carpelli duplici serie adfixa. Albumen tenue carnosum. Embryo rectus, radiculâ ad hylum directâ. — Herbæ foliis carnosis sæpius alternis, floribus sæpius cymosis.

§. I. ISOSTEMONÆ. (*DC.*)

1. *TILLEA Mich. gen. t. 20. DC. bull. phil. n. 49. p. 2.* — *Tillea sp. L.*

Calyx 3-4-partitus; petala 3-4; stam. 3-4; squamæ 0 aut minimæ; carpella 3-4 medio subconstricta disperma. — Herbæ parvæ annuæ.

1. *T. MUSCOSA* (*L. sp. 186.*) caule ramoso, foliis basi connatis, floribus sessilibus. ① in humidis provinc. australium, occidentalium et mediarum. — *Fl. fr. 4. p. 385. Lam. ill. t. 90. f. 2. (v. v.)*

II. BULLIARDA *DC. pl. grass. p. 7. Fl. fr. 4. p. 384.*

Calyx 4-fidus; petala 4; stamina 4; ovaria 4; squamæ 4 lineares calycis longitudine; carpella 4 polysperma. — Herbæ parvæ annuæ.

1. *B. VAILLANTII* (*DC. pl. grass. t. 74.*) caule erecto dichotomo, foliis oblongis acutis pedunculo brevioribus. ① in umbrosis humidis ad oras aquilegiorum Fontisbellaquei. — *Fl. fr. 4. p. 385. Tillæa aquatica Lam. ill. t. 90. f. 1. — Vaill. bot. t. 10. f. 1. (v. v. DC.)*

## §. II. DIPLOSTEMONEÆ. (DC.)

## † Gamopetalæ.

III. UMBILICUS DC. *bull. phil.* 1801. n. 49. *prod. t. 3. ined.* — *Cotyledonis* sp. *auct.*

Calyx 5-partitus; cor. gamopetala campanulata 5-fida, lobis tubi circiter longitudine; stamina 10 tubo corollæ inserta; squamæ 5 ovals aut cuneatæ obtusæ; carpella 5 apice attenuata, stylis subulatis.

§. 1. *Sediformes* (DC.), *foliis caulinis alternis.* — *Herbæ annuæ.*

1. U. SEDOIDES (DC. mss.) caule subradicante glabro, foliis oblongis convexis obtusis glabris, floribus ad apicem paucis subsessilibus. ① in summis Pyrenæis præsertim orientalibus propè nives æternas. — *Cotyledon sedoides*. DC. fl. fr. 5. p. 521. C. sediformis Lapeyr. — Herba parva simplex aut ramosa Sedi atrati ferè facie. (v. v.)

§. 2. *Cotyloideæ* (DC.), *corollis vix ad medium 5-fidis, foliis radicalibus petiolatis cucullatis plus minus peltatis.*

2. U. PENDULINUS (DC. pl. grass. t. 156.) radice tuberosâ, foliis inferioribus peltatis repando-crenatis, caule subsimplici, floribus racemosis pendulis, bracteis integris. ½ ad muros vetustos et rupes prov. australiorum et occidentaliū. — Fl. fr. 4. p. 382. *Cotyledon umbilicus* L. sp. 615. — *Clus. hist.* 2. p. 63. f. 1. — *Lob. ic.* t. 386. f. 2. (v. v.)

## †† Polypetalæ.

IV. SEDUM DC. *bull. phil.* n. 49. *prod. t. 3.* — *Sedum* et *Rhodiola* L.

Calyx 4-7-partitus sæpius 5-partitus, sepalis ovatis sæpius turgidis foliiformibus; petala et carpella totidem; stamina petalorum numero dupla, squamæ nectariferæ ovatæ integræ. — Surculi steriles folia conferta gerentes.

§. 1. *Foliis planis, floribus luteis.*

1. S. RHODIOLA (DC. pl. grass. t. 143.) radice tuberosâ, foliis lanceolatis denticulatis acuminatis, floribus abortu dioicis corymboso-capitatis. ½ in umbrosis et rupestribus Alpium Delphin. et Pyrenæorum. — Fl. fr. 4. p. 386. *Rhodiola rosea* L. sp. 1465. — *Cam. epit.* 169. ic. Sepala 4-5. Stam. 8-10. Caps. 4-5. (v. v. DC.)

§. 2. *Foliis planis, floribus albis aut purpurascensibus.*

2. S. TELEPHIUM (L. sp. 616.) caule erecto, foliis magnis sparsis aut oppositis ovatis serratis subintegræve sessilibus, corymbo folioso. ½ ad muros, vineas et inter lapides. — Fl. fr. 4. p. 387. — *Fuchs. hist.* 800. ic. (v. v.)

β. *latifolium* (L.) foliis latis sæpius oppositis, staminibus plerumque (respectu corollæ) longioribus. — *S. latifolium* Bert. *amœn.* 366. — *Clus. hist.* 2. p. 66. f. 1. (v. v.)

3. S. ANACAMPSEROS (L. sp. 616.) caulibus decumbentibus, foliis sparsis cuneiformibus integerrimis sessilibus, floribus corymbosis. ½ inter rupes apricas in Alpibus Galloprovinciæ et Pyren. orient. ex Lapeyr., propè Parisios ex Merat? — Fl. fr. 4. p. 387. — *Lob. ic.* t. 390. f. 2. (v. v.)

4. S. STELLATUM (L. sp. 617.) foliis sparsis rotundo-spathulatis ovatisve obtusè angulatis basi in petiolum subattenuatis, floribus lateralibus sessilibus solitariis, capsulis stellatum patentibus. ① in lapideis umbrosis provinc. australium, Arvernæ? ex Delarbre. — Fl. fr. 4. p. 388. — *Col. phyt.* 30. t. 11. (v. v. DC.)

5. S. CREPA (L. sp. 617.) caule ramoso paniculato pubescente, foliis sparsis

verticillatisque oblongo-lanceolatis spathulatisve integerrimis, petalis aristatis. ① in lapidosis secus colles et muros. — Fl. fr. 4. p. 389. — Clus. hist. 2. p. 68. ic. (v. v.)

β. *galioides* (DC. ined.) foliis superioribus suboppositis, infimis verticillatis spathulatis. — S. *galioides* All. ped. t. 65. f. 3. — Fl. fr. 4. p. 389. S. verticillatam Latourr. non L. (v. v.)

γ. *alsinaefolium* (DC. ined.) foliis alternis ovalibus. — S. *alsinaefolium* All. ped. t. 22. f. 2. Fl. fr. 3. p. 388. (v. s.)

§. 3. *Foliis teretibus, floribus albis purpurascentibus aut cœruleis.*

6. S. *CRUCIATUM* (Desf. cat. 162.) caulibus patulis basi ramosissimis in parte superiore glandulosis, foliis verticillatis semiteretibus oblongis obtusis glaberrimis, superioribus sparsis, floribus corymbosis, petalis acuminatis. ② in Corsicæ rupibus propè Corte reperiit Ph. Thomas. — Fl. fr. 4. p. 389. S. Monregalense Balb. misc. p. 23. f. 6. (v. s.)

7. S. *ALBUM* (L. sp. 619.) caulibus ascendentibus glabris, foliis sparsis teretiusculis oblongis obtusis patentibus glaberrimis, cymâ ramosâ, laciniis calycinis obovatis obtusis, petalis vix acutis. ② in muris et lapidibus frequens. — Fl. fr. 4. p. 390. S. micranthum Bast. Fl. fr. 5. p. 523. — Fuchs. hist. 35. ic. (v. v.)

β. *turgidum* (DC. ined.) foliis ovatis crassis, surculorum juniorum erectis, petalis obtusis. ② in Pyrenæis editioribus circâ Valletriam, usque ad aquas Bigerronum (Bagnères) descendens. — S. *turgidum* Ram. in DC. fl. fr. 4. p. 390. (v. v.)

8. S. *VILLOSUM* (L. sp. 620.) foliis sparsis oblongis suprà planiusculis pedunculis axillaribus 1-3-floris glanduloso-pubescentibus, floribus corymboso-paniculatis, laciniis calycinis lanceolatis subobtusis, petalis ovatis subobtusis. ① in pratis paludosis montosis. — Fl. fr. 4. p. 392. — Clus. hist. 2. p. 59. f. 3. (v. v.)

9. S. *MRAUTUM* (All. ped. t. 65. f. 5.) caule basi atolonifero, foliis villosis semiteretibus, inferioribus confertis, superioribus sparsis, floribus cymosis, laciniis calycinis lanceolato-ovatis obtusis, petalis lanceolato-ovatis aristatis. ② ② in rupestribus montosis Gallix australis, Arvernix, Pyrenæorum, propè Parisios ex Merat. — Fl. fr. 4. p. 392. (v. v.)

10. S. *CORSICUM* (Duby mss.) foliis glanduloso-villosis obovato-rotundis basi attenuatis obtusis, inferioribus confertis, superioribus sparsis, floribus cymosis laxis, laciniis calycinis lanceolato-ovatis obtusis brevibus, petalis lanceolato-ovatis subobtusis. ①? in Corsicâ propè Corte reperiit Ph. Thomas. (v. s.)

11. S. *AUBENS* (L. sp. 619.) caule erecto ramoso, foliis alternis oblongis obtusiusculis teretiusculis sessilibus patentibus glabris, cymâ ramosâ pubescente, floribus unilateralibus sessilibus, laciniis calycinis lanceolatis acutis, petalis acuminato-aristatis. ① in cultis et vineis passim. — Crassula rubens L. syst. veg. 253. Fl. fr. 4. p. 386. DC. pl. grass. t. 55. — Carpella matura puberula. (v. v.)

12. S. *CESPITOSUM* (DC. prod. t. 3. ined.) foliis alternis ovatis obtusis glabris, caule subsimplici glabro, floribus lateralibus subsessilibus, laciniis calycinis lanceolato-ovatis, carpellis stellatim patentibus. ① in apricis Monspeli, Avenionis, Antipolis, Corsicæ. — Crassula Magnolii DC. fl. fr. 5. p. 522. C. cespitosa Cav. ic. x. t. 69. f. 2. Tillæa rubra Gou. hort. 77. — Magn. bot. 237. ic. (v. v. DC.)

13. S. *ANDEGAVIENSIS* (DC. prod. t. 3. ined.) caule erecto basi simplici apice trichotomo, foliis subalternis rotundo-ovatis obtusis glabris, floribus pentandris in dichotomiâ et secus ramulos sparsis sessilibus, laciniis calycinis ovatis obtusis, petalis ovatis acutiusculis, capsulis erectis. ① in muris et rupibus schistosis Andegaviæ. — Crassula Andegaviensis DC. fl. fr. 5. p. 522. Sedum atratum Bast. ess. 167. — S. dasyphylli var. pentandra? (v. s.)

14. *S. NASTRHYLLUM* (L. sp. 618.) caule debili apice pubescente foliis oppositis ovatis obtusis pinguibus, floribus laxè racemosis sæpius hexapetalis, laciniis calycinis obovatis obtusis brevibus, petalis lanceolato-ovatis obtusiusculis.  $\mathcal{L}$  in aridis Galliz orientalis, australis et Pyrenæorum. — Fl. fr. 4. p. 391. Bull. herb. t. 11. (v. v.)
15. *S. BREVI-FOLIUM* (DC. rapp. 2. p. 79.) caule basi fruticuloso ramoso tortuoso, ramis erectiusculis glabris, foliis crassis globoso-ovatis obtusis brevibus, in aurculis oppositis, in caulibus florentibus sparsis, floribus corymboso-paniculatis, laciniis calycinis ovatis subacutis brevibus, petalis lanceolato-ovatis obtusiusculis.  $\mathcal{L}$  ad rupes apricas in Pyrenæis et in Corsicâ. — Fl. fr. 5. p. 524. *S. sphaericum* Lapeyr. (v. v.)
16. *S. ATRATUM* (L. sp. 1673.) caule ramisque erectis glabris, foliis sparsis teretibus obtusissimis glabris, floribus corymbosis fastigiatis, laciniis calycinis lanceolatis obtusis, petalis lanceolato-obovatis obtusiusculis. (1) in rupibus apricis Alpium, Pyrenæorum. — Fl. fr. 4. p. 391. All. ped. t. 65. f. 4. (v. v.)
17. *S. ANGELICUM* (Huds. ang. 196.) caulibus infirmis gracilibus pedicellis glabris, foliis subalternis ovatis gibbis pinguibus, cymâ bifidâ, laciniis calycinis lanceolato-ovatis subacutis, petalis lanceolatis acuminatis. (1) in rupibus Pyrenæorum et Galliz occidentalis. — Fl. fr. 4. p. 391. — Ray. syn. t. 12. f. 2. (v. v.)
18. *S. HEPTAPETALUM* (Poir. voy. barb. 2. p. 169.) caulibus erectis ramosis glabris, foliis sparsis ovato-oblongis planiusculis obtusis, floribus corymboso-paniculatis, laciniis calycinis obovatis obtusis, petalis 6-7 lanceolatis acuminatis. (1) ad rupes in Corsicâ. — Fl. fr. 4. p. 392. Fl. azurei. (v. s.)

§. 4. *Foliis teretibus basi solutis, floribus luteis.*

19. *S. SAXATILE* (L. sp. 619.) caule ramoso decumbente, ramis divaricatis, foliis sparsis semiteretibus obtusis, cymâ trifidâ, laciniis calycinis lanceolatis obtusis, petalis lanceolatis aristato-acuminatis. (1) in saxosis apricis Vogensium, Alpium, Pyrenæorum. — Fl. fr. 4. p. 394. All. ped. t. 65. f. 6. *S. divaricatum* Lapeyr. abr. 260. non Ait. — Lob. ic. 378. (v. v.)
20. *S. REPENS* (DC. fl. fr. 5. p. 525.) caulibus subsimplicibus basi repentibus ascendentibus, foliis ovato-teretibus adnato-sessilibus erectis sublaxis, floribus subcapitatis, laciniis calycinis lanceolatis obtusissimis, petalis lanceolatis subobtusis. (1) in summis Alpibus et Pyrenæis. — S. Guettardi Vill. dauph. 3. p. 678. t. 45. excl. syn. (v. v. DC.)
21. *S. ACER* (L. sp. 619.) foliis subovatis adnato-sessilibus gibbis erectiusculis alternis, cymâ trifidâ, laciniis calycinis ovalibus obtusis, petalis lanceolatis acuminatis.  $\mathcal{L}$  in apricis et muris frequens. — Fl. fr. 4. p. 393. Bull. herb. t. 30. DC. pl. grass. t. 117. (v. v.)
- β. glaciale*, floribus congestis terminalibus, caule ramisque radicanlibus. In Alpibus Galloprovinciæ ad nives. — *S. glaciale* Clarion in DC. fl. fr. 4. p. 393. (v. s.)
22. *S. SEXANGULARE* (L. sp. 620.) foliis teretibus adnato-sessilibus subpatulis ternatim verticillatis sparsisque, in surculis sexfariam imbricatis, cymâ trifidâ, laciniis calycinis lanceolatis obtusis, petalis lanceolatis acuminatis.  $\mathcal{L}$  in pratis siccis. — Fl. fr. 4. p. 394. — Cam. epit. 856. ic. *S. boloniense* Lois. not. 71. Fl. fr. 5. p. 525. (v. v.)
23. *S. REFLEXUM* (L. sp. 618.) foliis sparsis subulatis erectis, floribus breviter pedicellatis cymosis subcorymbosis, cymæ ramis virgineis reflexo-nutantibus, laciniis calycinis lanceolato-ovatis subacutis, petalis linearibus acutis.  $\mathcal{L}$  ad rupes et muros. — Fl. fr. 4. p. 394. — Fuchs. hist. 33. ic. (v. v.)
24. *S. SUPASTRÆ* (L. sp. 618.) caulibus basi ramosis, floriferis erectis, foliis teretibus subulatis basi solutis glaucis, surculis sterilibus cylindraceis densè imbricatis, floribus cymosis 5-7-petalis, laciniis calycinis ovato-lanceolatis obtusis, petalis lineari-lanceolatis obtusis.  $\mathcal{L}$  in muris et rupibus apricis provinc. aus-

traliū. — engl. bot. t. 170. *S. reflexum* DC. pl. grass. t. 116. — Dill. elth. 2. f. 333. (v. v.)

25. *S. ALTISSIMUM* (Lam. dict. 4. p. 634.) foliis sparsis teretibus acutis, inferioribus imbricatis, superioribus supra planis, floribus densè cymosis, laciniis calycinis ovatis acutis, petalis 6-8 lanceolatis subobtusis.  $\mathcal{L}$  inter saxorum fragmenta in provinciis australibus et in radicibus Pyrenæorum. — Fl. fr. 4. p. 394. DC. pl. gras. t. 40. *S. Nicæense* All. ped. t. 90. f. 1. — Jacq. vind. t. 81. (v. v.)

26. *S. ANOPETALUM* (DC. rapp. 2. p. 80.) foliis sparsis teretibus acutis erectis glaucis, surculorum imbricatis, floribus cymosis sessilibus, laciniis calycinis lanceolatis acuminatis, petalis 6-7 linearibus acuminatis erectis.  $\mathcal{L}$  in collibus et rupestribus provinc. australium. — Fl. fr. 5. p. 526. *S. Hispanicum* Fl. fr. 3. p. 395. non L. (v. v.)

27. *S. AMPLEXICAULE* (DC. rapp. 2. p. 80.) foliis sparsis teretibus acutis, surculorum basi in membranam amplexicantem dilatatis apice acuminatis, laciniis calyc. lanceolatis acutis, petalis lineari-lanceolatis subobtusis.  $\mathcal{L}$  in Cebennis et monte Ventoso. — Fl. fr. 5. p. 526. (v. s.)

#### V. SEMPERVIVUM *L. gen.* 612.

Calyx 6-12-fidus; petala et ovaria totidem; stamina petalorum numero dupla; squamæ ovatæ, latæ, emarginatæ aut laceræ.

##### §. 1. *Floribus ochroleucis.*

1. *S. GLOBIFERUM* (L. sp. 665.) foliis ciliatis, propaginibus globosis, floribus patentibus dodecagynis.  $\mathcal{L}$  in montibus Alsatiæ ex Mapp. — Fl. fr. 4. p. 397. Jacq. aust. t. 40. (v. s. ex Pedemontio.)

##### §. 2. *Floribus purpurascensibus.*

2. *S. ARACHNOIDEUM* (L. sp. 665.) propaginibus globosis villis intertextis, floribus cuneagynis, squamis oblongis truncato-dentatis.  $\mathcal{L}$  ad rupes apricas Alpium, Pyrenæorum, Cebennarum et montium Velaunorum. — Fl. fr. 4. p. 397. — Barr. ic. t. 391, f. 1 et 393. (v. v.)

3. *S. MONTANUM* (L. sp. 665.) foliis pubescentibus, propaginibus patentibus imberbibus, floribus subdodecagynis, squamis minimis.  $\mathcal{L}$  ad rupes apricas in Alpibus et Pyrenæis. — Fl. fr. 4. p. 396. Jacq. austr. app. t. 41. DC. pl. grass. t. 105. (v. v.)

4. *S. TECTORUM* (L. sp. 664.) foliis ciliatis, propaginibus patentibus, floribus dodecagynis, squamis cuneiformibus carunculatis.  $\mathcal{L}$  ad rupes Alpium et Pyrenæorum, in muris et tectis. — Fl. fr. 4. p. 396. — Fuchs. hist. 32. ic. (v. v.)



Ordo XLVI. NOPALEÆ. *Juss. Dict. sc. nat.* 35. p. 144. —  
*Cacti. Juss. gen.* p. 310.

Calyx monosepalus ovario adhærens,  $\infty$ -partitus. Petala numero indefinita. Stamina numero indefinita, cum petalis, disco ovarii apicem vestienti inserta. Stylus 1 tubulosus; Stigma multifidum. Bacca umbilicata 1-locularis polysperma. Semina in pulpâ nidulantia, integumento duplici; externo crustaceo, interno tenui membranaceo. Albumen nullum. Embryo incurvus aut spiralis. Radicula ad hylum versa. — Frutices caulibus carnosiss, sæpè folia æmulantibus, pilorum fasciculis aut aculeis quinquunciali ordine dispositis.

I. CACTUS *L. gen.* 613.

Vide caracterem ordinis.

1. *C. OPUNTIA* (*L. sp.* 669.) articulato-prolifer latus, articulis ovatis compressis, spinis setaceis, floribus luteis. Ex America orta spontè nunc crescit in Galloprovinciâ. — *Fl. fr.* 4. p. 405. — *Mill. dict.* 1. t. 191. (v. v. in Valesiâ.)

Ordo XLVII. FICOIDEÆ. *Juss. gen.* 315. *DC. fl. fr.* 5. p. 528.

Calyx monosepalus definitè partitus liber aut ovario adhærens. Corolla summo calyci inserta polypetala aut petalis basi coalitis monopetala, aut cum calyce coalita. Stamina numero indefinita summo calyci inserta; antheræ introrsæ. Styli plures. Capsula aut drupa multilocularis, loculis numero stylorum polyspermis. Semina angulo loculorum interiori affixa. Embryo incurvus. Albumen farinaceum centrale. — Flores axillares terminalesve. Folia succulenta.

I. MESEMBRYANTHEMUM *L. gen.* p. 628.

Calyx ovario adhærens persistens 5-fidus; petala plurima linearia basi coalita; styli 4-10 sæpius 5; capsula umbilicata carnosa.

1. *M. RODIFLORUM* (*L. sp.* 687.) foliis alternis teretiusculis obtusis basi ciliatis, petalis calyce brevioribus. (1) in arenosis maritimis Corsicæ circa Urcinium. — *DC. pl. gr.* t. 88. fl. fr. 5. p. 529. (v. s.)

Ordo XLVIII. GROSSULARIÆ. DC. *fl. fr.* 4. p. 406. *Berlandier diss. ined.*

Calyx ovario adhærens 4-5-partitus regularis coloratus. Petala 5-4 fauci calycis inserta, cum ejus laciniis alterna, æqualia. Stamina 4-5 rarissimè 6, ibidem inter petala inserta; filamenta libera; antheræ biloculares longitudinaliter internè dehiscentes. Ovarium uniloculare, placentis 2 parietalibus oppositis. Stylus unicus 2-3-4-fidusve. Fructus baccatus umbilicatus subglobosusve, calyce persistente coronatus, unilocularis. Semina plura arillata podospermis longis filiformibus suspensa. Integumentum exterius gelatinosum membranaceumve, interius tenuissimè membranaceum, albumini arcè adhærens. Albumen coæneum. Embryo minutus rectus in extremitate angustiore seminis situs. Radicula obtusa. — Fructus inermes vel spinosi, foliis alternis lobatis et incis.

I. RIBES L. *gen.* 281.

Cal. 5-lobus; pet. 5 parva; stam. 5 rarissimè 6; semina oblonga subcompressa.

Sect. I. GROSSULARIA *Tourn. inst.* 1. p. 639.

Caules aculeati, pedunculi 1-3-flori, calyx plus minùs campanulatus.

1. *R. UVA-CRISPA* (L. sp. 292.) ramis aculeatis, foliis obtusè 3-5-lobis villosulis, pedunculis 1-2-floris rarissimè 3-floris, sepalis reflexis glabris, petalis ad apicem subrotundis glabris, fauce barbatà, stylis puberulis. § ad sepes. — *Fl. fr.* 4. p. 408.

*α. sylvestre* (DC.) foliis parvis utrinquè villosopubescentibus, baccis glabris. — *Lob. ic.* 2. p. 206. (v. v.)

*β. sativum* (DC.) foliis amplioribus sæpè subtus glabris lucidis baccis majoribus glabris pilosisve. — In hortis culta. — *R. Grossularia* L. sp. 291. (v. v. c.)

Sect. II. RIBESIA *Berl. l. c.*

Flores racemosi, pedunculi 4-flori, calyx campanulatus aut cylindricus, canlis inermis.

2. *R. ALPINUM* (L. sp. 291.) foliis 3-5-lobis obtusis subtus lucidis suprà pilosis, bracteis lanceolatis sæpè flore longioribus, petalis minutis subabortientibus, antheris plus minùs subsessilibus, stylis coalitis. § ad sepes et in umbrosis montium. — *Fl. fr.* 4. p. 407. — *J. Bauh. hist.* 2. p. 98. *ic.* (v. v.)
3. *R. RUBRUM* (L. sp. 290.) foliis obtusè 3-5-lobis subtus pubescentibus, junioribus sæpè subtomentosis suprà glabris, racemis nutantibus, bracteis obtusis pedicellis brevioribus, calycibus plano-campanulatis, sepalis obtusis, petalis obcordatis. § in vallibus Jurassi, humiliorum Alpium; et in hortis cultam. — *Fl. fr.* 4. p. 406. — *Lob. ic.* 2. p. 202. f. 1. (v. v.)
4. *R. PRAZUM* (Wulf. in Jacq. misc. 2. p. 36.) foliis acuminatis 3-5-lobis subcordatis inciso-serratis longè pedunculatis suprà pilosis, racemis erectis, bracteis flore brevioribus, sepalis obtusis, petalis obcordatis. § in montium locis umbrosis, inter lapides juxta rivos. — *Fl. fr.* 4. p. 407. *Jacq. ic. rar.* 1. t. 49. *Lam. ill.* t. 146. f. 2. (v. v.)

5. *R. menum* (L. sp. 291.) foliis subtus punctato-glandulosis 3-5-lobis, racemis laxis, bracteis minutis subulatis obtusave pedicello multò brevioribus, petalis oblongis, calycibus campanulatis, sepalis reflexis. ♂ in montibus Delph. ex Vill., Arvernæ ex Delarb., Monspeii ex Gouan., in hortis cultum. — Fl. fr. 4. p. 407. — J. Banh. hist. 2. p. 99. f. 1. (v. v. c.)

Ordo XLIX. SAXIFRAGÆ. *Vent. tabl. t. 2. p. 277. Ser. in DC. prod. t. 3. ined.*

Calyx plus minùsve gamosepalus persistens ovario adhærens vel liber; sepala 4-5. Petala 4-5 libera apici tubi sæpè brevissimi calycis inserta et cum sepalis alterna, rariùs nulla. Stamina libera calyci inserta 8-10, petalis sepalisque opposita; antheræ longitudinaliter dehiscentes. Ovarium imo calyci adnatum, 2-loculare carpellis duobus margine seminiferis inter se coalitis constans. Styli 2 rariùs 4-5. Stigmata apice dilatata. Fructus 2-locularis, intrà stylos foramine, vel a basi ad apicem dehiscens, aut indehiscens baccatus 4-5-locularis. Semina minuta sæpiùs numerosa. Albumen carnosum, embryo teres, radícula infera. — Herbæ, foliis alternis aut radicalibus sæpè carnosis.

+ COROLLA POLYPETALA.

I. SAXIFRAGA L. gen. 559.

Calyx 5-sepalus, sepalis plus minùs inter se et cùm ovario coalitis; corolla 5-petala, petalis breviter unguiculatis integris; styli 2; capsula 2-locularis intrà stylos foramine dehiscens; semina numerosa rugosa vel lævia.

Sect. 1. PORPHYRION *Tausch. hort. canal. fasc. 1. Ser. in DC. prod. t. 3. ined.*

Calyx longus gamosepalus ovarium involvens erectus, semina angulosa. Folia opposita minima coriacea vel crassa perennantia.

1. *S. oppositifolia* (L. sp. 575.) caulibus ramosis prostratis, foliis inferioribus quadrifariam imbricatis ovatis carinatis ciliatis, caulinis cruciatim oppositis ovatis vel obovatis apice acutis unipunctatis, floribus solitariis terminalibus, calycis non adhærentis lobis obtusis ciliatis, staminibus corollâ brevioribus, seminibus oblongo-triquetris rugosis. ♀ in Alpibus, Monte-Ventoso et Pyrenæis. — Fl. fr. 4. p. 364. Lapeyr. fl. pyr. 36. t. 16. All. ped. t. 21. f. 3. (v. v.)
2. *S. retusa* (Gou. ill. 28. t. 18. f. 1.) glaucescens, caulibus ramosis prostratis, foliis inferioribus quadrifariam imbricatis triquetris crassis glabris apice retusis 3-punctatis, foliis caulinis obovatis planiusculis ciliatis subtruncatis, floribus 1-4, calycis tubo ovario adhærente lobis obtusis glabris, staminibus corollam æquantibus, seminibus lævibus nitidis. ♀ in Pyrenæis et Alp. Delphinatûs. — Fl. fr. 4. p. 365. Lapeyr. fl. pyr. t. 18. S. purpurea All. ped. t. 21. f. 2. (v. v.)
3. *S. biflora* (All. ped. t. 21. f. 1.) foliis obovatis subcoriaceis planis impunctatis

cruciatim oppositis subimbricatis nervosis ciliatis, floribus 2-5, calycis tubo ovario adhærente lobis obtusis, staminibus corollam æquantibus, seminibus obovato-triquetris apice truncatis.  $\mathcal{L}$  in Alpibus et Pyrenæis propè nives. — Fl. fr. 4. p. 365. Lapeyr. pyr. t. 18. (v. s.)

Sect. II. *AIZONIA* Tausch. hort. canal. fasc. 1. Ser. l. c.

Calyx longus gamosepalus adhærens ovarium involvens erectus, semina ovato-triquetra rugosa. — Folia alterna sessilia crassa coriacea glaucescentia cartilagineo-marginata.

§. 1. *Foliis ligulatis obtusis.*

4. *S. CORTEXLEON* (L. sp. 570.) caule pedunculis calycibusque pilis glandulosis viscidis obtectis, foliis rosularum obovato-oblongis obtusissimis cartilagineo-serratis, floribus pyramidato-paniculatis, lobis calycinis lineari-lanceolatis obtusis tubo campanulato longioribus, petalis cuneato-spathulatis obtusis, stylis subparallelis apice deflexis, stigmatibus subcapitatis.  $\mathcal{L}$  in rupibus humidis Alpium, Pyrenæorum. — *S. pyramidalis* Lapeyr. pyr. 32. Fl. fr. 4. p. 360. Stern. sax. 2. t. 2. (v. v.)
5. *S. LONGIFOLIA* (Lapeyr. fl. pyr. t. 11.) caule pedunculis calycibus pilis glandulosis viscidis obtectis, rosularum foliis spathulato-oblongis obtusis cartilagineo-crenatis, floribus paniculatis, lobis calycinis subovatis obtusis tubo hemisphærico vix longioribus, petalis obovato-spathulatis, seminibus ovato-triquetris tenuissimè rugulosis.  $\mathcal{L}$  in rupibus editiorum Pyrenæorum. — Fl. fr. 4. p. 359. (v. v.)
6. *S. AIZOON* (Jacq. austr. t. 438.) pedunculis calycibusque piloso-glandulosis viscidis, foliis rosularum ovato-spathulatis obtusis cartilagineo-serratis, floribus paniculato-corymbosis, lobis calycinis ovatis obtusis, petalis obovatis, stylis divergentibus calyce longioribus, seminibus ovato-triquetris punctato-rugosis.  $\mathcal{L}$  ad saxa et rupes montium. — Fl. fr. 4. p. 360. — Barr. ic. t. 1309, 1311, 1312. (v. v.)
7. *S. MUTATA* (L. sp. 570.) foliis rosularum elongato-lanceolatis obtusissimis basi ciliatis, caulinis spathulatis ciliato-glandulosis, floribus pyramidato-paniculatis, pedunculis calycibusque piloso-glandulosis viscosissimis, lobis calycis ovatis obtusis petalis linearibus oblongis acutis duplò brevioribus.  $\mathcal{L}$  in Pyrenæis ex Lapeyr. — Fl. fr. 5. p. 516. Sims. bot. mag. t. 351. Jacq. ic. rar. 3. t. 466. — Hall. helv. t. 16. f. 2. (v. v. in Sabaudia.)

§. 2. *Foliis ligulatis plus minùsve triquetris.*

8. *S. MEDIA* (Gou. ill. 27.) caulibus pedunculis calycibusque glanduloso-pilosis viscidis, fol. rosularum glabris glaucis ovato-spathulatis margine membranaceis submucronatis, caulinis lanceolatis glanduloso-ciliatis acutis, laciniis calycinis ovato-lanceolatis obtusis petala obovata superantibus.  $\mathcal{L}$  in Pyrenæorum rupibus calcareis. — Fl. fr. 4. p. 361. Lam. ill. t. 273. f. 6. *S. calyciflora* Lapeyr. fl. pyr. t. 12. *S. diapsenioides* Lapeyr. abr. supp. 54 non Bell. ex cl. Dufour. — Petala calycesque purpurei. (v. v.)
- $\beta$ . *ambigua*, petalis purpureis oblongis calycem superantibus. — In Pyrenæis propè St.-Beat. Hybrida inter *S. mediam* et *aretioidem*. — *S. ambigua* DC. fl. fr. 5. p. 517. (v. s.)
9. *S. ARETIOIDES* (Lapeyr. fl. pyr. t. 13.) caulibus pedunculis calycibusque glanduloso-viscidis, foliis rosularum lineari-spathulatis subtriquetris, caulinis linearibus obtusis glanduloso-ciliatis, floribus corymbosis densis, calycis laciniis ovatis obtusis petalis obovatis brevioribus.  $\mathcal{L}$  in summis Pyren. — Fl. fr. 4. p. 362. Petala lutea. (v. v.)
- $\beta$ . *luteo-purpurea*, foliis glaucis longioribus subacutis. — In Pyrenæis propè St.-Beat. — *S. luteo-purpurea* Lapeyr. fl. pyr. t. 14. Fl. fr. 4. p. 362. Petala lutea, calyx purpureus. Hybrida inter *S. aretoidem* et *mediam* monente cl. Arnott. (v. s.)

10. *S. cæsia* (L. sp. 571.) caule pedunculisque glabriusculis, foliis lineari-spathulatis plus minùve triquetris punctorum duplici serie subtùs notatis, rosularum recurvis basi ciliatis, calycis laciniis lanceolato-ovatis obtusis, petalis ovato-spathulatis.  $\mathcal{U}$  in Alpihus et Pyrenæis. — Fl. fr. 4. p. 363. Scop. carn. t. 15. Jacq. aust. t. 374. Fl. albi. (v. v.)

§. 3. *Foliis triquetris mucronatis subpungentibus.*

11. *S. VANDERLII* (Sternb. sax. 34. t. 10. f. 3. excl. syn. Jacq.) caulibus calycibusque piloso-glandulosis, foliis firmis ciliatis, radicalibus aggregatis densis glabris pungentibus, floribus corymbosis, lobis calycis ovatis acutiusculis petalis obovato-cuneatis brevioribus.  $\mathcal{U}$  in Pyrenæis propè l'*Étang d'Amour* ex Lapeyr. — *S. burseriana* Lapeyr. fl. pyr. 35. non L. (v. v. in Italià D.)

Sect. III. *DACTYLITES Tausch. l. c. Ser. l. c.*

Calyx longus gamosepalus ovario adhærens, erectus persistens. — Folia alterius plana margine nec cartilaginea nec punctata integra aut lobata.

\* *Radice perenni stolonifera.*

12. *S. SEDOIDES* (L. sp. 572.) foliis lineari-spathulatis integris obtusis inferioribus patulis, superioribus villosulis ciliatis, pedunculis piloso-glandulosis 1-4-floris, calycis pilosi sepalis lanceolato-ovatis subacutis petala lanceolata acuta superantibus.  $\mathcal{U}$  in Pyrenæis orientalibus ex Lapeyr. — Fl. fr. 5. p. 518. Jacq. misc. 2. t. 21. f. 22. *S. trichodes* Scop. carn. t. 15. Fl. lutei. (v. s. ex Alp. Tyrol.)
13. *S. PLANIFOLIA* (Lapeyr. fl. pyr. p. 31.) cespitosa, foliis lineari-oblongis obtusissimis planis integris piloso-glandulosis, inferioribus confertis imbricatis, caulibus 1-3-floris piloso-glandulosis, calycis laciniis lanceolatis obtusis petalis obovatis submarginatis duplò brevioribus.  $\mathcal{U}$  ad rupes humidæ editiorum Alpium et Pyren. — Fl. fr. 4. p. 367. *S. muscoides* All. ped. t. 61. f. 2. excl. syn. Folia antiquiora cinerascens. Fl. ochroleuci. (v. v.)
14. *S. ANDROSACHA* (L. sp. 571.) foliis obtusis, radicalibus obovato-spathulatis integris rariùs apice 3-dentatis in petiolum basi subattenuatis ciliatis, caulibus 1-3 lanceolato-ovatis lanceolatisve, caulibus 1-4-floris, calycis laciniis ovatis obtusissimis petalis obovatis obtusis brevioribus.  $\mathcal{U}$  in Alpihus et Pyrenæis. — Fl. fr. 4. p. 367. *S. Pyrenaica* Scop. carn. t. 16. — Pluck. t. 222. f. 2. Fl. albescentes. (v. v.)
15. *S. MUSCOIDES* (Wulf. in Jacq. misc. 2. p. 125.) cespitosa plus minùve glanduloso-viscosa, caule erecto 1-8-floro, foliis linearibus obtusissimis integris aut apice 3-lobis, inferioribus confertis ciliatis, laciniis calycis lanceolatis obtusissimis petalis obtusis elliptico-linearibus subbrevioribus.  $\mathcal{U}$  in rupibus Alpium, Montis-Aurei, Pyrenæorum. — Fl. fr. 4. p. 376. *S. cespitosa* Scop. carn. t. 14. *S. moschata* Jacq. misc. 2. t. 21. f. 1. Lapeyr. fl. pyr. t. 37. Fl. fr. 5. p. 520. — Seg. ver. 1. t. 9. f. 4. Variat petalis flavescentibus aut purpureis. (v. v.)
- β. elatior* (Ser.) caule elongato majore, foliis longius petiolatis, laciniis calycis ovatis petalis brevioribus. — In Jura (Ph. Thomas) et Pyrenæis. — *S. cespitosa* Lapeyr. fl. pyr. t. 35. (v. s.)
16. *S. EXARATA* (Vill. damp. 4. t. 45. non All.) plus minùve piloso-viscidula, foliis nervosis obtusis cuneatis 3-5-fidis, caulibus rariùs integris, radicalibus petiolatis, antiquioribus reflexis, calycis laciniis ovato-lanceolatis obtusis petalis ovatis obtusis nervosis plus minùve brevioribus.  $\mathcal{U}$  in rupibus editiorum Alpium et Pyrenæorum. — Fl. fr. 4. p. 374. *S. hypnoides* All. ped. t. 21. f. 4. non L. *S. intricata* Lapeyr. fl. pyr. t. 33. Fl. fr. 4. p. 374. *S. nervosa* Lapeyr. pyr. t. 39. (v. v.)
17. *S. GROENLANDICA* (L. sp. 578.) viscosa cespitosa, foliis pubescenti-viscosis cuneatis apice 3-lobis segmentis obtusis, radicalibus densissimè imbricatis, caulibus 1-2-phyllis 2-5-floris floribus subcapitatis, laciniis calycis ovatis

- obtusis petalis obovatis obtusis multò brevioribus, staminum filamentis persistentibus demum purpureis.  $\frac{1}{2}$  in editioribus Pyrenæis. — Fl. fr. 4. p. 376. Lapeyr. fl. pyr. t. 19. Fl. albi, calyces purpurei. (v. v. DC.)
18. *S. PUBESCENS* (DC. fl. fr. 4. p. 375.) pubescenti-viscida, foliis cuneatis apice 3-5-lobis segmentis linearibus obtusis divaricatis, radicalibus petiolatis, antiquioribus reflexis, caulibus oligophyllis, floribus subcorymbosis, laciniis calycis lanceolatis obtusis petalis rotundato-ovatis multò brevioribus.  $\frac{1}{2}$  in rupibus Alpium Galloprovinciæ, Montis-Ventosi, Cebenn., Pyren. — *S. mixta* Lapeyr. fl. pyr. t. 20 et 21. (v. v.)
19. *S. AJUGÆFOLIA* (L. sp. 578.) surculis prostratis foliosis, foliis glabris cuneato-petiolatis 3-5-fidis, segmentis lanceolatis acutis, pedunculis axillaribus ascendentibus paucifloris oligophyllis, foliis superioribus linearibus indivisis, calycis laciniis lanceolatis subacutis petalis ovato-ellipticis obtusis duplò brevioribus.  $\frac{1}{2}$  in saxosis humidis secus nives in Pyrenæis. — Fl. fr. 4. p. 571 excl.; var.  $\beta$ . Lapeyr. fl. pyr. t. 31. (v. v.)
20. *S. CAPITATA* (Lapeyr. fl. pyr. t. 30.) surculis basi nudis procumbentibus dein erectis foliosis, foliis subcapitatis coriaceis cuneatis basi ciliatis 3-5-fidis, laciniis latis cuneatis 3-lobis, segmentis ovatis acutis obtusisve, pedunculis axillaribus erectis, foliis superioribus lineari-lanceolatis acutis, laciniis calycis ovato-lanceolatis acutis petalis rotundato-ovatis duplò brevioribus.  $\frac{1}{2}$  in editioribus Pyrenæis. — *S. ajugæfolia*  $\beta$ . Fl. fr. 4. p. 371. (v. v.)
21. *S. ASCENDENS* (L. sp. 579.) caule ascendente apice pubescente ramoso-paniculato, foliis crassis basi cuneatis palmato-3-5-partitis, lobis sub 3-fidis, segmentis inciso-dentatis acutis, floribus paniculatis, calycis glanduloso-pubescentia laciniis lanceolatis acutis petalis obovatis multò brevioribus.  $\frac{1}{2}$  secus rivulos Pyrenæorum et montium Corsicæ. — Fl. fr. 4. p. 370. *S. aquatica* Lapeyr. fl. pyr. t. 28 et 29. Fl. magni albi. (v. v.)
22. *S. GERANIODES* (L. sp. 578.) viscida, caulibus erectis subnudis, foliis radicalibus reniformibus longè petiolatis sub 5-palmatis laciniis 3-lobis lobis lanceolatis ovatisve acutis, caulinis subsessilibus 5-fidis laciniis lineari-lanceolatis, superioribus linearibus, floribus paniculato-corymbosis, calycis 5-fidi laciniis linearibus lanceolatisve acutis petalis oblongo-ovatis triplò brevioribus.  $\frac{1}{2}$  ad rupes umbrosas Pyrenæorum. — Fl. fr. 4. p. 372. Gou. ill. t. 18. f. 2. Lapeyr. fl. pyr. t. 43. (v. v.)
23. *S. LADANIFERA* (Lapeyr. fl. pyr. t. 42.) viscida, foliis radicalibus gemmiferis longè petiolatis 5-7-palmato-partitis laciniis integerrimis linearibus, caulinis subpetiolatis 3-fidis, summis linearibus, floribus paniculatis, calycis 3-fidi laciniis lineari-lanceolatis acutis petalis ovato-spathulatis duplò triplòve brevioribus.  $\frac{1}{2}$  ad rupes Pyren. orientaliū. — Fl. fr. 4. p. 373. *S. pedatifida*  $\beta$  ladanifera Don in trans. linn. 13. p. 414. (v. v.)
- $\beta$ . *pedatifida*, calycis laciniis linearibus angustis petalis ovato-spathulatis longè unguiculatis brevioribus. — In editioribus montibus agri Ligericini (Prost), Corsicæ. — *S. pedatifida* Sm. eng. bot. t. 1276. *S. cervicornis* Viv. cors. app. p. 2. (v. s.)
24. *S. PENTADACTYLIS* (Lapeyr. fl. pyr. t. 44.) glabra, surculis basi suffrutescentibus erectis coronatis foliis congestis rigidis plus minùsve petiolatis 3-5-palmatis laciniis linearibus obtusis, antiquioribus gemmiferis, floribus corymboso-paniculatis aut subcapitatis, laciniis calycinis lanceolatis acutis petalis ovato-ellipticis duplò triplòve brevioribus.  $\frac{1}{2}$  ad rupes in Pyren. orient. — Fl. fr. 4. p. 374. An satis à *S. ladanifera*  $\alpha$  distincta? (v. v.)
25. *S. HYFNOIDES* (L. sp. 579.) caulibus erectis subnudis paucifloris, ramis sterilibus abortu sæpè gemmiferis, foliis flaccidis ciliatis integris vel 3-7-sectis, laciniis linearibus acuminatis, calycis laciniis lineari-lanceolatis acutis petalis oblongo-obovatis duplò triplòve brevioribus.  $\frac{1}{2}$  in montibus calcareis Galliæ australis, Cebenn., Arvernæ. — Fl. fr. 4. p. 376. Vill. dauph. 4. t. 45. Planta valdè polymorpha. (v. v.)

\*\* *Annua astolonosa.*

26. *S. PETREA* (L. sp. 578. excl. syn.) caulibus erectis aut ascendentibus ramosis foliosis, foliis cuneiformibus 3-5-inciso-dentatis aut integris spathulatis lanceolatisve villosis-viscidis, inferioribus confertis, lacinii calycis ovato-lanceolatis obtusis petalis obovatis duplò brevioribus. ① in rupestribus Alpium Delph. et Pyrenæorum. — Fl. fr. 4. p. 370. S. ascendens All. ped. t. 22. f. 3. non Gouan. S. controversa Sternb. sax. 43. t. 16. f. 1. — 5. (v. v.)
27. *S. TRIDACTYLITES* (L. sp. 578.) caule erecto paucifloro glutinoso, foliis crassiusculis inferioribus integris obovato-spathulatis petiolatis, mediis 3-fidis alternis lacinii ovatis obtusis, summis linearibus, cal. lacinii brevibus ovatis obtusis petalis obovatis brevioribus. ① in muris et arenosis. — Fl. fr. 4. p. 369. — Black. t. 212. (v. v.)

\*\*\* *Perennes astolonosæ radice bulbifera.*

28. *S. GRANULATA* (L. sp. 576.) pubescenti-viscosa, caule erecto stricto, foliis reniformibus, inferioribus longè petiolatis inciso-crenatis, superioribus sessilibus lobatis, floribus paniculato-corymbosis, calycis lacinii oblongo-lanceolatis obtusis petalis obovato-oblongis triplò brevioribus. 2/ in pratis et ad oras sylvarum passim. — Fl. fr. 4. p. 368. — Cam. epit. 719. ic. (v. v.)
- β. *Corsica* (Ser. ined.) pygmaea, caule 1-4-floro, foliis radicalibus longè petiolatis cordatis 5-7-lobis lobis rotundatis obtusis, caulinis 3-fidis. — In Corsicâ. (Ph. Thomas.)
- γ. *penduliflora* (Ser.) foliis omnibus petiolatis, inferioribus profundè cordatis 7-9-lobis lobis rotundis obtusis, floribus subumbellatis cernuis, calycis lacinii ovato-lanceolatis. — In Monte-Aureo. S. penduliflora Bast. in Journ. bot. 1814. p. 17. Fl. fr. 5. p. 519. (v. s.)

Sect. IV. *ARABIDIA Tausch. l. c. Ser. l. c.*

Calyx basi vix gamosepalus ovario non adhærens post anthesin reflexus, petala longè unguiculata bimaculata, semina oblonga striata, folia alterna cuneiformia vel obovata in stolonibus rosulata.

29. *S. CLUSII* (Gou. ill. p. 28. excl. plur. syn.) caulibus erectis ramosis foliosis, foliis radicalibus spathulato-obovatis flaccidis basi attenuatis pilosis dentatis obtusis, caulinis cuneatis spice dentatis, superioribus linearibus, floribus paniculatis, pedicellis post anthesin divaricatis, sepalis ovatis obtusis petalis lanceolatis basi unguiculatis subinæqualibus brevioribus. 2/ in humidis umbrosis Pyren. et Cebenn. — Fl. fr. 4. p. 380. S. leucanthemifolia Lapeyr. fl. pyr. t. 25. (v. v.)
30. *S. STELLARIS* (L. sp. 572.) caulibus erectis subsimplicibus nudis, foliis rosulatis obovato-spathulatis crassiusculis dentato-serratis obtusis basi in petiolum subattenuatis glabrisculis, floribus paniculato-corymbosis, pedicellis longis, sepalis ovatis obtusis petalis lineari-lanceolatis basi unguiculatis duplò brevioribus. 2/ in humidis montium. — Fl. fr. 4. p. 379. — Pluck. t. 58. f. 2. (v. v.)

Sect. V. *HYDATICA Neck. el. bot. 2. p. 387. Tausch. l. c. Ser. l. c.*

Calyx basi vix gamosepalus ovario non adhærens, semina subæpærica aspera. — Folia cuneata orbiculatave.

\* *Calyce post anthesin reflexo.*

31. *S. CUNEIFOLIA* (L. sp. 574.) foliis rosulatis coriaceis cuneato-spathulatis obtusissimis dentato-crenatis glabris basi in petiolum attenuatis, scapo erecto nudo, floribus laxè paniculatis, calycis sepalis lanceolatis acutis petalis oblongo-lanceolatis obtusis brevioribus. 2/ in umbrosis humidis Alp., Cebenn., Pyren. — Fl. fr. 4. p. 377. Scop. carn. t. 13. — J. Bauh. hist. 3. p. 684. f. 2. (v. v.)
32. *S. UMBROSA* (L. sp. 574.) foliis rosulatis rotundato-spathulatis obovatisve

subretusis cartilagineo-crenatis in petiolum dilatatum ciliatum subattenuatis, scapo erecto stricto nudo viscido-glanduloso, floribus paniculatis, sepalis obovatis obtusis petalis ovato-oblongis duplò brevioribus.  $\mathcal{L}$  in Pyrenæis. — Fl. fr. 4. p. 378. Lapeyr. fl. pyr. t. 22. (v. v.)

33. *S. hirsuta* (L. sp. 574.) foliis rosulatis ovalibus basi subcordatis cartilagineo-crenatis retusis subglabris longè petiolatis, petiolis hirsutis, floribus paniculatis, sepalis ovato-lanceolatis obtusis petalis oblongo-ovatis brevioribus.  $\mathcal{L}$  in Pyrenæis. — Fl. fr. 4. p. 378. Lap. fl. pyr. t. 23. — Magn. hort. 87. ic. (v. s.)

$\beta$ . *Geum*, foliis dentatis utrinquè plus minùs hirsutis. — *S. Geum* L. sp. 574. Lapeyr. fl. pyr. t. 24. Fl. fr. 4. p. 379. (v. s.)

\*\* *Calyce post anthesin non reflexo.*

34. *S. rotundifolia* (L. sp. 576.) caulibus erectis foliosis, foliis reniformi-cordatis inæqualibus dentatis crenatisve, inferioribus longè petiolatis petiolis hirsutis, superioribus sessilibus, floribus paniculatis, sepalis lineari-lanceolatis acutis petalis lineari-ellipticis duplò brevioribus.  $\mathcal{L}$  in umbrosis montium. — Fl. fr. 4. p. 368. — Cam. epit. 764. ic. (v. v.)

#### Sect. VI. *HIACULUS* Clus. *Tausch. l. c.*

Calyx basi vix gamosepalus ovario non adhærens; sem. oblonga. — Caulis foliosus, folia alterna linearia aut lanceolata integra.

35. *S. Hiaculus* (L. sp. 576.) caule simplici erecto folioso, foliis obtusis, radicalibus spathulato-lanceolatis basi in petiolum ciliatum angustatis, caulinis oblongo-lanceolatis, sepalis oblongo-ovatis obtusissimis petalis ovatis obtusis duplò brevioribus.  $\mathcal{L}$  in paludosis turfosissimis Jurassi. — Fl. fr. 4. p. 377. — Hall. helv. t. 11. Fl. lutei. (v. v. DC.)

36. *S. aspera* (L. sp. 575.) caulibus foliosis basi geniculatis procumbentibus, ramulis sæpè abortivis gemmiformibus, foliis lineari-lanceolatis mucronatis lævibus margine sæpius ciliis asperis instructis, floribus corymboso-paniculatis, sepalis ovatis obtusis petalis obovato-oblongis duplò brevioribus.  $\mathcal{L}$  in Alpibus, montibus Arvernæ, Pyren. — Fl. fr. 4. p. 363. *S. bryoides* L. sp. 572. — Scheuchz. itin. 2. t. 20. f. 3 et t. 21. f. 2. (v. v.)

37. *S. aizoides* (L. sp. 576.) caulibus basi decumbentibus foliosis, foliis linearibus basi plus minùsve ciliatis crassiusculis submucronatis, floribus laxè paniculatis solitariisve, sepalis ovatis mucronatis petalis oblongo-ellipticis obtusis brevioribus.  $\mathcal{L}$  in umbrosis et humidis secùs rivulos montium. — Fl. fr. 4. p. 366. *S. autumnalis* L. sp. 575. — Fl. lutei sæpè croceò maculati. (v. v.)

### ++ COROLLA NULLA.

#### II. *CHRYSOSPLENIUM* Tourn. inst. t. 60. L. gen. 558.

Calyx ovario adhærens 4-5-sepalus sepalis apice liberis; petala nulla; stam. 8-10; caps. 1-locularis 2-valvis.

1. *C. alternifolium* (L. sp. 569.) foliis alternis reniformi-cordatis subduplicato-crenatis.  $\mathcal{L}$  in humidis umbrosis imprimis montium. — Fl. fr. 4. p. 381. — Dalech. lugd. 1114. f. 1. (v. v.)
2. *C. oppositifolium* (L. sp. 569.) foliis oppositis reniformibus subcuneatis irregulariter crenatis.  $\mathcal{L}$  in humidis montanis. — Fl. fr. 4. p. 381. — Dalech. lugd. 1114. f. 2. (v. v.)

#### III. *ADOXA* L. gen. 501.

Calyx 4-5-sepalus sepalis inter se et cum ovario adnatis extùs 2-4-squamosus; petala nulla; stam. 8-10; styli 4-5; bacca globosa 4-5-locularis; semina margine membranacea, radícula supera.

1. *A. Moschatellina* (L. sp. 527.) foliis biternatisectis glaucescentibus, floribus capitatis sessilibus.  $\mathcal{L}$  in sepibus, vere. — Fl. fr. 4. p. 382. — Lob. ic. 674. f. 2. (v. v.)



Ordo L. UMBELLIFERÆ. *Juss. gen.* 218. *Koch Umbellif. nov. disp. in novis actis nat. cur. t. 12. pars 1. p. 55.*

Calyx ovario adhærens, limbo 5-dentato vel edentulo. Petala 5 ovario aut glandulæ ovarium tegenti inserta, nunc integra in acumen sensim attenuata aut in circulum involuta apice brevi lato retuso aut in lacinulam inflexam subito coarctata, nunc emarginata bifidave. Stamina 5 petalis alterna et cum iis inserta. Styli 2 sæpius persistentes. Fructus constans 2 carpellis superficie plus minus latâ conglutinatâ et extus calyce adhærente obductis, maturitate à basi ad apicem segregatis, et ex apice axis filiformis *carpophori* dicti suspensis. Carpello cuique superficies duo, altera convexior *dorsalis*, altera quâ gemina sibi incumbunt carpella planior *commissuralis*. Superficies convexior striis quinque nunc elevatis nunc depressis nunc tenuibus *juga* dictis notatur. Horum jugorum unum medium dorsi tenens *carinale*, unum utrinque marginem occupans vel non longè ab illo remotum *juga lateralia*, unum utrinque inter carinam et marginem situm *juga intermedia*, et hæc 5 *primaria* vocantur cum semper saltem leniter indicata adsint. Inter juga primaria non raro quatuor alia *secundaria* in cujusvis *valleculæ* ( id est spatii inter 2 juga interpositi plerumque depressi ) existunt, quorum duo valleculas interiores *juga secundaria interiora*, duo valleculas exteriores *juga secundaria exteriora* occupant. Cujusque carpelli pericarpium sæpissimè *vittis* id est canalibus verticalibus rectis filiformibus vel clavulatis frequenter discoloribus oleum vel resinam continentibus gaudet; carpellum *paucivittatum* si vittæ solitariæ in medio vallecularum vel et binæ in valleculis exterioribus adsunt, si majori sunt vittæ numero *carpellum multivittatum*, si verò nullæ *evittatum*, dicitur. Carpella in quibus seminis superficies commissuralis plana est *orthosperma*, in quibus verò seminis superficies commissuralis plus minusve inflectitur *campylosperma* dicuntur. Embryo minimus ad apicem albuminis lignosi; radícula supera. — Herbæ foliis alternis, sæpius decompositis, floribus umbellatis aut rarissimè capitatis, involucro et involucello sæpius cinctis.

Subordo I. UMBELLIFERÆ PERFECTÆ MULTIJUGATÆ, id est carpello jugis 5 primariis et 4 secundariis instructo.

Trib. I. *THAPSIEÆ*. *Koch umb. p. 73.*

Carpella multijugata, jugis primariis 5 filiformibus, secundariis

4, interioribus filiformibus alatisve, exterioribus alatis alâ inermi. Semen (albumen) anticè planiusculum et idèò carpellum orthospermum.

† *Fructus octo-alatus, jugis secundariis in alam insignem protractis.*

I. LASERPITIUM L. gen. 344.

Calycis margo quinquedentatus; petala obovata emarginata cum lacinulâ inflexâ; fructus a dorso compressus vel sectione transversali subteres; carpella jugis primariis filiformibus, 2 lateralibus plano commissurali impositis, secundariis 4 alatis alâ integrâ; valleculæ sub jugis secundariis 1-vittatæ; semen anticè planiusculum. — Involucrum et involucellum polyphyllum.

\* *Fructu maturo glabro vel in jugis primariis pilis brevissimis adpressis obsito.*

1. L. SILEX (L. sp. 357.) foliis bipinnatis, foliolis lanceolatis integerrimis mucronatis simplicibus ternatisque, involucri foliolis lineari-lanceolatis aristato-acuminatis glabris, æminum alis brevibus. ½ in montibus provinc. australium. — Fl. fr. 4. p. 313. Jacq. austr. t. 145. (v. v.)
2. L. AQUILEGIVOLVUM (Murr. syst. 228.) foliis bis trifidis, foliolis ovatis basi subcordatis glabris subtus cinereis sæpè trilobis apice dentatis dentibus mucronatis, jugorum alis subæqualibus. ½ in umbrosis humidis Pyren. et Cebennarum. — Fl. fr. 5. p. 510. L. Alpium Wald. et Kit. hung. t. 253. (v. v.)
3. L. ASPÆNUM (Crantz austr. 8°. p. 50. t. 1. f. 2. et n. 6.) foliis bipinnatis tripinnatisve, foliolis ovali-oblongis obliquè cordatis subtus pubescentibus dentatis, dentibus mucronatis, involucri foliolis lineari-setaceis, seminum alis crispis. ½ in montibus. — Fl. fr. 5. p. 509. L. latifolium Fl. fr. 4. p. 312. — Lob. ic. 704. f. 2. (v. v.)
4. L. GLABRUM (Crantz austr. 3. p. 54.) foliis bipinnatis glaberrimis vaginis amplis, inferiorum foliolis rotundo-ovatis obliquè cordatis dentatis dentibus mucronatis, superiorum foliolis lanceolatis integerrimis, involucri foliolis setaceis glaberrimis, alis magnis subæqualibus crispis. ½ in locis montosis siccis Alsatiz, Jurass, Massiliæ, Pyrenæorum ex Lapeyr. — Fl. fr. 5. p. 509. L. latifolium L. sp. 356. — Moris. s. 9. t. 19. n. 1. (v. v.)
5. L. GALLICUM (L. sp. 357.) foliis 3-pinnatis, foliolis cuneatis plus minùs 3-5-fidis glabris, laciniiis oblongis apice calloso-mucronatis, involucri foliolis lineari-lanceolatis glabris acuminatis, alis subæqualibus. ½ in montibus aridis provinc. australium orientaliumque. — Fl. fr. 4. p. 312. — Pluck. t. 198. f. 6. (v. v.)
6. L. HIRsutum (Lam. fl. fr. 3. p. 648.) foliis supra-decompositis hirsutis, involucri foliolis lineari-lanceolatis aristato-acuminatis margine scarioso ciliatis. ½ in Alpibus Delphin. et Galloprovinciæ. — Fl. fr. 4. p. 313. L. Halleri All. — Hall. helv. t. 19. (v. v. in Helv.)

\*\* *Fructus maturi jugis primariis setulis patentibus obsitis.*

7. L. PRUTHENICUM (L. sp. 357.) foliis 3-pinnatis, foliolis lanceolatis integerrimis acutis extimis coalitis, involucri foliolis lineari-lanceolatis aristato-acuminatis, jugorum exteriorum alis majoribus. ½ in pascuis et dametis Galliz boreali-orientalis. — Fl. fr. 4. p. 312. — Brey. cent. t. 48. (v. v.)

†† *Fructus 4-alatus, jugis secundariis exterioribus tantum alatis.*

II. THAPSIA Tourn. inst. t. 171. L. gen. n. 361.

Calyx 5-dentatus; petala elliptica integra acuminata acumine inflexo vel involuto; fructus a dorso compressus; carpella jugis primariis 5 filiformibus, 2 lateralibus plano commissurali impositis, secundariis 4.

interioribus filiformibus tenuissimis, exterioribus membranaceo-alatis alâ integrâ; valleculæ sub jugis secundariis univittatæ; semen complanatum. — Invol. universale et parziale oligophyllum caducum, vittæ commissurales duæ.

1. *T. villosa* (L. sp. 575.) foliis ternato-decompositis, foliolis dentatis villosis basi cœdunatis. ½ in locis sterilibus umbrosis provinc. mediterranearum. — Fl. fr. 4. p. 342. Koch umb. f. 3. 4 et 5. — Clus. hist. 2. p. 192. ic. (v. v. DC.)

Trib. II. *DAUCINÆ*. Koch l. c. p. 76.

Fructus a dorso lenticulari-compressus aut sectione transversali subteres. Carpella multijugata, jugis primariis 5 filiformibus setulosis, secundariis 4 magis prominulis aculeatis. Semen anticè planiusculum.

III. *DAUCUS* L. gen. n. 333.

Calyx 5-dentatus; petala obovata emarginata cum lacinulâ inflexâ, exteriora sæpè radiantia et profundè bifida; fructus a dorso compressus; carpella jugis lateralibus plano commissurali impositis, secundariis 4 æqualibus magis prominulis unifariam aculeatis, aculeis basi tenuissimè connexis; valleculæ sub jugis secundariis 1-vittatæ. — Invol. universale et parziale polyphyllum; vittæ commissurales duæ.

1. *D. MAURITANICUS* (All. ped. t. 61. f. 1.) foliis multipinnatis glaberrimis, superiorum foliolis acuminatis linearibus trifidis glabris, inferiorum foliolis lanceolatis inciso-dentatis, involucri foliolis lineari-pinnatifidis striatis, aculeis diametrum fructûs oblongo-ovalis subsuperantibus. ② circa Nicæam et in agro Ruscinonensi propè Prades. — Fl. fr. 5. p. 512 an L. sp. 348? (v. v. DC.)
2. *D. MARITIMUS* (Lam. dict. 2. p. 634.) caule elongato lævi basi glabro superne scabrinusculo, foliis glabris, inferioribus 2-pinnatis foliolis laciniaüs laciniiis linearibus acuminatis, superioribus pinnatis foliolis linearibus acuminatis, involucri foliolis glabris lineari-pinnatifidis striatis, aculeis fructûs ovati diametrum subæquantibus. ① in arenosis littoribus Occitanie, Narbonæ et Vindilis (*Belle-Île en mer*). — Fl. fr. 4. p. 329. (v. v.)
3. *D. GUMMIFERA* (Lam. dict. 1. p. 634.) caule erecto basi hispido ramoso striato, foliis bipinnatis, foliolis glabrinusculis ovatis dentatis incisive crassis obtusis mucronulatis, involucri foliolis striatis ciliatis pinnatifidis, laciniiis lanceolato-linearibus acuminatis, aculeis fructûs ovati diametro minoribus. ½ in littoribus saxosis maris Mediterranei. — Fl. fr. 4. p. 328. et 5. p. 512. D. Gingidium L. sp. 348? D. lucidus L. f. supp. 179. (v. v.)
4. *D. CAROTA* (L. sp. 348.) caule hispido, foliolis 2-3-pinnatis villosis, foliolis incisè lacinia linearibus acutis, involucri foliolis lineari-pinnatifidis lævibus ciliatis, aculeis fructûs oblongo-ovalis diametro minoribus. ② in pascuis arvisque. — Fl. fr. 4. p. 327. Koch umb. f. 11. — Lob. ic. 723. f. 1. — J. Baub. 3. p. 2. p. 64. (v. v.)
5. *D. HERNANDUS* (Desf. atl. 1. t. 63.) caule hispido ramoso, foliis inferioribus hispidis 3-pinnatis foliolis ovatis incisè dentatisque, superioribus glabrinusculis 2-pinnatis, foliolis 3-5-fidis, lacinia linearibus lanceolatis acuminatis, involucri foliolis lineari-pinnatifidis ciliato-scabris basi scarioso-dilatatis, aculeis fructûs ovati diametrum subsuperantibus. ② in saxis maritimis propè Dieppe, Cherbourg et St.-Malo ex cl. de Brébisson. — Fl. fr. 4. p. 328. (v. v. DC.)

IV. *ORLAYA* Hoffm. umb. ed. 2. t. 1. p. 58. — *Caucalidis* sp. L.

Calyx 5-dentatus; petala obovata emarginata cum lacinulâ inflexâ,

exteriora radiantia profundè bifida; fructus a dorso lenticulari-compressus; carpella jugis lateralibus plano commissurali impositis, secundariis 4 bi-trifariam aculeatis, exterioribus magis prominulis; vallecule sub jugis secundariis univittatæ; semen anticè concaviusculum. — Invol. univers. varium, parziale polyphyllum.

1. *O. GRANDIFLORA* (Hoff. l. c.) foliis bipinnatis, involucri involucellorumque foliolis integerrimis lanceolatis acuminatis margine scariosis, petalis exterioribus maximis, jugis secundariis interioribus 3-fariam, exterioribus 2-fariam aculeatis. ① in arvis. — Koch umb. f. 13. *Cnalcis grandiflora* L. sp. 346. Lam. ill. t. 192. f. 1. Fl. fr. 4. p. 330. *Aculei hamati*. (v. v.)
2. *O. MARITIMA* (Koch umb. p. 79.) hispido-hirsuta, foliis 3-pinnatifidis incisis, involucri involucellisque oligophyllis, foliolis linearibus acuminatis indivisis 3-fidisque, jugis secundariis interioribus bifariam aculeatis, exterioribus brevioribus satis alâ bifariam inæqualiter aculeatâ. ① in arenosis maritimis Corsicæ, Nicæ, Galloprovinc., Occitaniz. — *Cnalcis maritima* Gou. hort. 135. Fl. fr. 4. p. 331. — Ger. gallop. t. 10. *Aculei apice stellatum hamati*. (v. v.)
3. *O. PLATYCARPOS* (Koch umb. l. c. f. 12.) caule sparsim piloso, foliis 3-pinnatifidis incisis, involucri involucellisque oligophyllis, foliolis lanceolatis margine scariosis indivisis aut apice pinnatis, jugis secundariis bifariam aculeatis, interioribus exteriores satis alâ bifariam inæqualiter aculeatâ æquantibus. ① in provinciis australioribus. — *Cnalcis platycarpus* L. sp. 347. Fl. fr. p. 331. — J. Bauh. p. 2. p. 80. f. 1. *Aculei purpurascens apice hamati, majores quam in præcedente*. (v. v.)

### Trib. III. CAUCALINEÆ. Koch umb. 79.

Fructus a latere contractus vel subteres. Carpella multijugata, jugis primariis 5 filiformibus setosis vel aculeatis, secundariis 4 magis prominulis aculeatis vel oblitteratis copiâ aculeorum totam valleculam obtegentium. Semen involutum vel margine inflexum et idcirco carpella campylosperma.

#### V. CAUCALIS Hoffm. umb. ed. 2. 1. p. 55. — *Cnalcidis* sp. L.

Calyx 5-dentatus; petala obovata emarginata cum lacinulâ inflexâ, exteriora radiantia profundè bifida; fructus a latere subcompressus; carpella jugis lateralibus plano commissurali impositis, secundariis 4 magis prominulis uni-bifariam aculeatis; vallecule sub jugis secundariis 1-vittatæ; semen involutum vel margine inflexum. — Invol. universale et parziale varium; vittæ commissurales 2.

1. *C. DAUCOIDES* (L. sp. 346.) caule erecto ramoso glabresculo, foliis 3-pinnatifidis incisis, involucri involucellisque subnullis aut foliolis 1-3 subulatis minimis caducis, jugis primariis aculeatis, secundariis unifariam aculeatis, aculeis subæqualibus, semine involuto. ① in incultis et limitibus arborum. — Fl. fr. 4. p. 332. Koch f. 14. Umbellulæ 1-3-floræ. (v. v.)
2. *C. LEPTOPHYLLA* (L. sp. 347.) foliis bipinnatifidis, foliolis integris incisive, involucri nullo, involucellis polyphyllis, foliolis hispidis lanceolato-linearibus acuminatis, jugis primariis setosis, secundariis bifariam aculeatis aculeis æqualibus, semine anticè profundo sulco exarato sed non involuto. ① in arvis, locis aridis. — *C. parviflora* Lam. dict. 1. p. 657. DC. fl. fr. 4. p. 332. — Jacq. vind. 2. t. 195. Umbellæ 1-3-radiatæ. (v. v.)

#### VI. TURGENIA Hoffm. umb. ed. 2. 1. p. 59. — *Tordylii* sp. L.

Calyx 5-dentatus; petala obovata emarginata cum lacinulâ inflexâ, exteriora radiantia bifida; fructus a latere contractus; carpella jugis

primariis lateralibus plano commissurali impositis muriculatis, reliquis omnibus bi-trisariam aculeatis, aculeis æqualibus; vallecule sub jugis secundariis univittatæ; semen involutum. — Involucrum univ. et partiale è foliolis pluribus.

1. *T. LATIFOLIA* (Hoffm. l. c.) foliis pinnatis, foliolis lanceolatis inciso-dentatis decurrentibus, involucri involucrellisque polyphyllis, foliolis ovato-lanceolatis scariosis acuminatis. ① in arvis inter segetes. — Koch f. 16. *Caucalis latifolia* L. syst. 205. Fl. fr. 4. p. 330. — Garid. aix. t. 22. (v. v.)

VII. *TORILIS* *Adans. Hoffm. gen. umb. ed. 2. 1. p. 53.* — *Tordylii* et *Scandicis* sp. L.

Calyx 5-dentatus; petala obovata emarginata cum lacinulâ inflexâ, exteriora majora bifida; fructus a latere contractus; carpella jugis primariis setulosis, lateralibus plano commissurali impositis, secundariis copiâ aculeorum totam valleculam occupantium oblitteratis; vallecule sub aculeis univittatæ; semen margine inflexum. — Involucrum varium, involucrellum polyphyllum.

1. *T. NODOSA* (Gært. fruct. t. 20. f. 6.) foliis tenuiter 2-3-pinnatifidis, umbellis lateralibus breviter pedicellatis, involucri involucrellorumque foliolis linearisetaceis hispidis, jugorum secundariorum aculeis caducis. ① in locis aridis, ad margines arborum. — *Caucalis nodiflora* Fl. fr. 4. p. 334. *C. nodosa* Huds. non All. et DC. — J. Bauh. 3. p. 2. p. 83. f. 2. Fructus aculeorum caducorum basi permanente tuberculatus. (v. v.)

2. *T. ANTHRISCUS* (Gmel. bad. 1. p. 613.) caule ramoso elato, foliis pinnatis foliolis pinnatifidis lacinii incisâ dentatisque decurrentibus, involucri involucrellisque foliolis 4-5 linearisubulatis hispidis, umbellis longè pedunculatis, fructus aculeis incurvis persistentibus. ① in locis incultis et ad sepes. — Koch umb. f. 15. *Caucalis Anthriscus* Fl. fr. 4. p. 333. — Bauh. prod. p. 80. ic. (v. v.)

3. *T. INFESTA* (Hoffm. l. c.) caule ramosissimo diffuso, foliis pinnatis foliolis pinnatifidis lacinii incisâ, involucri subnullo, involucrellis foliolis 3-5 linearisubulatis hispidis, umbellis longè pedunculatis, carpellis aculeatis aculeis breviter hamatis persistentibus. ① in arvis et ad vias. — *Cauc. arvensis* Fl. fr. 4. p. 333. *Scapdix infesta* L. sp. 237. *T. Helvetica* Röm. et Sch. — Jacq. vind. 3. p. 12. t. 16. An satis à præcedente distincta? (v. v.)

Trib. IV. *CORIANDRÆ*. Koch umb. p. 82.

Fructus globosus, vel è carpellis duobus subglobosis didymus. Carpella multijugata, jugis primariis 5 depressis flexuosisque vel sulcum obsoletum tantum formantibus, lateralibus antè marginem (id est locum ubi binæ carpellorum superficies sese contingunt) accessorium (id est à jugis lateralibus dictinctum) positum; secundariis 4 magis prominulis, omnibus apteris. Semen a basi ad apicem involutum vel curvatum, hinc anticè excavatum.

VIII. *CORIANDRUM* *Hoffm. umb. gen. ed. 2. 1. p. 189.* — *Coriandri* sp. L.

Calyx 5-dentatus; petala obovata emarginata cum lacinulâ inflexâ, exteriora radiantia bifida; fructus globosus; carpella jugis primariis 5 depressis flexuosis, secundariis 4 magis prominulis carinatis; vallecule evittatæ, commissura 2-vittata; semen anticè excavatum membranâ solutâ tectum. — Invol. nullum, involucrellum dimidiatum; carpella cohærentia difficilè separanda.

1. *C. SATIVUM* (L. sp. 367.) ① in arvis. — Koch f. 72 et 73. Fl. fr. 4. p. 292. — Cam. epit. 523. ic. Caulis teres, folia decomposita, foliolis linearibus. (v. v.)

IX. BIFORA *Hoffm. umb. ed. 2. 1. p. 191.*

*Calycis* margo obsoletus; *petala* obovata emarginata cum lacinulâ inflexâ; *fructus* carpellis subgloboso-ventricosus didymus; *carpella* granulato-rugulosa (juga secundaria), striis 5 impressis obsoletis notata (jugis primariis), lateralibus duobus semicircularibus antè marginem accessorium positis evittatis; commissura biforata; semen a basi ad apicem involutum. — *Invol.* univers. et partiale oligophyllum vel nullum.

1. *B. TESTICULATA* (Spreng. ap. Röm et Sch. t. 6. p. 448.) (1) inter segetes in provinc. mediterraneis. — *B. flosculosa* Bieb. taur. 3. p. 234. *Coriandrum testiculatum* L. sp. 367. Fl. fr. 4. p. 293. — Pluck. t. 169. f. 2. *Caulis* angulatus, folia decomposita, foliolis lineari-lanceolatis (v. v.)

Trib. V. *SILERINÆ. Koch umb. 84.*

*Fructus* a dorso lenticulari-compressus. *Carpella* multijugata, jugis primariis 5, lateralibus marginantibus, secundariis 4, omnibus filiformibus apteris. Semen antice planiusculum.

- X. *SILER Scop. carn. 1. p. 217. Gerth. fruct. 1. p. 92. t. 21. — Laserpitii sp. L.*

*Calyx* 5-dentatus; *petala* obovata emarginata cum lacinulâ inflexâ; *fructus* a dorso lenticulari-compressus; *carpella* jugis filiformibus elevatis obtusis, lateralibus marginantibus, secundariis minus prominulis; vallecule sub jugis secundariis univittatæ. — *Invol.* univ. et partiale oligophyllum caducum.

1. *S. TRILOBUM* (Scop. carn. 1. p. 217.) foliis biternatis, foliolis trilobis obtusis rotundo-ovatis crenatis. 2. in pascuis montanis Galloprovinciæ ex Ger, Pyren. ex cl. Benth. — Koch l. c. f. 34 et 35. *Laserpitium trilobum* L. *Angelica aquilegifolia* Fl. fr. 4. p. 306, et 5. p. 508. — Moris. a. g. t. 3. n. 3. *Civitas dubia.* (v. s. c.)

Subordo II. *UMBELLIFERÆ PERFECTÆ PAUCIJUGATÆ*, id est carpello jugis 5 primariis, secundariis nullis, vel et primariis basi et dorso oblitteratis apice tantum jugato.

Trib. VI. *TORDYLINEÆ. Koch 85.*

*Fructus* a dorso lenticulari-vel plano-compressus, margine dilatato incrassato noduloso vel plicato cinctus. *Carpella* jugis primariis 5 subindè tenuissimis vel obsoletis, lateralibus margini dilatato contiguis velleumdem formantibus. Semen complanatum et indè carpellum orthospermum.

XI. *CONDYLOCARPUS Hoffm. 1. p. 202. — Tordylii sp. L.*

*Calyx* 5-dentatus; *petala* obovata emarginata cum lacinulâ inflexâ, exteriora radiantia bifida; *fructus* a dorso plano-compressus, margine accessorio incrassato noduloso-crenato cinctus; *carpella* jugis tenuissimis tribus dorsalibus æquidistantibus, 2 lateralibus margine incrassato contiguis remotioribus, vallecule 3-vittatæ; vittæ striis separatæ; semen complanatum. — *Involuc.* et involucellum foliolis pluribus; commissura multi-striata 10-12-vittata.

1. *C. OFFICINALIS* (Koch umb. 87.) foliorum inferiorum foliolis ovatis incisiss

oreinato-dentatis, superiorum linearibus dissectis, fructu glanduloso, commissurâ 10-vittatâ. ① in provinciis australibus ad arva. — *Tordylium officinale* L. sp. 345. Fl. fr. 4. p. 335. — J. Bauh. hist. 3. p. 2. p. 84. f. 1. (v. v.)

XII. *TORDYLIUM* Hoffm. umb. 1. p. 198. — *Tordylii* sp. L.

Calyx 5-dentatus; petala obovata emarginata cum lacinulâ inflexâ, exteriora radiantia bifida; fructus a dorso plano-compressus, margine accessorio incrassato rugoso-tuberculato cinctus; carpella jugis tenuissimis, tribus dorsalibus æquidistantibus, 2 lateralibus margini incrassato contiguis vel ab eodem obtectis; valleculæ 1-vittatæ, vittis filiformibus; semen complanatum. — Involucrum et involucellum polyphyllum; commissura bivittata.

1. *T. maximum* (L. sp. 345.) foliis pinnatis, inferiorum foliolis lanceolato-ovatis inciso-crenatis, superiorum foliolis lanceolatis crenatis, terminali majore, fructu hispido. ① in incultis et marginibus arborum. — Fl. fr. 4. p. 335. Koch umb. f. 24 et 25. — Clus. hist. 2. p. 291. f. 1. Juga obsoleta. (v. v.)

Trib. VII. *PEUCEDANEÆ*. — *Selineæ* Koch umb. p. 88.

Fructus a dorso plano-vel lenticulari-compressus, margine dilatato lævigato alato complanato vel convexiusculo cinctus. Carpella jugis primariis 5 filiformibus subindè tenuissimis, lateralibus margini dilatato contiguis vel in eundem abeuntibus, secundariis nullis, totâ superficie commissurali connexa et sic fructus utrinquè unialatus. Semen complanatum vel dorso convexiusculum et hinc carpellum orthospermum.

XIII. *HERACLEUM* L. gen. n. 345.

Calyx 5-dentatus; petala obovata emarginata cum lacinulâ inflexâ, exteriora sæpè radiantia bifida; fructus a dorso plano-compressus, margine dilatato complanato cinctus; carpella jugis tenuissimis, tribus dorsalibus æquidistantibus, 2 lateralibus margini dilatato contiguis remotis; valleculæ univittatæ, vittis clavulatis; semen complanatum. — Involucrum caducum, involucellum polyphyllum.

Sect. I. *SPHONDYLIIUM* Hoffm. gen. umb. 1. p. 129.

Flores radiantes, commissura 2-vittata.

1. *H. SPHONDYLIIUM* (L. sp. 1. p. 358.) foliis pinnatis aut dilatatione petiolorum pinnatifidis, foliolis 3-5-lobatis lobis incisè crenatis, involucellorum foliolis lineari-setaceis, carpellis glabris orbicularibus. ② in pratis et ad sepes frequens. — Fl. fr. 4. p. 315. — Lob. ic. 703. f. 2. (v. v.)
2. *H. PYRENAICUM* (Lam. dict. 1. p. 403.) foliis subitè cano-tomentosis simplicibus rarissimè pinnatis cordato-palmatis 7-5-lobis acutis incisè crenatis, involucellorum foliolis lineari-subulatis, carpellis pubescentibus ellipticis. ② in pascuis Pyrenæorum præsertim orientalium. — Fl. fr. 4. p. 315. *H. setosum* Lapeyr. abr. 153 et *H. amplifolium* Lapeyr. abr. suppl. 44 ex cl. Benth. (v. v.)
3. *H. ANGUSTIFOLIUM* (L. mant. 57.) foliis tripartitis, lobis pinnatifidis lacinii confluentibus lanceolato-elongatis remotè dentatis acuminatis, involucellorum lacinii setaceis. ② in petrosis montium Lugdunensium ex Latour. Delph. propè Gratianopolim ex Vill. — Fl. fr. 5. p. 510. — Pluck. 1. 63. f. 3. (v. s. ex Valesiâ.)

4. *H. P. minimum* (Lam. fl. fr. 3. p. 413.) caule prostrato humili, foliis glaberrimis 2-pinnatis, foliolis incisis lacinii lanceolatis acutis, involuicellorum foliolis setaceis.  $\frac{1}{2}$  in lapidosis Alpibus Delph. — Fl. fr. 4. p. 316. H. pumilum Vill. dauph. a. t. 14. (v. s. sine fructu.)

Sect. II. *WENDIA Hoffm. l. c.*

Commisura evittata aut vittis 2 minimis.

5. *H. ALPIMUM* (L. sp. 359.) foliis simplicibus, subtus glabris cordato-palmatis lobis obtusiusculis subincisis crenato-dentatis, involuicellorum foliolis setaceis, carpellis obovato-orbicularibus.  $\frac{1}{2}$  in pascuis et ad oras sylvarum Jurassi, Alpium Delphin. ex Vill., Galloprov. ex Ger. — Fl. fr. 4. p. 316. — C. Bauh. prod. p. 83. ic. Vittæ commissurales 2 minimæ. (v. s.)

XIV. *PASTINACA L. gen. n. 362.*

Calycis margo obsoletus vel minutè denticulatus; petala subrotunda integra involuta lacinulâ latâ retusâ; fructus a dorso plano-compressus margine dilatato complanato cinctus; carpella jugis tenuissimis, tribus intermediis æquidistantibus, 2 lateralibus remotis margini accessorio contiguus sed ab illo sejunctis; valleculæ 1-vittatæ; semen complanatum. — Involucrum universale et parziale nullum vel oligophyllum.

1. *P. SATIVA* (L. sp. 376.) caule profundè sulcato, foliis simpliciter pinnatis, foliolis 5-11 lanceolatis incisis grossè serratis, superioribus sæpè decurrentibus, carpellis glaberrimis ovato-orbicularibus, commissurâ 2-vittatâ. ② in locis incultis ad sepea et vias. — Fl. fr. 4. p. 341. Lam. ill. t. 206. Koch umb. f. 26 et 27. (v. v.)
2. *P. KOCNII* (Duby mas.) caule striato foliisque simpliciter pinnatis utrinquè viridibus breviter pubescentibus, radicalibus 9-11 caulinis 3-5-foliolatis, foliolis ovatis basi cuneatis cordatisve indivisis argutè dentatis dentibus mucronatis, terminali majore subtrilobo, involuicellis nullis, carpellis glabris ovato-orbicularibus, commissurâ 4-6-vittatâ, vittis 2 majoribus.  $\frac{1}{2}$  in Corsicâ. — *P. graveolens* Salzmann. non Bieb. (v. s.)

*β. latifolia*, foliolis radicalibus dilatatis amplis subincisis. (v. s.)

XV. *ANETHUM Hoffm. umb. 1. p. 117. — Anethi sp. L.*

Calycis margo obsoletus; petala subrotunda integra involuta lacinulâ subquadratâ retusâ; fructus a dorso lenticulari-compressus, margine dilatato complanato cinctus; carpella jugis filiformibus æquidistantibus, tribus intermediis acutè carinatis, 2 lateralibus obsoletioribus in marginem abeuntibus; valleculæ 1-vittatæ; semen convexiusculum anticè planum. — Involucrum universale et parziale nullum; vittæ commissurales duæ.

1. *A. GRAVEOLENS* (L. sp. 377.) caule tereti glaberrimo striato, foliis decompositis glabris glaucis, lacinulis lineari-filiformibus integerrimis, carpellis glaberrimis ellipticis. ① in Aquitaniâ, Occitaniâ, Galloprovinciâ et circâ Parisios colitur et inde inter segetes subspontanea aliquandò occurrit. — Lob. ic. 776. (v. s.)

XVI. *PEUCEDANUM Koch umb. 1. p. 92. — Pencedani et Selini sp. L.*

Calycis margo 5-dentatus; petala obovata in lacinulam inflexam coarctata emarginata vel subintegra, fructus a dorso plano-vel lenticulari-compressus, margine dilatato complanato cinctus; carpella jugis subæquidistantibus, 3 intermediis filiformibus, 2 lateralibus obsole-



tioribus conformibusve margini dilatato contiguïs vel in eundem abeuntibus; semet anticè planum; vallecule 1-3-vittatæ. — Involucrum varium, involucellum polyphyllum.

## Sect. I. EUPEUCEDANUM.

Involucrum nullum aut oligophyllum; margo carpellorum minùs latus.

1. *P. PANICULATUM* (Lois. gall. 722.) caule glaberrimo tereti striato in parte superiore ramosissimo, foliis ternato-decompositis laciniis filiformibus, involucri foliolis 1-2 setaceis caducis, involucellis polyphyllis foliolis setaceis.  $\frac{1}{2}$  in Corsicâ. — Fl. fr. 5. p. 513. Umbellæ numerosæ paniculatim dispositæ. (v. s. sine fructu.)
2. *P. OFFICINALE* (L. sp. 353.) caule glaberrimo striato, foliis 3-4-pinnatis, foliolis lineari-ensiformibus, involucri foliolis 2-3 setaceis caducis minimis, involucelli polyphylli foliolis setaceis equalibus, carpellis glaberrimis ovato-ellipticis, commissurâ 2-vittatâ.  $\frac{1}{2}$  in umbrosis subhumidis. — Fl. fr. 4. p. 336. — Lob. ic. 782. f. 1. et 2. Fl. lutei. (v. v.)
3. *P. PARISIENSE* (DC. fl. fr. 4. p. 336.) caule glaberrimo striato, foliis ter aut quater pinnatis, foliolis ensiformibus, involucro 8-10-phylo foliolis lineari-subulatis, involucellis setaceis, commissurâ 2-vittatâ.  $\frac{1}{2}$  in sylvis Gallie mediæ et Occitanie ex cl. Benth. — *P. Gallicum* Pers. Fl. albi. (v. s. sine fructu.)
4. *P. CARVIFOLIUM* (Vill. dauph. 2. p. 630.) caule tereti glaberrimo striato, foliis pinnatis, foliolis pinnatifidis lobis decussatis 3-5-fidis laciniis linearibus acutis, involucro nullo, involucellis 3-4-phyllis foliolis lineari-subulatis, commissurâ 4-6-vittatâ, valleculis 2-3-vittatis.  $\frac{1}{2}$  in pratis sylvis et dumetis humidis Gallie mediæ, borealis orientalisque. — *Selinum* Chabreï Jacq. austr. 1. t. 72. Fl. fr. 4. p. 322. — Crantz. austr. 3. p. 22. t. 3. f. 2. (v. v.)

## Sect. II. THYSSELINUM Hoffm. umb. 1. p. 134.

Involucrum polyphyllum, vittæ commissurales tectæ, margo carpellorum minùs latus.

5. *P. PALUSTRE* (Mœnch. meth. p. 82.) caule tereti glaberrimo sulcato, foliis 2-3-pinnatis foliolis pinnatifidis, laciniis lineari-lanceolatis obtusiusculis, involucro involucellisque polyphyllis, foliolis lanceolato-linearibus apice subulatis, pedunculis puberulis, fructu lanceolato-elliptico.  $\frac{1}{2}$  in paludibus et pratis montanis Arvernæ ex Delarbre, Alsatiæ ex Mapp., Lotharingiæ ex Will. — Koch umb. p. 94. excl. Lin. syn. *Selinum* sylvestre L. hort. ups. 59. Fl. fr. 4. p. 319. Fl. dan. t. 412. Juga æqualia. (v. s. ex Daniâ.)

## Sect. III. CERVARIA.

Involucrum polyphyllum, vittæ commissurales superficiales, margo carpellorum minùs latus.

6. *P. ALSATICUM* (L. sp. 354.) caule tereti striato, foliis tripinnatis, foliolis pinnatifidis laciniis mucronatis in foliis superioribus lanceolato-ovatis, in radicalibus lineari-lanceolatis. involucris involucellisque polyphyllis, foliolis lanceolato-linearibus apice subulatis, fructu lanceolato-elliptico.  $\frac{1}{2}$  in sylvis subhumidis Alsatiæ, Galloprovinc., Pyren. orient. — Fl. fr. 4. p. 337. Jacq. austr. t. 70. non Poia. (v. s.)
7. *P. CERVARIA* (Lapeyr. abr. 149.) caule tereti striato, foliis bipinnatis, foliolis lanceolato-ovatis incisus inæqualiter argutè serratis basi obliquè cordatis dilatatis, involucro involucellisque polyphyllis, foliolis lineari-subulatis, fructu ovato.  $\frac{1}{2}$  in lapidosis montanis Alpium, Jurassi, Alsatiæ, Andegaviæ ex Bast. — *Selinum* Cervaria Crantz austr. 167. t. 3. f. 1. Fl. fr. 4. p. 319. — Fuchs, hist. 233. ic. (v. v. DC.)
8. *P. OROSELITUM* (Mœnch. meth. p. 82.) caule tereti striato, foliis 3-pinnatis, foliolis divaricatis cuneiformibus incisus 3-fidis pinnatifidisve acutis, invo-

luero involucrellisque polyphyllis, foliolis linearibus, fructu elliptico glabro.  $\frac{1}{2}$  in sylvis et montanis. — *Selinum Oreoselinum* Fl. fr. 4. p. 319. — *Clas. hist.* 2. p. 195. f. 2. (v. v.)

*ℓ. angulatum*, caule angulato pubescente. In Pyrensis. — Fl. fr. 5. p. 511. (v. v. DC.)

Sect. IV. *OREOSELINUM Hoffm. umb. gen. p. 155.*

Involucrum polyphyllum, vittæ commissurales superficiales, margo carpelorum latus.

9. *P. AUSTRIACUM* (Koch. umb. p. 94.) caule subangulato sulcato glabro, foliis 3-pinnatis, foliolis ovatis 3-5-lobis lobis cuneiformibus incisive lacinias mucronatis, involucris involucrellisque polyphyllis, foliolis lanceolatis lanceolato-linearibus mucronatis.  $\frac{1}{2}$  in Monte-Aureo ex Lam., Delph. montosis ex Vill. — *Selinum Austriacum* Jacq. aust. t. 71. Fl. fr. 4. p. 321. Commissura sæpe 4-vittata, vallecule exteriores subinde 2-vittatæ ex Koch. (v. s. c. sine fructu.)

10. *P. MONTANUM* (Koch l. c.) caule striato glabro subangulato, foliis 3-pinnatis, foliolis pinnatifidis lacinias mucronatis, involucro involucrellisque polyphyllis, foliolis lanceolato-linearibus acuminatis, stylis divaricatis reflexis, carpellis obovatis basi retusis, commissurâ 2-vittatâ.  $\frac{1}{2}$  in paludibus et pratis montanis Gallie mediæ, occident., borealis orientalisque. — *Selinum palustre* L. sp. 350. Crantz austr. p. 172. t. 4. f. 1. Fl. fr. 4. p. 320 excl. syn. Crantz. Jaga lateralia obsoleta. (v. s.)

XVII. *IMPERATORIA L. gen. n. 359.*

Calycis margo obsoletus; petala obovata in lacinulam inflexam coarctata emarginata vel subintègra; fructus a dorso plano-vel lenticulari-compressus margine dilatato latissimo complanato cinctus; carpella jugis subæquidistantibus, 3 intermediis filiformibus, 2 lateralibus obsoletioribus margini contiguis; semen anticè planum. — Involucrum nullum.

1. *I. OSTRUTRIUM* (L. sp. 372.) caule tereti glabro striato, foliis ternatis, foliolis rotundo-ovatis 2-3-lobis dentatis dentibus mucronatis.  $\frac{1}{2}$  in pascuis montanis. — Fl. fr. 4. p. 286. — *Garid. aix.* t. 55. (v. v.)

XVIII. *OPOBANAX Koch umb. p. 96.*

Calycis margo obsoletus; petala subrotunda integra involuta cum lacinulâ acutiusculâ; fructus a dorso plano-compressus, margine dilatato convexo cinctus; carpella jugis tribus dorsalibus filiformibus tenuissimis, lateralibus nullis (præter marginem incrassatam); vallecule trivittatæ; semen complanatum. — Involucrum et involucrellum oligophyllum; commissura 8-10-vittata.

1. *O. CUTRARIUM* (Koch l. c.) caule tereti glabro striato, foliis pinnatis bipinnatisque, foliolis serrato-crenatis sinuatis basi anticè excisis.  $\frac{1}{2}$  ad margines arborum in provinciis mediterraneis. — *Pastinaca Opopanax* L. mant. 357. *Con. ill.* t. 13 et 14. Fl. fr. 4. p. 341. (v. v.)

XIX. *FERULA Tourn. t. 170. L. gen. n. 343.*

Calycis margo 5-dentatus; petala ovata integra acuminata, acumine ascendente vel incurvato; fructus a dorso plano-compressus, margine dilatato complanato cinctus; carpella jugis 3 dorsalibus filiformibus tenuissimis, duobus lateralibus obsoletis in marginem dilatatum abeuntibus; vallecule 3-vittatæ; semen complanatum. — Involucrum nullum aut ut involucrellum oligophyllum; commissura 4-vittata.

1. *F. GLAUCA* (L. sp. 355.) foliis suprà-decompositis, foliolis lanceolato-linearibus planis, superioribus limbo abortivo subsimplicibus, involuero nullo, carpellis elliptico-ovatis.  $\mathcal{L}$  Monspelii et in Galloprovinciæ insulâ Sanctæ-Margaritæ (Stire). — Fl. fr. 5. p. 514. Koch umb. f. 3o et 31. — J. Bauh. hist. 3. p. 2. p. 45. f. 2. (v. v. DC.)
2. *F. communis* (L. sp. 355.) foliis suprà-decompositis, foliolis lineari-filiformibus mucronatis, umbellis ternatis, involucri sæpè caduci foliolis 4-5 linearilanceolatis acuminatis, carpellis ovatis.  $\mathcal{L}$  in montosis maritimis Corsicæ, Niocæ ex All. Galloprovinciæ ex Ger., Occitanis? et agri Ruscinon. ? — Fl. fr. 4. p. 343. — Lob. ic. 778. f. 2. (v. s.)

XX. FERULAGO Koch l. c. p. 97.

Calyx 5-dentatus, petala subrotunda integra curvata lacinulâ brevi inflexâ; fructus a dorso plano-compressus, margine dilatato complanato cinctus; carpella jugis tribus dorsalibus latiusculis obtusis, lateralibus duobus obsoletis in marginem dilatatum abeuntibus; semen complanatum undiquè vittis copiosissimis coopertum, dorsalibus pericarpio crassiusculo tectis; commissuralibus superficialibus. — Invol. et involucellum polyphyllum.

1. *F. nodiflora* (Koch l. c.) caule striato subangulato, foliis suprà-decompositis, foliolis filiformibus acutis divaricatis, umbellis pedunculatis ad nodos verticillatis, involucri foliolis 7-10 linearilanceolatis acutis, carpellis ellipticis.  $\mathcal{L}$  in vineis Niocæ ex All., Corsicæ. — Ferula nodiflora L. sp. 356. Fl. fr. 4. p. 343. — Barr. ic. 835. (v. s.)

Trib. VIII. ANGELICÆ. Koch l. c.

Fructus a dorso compressus, margine dilatato alato et ob superficies commissurales centro tantum connexas duplicato cinctus, hinc utrinquè hialatus. Carpella jugis primariis quinque, dorsalibus tribus filiformibus vel in alam expansis, lateralibus semper in alam abeuntibus duplò saltem latiore quam dorsalium jugorum; secundariis nullis. Semen dorso convexiusculum et ideo carpellum orthospermum.

Sect. I. ANGELICÆ NUCLEATÆ.

Semina in pericarpio libera nec nisi latere commissurali adnata.

XXI. ARCHANGELICA Hoffm. umb. 1. p. 166. — Angelicæ sp. L.

Calyx 5-dentatus; petala elliptica integra acuminata acumine incurvato; fructus a dorso subcompressus, utrinque 2-alatus; carpella jugis crassiusculis carinatis, tribus dorsalibus elevatis, duobus lateralibus in alam duplò latiore dilatatis; semen vittis copiosis undique tectum. — Involucrum subnullum, involucellum dimidiatum polyphyllum.

1. *A. officinalis* (Hoffm. l. c.) foliis bipinnatis, foliolis ovato-lanceolatis inæqualiter grossè dentatis, impari lobato.  $\mathcal{L}$  in montanis Galloprov. ex Ger., Alsatia ex Nestler. — Koch umb. t. 17. 18 et 19. Angelica Archangelica L. sp. 360. Fl. fr. 4. p. 305. (v. s. c.)

Sect. II. ANGELICÆ SOLIDÆ.

Calyx pericarpium et semen in unum coadunata.

XXII. ANGELICA Hoffm. umb. 1 p. 158. — Angelicæ sp. L.

Calycis margo obsoletus; petala lanceolata integra acuminata acu-

mine recto vel incurvato; fructus a dorso compressus, utrinque bialatus; carpella jugis tribus dorsalibus filiformibus elevatis, lateralibus in alam membranaceam multò latiore dilatatis; valliculæ 1-vittatæ; semen dorso convexum anticè planiusculum. — Involucr. oligophyllum aut nullum, involucellum polyphyllum; vittæ commissurales 2-4; carpella carinâ tantum angustâ cohærentia.

1. *A. RAZOULII* (Gou. ill. p. 13. t. 6.) caule striato apice pubescente, foliis 2-3-pinnatis, foliolis lanceolatis serratis acutis subtus scabris basi decurrentibus et sæpè 2-partitis, involucri foliolo 1 filiformi brevi, pedunculis pubescentibus, carpellis ovatis, vittis pericarpio tectis.  $\frac{1}{2}$  in Pyrenæis. — Fl. fr. 4. p. 306 et 5. p. 508. *A. ebullifolia* Lapeyr. — Bocc. mus. t. 99. Stamina longissima, petalorum acumen incurvum. (v. s.)
2. *A. MONTANA* (Schleich. exsic. Spreng. umb. prod. 16.) caule striato apice pubescente, foliis 3-pinnatis, foliolis lanceolatis ovatisve acuminatis glaberrimis basi decurrentibus argutè serratis dentibus mucronatis, pedunculis pubescentibus, involucri foliolo unico filiformi, carpellis ovatis latis, vittis superficialibus.  $\frac{1}{2}$  in Jurasso, Alpibus et montibus Ligericinis (Prost). — *Imperatoria montana* DC. fl. fr. 5. p. 504. Stam. longissima; petalorum acumen breve subrectum. (v. s.)
3. *A. SYLVESTRIS* (L. sp. 361.) caule striato puberulo apice pedunculisque tomentosis, foliis 2-3-pinnatis foliolis lanceolatis argutè serratis acutis glabris basi non decurrentibus, involucri foliolis 1-2 sæpius abortientibus.  $\frac{1}{2}$  in locis humidis et ad rivulos frequens. — Koch umb. f. 20 et 21. *Imperatoria sylvestris* Fl. fr. 4. p. 286. — Lob. ic. 699. Petalorum acumen subrectum. (v. v.)
4. *A. PYRÆNÆA* (Spreng. umb. gen. p. 62.) caule striato subsimplici, foliis pinnatis 2-pinnatisve foliolis 2-pinnatifidis lacinii linearibus acutis, involucri foliolo unico setaceo, umbellâ 4-7-radiatâ, carpellis elliptico-ovatis, vittis superficialibus.  $\frac{1}{2}$  in pascuis montanis Pyrenæorum, Arvernæ et Vogesorum. — *Selinum Pyrenæum* Gou. ill. p. 11. t. 5. Fl. fr. 4. p. 323. — Moris. s. g. t. g. n. 2. Petalorum acumen brevissimum. (v. v.)

XXIII. *SELINUM Hoffm. umb. gen. 1. p. 150. — Selini sp. L.*

Calycis margo obsoletus; petala obovata emarginata cum lacinulâ inflexâ; fructus a dorso compressus, carpellis centro tantum adnexis utrinquè bialatus; carpella jugis 5 membranaceo-alatis, alâ jugorum lateralium duplò latiore; valliculæ uni-vittatæ; semen subsemiteres. — Involucrum oligophyllum; involucellum polyphyllum; valliculæ exteriores sæpè bivittatæ; vittæ commissurales duæ.

1. *S. CARVIPOLEA* (L. sp. 350.) caule sulcato acutangulo glabro, foliis 3-pinnatis, foliolis pinnatifidis incisive, lacinii lanceolatis mucronatis, carpellis ovato-orbicularibus.  $\frac{1}{2}$  in sylvis humidis et stagnis umbrosis. — Fl. fr. 4. p. 322. Jacq. austr. t. 16. Koch umb. f. 22 et 23. (v. v.)

XXIV. *LEVISTICUM Koch umb. p. 101. — Ligustici sp. L.*

Calycis margo obsoletus; petala incurvata subrotunda integra cum lacinulâ brevi; fructus a dorso compressus, carpellis versùs marginem hiantibus utrinquè bialatus; carpella jugis quinque alatis, alâ jugorum lateralium duplò latiore; valliculæ uni-vittatæ; semen convexum anticè planiusculum. — Involucrum universale et parziale polyphyllum.

1. *L. OFFICINALE* (Koch l. c.) caule striato, foliis 2-pinnatis, foliolis cuneatis indivisis trilobisve apice incis.  $\frac{1}{2}$  in pascuis montanis Delph. ex Vill., Galloprov. ex Ger. Cebennarum ex Gouan, Pyr. orient. — *Angelica Levisticum* All. Fl. fr. 4. p. 306. Commissura quadrivittata. (v. s.)

Trib. IX. *SESELINEÆ*. — *Seselineæ* et *Ammineæ*. Koch  
umb. p. 102 et 114.

Fructus sectione transversali teres vel a latere compressus et aliquando latusculis contractis didymus. Carpella jugis primariis quinque filiformibus vel alatis, lateralibus marginantibus, omnibus æqualibus vel lateralibus paulò tantum latioribus, secundariis nullis. Semen dorso semiteres vel gibbo-convexum, anticè planiusculum vel undiquè teres et ideò carpella orthosperma.

Subtrib. I. *SESELINEÆ INTEGRIFOLIÆ*.

Petala arcte involuta, folia non dissecta.

XXV. *BUPLEVRUM* L. gen. n. 328.

Calycis margo obsoletus; petala subrotunda integra arcte involuta lacinulà latà retusà; fructus a latere compressus vel latusculis contractis subdidymus, stylopodio depresso coronatus; carpella jugis quinque æqualibus alatis vel filiformibus argutis, vel tenuissimis et obsoletis, lateralibus marginantibus; valleculæ vittatæ vel evittatæ; semen tereti-convexum, anticè planiusculum. — *Involucrum varium*.

Sect. 1. *ODONTITES* Hoffm. umb. p. 116.

Juga tenuissima depressa, valleculæ plerumque granulatæ 1-0-vittatæ, involuella umbellulas superantia. — *Species annuæ*.

1. *B. tenuissimum* (L. sp. 342.) caule erecto substriato ramoso, ramis elongatis patulis, foliis lineari-lanceolatis acutis basi attenuatis, floribus subsessilibus, umbellulis terminalibus lateralibusve, involucellis umbellulà longioribus involucriisque 3-5-phyllis, foliolis brevibus lanceolato-linearibus acuminatis, fructibus globosis, jugis tenuibus valleculisque evittatis granulatis. (I) in sterilibus herbosis Galliæ mediæ, australis, occidentalisque. — Fl. fr. 4. p. 350. — Barr. ic. t. 1248. (v. v.)
2. *B. elaucum* (DC. fl. fr. 5. p. 515.) caule diffuso obsoletè angulato à basi ramosissimo, foliis linearibus acuminatis ad basim subincrassatis, umbellis lateralibus terminalibusve, involucellis umbellulà longioribus involucriisque 5-phyllis, foliolis linearibus aristato-acuminatis, fructibus granulatis, jugis oblitteratis. (I) in arenosis maritimis Corsicæ, Narbonæ, Massiliæ, Telonis, Niceæ. (v. v.)
3. *B. semicompositum* (L. sp. 342.) caule erecto substriato ramoso, foliis lanceolatis obtusis mucronatis basi attenuatis subsemialexicaulibus, involucellis umbellulà longioribus involucriisque 5-phyllis, foliolis lineari-lanceolatis aristato-acuminatis, jugis subobsoletis valleculisque evittatis granulatis. (I) in sterilibus Narbonæ ex Lam., Niceæ ex All. — Fl. fr. 4. p. 350. Gouan ill. p. 9. t. 7. f. 1. (v. v. c.)
4. *B. Odontites* (L. sp. 342.) caule tereti striato ramoso ramis divaricatis, foliis lineari-ensiformibus acuminatis glabris, involucellis umbellulam superantibus involucriisque 5-phyllis, foliolis lanceolatis venosis acuminatis, jugis tenuissimis depressis, valleculis lævibus 1-vittatis. (I) in messibus et collibus provinc. austr., Arvernæ, Andegaviæ, Pictaviæ, et propè Cherbourg. — Fl. fr. 4. p. 349. — J. Bauh. hist. 3. part. 2. p. 201. f. 1. (v. v.)
5. *B. rotundifolium* (L. sp. 340.) caule erecto læviusculo apice ramoso, foliis obtusis mucronatis inferioribus lanceolato-ovatis superioribus ovatis perfoliatis, invol. nullo, involucellis 5-phyllis, foliolis rotundo-ovatis acuminato-aristatis, jugis depressis tenuissimis, valleculis granulatis subevittatis.

① inter segetes Galliz australis, mediz occidentalisque. — Fl. fr. 4. p. 345. Lob. ic. t. 396. f. 1. (v. v.)

# Sect. II. EURUPLEVUM.

Juga elevata subalata, vallecule læves 1-3-vittatæ. — Species plerumque ♀.

† *Involucellis umbellulæ superantibus.*

6. *B. LONGIFOLIUM* (L. sp. 341.) caule erecto substriato subsimplici, foliis acutis, radicalibus lanceolatis in petiolum coarctatis, superioribus cordato-ovatis amplexicaulibus, involucri inæqualibus sub-5-phyllis, involucellis umbellulâ longioribus, foliolis lanceolato-ovatis acutis, jugis membranaceo-alatis, valleculis lævibus, vittis 3 obsoletis. ♀ in lapidosis montanis Alpium, Jurassi, Vogesorum, Arverniz. — Fl. fr. 4. p. 345. — J. Bauh. hist. 3. p. 199. f. 1. (v. v.)
7. *B. PYRENEUM* (Coul. ill. p. 8. t. 4. f. 1. 2.) caule simplici lævi erecto flexuoso folioso, foliis lanceolatis acuminatis, radicalibus in petiolum attenuatis, superioribus cordato-ovatis amplexicaulibus, involucri 3-phyllis, foliolis ovato-lanceolatis obtusis umbellarum radios æquantibus, involucellis 5-phyllis, foliolis rotundis obtusissimis umbellulam superantibus, jugis alatis, valleculis 3-vittatis. ♀ in rupibus editiorum Pyrenæorum. — Fl. fr. 4. p. 346. (v. v.)
8. *B. STELLATUM* (L. sp. 340.) caule substriato simplici subnudo, foliis radicalibus lanceolato-linearibus linearibusve acutis in petiolum attenuatis, caulino nullo aut unico sessili, involucri 1-3-phylo, foliolis lanceolato-ovatis amplexicaulibus acutis, involucellis 9-10-phyllis, foliolis cœdunatis obovatis macronulatis umbellulam superantibus, jugis membranaceo-alatis, valleculis 1-vittatis. ♀ in rupestribus Alpium et Corsicæ montium. — Fl. fr. 4. p. 346. — Hall. herb. t. 18. (v. v.)
9. *B. GRAMINIFOLIUM* (Vahl. symb. 3. p. 48.) caule substriato simplici stricto subnudo, foliis radicalibus numerosis graminæis linearibus longè acuminatis basi dilatatis, caulinis 1-2 ensiformibus acuminatis semi-amplexicaulibus, involucri 3-6-phylo, foliolis lanceolato-elongatis acuminatis, involucellis 7-9-phyllis, foliolis lanceolatis cuspidatis umbellulam superantibus, jugis elevatis subalatis, valleculis lævibus. ♀ in Alpibus Galloprovinciæ. — Fl. fr. 4. p. 347. *B. petræum* Jacq. rar. 1. t. 56. Vill. dauph. 2. t. 14. non L. (v. v. D. in Italiâ.)
10. *B. RANUNCULOIDES* (L. sp. 342.) caule substriato subsimplici, foliis linearilanceolatis acuminatis, caulinis basi dilatatis amplexicaulibus, involucri 5-phyllis, foliolis inæqualibus lanceolatis ovatisve acutis, involucellis 5-6-phyllis, foliolis obovatis cuspidatis umbellulam subsuperantibus, jugis elevatis argutis subalatis, valleculis lævibus 1-vittatis. ♀ in editioribus montibus. — Fl. fr. 4. p. 348. *B. angulosum* L. *B. repens* et *B. obtusatum* Lapeyr. ex cl. Benth. — J. Bauh. hist. 3. p. 199. f. 1. (v. v.)  
*β. caricifolium*, foliis linearibus angustioribus, involucri 1-3-phylo caduco. — Inter lapides et saxa Alpium et Pyrenæorum. — *B. caricifolium* Fl. fr. 4. p. 348. an Wild? Vix varietas. (v. v.)
11. *B. JUNCUM* (L. sp. 342.) caule substriato ramoso, foliis linearibus acuminatis, involucellis 5-phyllis umbellulâ subbreioribus, involucri 2-5-phyllis, foliolis lineari-lanceolatis acuminatis, fructibus globosis, jugis tenuibus argutis subalatis, valleculis lævibus, vittis 3 aut 0. ① in margine arvorum et sepibus Galliz austr., orientalis, propè Parisios ex Thuil., in Arverniâ ex Del., Andegaviâ ex Bast. — Fl. fr. 4. p. 351. — Moris. s. 9. t. 12. n. 3. (v. v.)
12. *B. GERARDI* (Murr. syst. 274.) caule lævi ramoso flexuoso, foliis linearibus acuminatis sessilibus, involucellis umbellulam superantibus involucrique foliolis 5 lineari-subulatis inæqualibus, fructû cylindrico, jugis tenuibus acutiusculis subelevatis, valleculis lævibus 3-vittatis. ① in sterilibus provinciarum mediterranearum. — Fl. fr. 4. p. 351. — Ger. gallop. t. 9. (v. v.)

†† *Involucellis umbellulis brevioribus.*

13. *B. FALCATUM* (L. sp. 341.) caule substriato flexuoso folioso ramoso, ramis erectis, foliis lanceolatis nervosis mucronulatis utrinque attenuatis basi amplexicaulibus, inferioribus petiolatis, superioribus linearibus, involucri foliolis 3-5 inæqualibus lanceolatis linearibusve, involucellis foliolis 5-6 lanceolato-linearibus acuminatis umbellulâ brevioribus, jugis elevatis subulatis, valleculis lævibus 3-vittatis. ½ in lapidosis aridis, ad sepes et inter dumeta. — Fl. fr. 4. p. 347. — Lob. ic. 456. f. 1. (v. v.)

β. *petiolare* (DC.) foliis radicalibus ovalibus longè petiolatis. — B. petiolare Lapeyr. (v. v.)

14. *B. ARIIDUM* (L. sp. 342.) caule stricto substriato ramoso ramis divaricatis, foliis inferioribus lanceolatis acutis nervosis in petiolum basi amplexicaulem coarctatis, superioribus lineari-lanceolatis sessilibus, involucellis umbellulâ multò minoribus involucrisque minimis 5-phyllis, foliolis linearibus acutis, jugis elevatis subulatis, valleculis lævibus 1-vittatis. ½ in lapidosis aridis Galloprov., Occitaniz. — Fl. fr. 4. p. 349. — Lob. ic. t. 456. f. 2. (v. s.)

Sect. III. *TENORIA Spreng. in Rœm. et Schult. vi. p. xxxiv.*

Juga elevata subelevatave, valleculæ læves 1-vittatæ, involucella umbellulis breviora. — Species suffruticosæ.

15. *B. SUFFRUTICOSUM* (L. amæn. 4. p. 269.) suffrutescens, caule tortuoso ramoso ramis erectis strictis, foliis linearibus rigidis convolutis acuminatis, involucellis umbellulâ brevioribus involucrisque 5-phyllis, foliolis brevibus reflexis linearibus acutis, fructibus glaucis, jugis vix elevatis, valleculis lævibus. ½ in agro Ruscinonensi inter Narbonam et Perpignanum reperiit Ph. Thomas. — Cav. ic. 2. t. 106. — Barr. ic. t. 1255. (v. s.)

16. *B. SPINOSUM* (L. supp. p. 178.) suffrutescens, caule recto lævi folioso ramoso, ramis tortuosis divaricatis patulis subnudis, foliis linearibus subcoriaceis basi dilatatis amplexicaulibus, involucris involucellisque minimis, jugis subelevatis, valleculis lævibus. ½ in Corsicâ reperiit Ph. Thomas. — Gou. ill. p. 8. t. 2. f. 3. (v. s.)

17. *B. FRUTICOSUM* (L. sp. 343.) suffrutescens, caule stricto ramoso, foliis lanceolato-obovatis venosis mucronulatis, involucris involucellisque umbellulâ duplò brevioribus polyphyllis, foliolis caducis lanceolato-obovatis obtusiusculis, jugis elevatis subulatis, valleculis læviusculis 1-vittatis. ½ in provinciis mediterraneis. — Fl. fr. 4. p. 345. — Moris. s. 9. t. 6. n. 1. Vittæ tenuissimæ. (v. v.)

## Subtrib. II. SESELINEÆ SECTIFOLIÆ.

Subdiv. I. SESELINEÆ SOLIDÆ, id est semine pericarpio calyceque in unum coadunatis.

† *Multivittatæ.*

XXVI. *BRIGNOLIA Bertol. amæn. ital. p. 97. — Sii sp. L.*

Calycis margo 5-dentatus; petala subrotunda integra involuta lacinulâ latâ retusâ; fructus subteres a latere subcompressus lineari-elongatus; carpella jugis 5 filiformibus obtusis æqualibus, lateralibus marginantibus; valleculæ 1-3-vittatæ; semen subsemiteres. — Involucrum et involucellum polyphyllum, commissura 4-6-vittata.

1. *B. PASTINACÆFOLIA* (Bertol. l. c.) foliis bipinnatis, foliolis ovatis basi obliquis serratis serraturis mucronatis, superioribus decurrentibus, involucri involucellisque polyphyllis, foliolis setaceis, carpellis glaberrimis. ½ in arvis circâ Bonifacio. — Sium Siculum L. sp. 362. Fl. fr. 5. p. 507. — Zanoni. hist. 78. t. 30. Fl. aurei. (v. s.)

## XXVII. ATHAMANTA Koch umb. p. 105. — Athamante sp. L.

Calycis margo 5-dentatus; petala obovata emarginata cum lacinulâ inflexâ, brevissimè unguiculata; fructus in collum attenuatus sectione transversali subteres vel a latere parum compressus; carpella jugis 5 filiformibus apteris æqualibus, lateralibus marginantibus; valleculæ 2-3-vittatæ; semen subsemiteres. — Invol. et involucellum polyphyllum.

1. A. MACEDONICA (Spreng. in Rœm. et Schult. vi. p. 491.) caule pubescente ramosissimo, foliis glabriusculis 2-pinnatis superioribus pinnatis, foliis rhombo-ovatis incisive crenatis, involucri involucellisque polyphyllis, foliolis linearibus hirsutissimis, fructu hispidio. (2) in pratis sterilibus circa Niceam ex All. et in hortis sæpè cultus. — Bubon Macedonicum Fl. fr. 4. p. 298. — Lob. ic. t. 708. f. 1. Semen vittis 10-12 undiquè obtectum, juga tenuissima setacea. (v. v. c.)
2. A. CRETENSIS (L. sp. 352.) caule hirsuto tereti sublevi, foliis hirsutis 3-pinnatis, foliolis 3-5-lobis lineari-lanceolatis acutis, involucri caduco involucellisque polyphyllis, foliolis scariosis linearibus aristatis, fructu villosissimo. 2½ in montibus editioribus. — Fl. fr. 4. p. 318. — Cam. epit. 536. ic. Juga planiuscula latiuscula, valleculæ 2-3-vittatæ, commissura 6-vittata. (v. v.)  
*♀. decipiens*, foliis glabriusculis, foliolis linearibus. — In rupestribus montium Delph. et Galloprov. — A. Matthioli Fl. fr. 4. p. 318. non Wulf. nec Jacq. A. rupestris Vill. non Scop. (v. s.)

## XXVIII. CONOPODIUM Koch umb. p. 118. — Bunii sp. DC.

Calycis margo obsoletus; petala obovata emarginata vel subemarginata cum lacinulâ inflexâ; fructus a latere compressus lineari-oblongus, stylopodio conico immarginato stylisque rectis coronatus; carpella jugis 5 æqualibus filiformibus obtusis, multivittata; semen semitereti-convexum anticè planiusculum. — Involucrum et involucellum oligophyllum; carpophorum (axis fructus) liberum apice bifidum.

1. C. DENUDATUM (Koch l. c.) radice bulbosâ, caule erecto simplici, foliis 2-pinnatis, foliolis linearibus integris, involucri foliolis linearibus, jugis tenuissimis, valleculis 2-3-vittatis. 2½ in pratis montosis Delph., Lugd., Arvernæ, Cebenn., Pyren., Burdigalæ (Des Moul.), Andegaviæ (Bast.), Aureliæ, Bellovaci ex cl. Graves. — Bonium denudatum Fl. fr. 4. p. 325. (v. v.)

## XXIX. PIMPINELLA L. gen. n. 366.

Calycis margo obsoletus; petala obovata emarginata cum lacinulâ inflexâ; fructus a latere compressus ovatus, stylopodio pulvinato stylisque reflexis coronatus; carpella jugis 5 filiformibus æqualibus, lateralibus marginantibus; valleculæ multivittatæ; semen gibbo-convexum anticè planiusculum. — Invol. et involucell. 0; carpophorum liberum bifidum.

## Sect. I. TRAGOSELINUM Mœnch. meth. p. 99.

Fructus glabri.

1. P. MAGNA (L. mant. p. 219.) caule erecto sulcato glabro, foliis pinnatis, radicalium foliolis inciso-dentatis lobatisve impari trilobo, jugo inferiore pedicellato, superiorum foliolis pinnatifidis laciniis linearibus. 2½ in locis incultis et ad oras sylvarum. — Fl. fr. 4. p. 281. — Barr. ic. 243. (v. v.)
2. P. DISSECTA (Retz obs. 3. p. 30. t. 2.) caule erecto striato simplici glabro, foliis pinnatis, foliolis pinnatifidis sessilibus, segmentis sublinearibus acutis. 2½ in sabulosis aridis passim. — Fl. fr. 4. p. 282. An præced. var? (v. v.)



## UMBELLIFERÆ.

229

3. *P. SAXIFRAGA* (L. sp. 378.) caule erecto striato infernè puberulo, foliis pinnatis, foliolis sessilibus infernè puberulis, radicalium ovatis grossè dentatis, superiorum lanceolatis linearibusve incis. ♀ in pascuis siccis. — Fl. fr. 4. p. 281. Jacq. aust. t. 395. Koch umb. f. 55 et 56. (v. v.)

### Sect. II. *TRAGIUM Spreng. umb. prod. p. 26.*

Fructus villosi.

4. *P. PERGRINA* (L. sp. 378.) caule erecto læviusculo, foliis pinnatis, radicalium petiolo pubescente, foliolis crenatis obtusis inferioribus ovatis subpedicellatis summo cordato cuneatove, superiorum foliolis linearibus lanceolatis incis. acutis, umbellis 15-20-radiatis, fructibus hispidis. (1) (2) in sepibus et agris Corsicæ, Nicæ, Galloprov.; Occitanicæ. — Fl. fr. 5. p. 502. *P. hispida* Lois. not. 48. Fl. fr. 5. p. 502. — Moris. hist. 2. s. 9. t. 9. n. 13. (v. v.)
5. *P. TRAGIUM* (Vill. dauph. 2. p. 606.) caulibus erectis diffusivè lævisculis, foliis tenuissimè puberulis pinnatis, foliolis pinnatifidis segmentis inciso-dentatis acutiusculis, umbellis 4-7-radiatis, fructibus velutino-incanis. ♀ in siccis lapideis Galloprovinciæ, Delph. austral., Occitanicæ et Pyren. orientali-um. — Fl. fr. 5. p. 501. *P. canescens* Lois. not. t. 4. — Col. phyt. 76. (v. v.)

### XXX. *SIUM Koch umb. 117. — Sü sp. L.*

Calycis margo 5-dentatus, subindè obsoletus; petala obovata emarginata cum lacinulâ inflexâ; fructus a latere compressus vel contractus et subdidymus, stylopodio margine depresso stylisque reflexis coronatus; carpella jugis quinque æqualibus filiformibus obtusiusculis, multivittata; semen subteres. — Involucrum varium, involucellum polyphyllum; carpophorum bipartitum.

\* *Jugis lateralibus marginantibus; valleculis 3-4-vittatis vittis superficialibus; semine tereti-convexo anticè planiusculo.*

1. *S. SISARUM* (L. sp. 361.) caule erecto, foliis pinnatis superioribus ternatis, foliolis ovato-lanceolatis lineari-lanceolatisque serratis acuminatis, involu- cro caduco involucellisque polyphyllis, foliolis æqualibus lineari-lanceolatis, commissurâ 4-vittatâ. ♀ in hortis colitur. — Fl. fr. 4. p. 301. — Lob. ic. 710. f. 1. (v. v. c.)
2. *S. LATIFOLIUM* (L. sp. 361.) foliis pinnatis, foliolis oblongo-lanceolatis æqualiter argutè serratis mucronatis, involu- cro inæquali involucellisque polyphyllis, foliolis linearibus acuminatis, commissurâ 6-vittatâ. ♀ ad sepes et in paludosis. — Fl. fr. 4. p. 299. — Lob. ic. 208. (v. v.)
- \* \* *Jugis lateralibus paulò antè marginem carpelli positus; seminibus teretibus multivittatis et pericarpio corticoso arcuè adnato tectis.*
3. *S. ANGUSTIFOLIUM* (L. sp. 1672.) foliis pinnatis, foliolis lanceolatis inæqualiter lobatis serratisque acutis, umbellis pedunculatis oppositifoliis, involu- cro involucellisque polyphyllis, foliolis lineari-lanceolatis acutis. ♀ in rivulis fossisque. — Fl. fr. 4. p. 299. (v. v.)

### XXXI. *LIGUSTICUM Tourn. 171. L. gen. n. 346.*

Calycis margo 5-dentatus vel obsoletus; petala obovata aut obovato-oblonga in lacinulam inflexam coarctata brevissimè unguiculata; fructus sectione transversali subteres vel a latere parum compressus; carpella jugis 5 argutis subalatis æqualibus, lateralibus marginantibus; valleculæ multivittatæ; semen subsemiteres. — Involucrum varium, involu- cellum polyphyllum.

#### Sect. I. *EULIGUSTICUM.*

Petala acutè emarginata brevissimè unguiculata.

\* *Calycis margine minute 5-dentato; involucri universali polyphylo.*

1. *L. FRUTULACRUM* (All. ped. t. 60. f. 1.) caule tereti folioso sulcato apice ramoso, foliis 2-3-pinnatis, foliolis pinnatifidis lacinii linearibus subcarnosis, involucri foliolis pinnatifidis lacinii linearibus.  $\mathcal{L}$  in Jurasso suprâ Gex, Alpibus Delphin., Galloprovinc., et Pyrenæis editioribus ex Lapeyr. — Fl. fr. 4. p. 308. (v. s.)
2. *L. SIMPLEX* (All. ped. t. 71. f. 2.) caule nudo simplici striato, foliis pinnatis, foliolis multifidis lacinii acutis linearibus, involucri foliolis pinnatifidis, lacinii linearibus.  $\mathcal{L}$  in pascuis Alpinis Delph., Pyren. — *Laserpitium simplex* L. mant. 56. Fl. fr. 4. p. 314. — Vill. dauph. 2. t. 14. Juga alata. (v. v.)

\* \* *Calycis margine obsolete; involucri nullo aut 1-phylo.*

3. *L. PYRENAICUM* (Gou. ill. t. 7. f. 4.) caule striato folioso, foliis tripinnatis, foliolis pinnatifidis lacinii lineari-lanceolatis mucronatis, involucri 1-phylo lineari brevi caduco.  $\mathcal{L}$  in siccis petrosis Pyrenæorum. — Fl. fr. 4. p. 309. — Seg. ver. 41. t. 13. Koch umb. f. 46 et 47. (v. s.)

#### Sect. II. *SILAUS* Koch umb. p. 105.

Petala subintegra basi truncata sessilia.

4. *L. SILAUS* (Duby mss.) caule erecto striato, foliis 3-2-pinnatis, foliolis 3-5-lobis lanceolatis acutis, involucri nullo aut raro 1-2-phylo, involucri polyphyllis, foliolis indivisis linearibus acuminatis.  $\mathcal{L}$  in pratis. — *Pencœdanum* Silaus L. sp. 354. Fl. fr. 4. p. 337. — Lob. ic. 738. f. 1. (v. v.)

XXXII. *MEUM* Jacq. Gärtn. fruct. 1. p. 106. t. 23. Koch umb. p. 103. — *Athamantæ* et *Phellandrii* sp. L.

Calycis margo obsoletus; petala integra ovato-elliptica basi apiceque acuta; fructus sectione transversali a latere subcompressus; carpella acuta jugis 5 prominulis carinatis æqualibus, lateralibus marginantibus; vallecule multivittatæ; semen subsemiteres. — Involucrum oligophyllum aut nullum, involucellum polyphyllum; vittæ commissurales 6-8.

1. *M. MUTELLINA* (Gärtn. fruct. 1. p. 106. t. 23.) caule simplici lævi subnudo, foliis 2-pinnatis, foliolis 2-pinnatifidis lacinii lineari-lanceolatis, involucri caduco, involucellorum foliolis lanceolatis acuminatis, carpellis lanceolatis utrinque acutis.  $\mathcal{L}$  in pascuis editioribus Vogesorum, Alpium, Montis-Aurei. — *Ligusticum Mutellina* Fl. fr. 4. p. 310. All. ped. t. 60. f. 2. — Cam. epit. p. 8. ic. (v. s.)
2. *M. TENUIFOLIUM* (Duby mss.) caule simplici subnudo lævi, foliis radicalibus decompositis, caulinis 3-fidis, foliolis lineari-filiformibus, involucri foliolis 1-2 involucellorum 5-8 lineari-lanceolatis acutis.  $\mathcal{L}$  in fissuris rupium editiorum Pyrenæorum. — *Ligusticum tenuifolium* DC. fl. fr. 4. p. 309. L. splendens Lapeyr. Wallrothia splendens Spreng. in Rœm. et Schult. 6. p. 557. (v. s.)
3. *M. ATHAMANTICUM* (Moris. hist. 2. s. 9. t. 2. n. 2.) caule simplici subnudo striato, foliis decompositis, foliolis capillaribus, involucri foliolis 3-5, involucellorum foliolis 10-12 basi lanceolato-linearibus apice subulato-acuminatis, carpellis lanceolatis angustis utrinque acutis.  $\mathcal{L}$  in pascuis montium editiorum. — Jacq. aust. t. 303. *Ligusticum Meum* Fl. fr. 4. p. 310. (v. v.)

#### XXXIII. *TROCHISCANTHES* Koch umb. p. 103.

Calycis margo 5-dentatus; petala spathulato-obovata subintegra cum lacinulâ triangulari inflexâ, longè unguiculata; fructus a latere subcompressus; carpella jugis quinque argutis subulatis æqualibus, lateralibus marginantibus; vallecule multivittatæ; semen subsemiteres. — Involucrum nullum, involucellum 3-5-phyllum.

1. *T. nodiflorus* (Koch l. c.) caule tereti striato, foliis ter aut quater ternatis, foliolis lanceolato-ovatis acutis grossè dentatis, ramis verticillato-paniculatis bracteatis, bracteis lineari-lanceolatis integerrimis.  $\frac{1}{2}$  in sylvis umbrosis montosis Delph. et Galloprovinciæ superioris. — *Imperatoria nodiflora* Fl. fr. 4. p. 287. Lig. nodifl. Vill. 2. t. 13. (v. s.)

† † *Paucivittatæ.*

XXXIV. *MOLOSPERMUM* Koch umb. p. 108. — *Ligustici* sp. L.

Calyx quinquedentatus foliolosus; petala lanceolata integra longè acuminata acumine ascendente; fructus sectione transversali a latere parum compressus; carpella jugis 5 aut abortu 3 argutis membranaceo-alatis æqualibus, lateralibus marginantibus; vallecule 1-vittatæ; semen semiteres. — Involucrum universale et partiale polyphyllum.

1. *M. Peloponnesiacum* (Koch l. c.) foliis multiplici-pinnatis, foliolis pinnatis incisim acuminatis, involucris involucellisque polyphyllis, foliolis lanceolatis acuminatis, carpellis lanceolatis utrinquè attenuatis.  $\frac{1}{2}$  in Alp. Delph., Galloprov., Cebenn., Pyren. orient. — *Ligusticum Peloponnesiacum* L. sp. 360. Fl. fr. 4. p. 307. — Cam. epit. 514. ic. (v. v. DC.).

XXXV. *ÆGOPIDIUM* L. gen. n. 368.

Calycis margo obsoletus; petala obovata emarginata cum lacinulâ inflexâ; fructus a latere compressus oblongus stylis reflexis coronatus; carpella jugis 5 filiformibus, lateralibus marginantibus; vallecule evittatæ; semen tereti-convexum, anticè planiusculum. — Invol. et involucellum nullum. Genus habitû et valleculis evittatis à *Caro* diversum sed tamen nimium affine.

1. *Æ. Podagraria* (L. sp. 379.) caule profundè striato, foliis 2-ternatis ternatisve, foliolis lanceolato-ovatis acuminatis.  $\frac{1}{2}$  ad sepes, in hortis et pomariis. — Fl. fr. 4. p. 280. — Lob. ic. t. 700. f. 2. (v. v.)

XXXVI. *CARUM* L. gen. n. 365.

Calycis margo obsoletus; petala regularia obovata emarginata cum lacinulâ inflexâ; fructus a latere compressus oblongus, stylis reflexis coronatus; carpella jugis 5 filiformibus æqualibus, lateralibus marginantibus; commissura plana; vallecule 1-vittatæ; semen tereti-convexum, anticè planiusculum. — Invol. et involucellum varium; carpophorum apice furcatum liberum.

\* *Involucro 1-phyllò aut nullo.*

1. *C. Carvi* (L. sp. 378.) caule erecto ramoso, foliis 2-pinnatis, foliolis decussatis pinnatifidis, laciniiis linearibus acutis, fructu oblongo-globose. ② in pratis montosis. — *Seseli Carvi* Fl. fr. 4. p. 285. — Cam. epit. 516. ic. (v. v.)

\*\* *Involucro et involucello polyphyllò.*

2. *C. Bulbocastanum* (Koch umb. p. 121.) radice tuberosâ, caule erecto striato, foliis 2-3-pinnatis, foliolis 2-3-lobatis laciniiis linearibus acutis, involucris involucellorumque foliolis linearibus brevibus acutis, fructibus elongato-oblongis, carpellis sectione verticali sublinearibus.  $\frac{1}{2}$  in pascuis subhumidis Galliæ orientalis, mediæ, Pyrenæorum ex Lapeyr., Burdigalæ ex Laterr. — *Bunium Bulbocastanum* L. sp. 349. Fl. fr. 4. p. 325. — Lob. ic. 745. f. 1. (v. v. DC.)
2. *C. verticillatum* (Koch umb. 122.) radice fasciculatâ, caule erecto substriato, foliis pinnatis, foliolis multifido-capillaceis subverticillatis, involucris involucellisque brevibus lineari-lanceolatis, fructu oblongo, carpellis

linearibus.  $\mathcal{L}$  in pratis humidis passim. — *Sium verticillatum* Lam. enc. 1. p. 407. DC. fl. fr. 4. p. 302. — Dalech. lugd. 718. ic. (v. v.)

XXXVII. DREPANOPHYLLUM *Moench. supp. 2. Hoffm. umb. 1. p. 109. — Sii sp. L.*

Calycis margo 5-dentatus; petala obovata curvata sinu lato emarginata cum lacinulâ inflexâ, fructus teres elongato-oblongus stylis reflexis coronatus; carpella jugis 5 filiformibus æqualibus, lateralibus marginantibus; vallecule 1-vittatæ vittis filiformibus; semen tereti-convexum, anticè planiusculum. — Involucrum polyphyllum, involucellum submediatum, foliolis interioribus brevissimis.

1. *D. FALCARIA* (Moench. l. c.) caule striato, foliis pinnatis, foliolis linearibus argutè serratis decurrentibus.  $\mathcal{L}$  in arvis et ad sepes passim. — *D. agreste* Hoffm. *Sium Falcaria* L. sp. 362. Fl. fr. 4. p. 301. — Lob. ic. 2. t. 24. f. 1. (v. v.)

XXXVIII. PETROSELINUM *Hoffm. umb. 1. p. 78. — Apii sp. L.*

Calycis margo obsoletus; petala subrotunda incurvata integra vix emarginata, in lacinulam inflexam coarctata; fructus ovatus a latere contractus subdidymus; carpella jugis quinque filiformibus æqualibus, lateralibus marginantibus; vallecule 1-vittatæ; semen gibbo-convexum anticè planiusculum. — Involucrum oligophyllum, involucellum polyphyllum; carpophorum bipartitum.

1. *P. SATIVUM* (Hoffm. l. c.) foliis inferioribus 2-pinnatis, foliolis ovatis incisim dentatis, caulinis linearibus integrisculis, umbellis pedunculatis, involacris 1-3-phyllis, involucellis polyphyllis foliolis setaceis. ② in umbrosis Galloprovinciæ, in hortis cultum. — *Apium Petroselinum* L. sp. 379. Fl. fr. 4. p. 339. — Black. t. 172. a. J. Bauh. hist. 3. part. 2. p. 99. ic. et 97. f. 2. (v. v. c.)

2. *P. SEGETUM* (Koch umb. p. 128.) foliis omnibus pinnatis, foliolis lanceolato-ovatis incisim serratis, umbellis longè pedunculatis 2-3-radiatis, invol. 2-phyllis, involucellis 4-6-phyllis foliolis lanceolatis brevissimis, fructu ovato-globoso, stylis brevissimis. ① in agris subhumidis et ad sepes Galliæ mediæ et occidentalis. — *Sium segetum* Fl. fr. 4. p. 303. *Sison segetum* L. Jacq. vind. t. 134. (v. s.)

XXXIX. APIUM *Hoffm. umb. 1. p. 75. — Apii sp. L.*

Calycis margo obsoletus; petala subrotunda integra cum apiculo arcte involuto; fructus subrotundus a latere contractus didymus; carpella jugis 5 filiformibus æqualibus, lateralibus marginantibus; vallecule univittatæ, exteriores sæpè 2-3-vittatæ; semen gibbo-convexum anticè planiusculum. — Invol. involucellumque nullum; carpophorum indivisum.

1. *A. GRÆVROLENS* (L. sp. 379.) foliis pinnatis ternatisve, foliolis cuneiformibus incisim apice dentatis, umbellis sessilibus pedunculatisve, stylis in fructu incurvis. ② in paludibus et ad rivulos, in hortis cultum. — Fl. fr. 4. p. 339. — Cam. epit. 727. ic. (v. v. c.)

XL. ÆTHUSA *L. gen. n. 355.*

Calycis margo obsoletus; petala obovata emarginata cum lacinulâ inflexâ; fructus ovato-globosus; carpella jugis 5 elevatis crassis acutè carinatis, lateralibus marginantibus et paulò latioribus, carinâ sub-

latâ cinctis; valleculæ evittatæ; semen semiglobosum. — Involucrum nullum aut 1-phyllum, involucellum triphyllum.

1. *Æ. CYNAPIUM* (L. sp. 367.) caule ramoso tereti lævi, foliis 2-pinnatis, foliolis pinnatifidis lacinii linearibus indivisis incisive, invol. o aut 1-phylo caduco, involuclis 3-phyllis foliolis lineari-subulatis. ① in cultis frequens. — Fl. fr. 4. p. 293. — J. Bauh. hist. 3. p. 2. p. 180. ic. (v. v.)

XL. *TRINIA Hoffm. umb. 1. p. 92. — Pimpinellæ sp. L.*

Dioica vel dioico-polygama. Calycis margo obsoletus; petala masculæ plantæ lanceolata in lacinulam involutam coarctata subemarginata, fæminæ vel hermaphroditæ ovata breviter apiculata apiculo inflexo; fructus a latere compressus ovatus stylis reflexis coronatus; carpella jugis 5 filiformibus prominulis æqualibus, lateralibus marginantibus; valleculæ evittatæ vel obsoletè vittatæ, juga vero vittata; semen gibbo-convexum anticè planiusculum. — Involucrum varium; carpophorum planissimum membranaceum bifidum.

1. *T. GLABERRIMA* (Hoffm. l. c.) radice fusiformi, caulibus diffusis lævisculis, foliis pinnatis foliolis multifidis lacinii linearibus, umbellis numerosis simplicibus compositisque, involucri nullo aut 1-phylo. ② inter saxa montosa Galliæ orientalis australisque, Andegaviæ ex Bast. — *Pimpinella dioica* L. mant. 357. Fl. fr. 4. p. 282. — Cus. hist. 2. p. 200. f. 1. (v. v.)

XLII. *SISON Koch umb. 123. — Sisonis sp. L.*

Calycis margo obsoletus; petala subrotunda curvata profundè emarginata cum lacinulâ inflexâ; fructus a latere compressus ovato-globosus, stylis brevissimis inclinatis coronatus; carpella jugis 5 filiformibus æqualibus, lateralibus marginantibus; valleculæ univittatæ vittis abbreviatis clavulatis; semen gibbo-convexum anticè planiusculum. — Involucrum universale et parziale oligophyllum; carpophorum bipartitum.

1. *S. ANOMUM* (L. sp. 362.) caule substriato, foliis inferioribus pinnatis foliolis ovato-lanceolatis inciso-dentatis incisive serrulatis, superioribus pinnatifidis lacinii linearibus integerrimis, umbellis lateralibus terminalibusque pauciradiatis. ② ad sepes et in glareosis passim. — *Sium Anomum* Roth. Fl. fr. 4. p. 304. — Barr. ic. t. 1190. (v. s.)

XLIII. *AMMI Tourn. 2. 159. L. gen. n. 334.*

Calycis margo obsoletus; petala obovata cum lacinulâ inflexâ emarginato-biloba, lobis inæqualibus irregularia; fructus a latere compressus ovato-oblongus stylis reflexis coronatus; carpella jugis 5 filiformibus æqualibus, lateralibus marginantibus; valleculæ 1-vittatæ; semen tereti-convexum anticè planiusculum. — Involucrum involuclumque polyphyllum; carpophorum liberum bipartitum.

1. *A. MAJUS* (L. sp. 349.) caule glauco striato, foliis inferioribus pinnatis 2-pinnatisve, foliolis lanceolatis argutè serrulatis acutis, superioribus multifidis lacinii lanceolato-linearibus argutè serratis, involucris dissectis lacinii lineari-capillaribus, involuclorum foliolis lineari-lanceolatis aristato-acuminatis. ① in arvis passim. — Fl. fr. 4. p. 326. — Lob. ic. t. 721. f. 1. (v. v.)
2. *A. GLAUCIFOLIUM* (L. sp. 349.) caule substriato, foliis pinnatis, foliolis omnibus multifidis lacinii linearibus integris acuminatis, involucris dissectis lacinii lineari-capillaceis, involuclorum foliolis lineari-subulatis.

① in arvis passim. — Fl. fr. 4. p. 326. — J. Bauh. 3. p. 2. p. 58. An prioris var? (v. v. DC.)

3. A. *VISHAGA* (Lam. dict. 1. p. 132.) caule valdè folioso striato, foliis 2-3-pinnatis, foliolis omnibus multifidis, laciniis linearibus integris acutis, umbellâ maturatone contractâ, radiis in receptaculum subcarnosum coactis, involucri dissecis laciniis lineari-filiformibus, involucellorum foliolis basi coalitis lineari-subulatis umbellulam superantibus. ① in pascuis provinc. austr. occidentaliumque usque ad Andegaviam ex Bast. — Fl. fr. 4. p. 327. — Lob. ic. 726. f. 1. (v. s.)

#### XLIV. CNIDIUM Hoffm. umb. gen. 1. p. 157.

Calycis margo obsoletus; petala obovata emarginata cum lacinulâ inflexâ; fructus sectione transversali subteres; carpella jugis 5 æqualibus alatis alâ submembranaceâ, lateralibus marginantibus; valliculæ 1-vittatæ; semen semiteres. — Involucrum varium, involucellum polyphyllum; commissura 2-vittata.

1. C. *MOUNIERI* (Spreng. umb. prod. p. 39.) caule subsulcato tereti, foliis pinnatis, foliolis pinnatifidis, laciniis linearibus, involucri involucellisque polyphyllis, foliolis lineari-setaceis, carpellis ovatis. ① in provinciis australibus. — Koch nmb. f. 48. Selinum Mounieri L. Fl. fr. 4. p. 321. Jacq. vind. t. 62. (v. v. c.)

#### XLV. SESELI L. gen. n. 360.

Calycis margo 5-dentatus; petala obovata in lacinulam inflexam coarctata emarginata vel subintegra; fructus ovatus vel oblongus sectione transversali subteres, stylis reflexis coronatus; carpella jugis 5 prominulis filiformibus vel elevatis crassis corticosis, lateralibus marginantibus et sæpè paulò latioribus; valliculæ univittatæ, semen subsemiteres. — Involucrum varium, involucellum polyphyllum; valliculæ exteriores 1-rarius 2-vittatæ; commissura 2-aut rarius 4-vittata.

##### Sect. I. LIBANOTIS Lam. fl. fr. 3. p. 427.

Involucrum universale polyphyllum.

1. S. *LIBANOTIS* (Koch umb. p. 111.) caule glabro striato sulcatove, foliis radicalibus 2-pinnatis, caulinis pinnatis foliolis pinnatifidis laciniis incisis lobis acutis lanceolatis, involucri involucellisque polyphyllis foliolis setaceis, carpellis ovalibus hirtis. 2/ ad sylvas in montanis. — Athamanta Libanotis L. sp. 351. Fl. fr. 4. p. 317. Koch umb. f. 49 et 50. — Liban. montana All. ped. t. 62. (v. v.)

Ⅎ. *pubescens* (DC. fl. fr. 4. p. 317.) caule foliisque pubescentibus. — In saxosis apricis Pyren. et Sequanæ propè Rothomagum. — Athamanta pubescens Retz obs. 3. p. 28. Fl. fr. 5. p. 511. (v. s.)

##### Sect. II. HIPPOMARATHRUM.

Involucrum universale nullum vel oligophyllum.

2. S. *TORTUOSUM* (L. sp. 373.) caule rigido tortuoso divaricato ramosissimo tereti, foliis glaucis 3-pinnatis, foliolis lanceolato-linearibus acutis, involucri involucellorum foliolis lineari-lanceolatis acuminatis, floribus subsessilibus congestis, carpellis subhirtis. 2/ inter saxa Gallie australis. — Fl. fr. 4. p. 285. — J. Bauh. hist. 3. p. 2. t. 16. f. 1. (v. s.)
3. S. *HIPPOMARATHRUM* (L. sp. 373.) caule lævisculo, foliis 2-3-pinnatis, foliolis linearibus angustis acutis, involucri subnullo, involucellis foliolis connatis 1-phyllis, floribus subsessilibus congestis, carpellis glabris ovalibus.

- ℥* in Alsatie collibus aridis ex Schauenb. — Fl. fr. 4. p. 283 et 5. p. 502. Jacq. aust. t. 143. — Crantz austr. p. 205. t. 5. f. 1 et 2. (v. s. c.)
4. *S. ANNUUM* (L. sp. 373.) caule substriato, foliorum 2-3-pinnatorum vaginis emarginatis, foliolis linearibus angustis acutis, invol. nullo, involu-cellis polyphyllis foliolis lanceolatis acutis umbellulis longioribus, carpellis glabris. *℥* ad rupes et in pratis siccis passim. — Fl. fr. 4. p. 283. Jacq. aust. t. 55. *Selinum dimidiatum* Fl. fr. 4. p. 323. (v. s.)
5. *S. MONTANUM* (L. sp. 372.) caule substriato, foliorum 3-pinnatorum vaginis integris, foliolis linearibus angustis glaucis mucronatis, superioribus 1-phyllis, involucri 1-3-phylli caducei involucellorumque polyphyll. foliolis linearibus subulatis, carpellis glaucis ellipticis. *℥* in montosis aridis. — Fl. fr. 4. p. 284. *S. glaucum* L. — Vaill. bot. t. 5. f. 2. (v. s.)
6. *S. ELATUM* (L. sp. 375.) caule elongato læviusculo ramoso, geniculis callosis, foliis pinnatis 2-pinnatisve, foliolis angustis linearibus, invol. o, involucellis polyphyllis, foliolis linearibus subulatis, carpellis glabris ovalibus. *℥* in montosis et ad oras sylvarum Mompelii, Delphin. ex Will., Aurelia ex Dub., Andegaviæ ex Bast. — Fl. fr. 4. p. 285. Gon. ill. 16. t. 8. (v. s.)

XLVI. PTYCHOTIS Koch umb. p. 124.

Calyx 5-dentatus; petala obovata bifido-emarginata, medio plicâ transversali notata lacinulam emittente; fructus a latere compressus ovatus vel oblongus stylis reflexis coronatus; carpella jugis 5 filiformibus æqualibus, lateralibus marginantibus; valliculæ 1-vittatæ; semen tereti vel gibbo-convexum anticè planiusculum. — Involucrum nullum, involucellum polyphyllum. Genus vix a *Seseli* sejungendum.

1. *P. HETEROPHYLLA* (Koch l. c.) caule ramosissimo striato divaricato, foliis radicalibus pinnatis, foliolis ovatis pinnatifidis laciniis inciso-dentatis, caulinis duplicato-ternatis foliolis linearibus angustissimis, invol. nullo aut 1-phylo foliolo lineari setaceo, involucellis foliolis 1-3 linearibus setaceis, fructu oblongo, carpellis glaberrimis. (2) in lapidosis provinc. austral. — *Seseli Saxifragum* L. sp. 374. Fl. fr. 5. p. 503. *Æthusa Bunius* Murr. syst. veg. 246. Fl. fr. 4. p. 293. *Juga vittata*? (v. v.)
- P. VERTICILLATA* (Duby mss.) caule erecto aut diffuso læviusculo, foliis pinnatis segmentis multipartitis lobis setaceis in foliis radicalibus brevioribus, invol. o, involucellis polyphyllis foliolis setaceis flores subæquantibus, fructu ovato, carpellis glaberrimis. (2) in Corsicâ. — *Seseli verticillatum* Desf. atl. 1. p. 260. Fl. fr. 5. p. 504. *Ses. ammoides* L? sp. 373 certò non Jacq. vind. t. 52. — Dalech. hist. 665. f. 2. (v. s.)

XLVII. HELOSCIADIUM Koch umb. p. 125. — *Sii* et *Sisonis* sp. L.

Calyx margo 5-dentatus subindè obsoletus; petala ovata integra acuta vel obtusiuscula et apiculata apice recto vel inflexo; fructus a latere compressus ovatus vel oblongus; carpella jugis 5 filiformibus prominulis æqualibus, lateralibus marginantibus; valliculæ 1-vittatæ; semen gibbo-vel tereti-convexum anticè planiusculum. — Involucrum varium, carpophorum integrum liberum.

1. *H. AULASTUM* (Koch l. c.) caule procumbente gracili, foliis pinnatis, foliolis profundè tripartitis subverticillatis, laciniis linearibus, umbellis longè pedunculatis multiradiatis, invol. involucellisque sub 5-phyllis foliolis lanceolatis. (1) in stagnis circâ Aquas Tarbellicas (Dax). — *Sium intermedium* DC. fl. fr. 4. p. 302. (v. s.)
2. *H. INUNDATUM* (Koch l. c.) caule repente fistuloso, fol. inferioribus multipartito-capillaceis superioribus pinnatis, foliolis cuneatis incisais lanceolatisve, umbellis pedunculatis 2-3-radiatis, involucris nullis, involucellis polyphyllis.

foliis lineari-lanceolatis brevibus, fructibus ovatis, stylis brevissimis incli-  
natis. (1) *P* in fossis et paludibus passim. — *Sium inundatum* Fl. fr. 4. p. 303.  
Sison inundatum L. — Pluck. t. 61. f. 3. (v. v.)

3. *H. repens* (Koch l. c.) caule repente striato fistuloso, foliis pinnatis foliolis  
dentatis incisive indivisisque, umbellis pedunculatis oppositifoliis, involuclis  
umbellulam subæquantibus involucrisque sub-6-phyllis foliolis lineari-lan-  
ceolatis, fructibus globosis, stylis reflexis. *℥* in paludosis et inundatis tur-  
fosis Galliz occident., mediz, borealis orientalisque. — *Sium repens* Fl. fr. 4.  
p. 300. Jacq. aust. 3. t. 260. (v. v. DC.)
4. *H. nodiflorum* (Koch l. c.) caule procumbente striato fistuloso, foliis pin-  
natis, foliolis lanceolatis serratis, umbellis oppositifoliis subsessilibus, invo-  
luclero 1-2-phylo, involucellis umbellulam subæquantibus foliolis lanceolatis,  
fructibus ovato-globosis, stylis reflexis. *℥* in rivulis et ad margines fluvio-  
rum. — *Sium nodiflorum* L. sp. 361. Fl. fr. 4. p. 299. — Moris. s. 9. t. 5.  
n. 7. (v. v.)

*ℓ. giganteum* (Des Moulins) caule 2-3-pedali, foliorum radicalium foliolis  
ovatis crenato-dentatis 2-pollicaribus. — Propè Burdigalam observavit  
cl. Des Moul. (v. s.)

**XLVIII. FENICULUM** *All. ped. 2. p. 25. Cært. fruct. 1. p. 105. t. 23. f. 5. —*  
*Anethi sp. L.*

Calycis margo obsoletus; petala subrotunda integra involuta lacinulâ  
subquadratâ retusâ; fructus sectione transversali subteres; carpella  
jugis 5 prominulis obtusè carinatis, lateralibus marginantibus et paulò  
lterioribus; vallecule univittatæ; semen subsemiteres. — Involucrum  
involucellumque nullum.

1. *F. officinale* (All. ped. n. 1359.) caule tereti ramoso striato lucido, foliis de-  
compositis, lacinii capillaceis, carpellis oblongo-ellipticis. (2) in lapidosis  
aridis. — *Anethum Feniculum* L. sp. 377. Fl. fr. 4. p. 340. — Moris. s. 9.  
t. 2. n. 1. (v. v.)

**XLIX. OENANTHE** *Lam. fl. fr. 3. p. 432. DC. fl. fr. 4. p. 295. — Oenanthe*  
*et Phellandrium L.*

Calycis margo 5-dentatus; petala obovata emarginata cum lacinulâ  
inflexâ; fructus sectione transversali subteres, stylis erectis coronatus; carpella  
jugis 5 convexiusculis obtusis, lateralibus marginantibus et  
paulò lterioribus; vallecule 1-vittatæ; semen tereti-convexum. — Invol-  
varium, involucellum polyphyllum. Flores radii pedicellati abortu ste-  
riles, disci sessiles vel brevissimè pedicellati fertiles; carpophorum  
nullum.

1. *OE. PHELLANDRIUM* (Lam. fl. fr. 3. p. 432.) caule fistuloso striato, foliis 3-pin-  
natis, foliolis decompositis lacinii divaricatis linearibus abbreviatis, invol. o,  
involucellis polyphyllis foliolis lineari-setaceis acuminatis umbellulâ breviori-  
bus, fructu ovato-elongato. *℥* in stagnis et fossis aquosis. — Fl. fr. 4. p. 295.  
*Phellandrium aquaticum* L. — Lob. ic. 735. f. 1. (v. v.)
2. *OE. FISTULOSA* (L. sp. 365.) radice repente, caule fistuloso striato lucido, fo-  
liis radicalibus decompositis lacinii linearibus abbreviatis, caulinis pinnatis,  
foliolis pinnatifidis indivisive lacinii lanceolatis acutis, involuclero nullo aut  
foliolo unico lineari-lanceolato umbellam superante, involucellis polyphyllis  
foliolis lineari-lanceolatis umbellulam subsuperantibus. *℥* in paludosis fre-  
quens. — Fl. fr. 4. p. 295. Cam. epit. p. 611. fig. ext. Juga connata pericar-  
pium crasse corticosum reddunt. (v. v.)
3. *OE. PIMPINELLOIDES* (L. sp. 365. non Poll.) radicem tuberculis oblongis,  
caule firmo sulcato, foliis radicalibus 2-pinnatis, foliolis pinnatifidis lacinii



integræ aut unidentatis lanceolatis acuminatis, caulinis pinnatis foliis linearibus integræ elongatis, involuclis umbellulam subæquantibus involucrique polyphyllis, foliolis lineari-setaceis.  $\mathcal{U}$  in pratis paludosis Galliæ mediæ, occidentalis australisque. — Fl. fr. 4. p. 297. — J. Bauh. hist. 3. p. 2. p. 191. f. 2. (v.  $\gamma$ .)

4. OE. *CHEEROPHYLLOIDES* (Pourr. aot. tol. 3. p. 323.) radicem fibris apicem versus tuberosis, caule sulcato, foliis radicalibus 2-pinnatis, foliolis ovato-cuneatis pinnatifidis laciniis mucronulatis, caulinis pinnatis 2-pinnatisve, foliolis linearibus integræ elongatis acuminatis, involuclis umbellulam subsuperantibus involucrique polyphyllis foliolis lineari-setaceis, fructibus cylindrico-elongatis.  $\mathcal{U}$  circa Narbonam, Valtetiam, Andegaviam. — Fl. fr. 5. p. 507. — Cam. epit. 610. f. III. OE. *pimpinelloides*  $\beta$ . Fl. fr. 4. p. 297. An satis à præcedente distincta? (v. s.)
5. OE. *LACHENALII* (Gmel. bad. 1. p. 678.) radicem fibris cylindræis, caule sublevi, foliis radicalibus pinnatis, foliolis ovalibus obovatisve sæpè 3-lobis, caulinis pinnatis 2-pinnatisve foliolis linearibus subacutis, involucri 4-6-phylli involuclorum polyphyll. foliolis linearibus umbellulâ brevioribus.  $\mathcal{U}$  in Alatiâ. — OE. *Rhenana* Fl. fr. 5. p. 506. Folia radicalia sæpè plantâ florente desunt. (v. s.)
6. OE. *APPROXIMATA* (Mer. par. 115.) radicem fibris cylindræis, caule sublevi, foliis radicalibus pinnatis, foliolis obtusè trilobis, caulinarum lobis linearibus, involuero nullo, involuclis polyphyllis foliolis lineari-lanceolatis acutis umbellulam subæquantibus.  $\mathcal{U}$  in humidis circa Parisios. — Fl. fr. 5. p. 507. (v. s.)
7. OE. *EXUCEDANIFOLIA* (Pol. pal. 1. p. 289. t. 2. f. 3.) tuberibus radicalibus ellipticis, caule sulcato, foliis pinnatis 2-pinnatisve, foliolis omnibus linearibus acutis, involuero subnullo, involuclis polyphyllis foliolis lineari-lanceolatis acutis umbellulâ duplò brevioribus, fructibus cylindricis.  $\mathcal{U}$  in fossis et paludibus Galliæ orientalis, mediæ et boreali-orientalis. — Fl. fr. 4. p. 297. — Dalech. lugd. 773. f. 2. (v. v. DC.)
8. OE. *GLOBULOSA* (L. sp. 365.) radice fusiformi ramosâ, caule vix striato, foliis pinnatis foliolis pinnatifidis radicalium laciniis lanceolatis subobtusis, caulinarum laciniis linearibus acutis, invol. o, involuclis polyphyllis foliolis lineari-setaceis umbellulam subsuperantibus, fructibus ovato-globosis.  $\mathcal{U}$  in stagis provinciarum australiorum. — Fl. fr. 4. p. 296. Gou. ill. t. 9. (v. s.)
9. OE. *CROCATA* (L. sp. 365.) caule striato, foliorum 2-3-pinnatorum foliolis omnibus cuneatis, caulinarum ovatis grossè dentatis incisive, superiorum lanceolatis, involucri involuclisque polyphyllis foliolis lineari-setaceis.  $\mathcal{U}$  in aquis et ad ripas fluviorum Galliæ præsertim mediæ et occidentalis. — Fl. fr. 4. p. 298. — Moris. s. 9. t. 7. n. 2. (v. v.)
10. OE. *APIFOLIA* (Brot. lusit. 1. p. 420.) caule sulcato, foliis radicalibus 2-pinnatis, foliolis pinnatis laciniis ovato-cuneatis incisè dentatis, caulinis pinnatis foliolis cuneatis pinnatifidis laciniis linearibus, invol. nullo, involuclis polyphyllis foliolis lineari-setaceis umbellulâ duplò brevioribus, fructibus elongato-cylindricis.  $\mathcal{U}$  ad aquas Corsicæ. (v. s.)

L. *CICUTA* L. gen. n. 354.

Calycis margo 5-dentatus foliolosus; petala obovata emarginata cum lacinulâ inflexâ; fructus subrotundus a latere contractus didymus; carpella jugis 5 planiusculis æqualibus lateralibus marginantibus; vallecule 1-vittatæ; vittæ (in sicco carpello) jugis elevatiores; semen teres. — Involucrium oligophyllum aut nullum, involuclum polyphyllum; carpophorum 2-partitum.

1. C. *VIROSA* (L. sp. 368.) caule fistuloso sulcato, foliis 2-pinnatis pinnatis simplicibusve foliolis 3-5-lobis, laciniis lineari-lanceolatis inciso-dentatis,

umbellis oppositifoliis, involucellis polyphyllis foliolis setaceis.  $\mathcal{U}$  ad stagna et fossas aquosas. — Koch umb. f. 53 et 54. Cicutaria aquatica Lam. DC. fl. fr. 4. p. 294. — Lob. ic. t. 208. f. 2. (v. v.)

Subdiv. II. SESELINEÆ NUCLEATÆ, id est semine libero nec pericarpio nisi latere commissurali adnato.

LI. CRITHMUM Tourn. t. 165. L. gen. n. 340.

Calycis margo obsoletus; petala subrotunda integra involuta lacinulâ obovatâ; fructus sectione transversali subteres; carpella jugis 5 elevatis argutis subalatis, lateralibus paulo latioribus marginantibus; semen semiteres liberum vittis copiosis undique tectum. — Involucrum et involucellum polyphyllum; pericarpium spongiosum.

1. C. MARITIMUM (L. sp. 354.) foliolis lanceolatis carnosius.  $\mathcal{U}$  inter saxa maritima. — Fl. fr. 4. p. 317. — Lob. ic. 39a. f. 4. (v. v.)

Trib. X. SCANDICINEÆ. Koch umb. p. 130.

Fructus a latere evidenter compressus vel contractus, lineari-elongatus, ut plurimum rostratus. Carpella paucijugata, jugis primariis quinque filiformibus subindè subalatis, lateralibus marginantibus, omnibus æqualibus, secundariis nullis; vel jugis oblitteratis lævigata et apice tantum jugata. Semen tereti-convexum, anticè profundè sulcatum vel margine involutum et idèd carpella campyloperma.

Sect. 1. SCANDICINEÆ SOLIDÆ EROSTRATÆ.

LII. CHÆROPHYLLUM L. gen. n. 358.

Calycis margo obsoletus; petala obovata emarginata cum lacinulâ inflexâ; fructus a latere compressus vel contractus; carpella jugis quinque obtusis 4-æqualibus lateralibus marginantibus; commissura sulco profundo exarata; valleculæ univittatæ; semen tereti-convexum sectione transversali semilunare. — Involucrum nullum vel oligophyllum, involucellum polyphyllum; carpophorum bifidum.

1. C. HIRSATUM (L. sp. 371.) caule striato petiolis nervisque hirsutis, foliis 2-pinnatis pinnatisve, foliolis pinnatifidis lacinulis lanceolatis incis, invol. nullo aut 1-2-phylo, involucelli polyphylli foliolis lanceolatis membranaceis ciliatis acuminatis, jugis obtusis depressis, stylis longissimis suberectis.  $\mathcal{U}$  in humidis montium. — Fl. fr. 4. p. 289. Jacq. austr. t. 148. (v. v.)

2. C. AUREUM (L. sp. 370.) caule striato, foliis 3-2-pinnatis, foliolis lanceolato-elongatis incis serratisque summis confluentibus, radicalium petiolis nervisque hirsutis, involucri nullo aut 1-phylo, involucellis polyphyllis foliolis hirsutis lanceolato-linearibus aristato-acuminatis, jugis obtusis subelevatis, stylis divergentibus.  $\mathcal{U}$  in umbrosis montium. — Fl. fr. 4. p. 289. — Lob. ic. 734. f. 2. (v. v.)

3. C. TEMULUM (L. sp. 370.) caule lævisculo hirto, geniculis tumidinasculis, foliis pubescentibus inferioribus 2-pinnatis superioribus pinnatis, foliolis ovatis lanceolatisve crenatis incis, summis confluentibus, invol. 0 aut 1-phylo, involucellis polyphyllis foliolis hirsutis lineari-lanceolatis acutis, stylis semidivergentibus brevibus.  $\mathcal{U}$  in locis incultis et sepibus. — Koch umb. f. 63. Fl. fr. 4. p. 290. — Tabern. ic. 94. (v. v.)

4. *C. suberosum* (L. sp. 370.) caule sub nodis inflato basi hirtio cæterum glabro fistuloso, foliis 3-2-pinnatis, foliolis pinnatifidis laciniis lineari-lanceolatis integerrimis, involucri nullo, involucellis polyphyllis foliolis linearibus acuminatis, stylis brevibus recurvis. ② in Alsatiæ dumetis ruderalisque ex J. Bauh. — Fl. fr. 5. p. 505. Koch umb. f. 61 et 62. — J. Bauh. hist. 3. p. 183. ic. (v. s. ex Germ.)
5. *C. nodosum* (Lam. enc. 1. p. 685.) caule hirtio scabro geniculis tumidiusculo, foliis inferioribus 2-pinnatis superioribus pinnatis, foliolis ovatis pinnatifidis incisive dentatis, umbellis 2-3-radiatis, invol. nullo, carpellis hirsutis, jugis obtusis valde depressis, stylis brevissimis erectis. ① in sepiibus et umbrosis circa Nicæam ex All., Parisios ex Lam., Andegaviam ex Bast. — *Caulis nodosa* Fl. fr. 4. p. 334. An potius sequentis generis? (v. s. c.)

## Sect. II. SCANDICINÆ SOLIDÆ ROSTRATÆ.

LIII. *ANTHRISCUS* Pers. syn. 1. p. 320. Hoffm. umb. 1. p. 38. — Scandicis et Chærophylli sp. L.

Calycis margo obsoletus; petala obovata truncata vel emarginata cum lacinulâ inflexâ, sæpè brevissima; fructus a latere contractus rostratus sectione verticali lanceolatus; carpella subteretia ejugata rostro solum quinquejugata; semen tereti-convexum anticè profunde sulcatum. — Involucrum nullum, involucellum polyphyllum; carpophorum apice bifidum.

\* *Carpellis lævibus.*

1. *A. sylvestris* (Hoffm. umb. gen. 1. p. 40.) caule sulcato ramoso, geniculis tumidis, foliis inferioribus 3-pinnatis, superioribus 2-pinnatis, foliolis lanceolatis incisim incisivo-dentatisve acutis, summis confluentibus, involucellis polyphyllis foliolis lanceolatis ciliatis, fructibus lucidis, stylis erectis subdivergentibus. ½ in pratis et ad sepes frequens. — Koch umb. f. 57 et 58. Chærophyllum sylvestre L. sp. 369. Fl. fr. 4. p. 288. — Dod. pempt. 701. (v. v.)
2. *A. torquata* (Duby mss.) caule striato ramoso glabro, foliis 2-pinnatis, foliolis lanceolato-ovatis grossè subinciso-dentatis, involucellis polyphyllis foliolis ciliato-hirsutis lanceolatis acutis, summis confluentibus, petalis valde inæqualibus exterioribus radiantibus, fructibus lucidis elongato-lanceolatis ciliorum minimorum seriè basi cinctis, stylis brevibus suberectis. ② in Alpibus Galloprovinciæ propè Colmars detexit cl. DC. — Chærop. torquatum Fl. fr. 5. p. 505. Myrrhis bulbosa All. ped. n. 1373. excl. syn. (v. v. DC.)
3. *A. ? Cicutaria* (Duby mss.) caule striato glabro, foliis inferioribus 2-pinnatis, superioribus pinnatis, foliolis elongato-lanceolatis incisim aut incisivo-dentatis acutis, summis confluentibus, involucellis lanceolato-linearibus ciliatis acuminatis, petalis valde inæqualibus exterioribus radiantibus, stylis divergentibus. ½ secus rivulos Alpium Delph. ex Vill. — Chærophyllum Cicutaria Vill. Fl. fr. 5. p. 506. Hujus generis, quantum videre licet per fructus nondum maturos. (v. s. ex Helv.)
4. *A. cerefolium* (Hoff. umb. 1. p. 41.) caule substriato glabro, foliis inferioribus 3-pinnatis, superioribus 2-pinnatis, foliolis ovatis incisivo-pinnatifidis laciniis lanceolatis, umbellis lateralibus subsessilibus, involucellorum foliolis lineari-lanceolatis acuminatis, fructibus elongato-linearibus, stylis brevibus erectis parallelis. ① in hortis colitur. — Chærophyllum sativum Lam. fl. fr. 3. p. 438. DC. fl. fr. 4. p. 291. (v. v. c.)

\* \* *Carpellis muricatis vel asuleolatis.*

5. *A. vulgaris* (Pers. syn. 1. p. 320.) caule striato glabro, foliis 3-4-pinnatis, foliolis pinnatifidis laciniis mucronulatis brevissimè ciliatis, umbellis lateralibus sessilibus pedunculatisque, involucellorum foliolis lineari-lanceolatis, fructibus aculeolatis oblongo-ovatis basi ciliorum seriè cinctis, sty-

lis brevissimis rectis. ① ad sepes et in marginibus arborum. — Koch umb. f. 59 et 60. *Caucalis scandinavica* Fl. dan. t. 863. Fl. fr. 4. p. 334. (v. v.)

LIV. SCANDIX *L. gen. n. 357.*

Calycis margo obsoletus; petala obovata truncata cum lacinulâ inflexâ; fructus a latere compressus longissimè rostratus; carpella jugis 5 obtusis æqualibus, lateralibus marginantibus; valleculæ evittatæ vel obsoletè vittatæ; semen tereti-convexum anticè profunde sulcatum. — Involucrum nullum vel oligophyllum, involucellum 5-7-phyllum; carpophorum apice furcatum.

Sect. I. PÆTEN.

Rostrum fructûs a dorso compressum.

1. *S. PÆTEN-VENERIS* (L. sp. 368.) foliis pinnatis, foliolis multifidis laciniiis linearibus acutis, umbellâ 2-3-radiatâ, invol. nullo aut umbellis folio superiore suffultis, involucellis erectis polyphyllis foliolis lanceolatis acuminatis, carpellis dorso rostrique margine brevissimè aculeolato-hispidis, stylis erectis parallelis. ① inter segetes frequens. — Fl. fr. 4. p. 291. — Cam. epit. 304. ic. (v. v.)

Sect. II. WYLIA *Hoffm. umb. 1. p. 5.*

Rostrum fructûs a latere compressum.

2. *S. AUSTRALIS* (L. sp. 369.) caulibus gracilibus, foliis pinnatis foliolis multifidis laciniiis capillaribus, umbellâ 2-3-radiatâ, involucro nullo, involucelli foliolis lanceolatis margine scariosis integris aut apice 2-3-fidis, carpellis attenuatis rostrique dorso aculeolato-hispidis, stylis erectis parallelis. ① in aridis provinc. mediterranearum. — Fl. fr. 4. p. 292. — Moris. s. g. t. 11. n. 4. (v. v.)

Sect. III. SCANDICINÆ, UTRICULOSÆ, id est pericarpio dorso inflato, anticè semine adnato.

LV. MYRRHIS *Scop. carn. ed. 2. sp. 207. Hoffm. umb. 1. p. 27.*

Calycis margo obsoletus; petala obovata emarginata cum lacinulâ inflexâ; fructus a latere compressus; semen involutum tectum pericarpio-è membranâ duplici formato, externâ in juga quinque æqualia arguta acutè carinata intus cava elevatâ, internâ arcte adnatâ; vittæ nullæ. — Involucrum nullum, involucellum polyphyllum; carpophorum apice fissum.

1. *M. ODORATA* (Scop. carn. n. 341.) caule striato, foliis pubescentibus 3-pinnatis, superioribus 2-pinnatis, foliolis pinnatifidis laciniiis ovatis lanceolatisve incisis, petiolis nervisque hirsutis, involucellorum foliolis lanceolato-linearibus membranaceis acuminatis. ② in pratis montanis. — Chærophyllum odoratum Lam. DC. fl. fr. 4. p. 290. — Cam. epit. 898. ic. (v. v.)

Trib. XI. SMYRNEÆ *Koch umb. p. 133.*

Fructus a latere compressus vel contractus turgidus. Carpella campylosperma paucijugata jugis primariis quinque, lateralibus marginantibus vel ante marginem positis, secundariis nullis; vel jugis ferè oblitteratis sublævigata. Semen involutum vel sulco profundo exaratum, semilunare vel complicatum.

Sect. 1. SMYRNEÆ SOLIDÆ MULTIVITTATÆ.

LVI. SMYRNIUM *Koch l. c. — Smyrni sp. L.*

Calycis margo obsoletus; petala lanceolata vel elliptica integra acu-

minata acuminis inflexo; fructus a latere contractus, carpellis reniformi-globosis didymus; juga 3 dorsalia prominula arguta, duo lateralia marginantia oblitterata; valliculæ multivittatæ; semen involutum. — Involucra varia; carpophorum bipartitum.

1. *S. OLUSATUM* (L. sp. 376.) caule striato, foliis 2-ternatis ternatisque, foliolis rotundo-ovatis basi cuneatis crenato-dentatis.  $\frac{1}{2}$  in pascuis humidis umbrosis Galliae australis, Aginni ex St.-Am., Bardigalæ ex Des Moulins, Andegavæ ex Bast., Parisiorum ex Mer. — Fl. fr. 4. p. 340. — Lob. ic. t. 708. f. 2. (v. v. DC.)
2. *S. FRAXOLATUM* (L. sp. 376.) radice tuberosâ, caule erecto sulcato in parte superiore 2-alato, foliis inferioribus ternatis foliolis rotundo-ovatis crenatis, superioribus simplicibus basi cordatis amplexicaulibus rotundo-ovatis crenatis integrisve.  $\frac{1}{2}$  in sylvis Corsicæ et Galloprovinciæ propè Telsonem. — Fl. fr. 5. p. 513. *S. Dioscordis* Spr. Koch umb. f. 38 et 39. — Dod. pempt. 698. f. 2. Moris. s. 9. t. 4. n. 7. (v. s.)

Sect. II. SMYRNEÆ SOLIDÆ PAUCIVITTATÆ VEL EVITTATÆ.

LVII. CONIUM L. gen. n. 336.

Calycis margo obsoletus; petala obovata emarginata cum lacinulâ inflexâ; fructus a latere compressus ovatus; carpella jugis 5 prominulis undulato-crenulatis æqualibus, lateralibus marginantibus; valliculæ multistriatæ evittatæ; semen sulco profundo angusto incisum. — Involucrum oligophyllum; involucellum triphyllum dimidiatum; carpophorum apice bifidum.

1. *C. MACULATUM* (L. sp. 349.) caule erecto fistuloso ramoso maculato, foliis 3-2-pinnatis, foliolis lanceolatis pinnatifidis incisive summis confluentibus. (2) ad sepes humidas. — Cicuta major Lam. DC. fl. fr. 4. p. 324. — Moris. s. 9. t. 6. n. 9. (v. v.)

LVIII. ECHINOPHORA Tourn. inst. t. 423. L. gen. n. 329.

Calycis margo 5-dentatus; petala obovata emarginata cum lacinulâ inflexâ, vel exteriora majora bifida; flores radii masculi longè pedicellati; femineus centralis solitarius; fructus ovatus subteres receptaculo cavo inclusus breviter rostratus rostro emerso; carpella jugis quinque depressis undulato-striatis æqualibus; valliculæ univittatæ, vittis membranâ arachnoideâ tectis. — Involucrum involucellamque polyphyllum.

1. *E. SPINOSA* (L. sp. 344.) caule ramoso diffuso, foliis crassis coriaceis 2-pinnatis foliolis subulato-spinosis.  $\frac{1}{2}$  in littoribus maris Mediterranei. — Fl. fr. 4. p. 352. — Lob. ic. 710. f. 1. (v. v.)

Sect. III. SMYRNEÆ NUCLEATÆ MULTIVITTATÆ.

LIX. CACHRYS Koch umb. 130. — Cachrydis sp. L.

Calycis margo obsoletus; petala ovato-lanceolata integra acuminata acuminis inflexo; fructus turgidus, sectione transversali subteres; carpella jugis 5 crassis obtusissimis æqualibus, lateralibus marginantibus; semen liberum vittis copiosis tectum involutum. — Involucrum involucellamque oligophyllum; pericarpium crassum suberosum.

1. *C. LEVIGATA* (Lam. dict. 1. p. 256.) foliis supra-decompositis glabris, lacinis setaceis brevibus, involucri involucelloræque foliolis lineari-setaceis.  $\frac{1}{2}$  in rupestribus maritimis provinciæ mediter. — Fl. fr. 4. p. 344. *C. Morisonii* All. — Moris. umb. t. 3. n. 1. (v. v. DC.)

## DICOTYL. seu EXOGENÆ.

## Sect. IV. SMYRNÆE UTRICULOSÆ.

## LX. PLEUOSPERMUM Hoffm. umb. p. 9.

Calycis margo 5-dentatus; petala obovata integra; fructus a latere compressus ovatus; pericarpium carpellorum è membranâ duplici formatum, externâ (seu epicarpio) in juga 5 alata cava inflatâ, internâ (endocarpio) arcte adnatâ et in juga 5 minora exterioribus opposita elevatâ; valliculæ endocarpii univittatæ; semen semilunare. — Involucrum involucellumque polyphyllum.

1. *P. AUSTRIACUM* (Hoffm. l. c.) caule erecto striato fistuloso, foliis ternatis pinnatisque foliolis pinnatifidis, laciniis lanceolatis incisissimis dentatis basi decurrentibus, involucris polyphyllis foliolis incisissimis. 2/ ad aquas iug montanis Galloprov., Delph. ex Vill. — *Ligusticum Austriacum* L. sp. 360. Fl. fr. 4. p. 307. L. Gmelini Vill. dauph. 2. t. 13. bis. — Clus. hist. 2. p. 193. f. 1. (v. s.)

## Subord. III. UMBELLIFERÆ IMPERFECTÆ.

## Trib. XII. SANICULÆE Koch umb. 138.

Fructus sectione transversali subteres. Carpella jugis quinque primariis æqualibus, secundariis nullis vel jugis oblitteratis tecta squamis paleiformibus aculeisve. Semen sectione transversali semiteres, antice planiusculum. Umbellulæ fasciculatæ vel capitatæ, vel tota umbella capitata.

## Sect. I. SANICULÆE UTRICULOSÆ.

## LXI. ASTRANTIA Tourn. t. 166. L. gen. n. 327.

Calycis margo 5-dentatus foliolosus; petala erecta conniventia oblongo-obovata a medio infracta in lacinulam longitudine ferè petali; fructus a dorso subcompressus; carpella jugis quinque elevatis obtusis plicato-dentatis inflatis; in cavitate juga minora fistulosa includentibus; vittæ nullæ; semen semiteres. — Involucrum varium, involucellum polyphyllum; carpophorum nullum.

1. *A. MAJOR* (L. sp. 339.) foliis radicalibus 5-lobis lobis trifidis acutis dentatis, involucris 3-fidis laciniis lineari-lanceolatis setoso-serratis, involucellorum foliolis lanceolatis integris aut apice 3-dentatis umbellulam superantibus. 2/ in pascuis montanis. — Fl. fr. 4. p. 353. Lam. ill. t. 191. f. 1. Koch umb. f. 42 et 43. (v. v.)
2. *A. MINOR* (L. sp. 340.) foliis radicalibus digitatis, foliolis septenis novenisve lanceolatis profundè dentatis, involucellorum foliolis lanceolatis integris acuminatis umbellulam subsequantibus. 2/ in pascuis Alpium et Pyren. — Fl. fr. 4. p. 353. Lam. ill. t. 191. f. 2. (v. v.)

## Sect. II. SANICULÆE SOLIDÆ.

## LXII. SANICULA Tourn. t. 173. L. gen. n. 326.

Calycis margo 5-dentatus foliolosus; petala erecta conniventia obovata emarginato-infracta in lacinulam longitudine petali; fructus sectione transversali teres subglobosum; carpella aculeis uncinatis dense tecta ejugata multivittata; semen semiglobosum. — Involucrum involucellumque polyphyllum; carpophorum nullum.

1. *S. EUROPEA* (L. sp. 339.) caule nudo, foliis palmatis 3-5-lobis lobis cuneatis 3-fidis dentatis, floribus sessilibus. 2/ in sylvis et æpibus. — Fl. fr. 4. p. 354. — Cam epit. 763. ic. (v. v.)

LXIII. ERYNGIUM *Tourn. t. 173. L. gen. 324.*

Calycis margo quinquedentatus foliolosus; petala erecta conniventia oblongo-obovata emarginato-infracta in lacinulam longitudine petali; fructus sectione transversali subteres obovatum; carpella squamis paleiformibus tecta ejugata evittata; semen subsemiteres. — Involucrum polyphyllum; flores paleis intermixti in capitulum collecti, carpophorum nullum.

1. *E. CAMPESTRE* (L. sp. 337.) caule ramosissimo, foliis spinosis, radicalibus bipinnatis foliolis decurrentibus ovatis variè contortis, involucri capitulum longe superantis foliolis linearibus spinosis.  $\frac{1}{2}$  in incultis. — Fl. fr. 4. p. 355. — Fuchs. hist. 297. ic. (v. v.)
2. *E. BOURGATI* (Gou. ill. p. 7. t. 3.) caule stricto, foliis radicalibus orbiculatis ternatis foliolis pinnatifidis lacinisque incisus divaricatis spinosis, involucri capitulum conicum superantis foliolis lanceolato-linearibus spinosis.  $\frac{1}{2}$  in herbosis Pyrenaeorum. — Fl. fr. 4. p. 355. Caulis amethystinus. (v. v.)
3. *E. SPIRA-ALBA* (Vill. dauph. 2. t. 17.) foliis omnibus digitatis, involucri capitulum ovatum maximum superantis foliolis linearibus pinnatifidis.  $\frac{1}{2}$  in lapidosis aridis montium Galloprovinciæ et Delph. — Fl. fr. 4. p. 356. — Dalech. Ingd. p. 1462. f. 1. (v. s.)
4. *E. ALPIMUM* (L. sp. 337.) foliis radicalibus profundè cordatis lanceolatis serratis, superioribus 5-3-lobatis, involucri capitulum oblongo-ovatum superantis foliolis linearibus pinnatifido-spinosis.  $\frac{1}{2}$  in pascuis Jurassi et Alpium Delph. et Gallopr. — Fl. fr. 4. p. 356. — Lob. ic. 2. p. 23. f. 2. (v. s.)
5. *E. MARITIMUM* (L. sp. 337.) foliis radicalibus subrotundis plicatis grosse dentato-spinosis, superioribus incisus, involucri capitulum ovatum superantis foliolis ovatis acuminatis spinosis.  $\frac{1}{2}$  in arenosis maritimis. — Fl. fr. 4. p. 355. — Cam. epit. 448. ic. (v. v.)
6. *E. PLANUM* (L. sp. 336.) foliis radicalibus elliptico-cordatis crenatis superioribus 5-6-lobatis, involucri capitulum rotundum superantis foliolis lanceolato-linearibus spinosis.  $\frac{1}{2}$  in Alpibus Galloprov. Italiæ vicinis. — Fl. fr. 4. p. 357. — Clus. hist. 2. p. 158. f. 1. (v. s. c.)
7. *E. MICHOTOMUM* (Desf. atl. 1. t. 55.) caule ramoso dichotomo, foliis radicalibus oblongis cordatis crenatis, superioribus 3-5-lobatis, involucri capitulum rotundum longe superantis foliolis linearibus acuminatis simplicibus tricuspidatisque.  $\frac{1}{2}$  Monspelii ex seminibus cum lanis Barbariæ allatum. — Fl. fr. 5. p. 515. (v. s.)

Trib. XIII. *HYDROCOTYLE* Spreng. in *Röm. et Sch.* 6. p. xxxii. Koch umb. p. 141.

Fructus a latere evidenter compressus. Carpella jugis primariis 5 subinde obsoletis, secundariis nullis. Semen antice planiusculum. Umbellæ simplices vel imperfectæ floribus capitatis vel verticillatis. Petala patentia integra acuta apice recto vel subinflexo.

LXIV. HYDROCOTYLE *Tourn. t. 173. L. gen. n. 325.*

Calycis margo obsoletus, petala ovata integra acuta apice recto, fructus a latere plano-compressus biscutatus; carpella jugis 5 filiformibus carinali et lateralibus sæpè obsoletioribus, intermediis duobus arcuatis, evittata; semen carinato-compressum.

1. *H. vulgare* (L. sp. 338.) foliis peltatis orbiculatis crenatis, umbellæ 5-8. florâ.  $\frac{1}{2}$  in paludibus. — Fl. fr. 4. p. 358. Koch umb. f. 64 et 65. — Lob. ic. 387. f. 1. (v. v.)

Ordo LI. CAPRIFOLIACEÆ Juss. gen. 210. excl. gen. *Loranthearum*.

**Calyx** 1-sepalus ovario adhærens sæpè basi bibracteatus, limbo integro aut diviso. **Corolla** regularis nunc 1-petala 4-5-fida; nunc 4-5-petala petalis basi latioribus. **Stamina** corollæ lobis numero æqualia, et iisdem alterna, in monopetalis corollæ, in polypetalis calyci inserta. **Stylus** 0 aut 1; **stigma** 1-3. **Bacca** aut capsula limbo calycino sæpiùs coronata uni-aut multilocularis, loculis 1-∞-spermis. **Sem.** spermodermis ossea. **Embryo** rectus in apice albuminis carnosus; **radicula** supera. — **Frutices** foliis oppositis alternisve exstipulatis, floribus sæpiùs corymbosis.

**Trib. I. HEDERACEÆ.** — *Hederaceæ* Ach. *Rich. bot. med.* p. 449.

**Calyx** ebracteatus. **Corolla** polypetala. **Stylus** 1. **Stigma** 1. **Drupa** aut **bacca** loculis 1-spermis.

I. *HEDERA* Tourn. t. 384. *L. gen. n.* 283.

**Calyx** 5-dentatus; **pet.** 5; **stam.** 5; **antheræ** basi bifurcatæ; **bacca** 5-locularis, sepimentis in maturitate oblitteratis.

1. *H. HELIX* (L. sp. 292.) foliis quinquangularibus 5-lobisque, floralibus ovatis, umbellâ erectâ. ♂ in sylvis, ad sepes et antiquos muros. — Fl. fr. 4. p. 278. Lam. ill. t. 145. (v. v.)

II. *CORNUS* Tourn. inst. t. 410. *L. gen. n.* 149.

**Calyx** 4-dentatus; **petala** 4; **stam.** 4; **drupa** non coronata fæta nuce 2-loculari.

1. *C. MAS* (L. sp. 171.) arborea, umbellis involucrium æquantibus antè folia nascentibus. ♂ in nemoribus et ad sepes. — Fl. 4. p. 277. Lam. ill. t. 74. f. 1. (v. v.)

2. *C. SANGUINEA* (L. sp. 171.) arborea, ramis erectis, foliis ovatis, cymis depressis. ♂ ad sepes et in sylvis. — Fl. fr. 4. p. 278. — Lob. ic. 2. p. 169. f. 2. (v. v.)

**Trib. II. SAMBUCINEÆ** A. *Rich. in dict. class.* 3. p. 173.

**Calyx** parvus bracteis cinctus. **Stylus** 0. **Stigmata** 3. **Cor.** 1-petala. **Bacca** 1-locularis 1-3 sperma.

III. *SAMBUCUS* Tourn. inst. t. 376. *L. gen. n.* 372.

**Calyx** 5-fidus; **cor.** rotata 5-loba; **stam.** 5; **bacca** 3-sperma.

1. *S. EUULUS* (L. sp. 385.) cymis 3-partitis, foliorum foliolis 7-9 oblongo-lanceolatis, stipulis foliaceis, caule herbaceo. ♀ ad margines viarum et fossarum humidarum. — Fl. fr. 4. p. 276. Blackw. t. 488. (v. v.)

2. *S. NIGRA* (L. sp. 385.) cymis 5-partitis, fol. foliolis 5-7 ovali-lanceolatis, caule arboreo. ♂ ad sepes in solo subhumido. — Fl. fr. 4. p. 276. fl. dan. t. 545. (v. v.)

3. *S. RACEMOSA* (L. sp. 386.) racemis compositis ovatis, fol. foliolis 5, caule arboreo. ♂ in montosis provinciarum orientalium borealiumque. Fl. fr. 4. p. 277. Jacq. rar. 1. t. 59. (v. v.)



IV. VIBURNUM *L. gen. n. 370.*

Calyx 5-fidus; corolla campanulata 5-loba; stam. 5; bacca trisperma.

§. 1. TINUS *Tourn. inst. t. 377.*

Bacca coronata.

1. V. TINUS (*L. sp. 383.*) foliis integerrimis ovatis ramificationibus venarum subtus villosoglandulosus.  $\text{\textcircled{J}}$  in lapidosis et umbrosis Galliae australis. — *Fl. fr. 4. p. 274.* — *Clus. hist. 1. p. 49. n. 1. II et III. (v. v.)*

§. 2. VIBURNASTRUM. — Viburnum et Opulus *Tourn.*

Bacca nuda.

2. V. LANTANA (*L. sp. 384.*) foliis cordatis serratis venosis subtus tomentosis.  $\text{\textcircled{J}}$  in dumetis ad sepes. — *Fl. fr. 4. p. 275.* — *Camer. epit. 122. ic. (v. v.)*
3. V. OPULUS (*L. sp. 384.*) foliis glabris 3-lobis acuminato-dentatis, petioliis glandulosus.  $\text{\textcircled{J}}$  in nemoribus ad sepes. — *Fl. fr. 4. p. 275.* — *Camer. epit. 977. ic. (v. v.)*

Trib. III. CAPRIFOLIEÆ *A. Rich. in dict. cl. l. c.*

Calyx bracteis cinctus; stylus 1; stigma trilobum; corolla monopetala; bacca 2-4-locularis, loculis 2- $\infty$ -spermis.

V. LONICERA *L. gen. n. 233.*

Calyx 5-dentatus urceolatus; corolla tubulosa campanulata vel infundibuliformis 5-fida subinæqualis; stam. 5; bacca 2-3-locularis, loculis polyspermis.

Sect. I. CAPRIFOLIUM *Linn. l. c.*

Baccæ solitariae; cor. tubulosa longa ringens.

1. L. BALCARICA (*DC. fl. fr. 5. p. 499.*) floribus verticillatis sessilibus terminalibus, corollis extus subglabris, foliis lanceolatis acuminatis persistentibus subtus glaucis, summis connato-perfoliatis.  $\text{\textcircled{J}}$  in agro Ruscitonensi usque ad Narbonam, Corsicâ (*herb. DC.*) et circâ Olbias et Massiliam reperiit. (*v. v. D.*)
2. L. ETREUSCA (*Senti viagg. 1. p. 113. t. 1.*) floribus in cespitula 3 pedicellata aggregatis, cor. extus pubescentibus, foliis subtus pubescentibus obovatis obtusissimis distinctis, summis subperfoliatis.  $\text{\textcircled{J}}$  frequens in sepibus et dumetis provinc. australiorum. — *Fl. fr. 5. p. 499. (v. v. DC.)*
3. L. CAPRIFOLIUM (*L. sp. 246.*) floribus verticillatis sessilibus terminalibus, cor. tubo extus pilosiusculo, foliis glabris ovatis, summis connato-perfoliatis.  $\text{\textcircled{J}}$  ad sepes et vineas in Galliâ australi. — *Fl. fr. 4. p. 270. Lam. ill. t. 150. f. 1. (v. v.)*
4. L. PERICLYMENUM (*L. sp. 247.*) floribus capitatis terminalibus, foliis omnibus distinctis.  $\text{\textcircled{J}}$  ad sepes et in nemoribus. — *Fl. fr. 4. p. 270. Black. t. 25. (v. v.)*

Sect. II. XYLOSTEUM *Linn. l. c.*

Baccæ geminae aut connatae; corolla infundibuliformis aut campanulata; pedunculi biflori.

\* *Baccis distinctis.*

5. L. XYLOSTEUM (*L. sp. 248.*) foliis ovatis acuminatis integerrimis pubescentibus.  $\text{\textcircled{J}}$  in umbrosis et sepibus. — *Fl. fr. 4. p. 271. Duham. arb. 2. t. 54. (v. v.)*
6. L. NIGRA (*L. sp. 24.*) bracteis calycinis subulatis calyce multo brevioribus, foliis ellipticis glabris integerrimis.  $\text{\textcircled{J}}$  in montibus Pyren. edit. ex cl. Benth.,

Arverniz, Galloprovinciz, Delphinatûs, Jurass. — Fl. fr. 4. p. 271. Jacq. austr. t. 214. (v. v.)

7. *L. PYRENAICA* (L. sp. 248.) foliis oblongo-obovatis glabris, bracteis cal. lanceolatis calyce 2-3-plôve longioribus, floribus subregularibus. ♂ in montibus Galloprovinciz? et Pyrenæorum. — Fl. fr. 4. p. 272. Duham. ed. 2. i. p. 53. t. 15. (v. v. DC.)

\* \* *Baccis coadunatis.*

8. *L. ALPIGENA* (L. sp. 248.) ovariis liberis, baccis coadunatis, foliis ovato-lanceolatis acuminatis subhirsutis. ♂ in montibus. — Fl. fr. 4. p. 272. Duham. ed. 2. p. 54. t. 16. (v. v.)
9. *L. CÆRULEA* (L. sp. 249.) ovariis baccisque coadunatis, foliis ovatis obtusiusculis, junioribus pubescentibus, antiquioribus glabris. ♂ in sylvis editioribus montium Galliæ ferè totius. — Fl. fr. 4. p. 272. Duham. ed. 2. i. p. 54. t. 17. (v. v.)

VI. LINNÆA Gron. L. gen. 774.

Cal. 5-fidus, basi cinctus caliculo persistente 4-partito; cor. campanulata 5-loba; stam. 4, 2 breviora; bacca sicca ovata 3-locularis, loculis 2-spermis.

1. *L. BORREALIS* (L. sp. 880.) ♂ in Alsatiâ?, Cebennis ex Gouan? — Fl. fr. 4. p. 269. fl. lapp. t. 12. f. 4. Dubia civis. (v. s. ex Helv.)

## Ordo LII. LORANTHÆ Juss. et Rich. ann. mus. 12. p. 292.

Calyx ovario adhærens 1-sepalus sæpè basi bibracteatus limbo subintegro. Corolla nunc 1-petala plus minùsve divisa, nunc polypetala petalis basi latioribus. Stamina corollæ lobis numero æqualia aut rariùs dupla illisque opposita. Stylus 1 aut nullus. Stigma 1. Bacca aut Capsula 1-locularis. Spermodermis membranacea. Semina albumine carnoso in parte superiori dehiscente impleta. Embryo cylindricus albumine longior. Radicula exserta infrà apicem coarctata claviformis supera. — Frutices plerumque parasitici monoïci, foliis oppositis rariùs alternis exstipulatis, floribus terminalibus axillari-busve solitariis corymbosis aut spicatis.

### I. VISQUM Tourn. 580. L. gen. n. 1105.

Diöicum. Calyx margine integro vix prominulo; petala 4 parva basi latiore juncta. Masc. Antheræ sessiles mediis petalis insertæ. Fem. Ovarium apice marginatum; stylus minimus, stigma capitatum, bacca parva non coronata 1-sperma.

1. *V. ALBUM* (L. sp. 1451.) foliis lanceolatis obtusis, caule dichotomo, spicis axillaribus. ♂ in truncis et ramis malorum, quercuum, ulmorum, pinorum, etc. — Fl. fr. 4. p. 273. ann. mus. t. 12. t. 27. (v. v.)
2. *V. OXYCEDRI* (DC. fl. fr. 4. p. 274.) aphyllum ramosum erectum carnosum hinc inde vaginulis obtectum. — In ramis Juniperi *Oxycedri* circa Narbonam reperit Broussonet, sed ex eo in Galliâ nuspiam repertum. — Lob. ic. 2. p. 222. f. 2. (v. s. ex Hispan.)

## Ordo LIII. RUBIACEÆ Juss. gen. 196.

Calyx monosepalus ovario adhærens, 4-5-rarius 6-lobus. Corolla 1-petala regularis summo calycis tubo inserta, 4-5-rarius 6-loba. Stamina corollæ inserta cum ejus lobis alterna, iisdem numero æqualia. Ovarium simplex. Stylus 1. Stigmata ovarii loculis numero æqualia. Fructus (in nostris) 2-locularis 2-spermus. Embryo rectus in albumine corneo; radícula infera; cotyledones foliaceæ. — Rubiaceæ nostræ herbæ radicibus rubescentibus, foliis integris verticillatis; floribus parvis rotatis aut tubulosis.

## I. RUBIA L. gen. n. 127.

Corolla campanulato-patens 4-5-loba; stam. 4-5; baccæ 2 glabræ subrotundæ.

1. *R. tinctorum* (L. sp. 158.) foliis annuis quaternis senisve ovato-lanceolatis lanceolatisve acutis subpetiolatis, corollæ lobis oblongis apice subcallosis, caule scaberrimo.  $\mathcal{Z}$  in arenosiâ Mospelii, Andegaviæ, etc.; colitur in Galloprovinciâ, Alsatia, etc. — Fl. fr. 4. p. 267. Mill. dict. n° 1. t. 1. (v. v.)
2. *R. perigrina* (L. sp. 158.) foliis perennantibus 4-6-nis lanceolatis acutis sessilibus, superioribus sæpius obcordatis, corollæ lobis mucronato-aristatis, caule scabro.  $\mathcal{Z}$  in Galliâ australi et occidentali. — Fl. fr. 4. p. 267. — Moris. s. 9. t. 21. n. 2. (v. v.)
3. *R. lucida* (L. syst. nat. 12. p. 732.) foliis perennantibus quaternis lanceolatis lucidis, inferioribus sæpius obcordatis mucronatis, corollæ lobis lanceolatis aristatis, caule læviusculo.  $\mathcal{Z}$  circa Parisios? in Galloprovinciâ, Occitaniâ, agro Ruscinonensi. — Fl. fr. 4. p. 268. (v. v. DC.)
4. *R. Requienii* (Duby mas.) foliis perennantibus senis octonisve linearibus angustis acuminatis sessilibus, corollæ lobis ovali-lanceolatis aristatis, caule scabro.  $\mathcal{Z}$  in Corsicâ circa Bonifacio reperiit cl. Requien. (v. s.)

## II. VAILLANTIA Tourn. a. g. 1706. t. 3. L. gen. 1151.

Corolla campanulata 3-4-fida, fructus florum ternorum calycibus demum basi coadunatis et limbo dilatatis tricornis.

1. *V. muralis* (L. sp. 1490.) glabra, foliis quaternis, floribus ternis polygamis, centrali fertili 4-fido, lateralibus sterilibus 3-fidis. ① ad rupes et muros in regione Olearum. — Fl. fr. 4. p. 266. 5. p. 499. — Mich. gen. p. 13. t. 7. (v. v.)

## III. GALIUM Scop. ed. 2. 1. p. 99. All. — Galii et Valantiæ sp. L.

Corolla rotata aut campanulata 4-fida; fructus calyce non coronatus.

§. 1. *Fructus glabro non tuberculoso.*

\* *Floribus luteis.*

1. *G. cruciata* (Scop. carn. 1. p. 100.) foliis quaternis trinerviis hirsutis subacutis, floribus polygamis 4-fidis, pedunculis diphyllis.  $\mathcal{Z}$  ad sepes et vias. — Fl. fr. 4. p. 250. — Lob. ic. t. 804. f. 2. (v. v.)
2. *G. verum* (Scop. carn. ed. 2. n. 144.) foliis quaternis subrigidis trinerviis ciliatis obtusissimis, floribus polygamis 4-fidis, pedunculis dichotom

aphyllis.  $\mathcal{Z}$  in montosis umbrosis Pyrenæorum, Occitanis et circa Dax. — Fl. fr. 4. p. 350.

*a. Bauhini*, foliis ovatis confertis, floribus luteis. — Scop. l. c. t. 2. G. Bauhini Röm. et Sch. 3. p. 218. Vaillantia crebrifolia St.-Am. ag. p. 424. (v. v.)

*β. Halleri*, fol. lanceolatis distantibus, flor. pallidioribus. — G. Halleri Röm. et Sch. 3. p. 218. (v. s.)

3. G. ARENARIUM (Lois. fl. 85.) foliis suboctonis oblongis crassiusculis glabris margine subrevolutis, pedicellis trifidis, caule ramosissimo prostrato.  $\mathcal{Z}$  in arenis maritimis à Bayona ad Venetam Armoracis. — Fl. fr. 5. p. 495. G. megalospermum  $\beta$  Fl. fr. 4. p. 249. (v. v.)

4. G. VERUM (L. sp. 155.) foliis suboctonis linearibus sulcatis, ramis floriferis brevibus, caule in parte superiori erecto.  $\mathcal{Z}$  in pratis, ad sepes et vias. — Fl. fr. 4. p. 248. — Cam. epit. 868. ic. (v. v.)

\*\* *Floribus rubris.*

5. G. PURPUREUM (L. sp. 156.) fol. 6-10-nis lineari-setaceis, floribus subterminalibus, pedunculis brevissimis, pedicellis sub-1-floris.  $\mathcal{Z}$  in Galloprovincia et circa Nicæam. — G. rubrum Fl. fr. 4. p. 251. — J. Bauh. hist. 3. p. 721. f. 2. (v. v. in Italiâ.)

6. G. RUBRUM (L. sp. 156.) fol. 6-8-nis inferioribus lanceolatis, superioribus lineari-lanceolatis linearibusve, pedunculis ramosis capillaribus folio longioribus, pedicellis multifloris.  $\mathcal{Z}$  circa Nicæam. — G. purpureum Fl. fr. 4. p. 251. — Clus. hist. 2. p. 175. f. 1. (v. v. DC.)

*β. pilosum*, caule basi piloso. — G. purpureum  $\beta$  Fl. fr. 5. p. 446. Circa Mimatem reperiit Prost. (v. s.)

\*\*\* *Floribus albis.*

G. CAMPESTRIS (Schousb. ex Wild. hort. ber. 1. p. 152.) foliis glaberrimis, infimis quaternis ovatis acutis, superioribus senis lineari-ellipticis acuminatis, caule decumbente tetragono. — In Corsicæ montibus reperiit Ph. Thomas — Flores sicci lutei, an vivi ochroleuci? (v. a.)

8. G. DIVARICATUM (Lam. dict. 2. p. 580.) foliis subseptenis linearibus mucronatis (oculo armato) hispidio-ciliatis, caule superne ramis capillaribus dichotomis et divaricatis paniculato. (1) in arenosis et lapideis provinciarum australiorum. — Fl. fr. 4. p. 259. ic. rar. t. 24. G. tenue Vill. 2. p. 322. t. 7. An à sequente satis diversum? (v. s.)

9. G. ANGULICUM (Huds. ang. 69.) foliis subsevis lineari-lanceolatis mucronatis margine cauleque scabris, floribus paniculatis terminalibus, pedunculis 2-aut 3-fidis. (1) in siccis lapideis vel arenosis. — Fl. fr. 4. p. 258. — Ray syn. t. 9. f. 1. (v. v.)

10. G. PYRÆNÆICUM (Lin. fil. sup. 121.) foliis senis linearibus acutissimis erectis nitidulis basi tumidis, floribus axillaribus solitariis foliis brevioribus.  $\mathcal{Z}$  in Pyrenæis præsertim orientalibus. — Fl. fr. 4. p. 260. Gou. ill. 5. t. 1. f. 4. (v. v.)

11. G. PUMILUM (Lam. dict. 2. p. 580.) foliis subseptenis linearibus subtus bisulcatis lævibus acutissimis basi tumidis, floribus subterminalibus subumbellatis foliis longioribus.  $\mathcal{Z}$  in saxosis Pyrenæorum præsertim centralium, montis Ventosi, montium Lugduni et Delphinatûs. — Fl. fr. 4. p. 260. Lam. ill. t. 60. f. 2.

*a. hypnoides*, pedunculis 2-3 floris. — G. hypnoides Vill. 2. p. 323. (v. v.)

*β. caespitosum*, ped. 1.-floris. — G. caespitosum Lam. ill. 1369. (v. v.)

*γ. pubescens*. — Circa Vallem-clausam et Massiliam. — G. pusillum L. sp. 154. (v. s.)

12. G. LÆVE (Thunb. fl. par. ed. 2. p. 77.) foliis subseptenis planis lineari-lanceolatis margine lævibus mucronatis glabris, floribus subumbellatis, ocellibus gla-

- bris diffusis ascendentibus.  $\mathcal{L}$  in pascuis montosis et sylvis siccis. — Fl. fr. 4. p. 256. (v. v.)
13. *G. supinum*. (Lam. fl. fr. 3. p. 379.) foliis subsenis lineari-lanceolatis mucronatis subasperis, caulibus prostratis ramosissimis, paniculâ 3-fidâ 3-choromâ.  $\mathcal{L}$  in aridis et lapidosis. — Fl. fr. 4. p. 259. — Juss. ac. par. 1714. t. 15. f. 2. An satis distincta sp? (v. s.)
14. *G. mucronatum* (Lam. dict. 2. p. 581.) foliis suboctonis scabris mucronatis, superioribus linearibus, inferioribus ovato-lanceolatis, pedunculis ramosis, pedicellis patulis, paniculâ divaricatâ, corollæ lobis acuminato-setaceis, caule ascendente infernè hirtò supernè glabro.  $\mathcal{L}$  in Pyrenæis, Delphinatû. — Fl. fr. 4. p. 258. *G. obliquum* Vill. dauph. 2. t. 8. Valdè var.  $\beta$  sequentis affine. (v. s.)
15. *G. Bocconi* (All. ped. 24.) foliis senis octonis deniave asperis mucronatis, inferioribus pubescentibus, pedunculis sub-3-choromis, caulibus 4-gonis prostratis.  $\mathcal{L}$  in montosis et umbrosis. — Fl. fr. 4. p. 257.
- a. *nitidulum*, floribus majoribus, pedicellis erectis, paniculâ coarctatâ. — *G. nitidulum* Thuill. ed. 2. t. p. 76. Circâ Parisios, Stampas, in Arvernâ montibus et Pyrenæis. (v. s.)
- $\beta$ . *micranthum*, flor. minimis, pedicellis patulis, paniculâ divaricatâ, in Galloprovinciâ et circâ Nicæam, in Pyrenæis reperi. — Boeck. mus. p. 145. t. 101. (v. v. D.)
16. *G. glaucum* (L. sp. 156.) foliis senis octomisve subtus glaucis linearibus submucronatis margine revolutis scabris glabris, floribus subcorymbosis, pedunculis dichotomis pedicellis erectis, paniculâ coarctatâ, caule lævi glabro subtereti ascendente.  $\mathcal{L}$  in lapidosis umbrosis Delphinatû, Galloprovinciæ, Occitaniæ, agri Ruscionensis. — Fl. fr. 4. p. 252. *G. campanulatum* Vill. dauph. 2. p. 326. t. 7. (v. v.)
17. *G. tenuifolium* (All. ped. 23.) foliis senis linearibus rigidis sulcatis acutissimis lucidis retrorsum scabris, floribus paniculatis pedicellis 3-choromis, caule subliguoso ramoso rigido 4-gono lævi.  $\mathcal{L}$  in provinciis australibus usquè ad Lugdunum. — Fl. fr. 4. p. 256. (v. v.)
18. *G. limpolium* (Lam. dict. 2. p. 578.) foliis lanceolato-linearibus glaberrimis subtus glaucis acuminatis margine lævibus inferioribus octonis, pedunculis paniculatis capillaribus, caule tereti.  $\mathcal{L}$  in umbrosis Delphinatû, Galloprovinciæ et Corsicæ (Ph. Thomas.) — Fl. fr. 4. p. 252. *G. atrovirens* Lapeyr. ex Benth. — Barr. ic. t. 356 et 583. (v. v.)
19. *G. sylvaticum* (L. sp. 155.) foliis ellipticis mucronatis lævibus subtus scabris inferioribus octonis, pedunculis capillaribus, caule lævi ramosissimo tereti.  $\mathcal{L}$  in Alatiâ, Delphinatû, agro Andegaviensi, Parisienaj. — Fl. fr. 4. p. 252. (v. v.)
20. *G. erectum* (Huds. ang. 68.) foliis senis octonisve lineari-lanceolatis margine aculeato-serrulatis mucronatis, paniculis 3-choromis, corollæ lobis acutis, caule lævinsculo flaccido.  $\mathcal{L}$  in pratis humidis montosis Pyrenæorum, Galloprovinciæ, Delphinatû. — Fl. fr. 4. p. 255. eng. bot. t. 2067. An satis distincta sp? (v. v. DC.)
21. *G. mollugo* (L. sp. 155.) foliis octonis ovatis lanceolatisve subserratis patentissimis mucronatis, corollæ lobis acutis, caule flaccido anguloso ramoso, ramis patentibus.  $\mathcal{L}$  ad sepes et vias humidas frequens. — Fl. fr. 4. p. 254. Bull. herb. t. 283. (v. v.)
- $\beta$ . *aristatum*, corollis acutiusculis. — *G. aristatum* L. sp. 152. Fl. fr. 4. p. 255. — Bar. ic. 356. Pro ipso *G. Mollugine* habet cl. Requien in herb. DC. (v. s.)
22. *G. palustre* (L. sp. 153.) foliis quaternis seniave obtusis inæqualibus glabris, inferioribus obovatis, superioribus lanceolatis, caule diffuso 4-gono lævinsculo.  $\mathcal{L}$  in humidis. — Fl. fr. 4. p. 254. fl. dan. t. 423. (v. v.)

*β. constrictum*, foliis linearibus. — *G. constrictum* Chaub. in *S. Am. fl. ag. bouq. t. 2.* (v. s.)

23. *G. ULIGINOSUM* (L. sp. 153.) foliis senis lanceolatis retrorsum serrato-aculeatis mucronatis rigidis, caulis angulis aculeato-scabris. ½ in turfosis paludosis circa Hunningam, Parisios et in Arvernâ. — *Fl. fr. 4. p. 259. eng. bot. t. 19.* An à priori satis distinctum? (v. s.)

24. *G. VILLARSII* (Req. in *Guer. vanc. ed. 2. p. 250.*) foliis senis linearibus glabris subacutis, caule ramosissimo procumbente, corollis obtusis, fructibus grossis sphaericis. ½ in saxosis Delphinatûs, montis Ventosi et Pyrenæorum. — *Fl. fr. 5. p. 497. G. megalospermum Vill. 2. p. 319. non All. G. cometerhizon La Peyr. ab. supp. 23.* (v. s.)

25. *G. SAXATILE* (L. sp. 154.) foliis senis obovatis obtusis ciliatis, caule ramosissimo procumbente. ½ in saxosis editiorum Pyrenæorum et Delphinatûs. — *Fl. fr. 4. p. 261. — Juss. acad. par. 1714. t. 15. f. 1. (v. v.)*

### §. 2. *Fructu glabro tuberculoso.*

26. *G. HANCYNICUM* (Weig. obs. p. 25.) foliis quaternis senisve lanceolato-obovatis acuminatis, caule procumbente, ramis confertis elongatis, pedunculis multifloris. ½ in montosis. — *Fl. fr. 4. p. 261. ic. rar. t. 25.* (v. s.)

27. *G. SPURUM* (L. sp. 154.) foliis senis lanceolatis carinatis scabris retrorsum aculeatis, pedunculis multifloris, fructiferis vix recurvis folio longioribus. ① in locis cultis frequens. — *Fl. fr. 4. p. 262.* (v. v.)

28. *G. TRICORNE* (With. brit. ed. 2. p. 153.) foliis suboctonis lanceolatis margine retrorsum aculeatis, pedunculis folio brevioribus trifloris, fructibus grossis nutantibus. ① in arvis et locis cultis frequens. — *Fl. fr. 4. p. 262. — Vill. bot. 14. n. 2. t. 1. f. 3. as.* (v. v.)

29. *G. SACCHARATUM* (All. ped. 39.) foliis subseptenis lineari-lanceolatis margine autrosam aculeato-scabris, floribus polygamis, pedunculis apice 3-floris, fructibus in pedunculo communi breviter pedunculatis. ① in locis cultis. — *Fl. fr. 4. p. 262. — Vaill. bot. p. 14. t. 4. f. 3. b.* (v. v.)

### §. 3. *Fructu hispido.*

30. *G. MURALE* (All. ped. t. 77. f. 1.) foliis inferioribus senis subreflexis, mediis quaternis, summis binis linearibus patulis, floribus verticillatis subsessilibus, fructibus lineari-ovoideis reflexis. ① ad muros et in sterilibus Gallie australioris. — *Fl. fr. 4. p. 264. G. minimum Röm. et Schult. (v. v. DC.)*

31. *G. VERTICILLATUM* (Lam. dict. 2. p. 585.) foliis lanceolatis pubescentibus reflexis, inferioribus senis, mediis quaternis, summis oppositis, floribus verticillatis sessilibus, fructibus globosis erectis. ① in Galloprovinciâ circa Arelatem, Vallem-clausam, Salun et ad radices montis Ventosi, in Occitaniâ et agro Ruscinonensi ex Benth. — *Fl. fr. 5. p. 498. Lois. not. 33. t. 2. G. verticilliflorum Pourr. (v. s.)*

32. *G. LITIGIOSUM* (DC. fl. fr. 4. p. 263.) foliis quaternis senisve linearibus erectis acuminatis glabris, pedunculis divaricatis capillaribus sub 3-chotomis. ① in locis siccis, ad muros et vias provinciarum australiorum. — *ic. rar. 1. t. 26. G. Parisiense L. sp. 157.* (v. v.)

33. *G. SETACEUM* (Lam. dict. 2. p. 584.) foliis subsenis lineari-setaceis erectis obtusiusculis (oculo armato) hispidulis, caule erecto filiformi apice ramoso, fructu subgloboso hispidissimo. ① in aridis Galloprovinciæ australis. — *Fl. fr. 5. p. 498. G. capillare Cav. ic. t. 191. f. 1. (v. s.)*

34. *G. MARITIMUM* (L. mant. 38.) foliis verticillatis lanceolato-ovatis villosis senis quaternis et binis, pedunculis unifloris, caule basi repente hirtio, geniculis glabris aut villosis. ½ in agro Ruscinonensi, Monspelienâ, Nicæensi? — *Fl. fr. 4. p. 263. — Barr. ic. t. 81. (v. v.)*

35. *G. APARINE* (L. sp. 157.) foliis octonis lanceolatis carinis scabris retrorsum

aculeatis, fructuum pilis uncinatis. ① in arvis, ad sepes et vineas. — Bull. herb. t. 315.

*α. vulgaris*, caule ramoso, geniculis villosis. — Fl. fr. 4. p. 263. (v. v.)

*β. Vaillantii*, caule simpliciusculo, geniculis subvillosis. — G. Vaillantii Fl. fr. 4. p. 263. — Vaill. bot. n. 1. t. 4. f. 4. (v. v.)

36. *G. boreale* (DC. fl. fr. 5. p. 498.) foliis quaternis lanceolatis trinerviis glabris obtusis, caule erecto, fructibus subsquamosis. ½ in montosis et lapidosis, ad sepes et in humidis.

*α. glabrum*, fructibus glabris. — G. rubroides β Fl. fr. 4. p. 253. G. orbibracteatum Chamb. in S.-Am. fl. ag. t. 1. (v. v.)

*β. scabrum*, fructibus subscabris. — Fl. Dan. t. 1024. (v. s.)

*γ. hispidum*, fructibus hispidis. — G. boreale L. sp. 156. Fl. fr. 4. p. 265. (v. s.)

37. *G. barrelieri* (Salzm. in bot. zeit. 1821. p. 107.) foliis quaternis ovatis 3-nerviis cauleque erecto hirsutis, paniculâ ramosissimâ, fruct. hispidis. ½ in Corsicâ collibus umbrosis. — Barr. ic. 324. (v. s.)

38. *G. rotundifolium* (L. sp. 156.) foliis quaternis subrotundo-ovatis trinerviis hispido-ciliatis, caule decumbente glabro, fructibus hispidis. ½ in sylvis umbrosis montium. — Fl. fr. 4. p. 266. — Barr. ic. 323. (v. v.)

#### IV. ASPERULA L. gen. n. 121.

Corolla infundibuliformis 3-vel sæpius 4-fida; baccae 2 siccae calycis limbo non coronatæ.

1. *A. levigata* (L. mant. 38.) foliis quaternis ellipticis uninerviis lævibus, fructibus scabris. ½ in montibus Lugdunensibus? circa Narbonam et Argentoratum? et in Corsicâ. — Fl. fr. 4. p. 246. Facies Galii rotundifolii sed corolla infundibuliformis et fructus glabri. (v. s.)

2. *A. taurina* (L. sp. 150.) foliis quaternis ovato-lanceolatis trinerviis, floribus polygamis fasciculatis vel verticillatis terminalibus. ½ in montosis et umbrosis Delphinatûs, Monspelii. — Fl. fr. 4. p. 245. — Lob. ic. t. 800. f. 1. (v. v. c.)

3. *A. odorata* (L. sp. 150.) foliis octonis lanceolatis, florum fasciculis pedunculatis, fructibus hirsutis. ½ in sylvis et umbrosis. — Fl. fr. 4. p. 245. — Lob. ic. t. 801. f. 1. (v. v.)

4. *A. hirta* (Ram. bull. phil. n. 41. p. 131. t. g. f. 1. 2. 3.) foliis senis, hirtis acutis internodio longioribus, floribus terminalibus aggregatis sessilibus bracteis superantibus. ½ in saxosis editiorum Pyrenæorum. — Fl. fr. 4. p. 244. Fructus glabri. (v. v.)

5. *A. cynanchica* (L. sp. 151.) foliis quaternis, infimis sæpius lanceolato-ovatis, mediis linearibus, superioribus oppositis, floribus 4-fidis, paniculâ contractâ. ½ in collibus lapidosis et pratis aridis. — Fl. fr. 4. p. 246. 5. p. 495. Admodum in decumbentiâ variabilis. (v. v.)

6. *A. tinctoria* (L. sp. 150.) foliis omnibus linearibus, inferioribus senis, intermediis quaternis, superioribus oppositis, floribus umbellatis distinctis petiolatis sæpius 3-fidis, paniculâ laxiusculâ. ½ in collibus aridis et lapidosis. — Fl. fr. 4. p. 245. — Tab. ic. t. 733. f. 1. Præcedenti nimium affinis. (v. v.)

7. *A. arvensis* (L. sp. 150.) foliis senis vel octonis obtusis glabris, floribus terminalibus sessilibus aggregatis, bracteis ciliatis. (X) in arvis. — Fl. fr. 4. p. 245. — Lob. ic. t. 801. f. 2. (v. v.)

#### V. SHERARDIA Gærtn. 1. p. 110. t. 24. DC. fl. fr. 4. p. 243.

Corolla infundibuliformis 4-fida; fructus calycinis deptibus persistentibus et accretis coronatus.

1. *S. arvensis* (L. sp. 149.) foliis senis acutis hispidis, floribus sessilibus terminalibus umbellatis, bracteis glabris. (X) in arvis. — Fl. fr. 4. p. 243. Lam. ill. t. 61. (v. v.)

VI. CRUCIANELLA L. *gen. n.* 126.

Corolla infundibuliformis, tubo gracili, limbo brevi 4-5-fido; calycis laciniae 2 oppositae profundae; fructus calyce non coronatus.

1. *C. MARTIMA* (L. sp. 158.) procumbens suffruticosa, foliis quaternis mucronatis, floribus oppositis 5-fidis. 3 in sabulosis maritimis Galloprovinciae et Occitaniae. — Fl. fr. 4. p. 248. — Barr. ic. t. 355. (v. v.)
2. *C. LATIFOLIA* (L. sp. 158.) procumbens, foliis quaternis lanceolatis, floribus spicatis. ① in siccis et sterilibus Nicææ, Galloprovinciae, Occitaniae. — Fl. fr. 4. p. 247. — Barr. ic. t. 520 et 549. (v. v.)
3. *C. MONSPELIACA* (L. sp. 158.) ramis procumbentibus erectivae, foliis acutis, caulinis quaternis ovatis, rameis quinis linearibus, floribus spicatis. ① in sabulosis circa Nicæam, Massiliam, Monspelium, Montem-Albanum, Viennam. — Fl. fr. 4. p. 247. An à præcedente diversa? (v. s.)
4. *C. ANGUSTIFOLIA* (L. sp. 157.) erecta, foliis senis linearibus, floribus spicatis. ① in siccis arenosis et lapideis Galliae australis. — Fl. fr. 4. p. 247. — Barr. ic. t. 550. (v. v.)

Ordo LIV. VALERIANEÆ DC. *fl. fr. ed.* 3. t. 4. p. 232.

*Dufr. val. mon.* 56.

Calyx 1-sepalus ovario adhærens, limbo nunc papposo intus revolutus, nunc dentato erecto. Corolla 1-petala tubulosa, apici ovarii inserta, 3-5-loba, interdum inæqualis aut calcarata. Stamina 1-5 tubo corollae inserta; Stylus 1. Stigmata 1-3. Capsula non dehiscens 1-3-localaris, loculis 1-spermis, 2 sæpissimè abortivis. Albumen 0; Embryo rectus; radícula supera. — Herbae oppositifoliae, floribus corymbosis, paniculatis aut subcapitatis.

I. VALERIANELLA *Tourn. inst.* 1. p. 132. *Dufr. val. p.* 56. \*

Corolla regularis 5-loba ecalcarata; stam. 3; capsula 3-localaris, limbo calycis recto nec involuto coronata.

1. *V. OLITORIA* (Moench. meth. 493.) caule debili, foliis lanceolatis integerrimis, capsula nudà globoso-compressa. ① in cultis ad vineas et agros. — Fl. fr. 4. p. 240. — Moris. 3. s. 7. t. 16. n. 36. (v. v.)
2. *V. CARINATA* (Lois. not. 149.) caule debili, foliis oblongis integris, capsula oblonga glabra edentula rimoso-carinata. ① inter segetes. — Fl. fr. 5. p. 492. *Dufr. val.* t. 2. Præcedentis varietas? (v. v.)
3. *V. FUMILA* (DC. fl. fr. 4. p. 242.) caule glabrinsculo, foliis inferioribus dentatis, summis linearibus multifidis, bracteis ciliatis, capsula turgida glabra lævi breviter 3-dentata. ① in Occitania et Galloprovincia. — Fl. fr. 5. p. 494. — Moris. 3. s. 7. t. 16. n. 37. (v. v. DC.)
4. *V. DENTATA* (DC. fl. fr. 4. p. 241.) caule glabro, foliis inferioribus lanceolatis integris, superioribus irregulariter basi dentatis, bracteis glabris, capsula glabra ovata dentibus 3-5 inæqualibus uno alio longiore coronata. ① inter segetes. — *Dufr. val.* t. 3. n. 5. — Moris. 3. s. 7. t. 16. n. 32. (v. v. DC.)
5. *V. AURICULA* (DC. fl. fr. 5. p. 492.) caule erecto glabro, foliis inferioribus lan-



- occolatis integris, superioribus irregulariter basi dentatis, bracteis glabris, capsulâ oblongâ glabrâ dente unico erecto acuto basi concavo coronatâ. ① Monspelii. — Hanc valdè suspicor tantum varietatem præcedentis esse. (v. s.)
6. *V. MIXTA* (Duf. val. 58 t. 3. f. 6.) caule anguloso, foliis integris, bracteis subciliatis, capsulâ minimâ ovatâ villosâ 3-4-denticulatâ. ① in arvis Andegaviæ, Occitaniz et Galloprovinciæ. — Fl. fr. 5. p. 492. — Moris. 3. s. 7. t. 16. n. 35. (v. s.)
7. *V. ASIOCARPA* (Desv. journ. 2. p. 314. ic.) caule angulato, foliis inferioribus oblongo-spathulatis, summis linearibus, bracteis glabris, capsulâ ovatâ angulosâ hirtâ irregulariter 6-dentatâ. ① inter segetes ex Aureliâ usque ad Galloprovinciam. — Fl. fr. 5. p. 493. Duf. val. t. 3. n. 4. — Moris. 3. s. 7. t. 16. n. 33. Staturâ admodum variabilis; folia intermedia aliquando basi dentata. (v. s.)
8. *V. ECHINATA* (DC. fl. fr. 4. p. 242.) caule glabro, foliis dentatis, capsulâ lineari 3-dentatâ, dente extimo majore recurvato. ① in arvis Galliz australis, in Limaniâ Arverniz. — Duf. val. t. 3. f. 10. — Moris. 3. s. 7. t. 16. n. 28. (v. s.)
9. *V. HAMATA* (DC. fl. fr. 5. p. 494.) foliis linearibus, bracteis ciliatis, capsulâ pubescente obovatâ angulatâ, limbo 6-dentato dentibus lineari-subulatis hamatis. ① Andegaviæ. (v. s.)
10. *V. DISCORDEA* (Lois. not. 148.) caule glabro, foliis integris dentatisve, capsulâ villosâ coronatâ limbo patente 6-12-radiato, dentibus lanceolatis setaceo-hamatis. ① inter segetes Galloprovinciæ, Occitaniz. — Fl. fr. 5. p. 493. Duf. val. t. 3. f. 3. — Moris. 3. s. 7. t. 16. n. 29. In speciminibus examini subjectis, capsulâ semper 6-dentata. An notæ characteristicæ satis constantes ad hanc speciem à sequenti rite sejungendum? (v. s.)
11. *V. CORONATA* (DC. fl. fr. 4. p. 241.) caule pubescente, foliis dentatis lobatisve, superioribus 3-5-partitis, capsulâ villosâ limbo campanulato 6-10-radiato, dentibus setaceo-hamatis. ① in arvis ex Aureliâ ad Niczam. — Duf. val. p. 60. t. 3. f. 2. — Moris. 3. s. 7. t. 16. n. 30. (v. s.)
12. *V. VESICARIA* (Mœnch. meth. 493.) caule subvillosa, foliis dentatis, superioribus lobatis, capsulâ ovatâ villosâ limbo calycis vesicario coronatâ. ① circâ Nions in Delphinatâ et Bourbonne-les-Bains ex Vill., propè Beauvais (cl. Graves). — Fl. fr. 4. p. 241. Duf. val. t. 3. f. 9. — Boerh. lûgd. r. t. 75. (v. s.)

#### II. FEDIA Mœnch. meth. p. 486. Duf. val. p. 54.

Corolla irregularis, vix 2-labiata ecalcarata; stamina 2; capsula 3-locularis, limbo calycis recto nec involuto coronata.

1. *F. CORUSCANS* (Gærtn. fruct. 2. t. 86. f. 3.) foliis ovato-obtusis, sessilibus subintegris. ① circâ Niczam ex All. — Fl. fr. 4. p. 240. — Moris. 3. s. 7. t. 16. n. 27. (v. s. ex Ital.)

#### III. CENTRANTHUS DC. fl. fr. 4. p. 238. — Valerianæ sp. L.

Corolla 5-loba regularis calcarata; stamen 1; capsula 1-locularis limbo calycis in pappum plumosum evoluta coronata.

1. *C. LATIFOLIUS* (Duf. val. p. 38.) foliis ovato-lanceolatis. ② in lapideis et maritimis Galliz australis. — C. ruber Fl. fr. 4. p. 239. — Moris. 3. s. 7. t. 14. n. 15. Variat floribus albis rubrisve. (v. v.)
2. *C. ANGUSTIFOLIUS* (DC. fl. fr. 4. p. 239.) foliis linearibus. ② in saxosis montium Galliz orientalis australisque. — Moris. 3. s. 7. t. 14. n. 16. (v. v.)
3. *C. CALCITRAPA* (Duf. val. 39.) foliis superioribus pinnatifidis. ① in saxosis Galliz australis. — Fl. fr. 5. p. 492. Valeriana Calcitrapa Fl. fr. 4. p. 238. — Clns. hist. 2. p. 54. f. 2. (v. v.)

## IV. VALERIANA. Neck. el. 1. p. 123. DC. fl. fr. 4. 233.

Corolla 5-loba regularis ecalcarata; stamina 3; cætera ut in *Cen-  
crantho*.

1. *V. OFFICINALIS* (L. sp. 45.) caule sulcato, foliis omnibus pinnatis, pinnis dentatis.  $\mathcal{L}$  in humidis et sylvis. — Fl. fr. 4. p. 233. — Lob. ic. t. 715. f. 1. (v. v.)
2. *V. PHU* (L. sp. 45.) caule glabro erecto, foliis radicalibus ovatis lanceolatis petiolatis integris obtusis, caulinis pinnatis, lobis inæqualibus lanceolatis acutis.  $\mathcal{L}$  in montibus Alsatiæ? circa Aginnum ex St.-Am. et Burdigalam ex Later. — Fl. fr. 4. p. 234. — Lob. ic. t. 714. f. 2. (v. v. c.)
3. *V. TRIPTERIS* (L. sp. 45.) caule glabro erecto, foliis petiolatis obtusis inæqualiter crenato-sinuatis dentatisve, radicalibus cordatis, caulinis ternatis ovato-oblongis, foliolis lateralibus lanceolatis.  $\mathcal{L}$  in montibus et Alpibus. — Fl. fr. 4. p. 234. — Moris. 3. s. 7. t. 14. n. 10. (v. s.)
4. *V. PYRÆNAÏCA* (L. sp. 46.) caule striato, foliis cordatis inæqualiter dentatis, inferioribus simplicibus, superioribus ternatis pinnatisque, foliolo terminali maximo.  $\mathcal{L}$  in umbrosis Pyrenæorum et montium Velaunorum propè *Chareniac* ex Arnaud. — Fl. fr. 4. p. 234. (v. v.)
5. *V. DIOÏCA* (L. sp. 44.) caule glabro erecto, foliis obtusis, radicalibus spathulato-ovatis integris subcordatis petiolatis, caulinis pinnatifidis lobis inæqualibus, floribus dioicis.  $\mathcal{L}$  in paludosis et pratis humidis. — Fl. fr. 4. p. 238. — Lob. ic. t. 715. f. 2. (v. v.)
6. *V. TUBEROSA* (L. sp. 46.) caule glabro erecto, foliis radicalibus lanceolato-oblongis in petiolum subdecurentibus integris obtusis, caulinis summisque pinnatifidis, lobis linearibus.  $\mathcal{L}$  in montosis Galliæ australis. — Fl. fr. 4. p. 235. — Moris. 3. s. 7. t. 15. n. 20 et 21. (v. v. DC.)
7. *V. GLOBULARIÆFOLIA* (DC. fl. fr. 4. p. 236.) caule glabro subprocumbente, foliis radicalibus ovatis petiolatis integris obtusis, infimis caulinisque pinnatifidis lobis oblongo-linearibus, summis linearibus.  $\mathcal{L}$  in saxosis editiorum Pyrenæorum. — V. heterophylla Lois. gall. 21. t. 2. V. glauca Lapeyr. abr. 19. (v. v.)
8. *V. MONTANA* (L. sp. 45.) caule erecto, foliis oblongis subdentatis, imis obtusis, summis acutis, floribus paniculatis.  $\mathcal{L}$  in montibus. — Fl. fr. 4. p. 235. — Moris. 3. s. 7. t. 15. n. 11. (v. v.)
9. *V. SALIUNCA* (All. ped. n. 9. t. 10. f. 2.) foliis spathulato-oblongis integris obtusis glaberrimis, caulinis lanceolato-linearibus, floribus umbellato-capitatis.  $\mathcal{L}$  in saxosis Alpium et Delphinatûs. — Fl. fr. 5. p. 492. V. supina Fl. fr. 4. p. 237. (v. v. in Helvetiâ.)
10. *V. SAXATILIS* (L. sp. 46.) caule recto, foliis indivisis, radicalibus ellipticis trinerviis ciliatis subdentatis petiolatis, caulinis lineari-lanceolatis sessilibus.  $\mathcal{L}$  circa Nicæam ex All., in Pyrenæis ex Lapeyr. — Fl. fr. 4. p. 237. — Moris. 3. s. 7. t. 14. n. 14. (v. v. D. in Italiâ.)
11. *V. CRUTICA* (L. sp. 46.) caule glabro ascendente, foliis integerrimis obtusis, radicalibus cuneato-oblongis, caulinis linearibus, floribus verticillato-racemosis.  $\mathcal{L}$  in saxosis Alpium Delphinatûs. — Fl. fr. 4. p. 236. Jacq. coll. 1. p. 24. t. 1. (v. s. ex Helv.)

Ordo LV. DIPSACEÆ Juss. gen. 194. excl. *Allioniâ et s. 2.*  
*Coulter mem. sur les Dips. in mem. soc. gen. t. 2. 1. pars p. 13.*

Calyx (internus) adhaerens, limbus pappiformis varie divi-  
 sus. Corolla 1-petala tubulosa supera calyci inserta, limbo  
 obliquo 4-5-fido; æstivatione cochleari. Stamina definita sæ-  
 pissimè 4-5 distincta corollæ adnata, nervisque ejusdem alter-  
 nantia. Stylus 1. Ovarium 1-loculare, ovulo unico ex apice  
 loculi pendulo. Akenium limbo calycis accreto coronatum.  
 Embryonis recti albumineque carnoso tecti radícula supera.  
 Involucellum (calyx externus) 1-phyllo sæpissimè 4-8-foveo-  
 latum, fructum arcuè cingens. — Herbæ aut suffrutices, foliis  
 oppositis verticillatisve. Flores in receptaculo communi aggre-  
 gati, involucri polyphyllo cincti.

I. SCABIOSA *Coul. l. c. p. 33.* — *Scabiosa sp. L.*

Calycis limbus setis 5 hinc inde abortientibus munitus; corolla  
 4-sæpius 5-loba; involucellum 8-foveolatum sæpius subcylindricum.

Sect. I. ASTEROCEPHALUS *Coul. l. c.*

Involucelli basis teres nec plicata.

1. *S. GRAMINIFOLIA* (L. sp. 145.) involucelli basi elongatæ foveolis linearibus  
 æqualibus, coronâ (1) patulâ involucelli basi vix æquali, nervis 24, calycis  
 limbi pedunculati setis coronæ æqualibus, foliis linearibus integerrimis holo-  
 sericeis. 2℥ in montibus Galloprovinciæ, Delphinatûs. — Fl. fr. 4. p. 232.  
 — *Moris. 3. s. 6. t. 15. n. 36. Fl. cærulescentes. (v. v. D. in Italiâ.)*
2. *S. STREILATA* (L. sp. 144.) involucelli basi elongatæ foveolis obovatis æqua-  
 libus, coronæ patulæ basi duplò longioris nervis 40, calycis limbi pedun-  
 culati setis discum versùs incrassatis coronæ æqualibus at ratione pedicelli  
 exsertis, foliis dissectis. (1) in maritimis Occitanix, Galloprovinciæ, Nicææ.  
 — Fl. fr. 4. p. 231. — *Clus. hist. 2. p. 1. ic. 1. (v. v.)*  
 β. *simplex*, caulibus subunifloris, foliis sæpius pinnatifidis. — *S. simplex*  
*Fl. fr. 4. p. 231. Desf. atl. t. 39. f. 1. (v. s.)*
3. *S. MONSPELIENSIS* (Jacq. misc. 2. p. 320.) involucelli basi vix elongatâ  
 foveolis linearibus apertis æquali, coronæ basi duplò longioris nervis 32-36,  
 calycis limbi pedunculati setis longè exsertis, foliis dissectis. (1) circâ Monspe-  
 lium, Arelatem, etc. — *Jacq. ic. rar. t. 24. Coul. dips. 34. S. stellata γ. Fl.*  
*fr. 5. p. 491. (v. s.)*
4. *S. UCRANICA* (L. sp. 144.) involucelli basi hemisphæricâ foveolis ovato-  
 linearibus æquali, coronæ nervis 20-24, calycis limbi in coronâ subsessilis  
 setis longè exsertis, foliis radicalibus pinnatifidis, caulinis linearibus inte-  
 gerrimis basi ciliatis. 2℥ circâ Fontainebleau. — *Fl. fr. 4. p. 230. 5. p. 490. S.*  
*Gmelini St.-Hil. bull. phil. n. 61. p. 149. t. 3. Fl. ochroleuci. (v. v. c.)*

Sect. II. VIDUA *Coul. l. c.*

Involucelli basis rotundata plicata.

5. *S. ATROFURFURÆ* (L. sp. 144.) capitulis ovatis, calycis longè pedunculati  
 pedunculo coronâ duplò longiore, setis patulis exsertis involucello duplò lon-  
 gioribus, involucri foliolis lineari-lanceolatis flores subæquantibus, foliis lan-  
 ceolato-ovatis indivisis dentatis pinnatifidisque lobo terminali maxime den-  
 tato. (1) in hortis culta. — *Fl. fr. 4. p. 231. — Moris. 3. s. 6. t. 14. n. 26. (v. v.)*

(1) Pars involucelli foveolis superior dilatata sæpius membranacea (limbus ca-  
 lycis externi).

6. *S. MARITIMA* (L. amœn. 4. p. 304.) capitulis globosis, calycis pedunculati pedunculo coronam vix aequante, setis erectis exsertis involucello 30 longioribus, involucri foliis linearibus fl. 20 brevioribus, foliis integris pinnatifidisque, radicalibus dentatis, superioribus integerrimis linearibus. ? in provinciis australioribus. — Fl. fr. 5. p. 490. Nondum pro certo habere possum hanc speciem at vult Cl. Coulter, à precedente non diversam esse. (v. v.)

### Sect. III. *Succisa* Coulter. l. c.

Involucelli basis nulla à tubo distinguenda.

7. *S. LUCIDA* (Vill. dauph. 2. p. 293.) foliis radicalibus lanceolato-ovatis obtusis crenatis, caulinis superioribusque glabris pinnatifidis lobis linearibus acutissimis, caule sub-1-floro prostrato, involucri flores superante, coronâ tubo 30 brevioris nervis 20-24, calycis setis nigris flores subæquantibus coronâ vero 4-5-plòve longioribus. ? in montibus Delphinatâ et Galloprovinciâ. — Fl. fr. 4. p. 228. — W. et K. hung. 2. t. 138 et 157. *S. columbaria* α Coulter. 38. (v. v. D. in Helv.)
8. *S. COLUMBARIA* (L. sp. 143.) foliis radicalibus lanceolatis acutis dentatis, caulinis superioribusque pubescentibus bipinnatifidis lobis acutis, caule multifloro erecto, involucri floribus brevioris, coronâ tubo dimidiò brevioris, nervis 20, calycis setis corollâ 3-6-plò minoribus coronâ vero subduplò longioribus. ? in siccis montosis. — Fl. fr. 4. p. 228. *S. gramuntia* L. sp. 143. Fl. fr. 5. p. 488. — Cam. ep. p. 711. ic. (v. v.)

fl. *ochroleuca*, floribus ochroleucis, setis coronâ 30 longioribus. — In provinciis australioribus. — *S. ochroleuca* L. sp. 146. Fl. fr. 4. p. 230. (v. s.)

9. *S. PYRENAICA* (All. ped. t. 25. f. 2.) foliis radicalibus spathulatis obtusissimis inæqualiter crenatis tomentosis, superioribus pinnatifidis lobis lanceolatis dentatis inæqualibus, capitulis erectis, involucri foliolis lanceolatis, coronâ tubo 30 brevioris, nervis 20, setis corollâ sub-3-plò minoribus coronâ vero 40 longioribus. ? in lapidosis montium Galloprovinciâ, Delphinatâ, Occitanîâ et Pyrenæorum. — Fl. fr. 4. p. 229. *S. columbaria* δ. Coulter. 38. Planta vix pedalis. (v. v.)

fl. *mollissima*, cano-tomentosa, foliis superioribus decompositis lobis linearibus integris. — Circâ Nicæam. — *S. mollissima* Fl. fr. 5. p. 490. (v. s.)

10. *S. MOLOSCHICA* (Bert. dec. 3. p. 49.) cano-tomentosa, foliis inferioribus lanceolatis obtusis crenatis, superioribus pinnatifidis lobis lanceolatis acutis lateralibus integris, capitulis nutantibus solitariis, involucri foliolis linearibus, coronâ tubo hirsuto 30 brevioris, nervis 20. ? in Pyrenæis. — Fl. fr. 5. p. 489. *S. columbaria* θ Coulter. 39. Culturâ perstat. Planta 2-4-pedalis. (v. v. DC.)

11. *S. SUAVEOLENS* (Desf. cat. h. par. p. 110.) foliis radicalibus integerrimis ovato-elongatis, caulinis pinnatifidis lobis linearibus longis, caulibus strictissimis, involucelli tubo subcylindrico, coronâ brevissimâ nervis 16, calycis setis patulis brevibus vix exsertis. ? in siccis Fontisbleaudi et Andegaviæ (Bastard.) — Fl. fr. 4. p. 229. — Tabern. ic. 160 et 161. (v. s.)

12. *S. URCOLATA* (Desf. atl. 1. p. 122.) foliis pinnatifidis, involucri foliolis coalitis, coronâ obsoletâ, calycis setis abortientibus. ? in Corsicâ. — Bocc. sic. 1. 52. (v. s.)

13. *S. SUCOSA* (L. sp. 142.) foliis integris, involucri foliolis 2-3-serratis, flosculis aequalibus, coronâ perbrevis undulatâ, setis exsertis connatis. ? in sylvis et collibus aridis. — Fl. fr. 4. p. 226. — Cam. epit. 397. ic. Moris. 3. s. 6. t. 13. h. 7. (v. v.)

### II. *KNAUTIA* Coulter. l. c. — Knautiæ et Scabiosæ sp. L.

Corollâ 4-loba; calycis limbus subcyathiformis; involucellum compressum 4-foveolatum semen arcte cingens, stipite brevi munitum.

1. *K. sylvatica* (Duby mss.) foliis in petiolos alatos basi dilatatos connatosque coarctatis, omnibus indivisis lanceolatis, radicalibus integris, caulinis crenatis, superioribus dentatis, invol. foliolis subacutis, coronæ dentibus minimis, calycis ciliis 8 aristatis.  $\mathcal{U}$  in sylvaticis montanis. — *K. arvensis*  $\gamma$  et  $\delta$  Coult. dips. 29. *S. sylvatica* L. sp. 142. Fl. fr. 4. p. 227. *S. integrifolia* L. sp. 142. non fl. fr. — Clus. hist. 2. p. 2. f. 3. (v. v.)

$\beta$ . *longifolia*, foliis glaberrimis. — In Jurasso. — *S. longifolia* W. et K. hung. 1. t. 5. Fl. fr. 5. p. 488. (v. s.)

2. *K. arvensis* (Coult. dips. 29.) foliis sessilibus, radicalibus inæqualiter pinnatifidis lobis lanceolatis, caulinis pinnatifidis lobis linearibus, superioribus integris lineari-lanceolatis, involucri foliolis obtusis, coronæ dentibus minimis, calycis ciliis 8-10 aristatis.  $\mathcal{U}$  ad vias, prata et agros. — *Sc. arvensis* L. sp. 142. Fl. fr. 4. p. 226. Fl. dan. t. 447. (v. v.)

$\beta$ . *collina*, foliorum radicalium lobis dentatis lobatisve. — *Sc. collina* Req. in Guér. vaubl. ed. 2. p. 248. Fl. fr. 5. p. 487. (v. v. DC.)

3. *K. hybrida* (Coult. dips. 30.) foliis radicalibus petiolatis inæqualiter pinnatifidis lobis obovatis rarius subintegris, caulinis sessilibus lobis lanceolatis, superioribus integris, involucri foliolis acuminatis, coronæ dentibus 12-15 perbrevibus, calycis ciliis 20-24 obsoletis. (1) in Occitaniâ, Galloprovinciâ et circâ Nicæam. — *S. hybrida* All. auct. p. 9. Fl. fr. 4. p. 227. *S. integrifolia* Fl. fr. 4. p. 227. non L. Savi fl. pis. 1. t. 2. f. 1. Flores rosei. (v. v.)

### III. CEPHALARIA Schrad. cat. sem. gott. 1814. Coult. l. c. — Scabiosæ sp. L.

Involucrum imbricatum bracteis brevius; corolla 4-loba; stigma longitudinale; calycis limbus subcyathiformis discoideusve; involucellum tetraëdrum 8-foveolatum interdum compressum.

1. *C. SYRIACA* (Schrad. l. c.) coronæ dentibus 8-4 aristatis, 4 iis alternantibus perbrevibus, bracteis aristatis, corollulis æqualibus, foliis lanceolatis integris dentatisve. (1) inter segetes propè Nemausim. — Coult. dips. 25. *Sc. Syriaca* L. sp. 141. Fl. fr. 5. p. 487. Fl. pallidæ cærulei. (v. s.)

2. *C. LEUCANTHA* (Schrad. l. c.) coronâ membranaceâ, bracteis exterioribus obtusis, interioribus vix acutis, foliis pinnatifidis.  $\mathcal{U}$  in montosis Gallie mediterraneæ. — *S. leucantha* L. sp. 142. Fl. fr. 4. p. 225. Fl. luteo-albidi. (v. v.)

3. *C. CENTAUROIDES* (Röm. et Schult. syst. 3. p. 49.) coronæ dentibus 4-8 obsoletisve, bracteis mucicis, exterioribus obtusis, interioribus acuminatis, foliis pinnatifidis.  $\mathcal{U}$  in Alpibus Galloprovinciæ. — *S. centauroides* Lam. ill. 1312. Fl. fr. 4. p. 224. Flores flavi. (v. s.)

4. *C. ALPINA* (Schrad. l. c.) involucelli coronæ dentibus 6-8 aristatis subæqualibus, bracteis acuminatis pilosis corollulis subæqualibus, foliis pinnatis, foliolis lanceolatis serratis.  $\mathcal{U}$  in alpibus et montibus. — Coult. dips. 24. *Sc. Alpina* L. sp. 141. Fl. fr. 4. p. 224. — Lob. ic. t. 537. f. 2. Flores flavi. (v. v.)

### IV. DIPSACUS L. gen. n. 114. Coult. dips. p. 21.

Calycis limbus subcyathiformis, involucellum 4-edrum 8-foveolatum; receptaculum paleaceum, paleis spinosis; stigma longitudinale; involucri foliola bracteas superantia.

1. *D. PILOSUS* (L. hort. ups. 25.) capitulo globoso piloso, bracteis rectis, involucri foliolis deflexis bracteas vix superantibus, foliis petiolatis appendiculatis. (2) ad sepes et fossas humidas. — Fl. fr. 4. p. 223. Lam. ill. t. 56. f. 2. (v. v.)

2. *D. LACINIATUS* (L. sp. 141.) capitulo ovato-cylindrico, bracteis rectis, involucri foliolis lineari-lanceolatis rigidis capitulo sub æd brevioribus, foliis sinuato-laciniatis. (2) ad vias et sepes rarius occurrit. — Fl. fr. 4. p. 222. — Moris. t. 3. s. 7. t. 36. n. 4. *D. sylvestris*  $\gamma$  Coult. dips. 22. (v. v.)

3. *D. VEROX* (Lois. fl. 719. t. 3.) capitulo ovato, bracteis rectis, involucri foliolis

triquetris patulis rigidis capitulum superantibus, caule foliisque aculeis horridis. ③ in maritimis Corsicæ. — Fl. fr. 5. p. 487. (v. s.)

4. *D. sylvestris* (Mill. dict. n. 2.) capitulo ovato-cylindrico, bracteis rectis, involucri foliolis linearibus incurvis capitulo longioribus, foliis connatis integris dentatisve. ② ad vias et sepes. — Fl. fr. 4. p. 222. Fl. dan. t. 965. — Moris. t. 3. s. 7. t. 36. n. 3. (v. v.)

5. *D. Fullonum* (Mill. dict. n. 1.) capitulo cylindrico, bracteis recurvis, involucri foliolis lanceolatis rigidis capitulo brevioribus, foliis connatis integris subcoriaceis. ② in agris passim cultus et provinciarum australium probabiliter indigenus. — Fl. fr. 4. p. 222. — Lob. ic. 2. t. 17. f. 2. Moris. l. c. n. 1. (v. v.)

---

Ordo LVI. COMPOSITÆ *Adans. fam.* 2. p. 103. *DC. fl. fr.* 4. p. 1. *Cass. dict. sc. nat.* 10. p. 131. — *Synanthereæ Rich.*

Calyx ovario adhærens, tubo nunc ovarii longitudini æquali et tunc *pappus* (seu calycis limbus abortivus) *sessilis*, nunc ovario longiore et tunc pappus *stipitatus* dicitur, limbo nunc vix ullo et tunc pappus *nullus* nunc integro nunc 5-dentato aut 5-lobo et tunc pappus *integer 5-dentatus 5-lobusve* dicitur, nunc sæpius evadente in pilos membranaceos innumeros nunc simplices et tunc pappus *pilosus*, nunc basi coalitos apice liberos et tunc pappus *ramosus*, nunc lateraliter barbato-plumosos et tunc pappus *plumosus* nuncupatur. Corolla summo calycis tubo inserta, constans petalis 5 basi plus minùsve inter se coalitis, nervos semper 2 simplicissimos marginales ad apicem confluentes et interdum tertium centralem gerentibus, per æstivationem valvatis, limbo vario in corollis stamineis naturali, in fæmineis neutrisve plus minùsve deformato, nempe in prioribus (stamineis) seu *regulari* nempe lobis 5 æqualibus, seu *subregulari*, seu *labiato* lobis extùs 3 intùs 2 in labia dispositis, seu *ringenti* lobis extùs 4 intùs 1 in labia duo dispositis, seu *obringenti* lobis extùs 1, intùs 4 in labia 2 dispositis, seu *palmato* lobo interno ad basin usque fisso cæteris vix ad medium divisus, seu tandem *fisso* lobo interiore brevissimo ad basin usque fisso, cæteris ad apicem usque ferè coalitis; in posterioribus fæmineis neutrisve (seu *ligulato*) nempe extùs in ligulam producto, seu *obligulato* nempe intùs in ligulam producto, seu *biligulato* nempe utrinquè in ligulam producto, seu *tubuloso* nempe limbo angusto et tubum simulante, seu *ampliato* nempe limbo ad basin in faucem latam dilatato. Stamina 5 cum corollâ inserta, cum ejus lobis alterna et tubo basi coalita, filamentis medio articulatis; antheræ oblongæ linearesve inter se in tubum coalitæ 4-valves valvis externis immobilibus, internis rimâ longitudinali dehiscenti-

bus semper ad apicem appendicem *terminalem*, sæpius ad basin appendices 2 *basilares* gerentibus. Ovarium 1 1-spermum supernè foveam nectariferam inter styli et corollæ staminum basin gerens. Stylus 1 in floribus androgynis teres cum apice nectararii articulatus, supernè divisus in ramulos 2 sæpius liberos interdum subadglutinatos, intus stigmatosos extus pilis pappulosis *collectores* dictis et pollen etenim colligentibus instructos; in femineis collectores nullos sed stigmatosas glandulas gerens, in masculis stigmatosas glandulas abortivas nullasque gerens sed collectores servans, in neutris rarissimis nec stigmata nec collectores gerens. Fructus a calycis tubo cum pericarpio proprio conflatus, unilocularis, evalvis, indehiscens, monospermus, sæpius sessilis, rarius pedicellatus ovario nempè suprâ basin calycis elevato stipitem tunc formante, tunc apice in collum post anthesin imprimis excrescens (pappi stipitem) attenuato, nunc in collum non attenuato, pappo nempè sessili. Placenta subcarnosa ad basin ovarii sita. Semen erectum. Albumen o. Embryo radiculâ subtereti inferâ. — Herbæ aut frutices. Flores aggregati in capitulos ramos caulemve terminantes, olim flores compositi dictos nunc seu *capitula*, seu *anthodia*, seu *calathides* nuncupatos. Cujusque capituli *receptaculum* seu pars pedunculi terminalis et plus minùsve dilatata, nunc membranaceum nunc carnosum, plus minùsve areolatum, flores gerens sæpius plurimos rarissimè solitarios. Circâ receptaculum est *involucrum* seu *Periclinium* constans squamis foliolisve plurimis nunc sæpius liberis, nunc rarius inter se coalitis, nunc uniseriatis nunc 2-∞-seriatis, seriebus internis sæpius longioribus; externâ (in biseriatis) interdum brevissimâ et tunc involucrum *calyculatum*. Super receptaculum et inter flores appendices nullæ aut variæ nempè squamellæ pili aut paleolæ. Cujusque capituli flores sunt inter se aut uniformes aut diffformes.

Subordo I. CORYMBIFERÆ *Vaill. Juss. gen. 177.*

Flores flosculosi aut radiati; receptaculum membranaceum aut vix carnosum; stigma super stylum non articulatum.

† SEMINA PAPPOSA, (excl. Bellide) PAPPO PLUS MINUS PILOSO.

I. EUPATORIUM *Tourn. t. 259. L. gen. 935.*

Involucrum cylindraceum, squamis imbricatis ovali-oblongis; flosculi pauci tubulosi hermaphroditi; receptaculum nudum; pappus pilosus.

x. E. CANNABINUM (L. sp. 1173.) foliis oppositis petiolatis 3-partitis, lacinii lanceolatis serratis, intermediâ longiore. *℥* in aquaticis et fossis humidis. — Fl. fr. 4. p. 129. — Lob. ic. t. 528. f. 2. (v. v.)

*β. indivium*, foliis superioribus indivisis.—Ray. syn. p. 180. — Moris. 3. s. 7. t. 13. n. 1. (v. s.)

## II. CACALIA L. gen. 933.

Involucrum simplex oblongum, squamis lineari-lanceolatis, basi subcalyculatum; flosculi omnes tubulosi hermaphroditi; receptaculum nudum; pappus pilosus.

1. *C. PETASITES* (Lam. dict. 1. p. 131.) caule herbaceo, foliis subtus incanis dentatis, radicalibus cordatis, inferioribus petiolatis, mediis acuminatis basi appendiculatis, superioribus sessilibus lanceolatis, corymbis fastigiatis, involucriis panicifloris.  $\mathcal{L}$  in Vogesis, Jurasso, Alpibus, monte-Aureo, et Pyrenæis. — Jacq. austr. t. 235. Fl. fr. 4. p. 127. excl. var. *β. C. albifrons* L. *Adenostyles albida* Cass. (v. s.)
2. *C. ALPINA* (Jacq. austr. t. 234.) caule herbaceo, foliis omnibus petiolatis cordatis dentatis glabris, superioribus acutis, corymbis fastigiatis, involucriis paucifloris.  $\mathcal{L}$  in saxosis humidis et umbrosis montium. — Fl. fr. 4. p. 127. *C. alpina β* L. sp. 1170. *Adenostyles viridis* Cass.—Lob. ic. t. 592. f. 1. (v. v.)
3. *C. LEUCOPHYLLA* (W. sp. 3. p. 1736.) caule herbaceo, foliis petiolatis utrinque tomentosis dentatis, inferioribus cordatis acuminatis, superioribus lanceolatis, floribus corymbosis, involucriis 15-20-floris.  $\mathcal{L}$  in petrosis montium Delphinatus, Galloprovincia. — Fl. fr. 4. p. 128. *C. tomentosa* Vill. dauph. 3. p. 171. *Adenostyles candidissima* Cass. (v. v. DC.)

## III. TUSSILAGO L. gen. 952.

Involucrum simplex, squamis margine membranaceis; flores flosculosi vel radiati; flosculi nunc omnes hermaphroditi, nunc feminei fertiles in radio, hermaphroditi in centro; pappus pilosus; receptaculum nudum.

### Sect. I. FARFARA DC. fl. fr. 4. p. 157.

Flores radiati, scapi uniflori.

1. *T. FARFARA* (L. sp. 1214.) scapo unifloro subnudo bracteato, foliis cordatis angulatis dentatis subtus pubescentibus.  $\mathcal{L}$  in argillosis subhumidis apricis. — Fl. fr. p. 4. p. 157. — Camer. epit. 590-591. ic. (v. v.)

### Sect. II. TUSSILAGO DC. l. c.

Flores flosculosi, caules uniflori basi foliosi.

2. *T. ALPINA* (L. sp. 1213.) caule unifloro subnudo, foliis reniformibus dentatis glabris.  $\mathcal{L}$  in pascuis editiorum montium. — Fl. fr. 4. p. 158. Lam. illust. t. 674. f. 7. *Homogyne Alpina* Cass. (v. v.)

### Sect. III. PETASITES DC. l. c.

Flores flosculosi, scapi multiflori.

3. *T. PETASITES* (Hop. tasch. 1803. p. 35.) thyrsio oblongo, foliis oblongo-cordatis inæqualiter denticulatis subtus pubescentibus, lobis approximatis.  $\mathcal{L}$  in humidis secus torrentes et fossas. — Fl. fr. 4. p. 158. Lam. ill. t. 674. f. 1. (v. v.)
4. *T. ALBA* (L. sp. 1214.) thyrsio fastigiato, foliis orbiculato-cordatis duplicato-argutè dentatis subtus villosa-pubescentibus.  $\mathcal{L}$  in humidis montanis Vogesorum, Jurassi, Burgundiæ, montis-Aurei. — Fl. fr. 4. p. 159. Fl. dan. t. 524. (v. v.)
5. *T. NIVKA* (Hop. tasch. 1803. p. 48.) thyrsio oblongo, foliis oblongo-cordatis inæqualiter dentatis subtus densè albo-tomentosis, lobis divaricatis.  $\mathcal{L}$  secus rivos in Alpibus editioribus Delphinatus, Galloprov., et summis Pyrenæis propè Benasque ex cl. Benth. — Fl. fr. 4. p. 159. — Moris. s. 7. t. 10. n. 4. (v. s.)
6. *T. FRAGRANS* (Vill. act. soc. hist. nat. par. 1. t. 12.) thyrsio fastigiato, floribus radiatis, foliis subrotundo-cordatis æqualiter denticulatis subtus pubescen-



tibus.  $\frac{1}{2}$  in Pyren. orient. ex Lois. Pourr. et Lapeyr. sed tamen dubia civis. — Fl. fr. 5. p. 472. *Nardosmia denticulata* Cass. (v. v. c.)

IV. CINERARIA *L. gen.* 957.

Involucrum simplex polyphyllum æquale; receptaculum nudum; flores radiati; pappus pilosus.

1. *C. CAMPESTRIS* (Retz obs. 1. p. 30.) floribus umbellatis, caule simplici, foliis tomentosis, radicalibus ovato-spathulatis subcrenulatis in petiolum coarctatis, caulinis sessilibus lanceolatis, superioribus integerrimis.  $\frac{1}{2}$  in collibus frondosis Parisiorum, Neustriæ, in Arvernæ montibus, Pyrenæis. — Fl. fr. 4. p. 169. — Clas. hist. 2. p. 22. f. 2. Fl. lutei. (v. v. DC.)

$\beta$ . *integrifolia*, pauciflora, foliis superioribus basi integerrimis. — *C. integrifolia* Jacq. aust. t. 179. Fl. fr. 4. p. 170. (v. s. ex Helv.)

2. *C. AURANTIACA* (W. sp. 3. p. 2081.) floribus subternis terminalibus, caule simplici, foliis sessilibus lanato-tomentosis, radicalibus ovatis obsolete crenatis, caulinis lanceolatis integerrimis, involucri sphaclatis.  $\frac{1}{2}$  in pratis siccis et editioribus Alpium. — Fl. fr. 4. p. 170. *C. Alpina* All. ped. t. 38. f. 2. non W. Fl. aurantiaci. (v. v.)

3. *C. LONGIFOLIA* (Jacq. aust. t. 181.) floribus umbellato-corymbosis, caule simplici, foliis radicalibus petiolatis spathulatis dentatis, caulinis oblongo-lanceolatis. ② in pascuis montanis irrigatis Delphinatûs, Galloprovinc., Pyren. orient. — Fl. fr. 4. p. 171. Fl. lutei. (v. s.)

4. *C. SIBIRICA* (L. sp. 1242.) racemo simplici folioso, foliis petiolatis glabris cordatis subcrenulatis, petiolo basi dilatato, caule simplici.  $\frac{1}{2}$  in paludosis montanis Pyrenæorum, Arvernæ, Ruthenensis (Rouergue) et Ligericini tractus (cl. Prost.). — Fl. fr. 4. p. 168. — Amm. ruth. t. 24. (v. s.)

5. *C. CORDIFOLIA* (Gou. ill. p. 69.) paniculâ pauciflorâ, caule simplici, foliis omnibus petiolatis cordatis duplicato-dentatis, petiolis basi subdentatis.  $\frac{1}{2}$  in Pyrenæis ex Lapeyr. — Fl. fr. 4. p. 171. — C. Bauh. prod. p. 69. f. 2. Dubia civis. (v. v. D. in Helv.)

6. *C. ALPINA* (W. sp. 3. p. 2084.) floribus subcorymbosis, caule simplici, foliis sessilibus subamplexicaulibus pinnatifidis, laciniis inæqualibus, lateralibus dentatis cuneatis, terminali inciso-dentatâ.  $\frac{1}{2}$  in Pyrenæis ex Lapeyr. — Jacq. austr. t. 177. (v. v. D. in Helv.)

7. *C. PALUSTRIS* (L. sp. 1243.) floribus corymbosis, foliis lato-lanceolatis dentato-sinuatis amplexicaulibus, caule villosa.  $\frac{1}{2}$  in arenosis aquosis et paludosis propè St.-Omer, Beauvais loco dicto *St.-Germer* (cl. Graves) et circâ Burdigalam ex Laterrade. — Fl. fr. 4. p. 169. — Lob. ic. t. 347. f. 2. (v. v. DC.)

8. *C. MARITIMA* (L. sp. 1244.) floribus paniculatis, involucri tomentosis, foliis pinnatifidis subtus cano-tomentosis, laciniis obtusis subtrilobis, caule suffrutescente.  $\frac{1}{2}$  in saxosis Galliæ Mediterraneæ et etiam in Delphinatû propè Di-gue. — Fl. fr. 4. p. 172. — Lob. ic. t. 227. f. 2. (v. v.)

V. SENECIO *L. gen.* 953.

Involucrum calyculatum, foliolis apice sphacelatis; flores nunc flosculosi, nunc radiati; receptaculum nudum; pappus pilosus, mollis.

§. 1. *Floribus radiatis, radio patente, foliis pinnatifidis.*

1. *S? LEUCOPHYLLUS* (DC. cat. 144.) foliis utrinquè lanatis niveis pinnatifido-lyratis, laciniis ovato-oblongis obtusis inferioribus distantibus mediis subincisis superioribus subcoarctatis, floribus corymbosis confertis.  $\frac{1}{2}$  in editioribus Pyrenæorum orientalium et montis Mesin. — Fl. fr. 5. p. 473. *S. incanus?* Lapeyr. abr. *S. palmatus* Lsp. suppl. An potius *Cinerariæ* species? (v. v.)

2. *S. INCANUS* (L. sp. 1219.) foliis utrinquè tomentosis niveis pinnatifidis, laciniis linearibus cuneatis obtusis subdentatis, corymbo coarctato.  $\frac{1}{2}$  in Alpibus Delphinatûs. — Fl. fr. 4. p. 162. — Barr. ic. t. 262. f. 1. (v. v. in Helv.)

3. *S. ARTEMISIAEFOLIUS* (Pers. ench. 2. p. 435.) foliis pinnato-multifidis, pinnis lineari-subulatis subincisis, involucri cylindricis demum longitudinaliter sulcatis, floribus corymbosis terminalibus.  $\mathcal{N}$  circa Parisios, Aureliam et in montosis Gallie mediz, australis et orientalis. — Fl. fr. 5. p. 472. *S. tenuifolius* Fl. fr. 4. p. 164. excl. syn. Jacq. Hoffm. et Huds. (v. v.)
5. *S. ASOTANFOLIUS* (L. sp. 1219.) foliis pinnatifido-multifidis linearibus nudis acutis, caule superne subundo subbifloro, squamis lanceolatis.  $\mathcal{N}$  in Alpibus Galloprovinciz ex Ger. — Fl. fr. 4. p. 164. — Clus. hist. 1. p. 334. f. 1. (v. s. ex Anstria.)
4. *S. JACOBÆ* (L. sp. 1219.) foliis radicalibus lyrato-pinnatifidis, caulinis pinnatifidis glabris lobis divaricatis lineari-lanceolatis subpinnatis, caule erecto, involucri subcylindrici squamis lanceolato-ovatis.  $\mathcal{N}$  in pratis, petrosis et ad vias frequens. — Fl. fr. 4. p. 163. — Fuchs. hist. 742. ic. (v. v.)
6. *S. AQUATICUS* (Huds. angl. 366.) foliis lyratis serratis glabris, inferioribus obovatis integris, involucri hemispharico, ligulis ellipticis.  $\mathcal{N}$  in paludibus et aquaticis. — Fl. fr. 4. p. 163. excl. syn. Clus. Fl. dan. t. 784. *S. Jacobæ* varietas? (v. v.)
7. *S. ZUCARFOLIUS* (Huds. angl. 366. non L?) foliis pinnatifidis subhirtis, lobis lanceolatis dentatis, terminali inciso-dentato, caule erecto, involucri squamis lanceolato-ovatis.  $\mathcal{N}$  in montosis dumetosis. — Fl. fr. 4. p. 164. (v. v.)  
*β. tenuifolius*, lacinii angustioribus. — In Delphinatū. — *S. tenuifolius* Jacq. austr. 3. t. 278. non Fl. fr. — Barr. ic. t. 153. (v. s.)
8. *S. SQUALIDUS* (L. sp. 1218.) foliis semiamplexicaulibus pinnatifidis, lacinii linearibus subdentatis distantibus, caule erecto, involucri squamis linearibus lucidis glaberrimis. ① in arvis, vineis, ad muros et vias Gallie mediterraneæ. — Fl. fr. 4. p. 162. *S. Gallicus* Vill. Daph. 3. p. 230. DC. syn. gall. p. 283. — Barr. ic. t. 262. f. 2. (v. v.)
9. *S. CRASSIFOLIUS* (W. sp. 3. p. 1982.) foliis carnosis amplexicaulibus indivisis partitisve crenatis, pedunculis squamosis, involucri squamis linearibus laxis glaberrimis. ① ex maritimis Massilie misit cl. Solier. — Fl. fr. 5. p. 472. — Barr. ic. 251. (v. s.)

### §. 2. *Floribus radiatis, radio revoluto.*

10. *S. SYLVATICUS* (L. sp. 1217.) foliis radicalibus indivisis dentatis, superioribus pinnatifidis lobis denticulatis, involucri cylindrici squamis subulatis. ① L. ② Vill. in sylvis submontanis. — Fl. fr. 4. p. 161. — Dill. elth. t. 258. f. 327. (v. v.)
11. *S. VISCOSUS* (L. sp. 1217.) foliis pinnatifidis viscidis, involucri hemispharici squamis laxis hirsutis. ① ad oras sylvarum in montosis. — Fl. fr. 4. p. 168. — Dill. elth. t. 258. f. 336. (v. v.)
12. *S. HUMILIS* (Desf. atl. 2. p. 271. t. 233.) caule procumbente, foliis subpaulatis obtusis duplicato-dentatis, floribus axillaribus terminalibusque subsolitariis, squamis glabris. ① in paludosis maritimis Corsicæ (Ph. Thomas). (v. s.)
13. *S. VERNICULACEUS* (Ten. fl. nap. t. 78.) foliis radicalibus petiolatis ovatis inciso-dentatis, superioribus subamplexicaulibus pinnatifidis lobis ovato-lanceolatis crenato-dentatis, floribus corymbosis, squamis lucidis glabris. ① in valle Asco Corsicæ reperiit Ph. Thomas. Folia fœniculum redolent ex Reyn. in h. DC. (v. s.)
14. *S. LIVIDUS* (L. sp. 1216.) caule recto simplici, foliis sinuato-dentatis, inferioribus ovatis petiolatis, superioribus oblongis amplexicaulibus, floribus subcorymbosis, squamis lucidis glabris. ① in maritimis Corsicæ, Galloprovinciæ. — Fl. fr. 5. p. 472. *S. nebrodensis* Fl. fr. 4. p. 162. excl. syn. (v. s.)

### §. 3. *Floribus radiatis, radio patente, foliis indivisis.*

15. *S. PALUDOSUS* (L. sp. 1220.) foliis semi-amplexicaulibus lanceolatis argutè

- serratis subtus adpressè villosis, corymbo terminali divaricato, caule stricto sublaesto.  $\mathcal{L}$  secus rivios et stagna in arundinetis. — Fl. fr. 4. p. 166. — Dalech. lugd. 1037. f. 2. (v. s.)
16. *S. TOURNEFORTII* (Lapeyr. abr. pyr. 516.) foliis oblongis utrinquè attenuatis subcrassis utrinquè glabris, dentibus semi-lanatis, corymbo paucifloro.  $\mathcal{L}$  in Pyrenæis editioribus. — *S. persicæfolius* Fl. fr. 4. p. 166. non L. Ram. bull. phil. n. 43. t. 11. f. 3. (v. v.)
17. *S. DORIA* (L. sp. 1221.) squamis involucri exterioribus patulis, foliis subdecurrentibus oblongo-lanceolatis glaucescentibus glabris serratis, corymbo multifloro.  $\mathcal{L}$  secus rivos in Galliâ australi. — Fl. fr. 4. p. 167. Jacq. austr. t. 185. (v. v. DC.)
18. *S. CACALIASTER* (Lam. fl. fr. 2. p. 132.) foliis lanceolatis argutè serratis, inferioribus decurrentibus, involucri squamis interioribus lineari-lanceolatis, floribus (ochroleucis) corymbosis aspièus flosculosis.  $\mathcal{L}$  in Vogesis et montibus Arvernæ. — Fl. fr. 5. p. 474. *Cacalis Sarraценica* L. sp. 1269. Fl. fr. 4. p. 128. (v. s.)
19. *S. SARRACENICUS* (L. sp. 1221.) foliis lanceolatis basi cuneatis argutè serratis subsessilibus glabriusculis, involucri squamis lineari-spathulatis, corymbo coarctato.  $\mathcal{L}$  in humidis umbrosis Alpium, Jurassi, agri Snessionensis et Forensis (le Forez). — Fl. fr. 4. p. 167. — Fuchs. hist. 728. ic. (v. v.)
20. *S. MEMORANSIS* (L. sp. 1221.) foliis sessilibus ovato-lanceolatis serratis margine ciliatis subtus tenuissimè pubescentibus, corymbo terminali folioso laxo.  $\mathcal{L}$  in montibus Galliæ australis, in Pyrenæis orientalibus. — Fl. fr. 4. p. 167. — Pluck. t. 236. f. 1. (v. s.)
21. *S. DORONICUM* (L. sp. 1222.) caule indiviso subtomentoso, foliis crassiusculis serratis, caulinis sessilibus, radicalibus petiolatis ovatis orbiculatisve subtus villosis.  $\mathcal{L}$  in pascuis subhumidis Alpium, Pyren., Cebennarum. — Fl. fr. 4. p. 168. — Clus. hist. 2. p. 17. f. 1. (v. v.)
- $\beta$ . *Barrelieri*, caule multifloro. — In Pyrenæis orient. — *S. Barrelieri* Gouill. 68? — Barr. ic. 146. 801. (v. v.)
- $\gamma$ . *tomentosus*, foliis subtus tomentosis. — *Lepicaune tomentosa* Lapeyr. abr. ex autopsiâ cl. Arnott. (v. s.)

#### §. 4. *Floribus flosculosis.*

22. *S. VULGARIS* (L. sp. 1216.) foliis amplexicaulibus pinnatifidis dentatis, floribus corymboso-coarctatis. ① in cultis frequens. — Fl. fr. 4. p. 161. Gærtn. fruct. 2. t. 166. f. 3. (v. v.)

#### VI. DORONICUM L. gen. n. 959.

Involucri squamæ duplcis ordinis æquales; flores radiati ligulis femineis; flosculorum semina pappo simplici coronata, ligularum nuda; receptaculum nudum.

1. *D. PARDALIANCHES* (L. sp. 1247.) foliis denticulatis, radicalibus cordatis longè petiolatis, mediis spathulato-cordatis, summis subrotundo-cordatis, caule hirsuto subsimplici.  $\mathcal{L}$  in sylvis montanis. — Fl. fr. 4. p. 173. — Cam. epit. 823. ic. (v. v.)
2. *D. SCORPIOIDES* (W. sp. 3. p. 2114.) foliis remotè denticulatis, superioribus oblongis amplexicaulibus, inferioribus ovatis petiolatis, petiolis alatis auriculatis, radicalibus ovatis subcordatis petiolatis, caule subglabro.  $\mathcal{L}$  in sylvis montanis Alpium, Pyrenæorum. — Fl. fr. 4. p. 174. — Clus. hist. 2. p. 16. (v. v.)
3. *D. PLANTAGINEUM* (L. sp. 1247.) foliis ovali-oblongis, inferioribus longè petiolatis, petiolis nudis, superioribus sessilibus amplexicaulibus, caule simplici.  $\mathcal{L}$  in Vogesis, Parisiis, Sinemuri, secus Gardanem (*Gardon*). — Fl. fr. 4. p. 174. — Dalech. hist. 1202. f. 2. (v. v. DC.)
4. *D. AUSTRIACUM* (Jacq. austr. t. 130.) foliis denticulatis, summis lanceolatis

amplexicanlibus, inferioribus spathulato-ovatis, radicalibus cordatis petiolatis.  $\mathcal{U}$  in Pyrenæis, montibus Ligerinæ (*Lozère*), Arvernæ, monte Pilato ex cl. Balbis, et ad ripas Druentis ex Lois.—Fl. fr. 5. p. 475.—Clus. hist. 2. p. 19. ic. (v. s.)

VII. ARNICA *L. gen. n. 958.*

Involucri squamæ duplicis ordinis æquales; flores radiati, ligulis femineis filamentis 5 sterilibus donatis; semina omnia pappo coronata; receptaculum nudum.

1. A. CORSICA (Lois. gall. 576. t. 20.) caule erecto supernè ramoso, foliis amplexicanlibus lanceolato-oblongis argutè denticulatis, floribus subcorymbosis.  $\mathcal{U}$  in Corsicà ad rupes aquis fluentibus madidas.—Fl. fr. 5. p. 475. Doronicum Corsicum Poir. enc. supp. 2. p. 517. Dor. Austriaco valdè similis. (v. s.)
2. A. DORONICUM (Jacq. aust. t. 82.) foliis remotè dentatis hirsutis, radicalibus petiolatis oblongis basi angustatis, caulinis alternis oblongo-lanceolatis obtusis.  $\mathcal{U}$  in petrosis Alpium editiorum Delphinatûs et Pyren. centr. ex cl. Benth.—Fl. fr. 4. p. 175. Grammarthron biligulatum Cass.—Clus. hist. 2. p. 17. f. 1. (v. v. DC.)
3. A. SCORPIOIDES (L. sp. 1246.) foliis dentatis dentibus acuminatis, radicalibus petiolatis elliptico-subrotundis, caulinis alternis oblongis acutis.  $\mathcal{U}$  in montibus Delphinatûs, Arvernæ, Pyrenæorum.—Fl. fr. 4. p. 176. Grammarthron scorpioides Cass.—Lob. ic. t. 649. f. 1. (v. v.)
4. A. MONTANA (L. sp. 1245.) foliis oppositis ovatis integris.  $\mathcal{U}$  in pascuis montanis.—Fl. fr. 4. p. 175.—Clus. hist. 2. p. 18. f. 1. (v. v.)  
*f. angustifolia*, foliis angustioribus, floribus sæpè cernuis.—In ericetis circà Dax.—Cineraria cernua Thore land. 344. (v. v.)

VIII. CHRYSOCOMA *L. gen. n. 939.*

Involucrum imbricatum, hemisphæricum vel ovatum squamis linearibus; receptaculum alveolatum; flosculi omnes hermaphroditi tubulosi; pappus pilosus, ciliatus.

1. C. LIXOSTRIS (L. sp. 1198.) herbacea, foliis linearibus glabris, involucris laxis, corollis sub-5-lobis.  $\mathcal{U}$  in argillosis apricis Gallis mediæ, orientalis, australis occidentalisque.—Fl. fr. 4. p. 141.—Lob. ic. 409. f. 1. (v. v.)
2. C. SAXATILIS (DC. fl. fr. 5. p. 468.) herbacea, foliis lanceolato-linearibus glutinosis pilosis, involucri multiseriati squamis exterioribus apice reflexis, corollis vix 5-dentatis.  $\mathcal{U}$  in rupestribus circà Massiliam et Telonem in Occitaniâ et agro Ruacinonensi ex cl. Benth.—*Isula saxatilis* Fl. fr. 4. p. 154. Erigeron glutinosum L. mant. 112. Chiliadenus camphoratus Cass.—Barr. ic. 158. (v. s.)

IX. ASTER *L. gen. n. 954.*

Involucrum imbricatum, squamis exterioribus patulis; receptaculum nudum; flores radiati, ligulis femineis fertilibus discoloribus oblongis; pappus pilosus.

1. A. ALPINUS (L. sp. 1226.) caule unifloro, foliis integerrimis radicalibus lanceolato-spathulatis, caulinis lanceolatis, invol. squamis subæqualibus lanceolatis obtusiusculis.  $\mathcal{U}$  in pascuis editiorum montium.—Fl. fr. 4. p. 144.—Clus. hist. 2. p. 15. f. 2. (v. v.)
2. A. AMELLUS (L. sp. 1226. non W.) caule corymboso, foliis oblongo-lanceolatis integris scabris, inferioribus obtusis, superioribus acutis, involucris subquarrosis, squamis obtusis, interioribus membranaceis apice coloratis.  $\mathcal{U}$  in Jurassi, Delph., Galloprov., Occit., Pyren. or. collibus, circà Aginnum et Corisopilum (Quimper).—Fl. fr. 4. p. 145. 5. p. 469.—Clus. hist. 2. p. 16. f. 1. (v. v.)

3. *A. PYRÆNEUS* (Desf. cat. par. 102.) caule hirsuto simplici suprà corymboso, foliis oblongo-lanceolatis subscabris subamplexicaulibus apice 4-5-dentatis, invol. squamis lineari-acuminatis hirsutis.  $\mathcal{U}$  in Pyrenæorum monte Esquierri. — Fl. fr. 4. p. 146. (v. s. c.)
4. *A. SALIGNUS* (W. sp. 3. p. 2040.) caule erecto glabro paniculato-corymboso, foliis lineari-lanceolatis acuminatis sessilibus integris glaberrimis impunctatis, inferioribus lanceolatis apice serratis, invol. squamis lineari-lanceolatis, interioribus acutissimis.  $\mathcal{U}$  in pratis humidis Argentorati, Mimatis. — Fl. fr. 5. p. 470. (v. s.)
5. *A. ACRIS* (L. sp. 1228.) caule subglabro corymboso, foliis lineari-lanceolatis basi sub 3-nerviis punctatis acutis, ramis foliosis, invol. squamis lineari-lanceolatis acutis.  $\mathcal{U}$  ad sepes et in herbosis Gallie mediterraneæ. — Fl. fr. 4. p. 146. — Carid. aix. t. 11. (v. s.)
6. *A. TAIFOLIUM* (L. sp. 1226.) caule glabro corymboso, foliis caulinis lineari-lanceolatis carnosius triplinerviis, inferioribus apice subserratis, invol. squamis laxis lanceolatis obtusis.  $\mathcal{U}$  in stagnosis maritimis et in salsis Lotharingie. — Fl. fr. 4. p. 145. — Gm. sib. 2. t. 80. f. 2. (v. v.)
7. *A. ANNuus* (L. sp. 1229.) caule subglabro supernè ramoso, foliis pilosiusculis, inferioribus subovatis serratis, superioribus lanceolatis, involucri hemisphæricis squamis subæqualibus lineari-lanceolatis acutis strigosis. (1) æcus Isaram ex Will., Rhenum ex Sch. — Fl. fr. 4. p. 146. Fl. dan. t. 486. *Diplopappus dubius* Cass. (v. s. ex Valesiâ.)
8. *A. CHINENSIS* (L. sp. 1232.) caule hispido, ramis unifloris, foliis ovatis grossè dentatis petiolatis, caulinis sessilibus lanceolato-acuminatis integris, involucri squamis maximis ciliatis. (1) ex Chinâ ortus, in hortis cultus. — Fl. fr. 4. p. 147. — Dill. elt. t. 34. f. 38. (v. v.)

X. ERIGERON *L. gen.* 951.

Involucrum oblongum imbricatum; receptaculum nudum; flores radiati, ligulis femineis paucis discoloribus linearibus angustissimis; pappus pilosus.

1. *E. CANADENSE* (L. sp. 1210.) caule hirtio, floribus paniculatis, foliis lanceolatis ciliatis. (1) in lapidosis et sylvis. — Fl. fr. 4. p. 144. — Moris. s. 7. t. 20. n. 29. (v. v.)
2. *E. ACRE* (L. sp. 1211.) caule multifloro, foliis radicalibus obtusis, caulinis acutis, involucri squamis linearibus acutissimis, pappo rufescente seminibus duplò longiore.  $\mathcal{U}$  in sterilibus et apricis frequens. — Fl. fr. 4. p. 142. *E. murale* Lapeyr. supp. — Col. ecphr. t. 26. f. 2. (v. v.)
3. *E. ALPINUM* (Lam. fl. fr. 2. p. 140.) caule paucifloro, foliis subtus villosis, caulinis subobtusis, invol. squamis lanceolatis acutis, pappo rufescente seminibus longitudine.  $\mathcal{U}$  in montosis frequens. — Fl. fr. 4. p. 142. (v. v.)  
 $\beta$ . *uniflorum*, caule 1-floro, foliis subglabris. — In editoribus. — *E. uniflorum* L. sp. 1211. (v. v.)
4. *E. VILLARSII* (Bell. act. tur. 5. p. 244. t. 9.) caule paniculato, foliis lanceolatis viscosis trinerviis scabris subdentatis sessilibus, squamis lineari-lanceolatis acutissimis, radio disco brevior.  $\mathcal{U}$  in editoribus Delphin., Galloprov., Pyren. — Fl. fr. 4. p. 143. (v. s.)

XI. SOLIDAGO *L. gen.* 955.

Involucrum imbricatum; receptaculum nudum; flores radiati, ligulis concoloribus subquinis; pappus pilosus.

1. *S. GRAVEOLENS* (Lam. fl. fr. 2. p. 145.) pubescenti-viscosa, foliis sublinearibus integerrimis, ramis lateralibus multifloris. (1) in arvis subhumidis Gallie australis et occidentalis. — Fl. fr. 4. p. 156. *Erigeron graveolens* L. sp. 1210. *Inula graveolens* Desf. Cass. — Barr. ic. t. 370. (v. v.)

2. *S. MINUTA* (L. sp. 1235.) caule simplicissimo, foliis lanceolatis acutis serratis glabris, pedunculis unifloris flore duplò longioribus.  $\mathcal{F}$  in Pyrenæis. — Fl. fr. 4. p. 156. — Herm. par. t. 245. (v. s.)
3. *S. VIRGA-AUREA* (L. sp. 1235.) caule supernè ramoso et pubescente, foliis caulinis lanceolatis utrinquè attenuatis serratis, inferioribus ellipticis pilosisculis, racemis erectis, pedicellis flore brevioribus.  $\mathcal{F}$  in petrosis sylvarum. — Fl. fr. 4. p. 156. Fl. dan. t. 668. (v. v.)
- $\beta$ . *ericetorum*, foliis subintegris lineari-lanceolatis.  $\mathcal{F}$  in sabulosis propè Dax. — *S. minuta* Thore non L. nec Vill. (v. v.)

XII. *BELLIDIASTRUM* Cass. dict. sc. nat. — Mich. gen. t. 29.

Involucrum 1-seriatum, squamis linearibus acutis; flores radiati, in disco hermaphroditi, in radio feminei, receptaculum conicum nudum; semina striata spinulosa.

1. *B. MICHELII* (Cass. in dict. sc. nat. 4. supp. p. 70.) scapo unifloro nudo, radio discolore, foliis petiolatis obovatis repandis.  $\mathcal{F}$  in sylvis montosis Galloprovinciæ, Delphinatûs. — *Doronicum Bellidiastrum* L. sp. 1247. Fl. fr. 4. p. 176. — Cam. epit. 654. ic. (v. v.)

XIII. *BELLIS* Tourn. t. 280. L. gen. n. 962.

Involucrum hemisphericum polyphyllum simplex, squamis lanceolatis; flores radiati; receptaculum nudum conicum; semina nuda.

1. *B. ANNUA* (L. sp. 1249.) caule subsimplici subfolioso, foliis ovali-spathulatis obtusis apice crenulatis, invol. squamis subobtusis glabriusculis apice ciliatis. (I) in pratis et collibus Galliæ mediterraneæ et Corsicæ. — Fl. fr. 4. p. 186. (v. v.)
2. *B. SYLVESTRIS* (Cyr. rar. 2. p. 22. t. 4.) scapo longo nudo unifloro, foliis oblongo-lanceolatis subacutis remotè denticulatis velutino-pubescentibus trinerviis, invol. squamis lanceolatis acutis hirtis.  $\mathcal{F}$  circà Telouem, Massiliam, Avenionem, in Occitaniâ et agro Rusciniensi ex Benth. — Fl. fr. 5. p. 478. — Dod. pempt. 265. f. 1. (v. s.)
3. *B. PERENNIS* (L. sp. 1248.) scapo subnudo unifloro, foliis spathulato-obovatis crenato-dentatis pubescentibus venosis, squamis lanceolato-ovatis obtusis hirtis.  $\mathcal{F}$  in graminosis ad vias. — Fl. fr. 4. p. 185. Lam. ill. t. 677. (v. v.)

XIV. *BELLIUM* L. gen. pl. ed. Schr. p. 561. DC. fl. fr. 4. p. 923.

Involucrum simplex polyphyllum, squamis lineari-spathulatis; receptaculum nudum conicum; flores radiati, flosculi 4-dentati tetrandri; semina paleis membranaceis 4-8 in pilos desinentibus coronata.

1. *B. BELLIDIOIDES* (L. mant. 285.) caule stolonifero, foliis spathulatis integris, scapis nudis filiformibus foliis triplò longioribus, seminibus paleis 4-8 coronatis. (I) in rupestribus maritimis Corsicæ. — Fl. fr. 4. p. 923 non Desf.; 5. p. 475. — Bocc. mns. p. 149. t. 107. (v. s.)
2. *B. NEVALE* (Req. in ann. sc. nat. 5. p. 383.) caule nudo, foliis spathulatis integris, scapis nudis villosis foliis vix duplò longioribus, seminibus paleis 4 coronatis. (I) in editioribus montibus Corsicæ. An satis à priore distincta? (v. s. in spec. non satis perfectis.)

XV. *CONYZA* L. gen. 950. DC. fl. fr. 4. p. 189.

Involucrum subrotundatum imbricatum; receptaculum nudum; flosculi omnes tubulosi, in centro hermaphroditi 5-dentati, in radio feminini steriles graciles 3-dentati; pappus pilosus ciliatus.

§. 1. *Involucro sub 1-seriato.*

1. *C. AMIGUA* (DC. fl. fr. 5. p. 468.) caule erecto ramoso piloso striato, foliis omnibus linearibus scabris planis hinc indè dentatis, capitulis paniculatis.

① in pratis circa Nemausim, Monspelium, Narbonem (Pourr.) cum *Erigone canadense*. — *Erigeron crispum* Pourr. chl. n. 470. (v. v. DC.)

§. 2. *Involucro multiseriato.*

2. *C. SICULA* (W. sp. 3. p. 1931) caule erecto lævi ramoso, foliis inferioribus lanceolatis, superioribus linearibus integerrimis margine revolutis, capitulis paniculatis. ① in fossis humidis et stagnis maritimis prov. mediterranearum. — Fl. fr. 4. p. 139. *Dimorphantes Sicula* Cass. — Magn. bot. monsp. t. 76. (v. s.)
3. *C. SQUARROSA* (L. sp. 1205.) caule erecto ramoso, foliis ovali-lanceolatis pubescentibus, inferioribus dentatis in petiolum attenuatis, capitulis corymbosis. ② ad sepes et oras sylvarum in siccis. — Fl. fr. 4. p. 139. Lam. ill. t. 697. f. 1. (v. v.)
4. *C. SAXATILIS* (L. sp. 1206.) caule suffruticoso prostrato, ramis cano-tomentosis, foliis linearibus margine revolutis subdentatis, pedunculis longissimis trifloris. ③ ad muros et inter saxa maritima Corsicæ, agri Ruscinonenſis et circa Nicæam, Olbiam. — Fl. fr. 4. p. 140. *Phagnalon subdentatum* Cass. — Barr. ic. t. 426. (v. v.)
5. *C. SORDIDA* (L. mant. 466.) caule suffruticoso prostrato, ramis cano-tomentosis, foliis linearibus integerrimis margine revolutis, pedunculis longis trifloris. ④ ad muros et saxa Gallie mediterraneæ, propè Nannetes? ex Bon. — Fl. fr. 4. p. 140. *Phagnalon tricephalum* Cass. — Barr. ic. t. 277 et 368. (v. v.)

XVI. *INULA* L. gen. 956.

Involucrum imbricatum; receptaculum nudum; flores radiati ligulis concoloribus; antheræ sæpius basi bisetosæ; pappus nunc pilosus simplex, nunc duplex, exterior membranaceus.

Sect. I. *COMVISARTIA*. Mer. fl. par. ed. 2. t. 2. p. 261.

*Squamæ* exteriores obovatæ, inter. *spatulatæ* coloratæ, pappus simplex.

1. *I. HELENIUM* (L. sp. 1236.) foliis amplexicaulibus subdentatis ovatis rugosis subtus tomentosis. ① circa Parisios, in Neustriâ, Pyrenæis ex Lapeyr. — Fl. fr. 4. p. 148. *Corvisartia Helenium* Mer. fl. par. 1. c. — Lob. ic. t. 574. f. 2. (v. s.)

Sect. II. *ENULA*.

*Squamæ* lineares lanceolataeve acutæ.

\* *Pappo simplici.*

2. *I. CRITHMOIDES* (L. sp. 1240.) foliis linearibus carnosis 5-cuspidatis, inv. squamis lineari-lanceolatis, seminibus hirtis. ② in paludosis maritimis ex Nicæâ ad Cherbourg usque (cl. de Breh.). — Fl. fr. 4. p. 154. *Limbarda tricuspidata* Cass. — Lob. ic. t. 395. f. 2. (v. v.)
3. *I. TUBEROSA* (Lam. dict. 3. p. 260.) caule suffruticoso, ramis unifloris, foliis linearibus rarò dentatis margine subhirsutis, involucri squamis linearibus, seminibus hirtis. ③ in siccis et maritimis Occitanie et in Cebennis. — Fl. fr. 4. p. 153. — Lob. ic. t. 350. f. 3. (v. v.)
4. *I. BRITANICA* (L. sp. 1237.) caule corymboso villosa, foliis amplexicaulibus lanceolatis basi serratis subtus pilosis, involucri squamis linearibus seminibusque hirsutis. ④ secus vias et fossas passim. — Fl. fr. 4. p. 149. — Moris. s. 7. t. 19. n. 8. *Britanica*. Dalech. hist. 1082. (v. v.)
5. *I. ALPINA* (L. sp. 1236.) foliis ovato-oblongis decurrentibus dentatis apice integerrimis, floribus corymbosis confertis, squamis linearibus ciliatis, seminibus glabris. ⑤ in montosis provinc. austral. — Fl. fr. 4. p. 155. — Garid. aix. t. 23. (v. v.)
6. *I. SQUARROSA* (L. sp. 1240.) foliis ovalibus rigidis tenuissimè serrulatis lucidis scabris reticulato-venosis, invol. squamis glabris exterioribus lanceolato-ovatis reflexis, seminibus glaberrimis. ⑥ in provinciis australibus. — Fl. fr. 4. p. 150. *I. Bubonium* Jacq. Vill. Lapeyr. — Pluckn. t. 16. f. 1. (v. v.)

- β. Germanica*, foliis ovato-oblongis. — In Delphinatu (h. DC.) et Santoniâ (Des Moulins.) — *I. Germanica*. L. Fl. fr. 4. p. 150. — Moris. s. 7. t. 19. n. 16. (v. s.)
7. *I. VAILLANTII* (Vill. dauph. 2. p. 216.) caule paniculato, foliis ovato-lanceolatis obsolete serratis subtus tomentosis, floribus subcorymbosis terminalibus, squamis exterioribus lanceolatis, interioribus linearibus ciliatis seminibusque glabris.  $\frac{1}{2}$  secus rivos et oras sylvarum Delphinatûs. — Fl. fr. 4. p. 152. — Hall. helv. n. 73. t. 2. (v. v. DC.)
8. *I. SALICINA* (L. sp. 1238.) foliis lanceolatis recurvis glaberrimis lucidis tenuissimè serrato-scabris, ramis angulatis glabriusculis, floribus inferioribus altioribus, squamis ciliatis exterioribus lanceolatis, seminibus glabris.  $\frac{1}{2}$  in pratis montosis. — Fl. fr. 4. p. 151. — Clus. hist. 2. p. 15. f. 1. Folia sæpè subamplexicaulia. (v. v.)
9. *I. HIRTA* (L. sp. 1239.) caule villosa subramoso, foliis lanceolato-oblongis integris rigidis hirtis, floribus paucis corymbosis, squamis exterioribus interioribus superantibus, seminibus glabris.  $\frac{1}{2}$  in pratis montosis et circà Parisios. — Fl. fr. 4. p. 151. — Clus. hist. 2. p. 14. f. 2. (v. s.)
10. *I. HYLENORDEZ* (DC. fl. fr. 5. p. 470.) foliis amplexicaulibus ovato-oblongis subhirsutis, radicalibus in petiolum angustatis subdentatis, floribus 3-4 corymbosis amplis, involucri hirsutissimi squamis externis lanceolatis foliaceis patulis.  $\frac{1}{2}$  secus agros Pyrenæorum orient. (v. v.)
11. *I. STAVIOLENS* (Ait. kew. 3. p. 224.) caule multifloro, foliis caulinis oblongis sessilibus subtus pilosis, radicalibus ellipticis subdentatis in petiolum longè angustatis, invol. squamis hirtis externis oblongis laxis, seminibus breviter pubescentibus.  $\frac{1}{2}$  in asperis Galloprovinciæ. — Fl. fr. 5. p. 471. I. oculus Christi Fl. fr. 4. p. 148. excl. syn. (v. s.)
12. *I. MONTANA* (L. sp. 1241.) caule unifloro, foliis lanceolatis hirsutis integerimis, involucri hirsuti squamis exterioribus oblongo-lanceolatis obtusis, interioribus linearibus acutis, seminibus hirtis.  $\frac{1}{2}$  in sterilibus montium provinc. austral. — Fl. fr. 4. p. 154. — Lob. ic. t. 350. f. 2. (v. v.)

\* \* *Pappo duplici.*

13. *I. ODORA* (L. sp. 1236.) caule paucifloro, foliis amplexicaulibus dentatis hirsutis, radicalibus petiolatis ovatis, caulinis sessilibus lanceolatis, superioribus basi auriculatis, pappo exteriori cupulari membranaceo 10-dentato.  $\frac{1}{2}$  in Corsicâ, Galloprov. et agro Ruscinonensi inter Collioure et Bagnols. — Fl. fr. 4. p. 148. — Barr. ic. t. 1145. (v. s.)
14. *I. VISCOSA* (Desf. atl. 2. p. 274.) hirsuto-viscosa, caule superne ramoso, foliis lanceolatis subserratis, pedunculis lateralibus unifloris foliosis, invol. squamis lineari-lanceolatis glabris, pappo externo brevissimo patulo.  $\frac{1}{2}$  in locis incultis, ad sepes et margines agrorum Galliæ australis. — Fl. fr. 4. p. 153. — Clus. hist. 2. p. 20. f. 1. (v. v.)
15. *I. DYSENTERICA* (L. sp. 1237.) caule pubescente paniculato, ramis lateralibus apice longioribus patulis, foliis amplexicaulibus cordato-oblongis nudis serratis subtus pilosis, invol. squamis setaceis, pappo externo brevi cupulari subintegro.  $\frac{1}{2}$  in fossis et humidis. — Fl. fr. 4. p. 149. Pulicaria dysenterica Cass. — Fuchs. hist. 436. ic. (v. v.)
16. *I. PULICARIA* (L. sp. 1238.) caule erecto paniculato, foliis amplexicaulibus oblongis undulatis villosis, pedunculis unifloris oppositifoliis, floribus subglobosis, radio minutissimo, inv. squamis linearibus, pappo externo profundè dentato subpiloso. ① in fossis humidis secus vias. — Fl. fr. 4. p. 150. Pulicaria vulgaris Cass. — Blackw. t. 103. (v. v.)

XVII. GNAPHALIUM Lam. DC. fl. fr. 4. p. 133.

Involucrum imbricatum, squamis margine sæpius coloratis, recep-



taculum nudum; flosculi omnes tubulosi, alii hermaphroditi, alteri feminini; pappus pilosus aut apice dentatus.

Sect. I. GNAPHALON. — Filago *Gærtn. non L.*

Flosculi exteriores feminei fertiles, pappus capillaris.

1. *G. LUTEO-ALBUM* (L. sp. 1196.) caule herbaceo simplici erecto, foliis semiamplexicaulibus lineari-lanceolatis subrepandis utrinquè pubescentibus, inferioribus obtusis, superioribus acutis, corymbo glomerato. ① in humidis. — Fl. fr. 4. p. 133. — Barr. ic. t. 367. (v. v.)
2. *G. SYLVATICUM* (L. sp. 1200.) caule herbaceo simplicissimo erecto tomentoso, foliis lanceolatis utrinquè attenuatis et lanatis, floribus terminalibus axillaribusque sessilibus spicatis. ② in sylvis. — Fl. fr. 4. p. 134.  
  - a. *fuscum*, squamis fuscis, foliis lanceolatis. — *G. fuscum* Fl. dan. t. 251. (v. v.)
  - β. *rectum*, foliis caulinis lineari-lanceolatis. — *G. rectum* Sm. fl. brit. 1. p. 870. Fl. dan. t. 1229. (v. v.)
3. *G. SUPINUM* (L. syst. veg. 623.) caule herbaceo simplici procumbente aut minimo, foliis linearibus acutis tomentosis, floribus capitatis. ② in pratis frigidioribus inter saxa et secus torrentes editiorum montium. — Fl. fr. 4. p. 133. — Bocc. mus. t. 85. (v. v.)
4. *G. ULIGINOSUM* (L. sp. 1201.) caule herbaceo ramoso diffuso lanato, foliis lineari-lanceolatis utrinquè angustatis tomentosis, floribus terminalibus congestis foliis brevioribus, squamis lanceolatis vix acutis. ① in paludibus et arvis humidis. — Fl. fr. 4. p. 135. *G. minimum* Lapeyr. non Smith (teste cl. Arnott.) — Lob. ic. t. 481. (v. v.)
5. *G. ARVENSE* (Lam. dict. 2. p. 759.) caule herbaceo erecto paniculato, foliis oblongo-lanceolatis lanuginosis, floribus congestis lateralibus terminalibusque cano-tomentosis. ① in campis sabulosis. — Fl. fr. 4. p. 136. (v. v.)
6. *G. GALLICUM* (Lam. l. c.) caule herbaceo erecto subdichotomo, foliis linearibus acuminatis canescentibus, floribus confertis axillaribus terminalibusque foliis semper brevioribus, squamis lanceolatis subobtusis. ① in arvis sabulosis. — Fl. fr. 4. p. 136. Logfia subulata Cass. (v. v.)
7. *G. GERMANICUM* (Lam. l. c.) caule herbaceo dichotomo, foliis lanceolatis tomentosis acutis, floribus globoso-capitatis axillaribus terminalibusque, squamis lucidis ovato-lanceolatis acuminatis. ① ad vias fossas et margines agrorum. — Fl. fr. 4. p. 135. Gifola vulgaris Cass. — Lob. ic. t. 480. f. 2. (v. v.)  
  - β. *pyramidatum*, foliis spathulatis. — *G. pyramidatum* W. sp. 3. p. 1895. Fl. fr. 5. p. 468. (v. s.)
  - γ. *lanuginosum*, caulis parte superiore foliis floribusque laxè lanuginosis. (v. s.)
8. *G. MONTANUM* (W. sp. 3. p. 1896.) caule herbaceo erecto subdichotomo, foliis lineari-lanceolatis adpressis tomentosis, floribus confertis axillaribus terminalibusque folia subsuperantibus, squamis linearibus acutis. ② in siccis montanis ad margines sylvarum. — Fl. fr. 4. p. 136. Logfia lanceolata Cass. (v. v.)
9. *G. MINIMUM* (Lob. ic. t. 481. f. 2.) caule herbaceo diffuso ramoso, foliis lanceolatis acutis planis canescentibus, floribus conicis subconfertis lateralibus terminalibusque, squamis subobtusis. ① in agro Borbonio et Aurelianiensi. — Fl. fr. 4. p. 137. Logfia brevifolia Cass. (v. s.)

Sect. II. ANTENNARIA *Gærtn. fruct. 2. p. 411. t. 167.*

Flosculi alii hermaphroditi, alii sterile; flosculorum sterilium pappus clavatus.

10. *G. DIOICUM* (L. sp. 1199.) sarmentis procumbentibus, caule simplicissimo, foliis radicalibus spathulatis, corymbo coarctato, floribus dioicis, squamis interioribus elongatis obtusis coloratis. ② in pratis montosis nudis. — Fl. fr. 4. p. 137. — Lob. ic. t. 483. f. 2. (v. v.)

11. *G. ALPINUM* (L. sp. 1199.) sarmentis nullis, caule simplicissimo, foliis radicalibus lanceolatis, floribus terminalibus aggregatis sessilibus, squamis interioribus elongatis acutis membranaceis. ② in pascuis apriciis Alpium editorum. — Fl. fr. 4. p. 138. Fl. dan. t. 332. (v. v.)
12. *G. LEONTOPODIUM* (Jacq. aust. t. 86.) caule herbaceo simplicissimo, foliis lineari-lanceolatis subtus tomentosis, floribus terminalibus sessilibus aggregatis, bracteis lanceolatis lanatis radiato-patentibus, involucris villosis. ① in Alpiis et Pyrenæis. — Fl. fr. 4. p. 138. *Leontopodium Alpinum* Cass. (v. v.)

XVIII. *ELYCHRYSUM* Gærtn. 2. t. 166. f. 7. DC. fl. fr. 4. p. 130.

Involucrum imbricatum, squamis inæqualibus obtusis scariosis sæpius coloratis; flosculi omnes tubulosi et hermaphroditi; receptaculum nudum; pappus pilosus vel subplumosus.

§. 1. *Involucris squamis albis.*

1. *E. FRIGIDUM* (W. sp. 3. p. 1098.) herbaceum ramosum procumbens, foliis quadrifariam imbricatis oblongis obtusis incanis, ramis unifloris, floribus sessilibus. ② in editoribus montibus Corsicæ. — Fl. fr. 4. p. 131. Gn. bellidiflorum Viv. fragm. p. 16. t. 19. (v. s.)

§. 2. *Involucris squamis flavis.*

2. *E. STÆCHAS* (DC. fl. fr. 4. p. 132.) fruticosum, foliis linearibus, corymbo composito, squamis ovatis obtusis. ③ in collibus aridis et arenosis Galliæ australis et in prov. occidentalibus ad Nannetes usque. — Gn. *Stæchas* L. sp. 1193. — Barr. ic. t. 278. 409. 410. (v. v.)
3. *E. ANGUSTIFOLIUM* (DC. fl. fr. 5. p. 467.) fruticosum, foliis linearibus, corymbo pluries composito, involucris oblongis, squamis ovato-spathulatis obtusissimis. ③ in Corsicæ, Cebennarum, Ruscinonensis agri locis aridis. — Gn. *angustifolium* Lam. dict. 2. p. 746. — Barr. ic. t. 1125. (v. v.)
4. *E. ARENARIUM* (DC. fl. fr. 4. p. 132.) herbaceum, foliis incanis tomentosis obtusis, radicalibus spathulato-lanceolatis, caulinis lineari-lanceolatis, corymbo composito. ② in arenosis Alsatiæ, Aureliæ, Lugduni, Bardigalæ ex Laterr., etc. — Gn. *arenarium* L. sp. 1195. — Barr. ic. t. 74. (v. s.)

† † SEMINA NUDA AUT ARISTATA.

XIX. *FILAGO* L. gen. 995.

Involucrum polyphyllum imbricatum, squamis ovalibus aristato-subulatis; flosculi tubulosi in disco masculi quadrifidi, in radio feminei fertiles, stigmatibus duobus; receptaculum oblongum in radio paleaceum in disco nudum; semina nuda compressa.

1. *F. PYGMÆA* (L. sp. 1311.) involucris squamis foliaceis lucidis glabris. ① in maritimis et stagnis exsiccatis Corsicæ et Galliæ mediterraneæ. — *Micropus pygmæus* Fl. fr. 4. p. 199. *Evax umbellata* Gærtn. 2. t. 165. f. 3. — Barr. ic. t. 127. n. 14. Hæc sola species omnibus gaudet characteribus a Linnæo huic generi assignatis. (v. v.)

XX. *MICROPUS*. L. gen. 996.

Involucrum 5-g-phyllum squamis laxis, flosculi tubulosi, in disco hermaphroditi steriles 5-dentati pentandri, stigmatibus simplicibus, in radio feminei fertiles stigmatibus duobus; receptaculum conico-subulatum, in radio paleaceum, in disco nudum; semina nuda compressa involucris foliolis involuta.

1. *M. SUPINUS* (L. sp. 1313.) foliis oppositis obovato-cuneatis, involucrio incano coriaceo echinato. ① in maritimis Galloprov., agri Nicæensis. — Fl. fr. 4. p. 200. Lam. ill. t. 694. f. 1. (v. s. ex Hisp.)

2. *M. erectus* (L. sp. 1313.) foliis alternis ovato-lanceolatis, floribus lanuginosis, involucri inermi. ① in campis sterilibus, imprimis australibus. — Fl. fr. 4. p. 199. — Clus. hist. 1. p. 339. f. 3. Barr. ic. t. 296. Caules sæpè decumbentes. (v. v.)

## XXI. CARPESIMUM L. gen. 948.

Involucrum hemisphæricum imbricatum squamis exterioribus patulis; flores flosculosi, flosculis in disco hermaphroditis 5-lobis, in radio femineis fertilibus 5-dentatis; receptaculum nudum; semina nuda.

1. *C. CRANUM* (L. sp. 1208.) floribus terminalibus solitariis cernuis. ½ in humidis et umbrosis Delph. ex Vill., Galloprov. ex Ger., Pyren. orientalium ex Lapeyr. — Fl. fr. 4. p. 187. Lam. ill. t. 696. f. 1. (v. s. ex Pedemontio.)

## XXII. BUPHTHALMUM L. gen. 977.

Involucrum imbricatum; flores radiati, flosculis hermaphroditis, ligulis femineis fertilibus; receptaculum paleaceum; semina pappo membranaceo obsoleto coronata, ligularum lateribus membranacea.

§. 1. *Involucro foliaceo bracteiformi.*

1. *B. SPINOSUM* (L. sp. 1274.) foliis alternis oblongo-lanceolatis amplexicaulis integerrimis hirsutis, involucri foliosis mucronatis. ① ad margines agrorum in Galliâ mediterraneâ, circâ Aginnum ex S.-Am. Burdigalam ex Lattier. — Fl. fr. 4. p. 217. — Clus. hist. 2. p. 13. f. 1. (v. v.)
2. *B. AQUATICUM* (L. sp. 1274.) involucri obtusè foliosis sessilibus axillaribus, foliis oblongis obtusis alternis subintegerrimis, caule dichotomo. ① secus aquas Occitaniz, Galloprov., Nicez. — Fl. fr. 4. p. 217. — Barr. ic. t. 552. (v. v.)
3. *B. MARITIMUM* (L. sp. 1274.) involucri obtusè foliosis pedunculatis, foliis alternis spathulatis, caule herbaceo. ½ in maritimis Nicez., Massiliz. — Fl. fr. 4. p. 217. — Barr. ic. t. 1151. (v. v.)

§. 2. *Involucro brevi subsquamoso.*

4. *B. SALICIFOLIUM* (L. sp. 1275.) foliis alternis oblongo-lanceolatis subserratis, involucri nudis. ½ ad radices montium Arverniz et Galliz australis. — Fl. fr. 4. p. 218. — Clus. hist. 2. p. 13. f. 3. (v. v.)

## XXIII. CHRYSANTHEMUM L. gen. 966. — Chrysanthemum et Pyrethrum Gertn. carp. 2. p. 420 et 430. DC. fl. fr. 4. p. 177 et 181.

Involucrum hemisphæricum imbricatum squamis margine scariosis; flores radiati; receptaculum nudum; semina nuda aut membranâ brevi coronata.

## Sect. 1. LEUCANTHEMUM.

Discus flavus, ligulæ discoloræ.

§. 1. *Involucro 1-seriato.*

1. *C. PAPPUSILLUM* (Lois. not. t. 6. f. 3.) caule pusillo subramoso stolonifero, foliis pinnatifidis, lobis 3-7 rotundis obtusis integerrimis, pedicellis axillaribus flore multoties longioribus. ① in insulis Sanguinariis propè Ajaccio Corsicæ et Lavezio propè Bonifacio. — Fl. fr. 5. p. 477. (v. s.)

§. 2. *Involucro multiseriato.*

\* *Receptaculo convexo.*

2. *C. TOMENTOSUM* (Lois. gall. 580. t. 18.) foliis palmatifidis tomentosis lobis obtusis, involucri squamis ovatis hirsutis, caule ascendente simplici uni-

- flo.  $\mathcal{L}$  in editoribus montibus Corsicæ. — *Pyrethrum tomentosum* Fl. fr. 5. p. 477. *P. minimum* Fl. fr. 4. p. 924. (v. s.)
3. *C. ALPINUM* (L. sp. 1253.) foliis inferioribus pinnatifido-dentatis, summis linearibus integerimis, involucri squamis lanceolatis glabris, caule 1-flo.  $\mathcal{L}$  in lapidosis Alpium et Pyrenæorum. — *Pyr. Alpinum* Fl. fr. 4. p. 182. — *Clas. hist.* 1. p. 335. f. 2. (v. v.)
- $\beta$ . *pubescens*, caule foliisque pubescentibus, squamis ciliatis. — *C. minimum* Vill. dauph. 3. p. 202. (v. v.)
4. *C. HALLERI* (Sut. helv. 2. p. 193.) foliis caulinis lanceolatis profundè tenuissimèque dentatis, radicalibus pinnatifidis, squamis lanceolatis glabris, caule 1-flo.  $\mathcal{L}$  in lapidosis Alpium Delphinatus, Galloprovinciæ. — *Pyr. Halleri* W. Fl. fr. 4. p. 182. — *Barr. ic. t.* 458. f. 2. (v. v.)
5. *C. CERATOPHYLLOIDES* (All. ped. t. 37. f. 1.) foliis pinnato-multifidis, pinnis linearibus acutis, caule erecto unifloro.  $\mathcal{L}$  in editoribus Delphinatus et Pyrenæorum orientalium. — *Fl. fr.* 4. p. 179. (v. v. DC.)
6. *C. MONSPELIENSE* (L. sp. 1252.) foliis imis palmato-multifidis, laciniis linearibus pinnatifidis, caule erecto unifloro.  $\mathcal{L}$  in Corsicæ et Occitanicæ locis montosis. — *Fl. fr.* 4. p. 180. *Jacq. obs. t.* 93. (v. s.)
7. *C. GRAMINIFOLIUM* (L. sp. 1252.) foliis linearibus subintegris aut tenuissimè inciso-dentatis, caule simplicissimo unifloro.  $\mathcal{L}$  in collibus Gallicæ australis. — *Fl. fr.* 4. p. 179. — *Magn. hort. monsp. t.* 31. (v. s.)
8. *C. LEUCANTHEMUM* (L. sp. 1251.) foliis amplexicaulibus lanceolatis serratis basi inciso-dentatis pinnatifidisve, radicalibus spatulatis in petiolum basi decurrentibus, caule erecto.  $\mathcal{L}$  in pratis frequens. — *Fl. fr.* 4. p. 178. — *Moris. 3. s. 6. t. 8. n. 2.* (v. v.)
- $\beta$ . *montanum*, foliis superioribus lineari-lanceolatis. — *C. montanum* L. sp. 1252. *All. ped. t.* 37. f. 2. — *Moris. l. c. n. 9.* Semina membranâ prominulâ dentatâ sæpè coronantur. (v. v.)
9. *C. MAXIMUM* (Ram. bull. phil. n. 42. p. 140.) foliis carnosio-fragilibus inferioribus spatulatis subintegris, mediis amplexicaulibus lanceolato-dentatis, summis linearibus integris, caule simplici unifloro.  $\mathcal{L}$  in Pyrenæis propè Valletriam in monte dicto *Pic de Lhéris*. — *Fl. fr.* 4. p. 178. — *Dodart ic. t.* 65. Flos duplò major quàm in præcedente. (v. s.)
- \* \* *Recept. hemisphærico, semine 4-gono.*
10. *C. PARTHENIUM* (Sm. brit. 2. p. 900.) foliis petiolatis compositis planis, pinnis ovatis incis, pedunculis ramosis corymbosis, caule erecto, involucri pubescente squamis lineari-spatulatis. (2)  $\mathcal{L}$  in incultis et lapidosis. — *Pyr. Parthenium* Fl. fr. 4. p. 183. *Matricaria Parthenium* Bull. herb. t. 203. (v. v.)
11. *C. CORYMBOSUM* (L. sp. 1251.) foliis pinnatifidis, pinnis lanceolatis pinnatifidis argutè serratis, summis confluentibus, pedunculis corymbosis, squamis lanceolato-ovatis.  $\mathcal{L}$  in montosis Gallicæ mediæ, orientalis, australis occidentalisque. — *P. corymbosum* Fl. fr. 4. p. 183. — *Clas. hist.* 1. p. 338. f. 1. *Barr. ic.* 781. 782. 785. 786. (v. v.)
- \* \* \* *Recept. conico, semine 3-gono.*
12. *C. INODORUM* (L. sp. 1253.) foliis sessilibus bipinnatifidis, pinnis linearifiliformibus bi-vel-tripartitis, caule ramoso patulo, seminibus membranulâ brevissimâ truncatâ coronatis. (1) in agris, ad vias. — *Pyr. inodorum* Fl. fr. 4. p. 184. — *Fuchs. hist.* 144. ic. (v. v.)
13. *C. MARITIMUM* (Pers. ench. 2. p. 462.) foliis bipinnatifidis, pinnis linearibus brevibus carnosis glabris trifidis, caule diffuso ramoso, seminibus membranâ brevissimâ obsolete 4-dentatâ coronatis.  $\mathcal{L}$  in arenosis maritimis ex Abbativillâ ad Nannetes. — *Pyr. maritimum* Sm. engl. bot. t. 970. *Fl. fr.* 5. p. 477. — *Matricaria maritima* L. sp. 1256. (v. v. DC.)

## Sect. II. CHRYSANTHOS.

Ligulæ disco concolores.

14. *C. SZOZTUM* (L. sp. 1254.) foliis amplexicaulibus supernè laciniatis, inferne profunde dentato-serratis. ① in arvis. — Fl. fr. 4. p. 181. — Clus. hist. 1. p. 334. f. 2. (v. v.)
15. *C. MYCONIS* (L. sp. 1254.) foliis ligulatis obtusis serratis, seminibus teretibus membranâ coriacea tubuloso-ligulatâ coronatis. ① in Corsicâ, Galloprovincia et circâ Niceam. — Fl. fr. 4. p. 180. — Moris. 3. s. 6. t. 4. n. 1. bis. (v. s.)
16. *C. CORONARIUM* (L. sp. 1254.) foliis bipinnatifidis acutis extrorsum latioribus, seminibus nudis alatis coronæ 3-dentatis disci dente 1 acuto terminatis. ① in hortis culta, et circâ Niceam ex All. indigena. — Fl. fr. 4. p. 181. — Clus. hist. 1. p. 335. f. 1. (v. v.)

XXIV. MATRICARIA *Tourn. t. 231. L. gen. 967.*

Involucrum hemisphæricum imbricatum squamis obtusis, flores radiati; receptaculum nudum conicum; semina nuda.

1. *M. CHAMOMILLA* (L. sp. 1256.) foliis bipinnatifidis, laciniis capillaribus, involucri subplano, squamis obtusiusculis. ① in hortis et cultis. — Fl. fr. 4. p. 184. — Lob. ic. 770. f. 1. (v. v.)
2. *M. SUAVOLIX* (L. sp. 1456.) foliis triplicato-pinnatifidis laciniis tenuissimis, involucri squamis subacutis obtusiusculis. ① circâ Narbonam, Abbavillam, in Arvernâ. — Fl. fr. 4. p. 185. Flores dimidiò minores quàm in præcedente, a quâ tamen non satis distincta. (v. s.)

XXV. ANTHEMIS *L. gen. 970.*

Involucrum hemisphæricum squamis subæqualibus margine scariosis imbricatum; flores radiati, flosculis hermaphroditis, ligulis lanceolatis femineis fertilibus; receptaculum conicum paleaceum; semina cylindrica aut 3-gona non marginata, nuda aut membranulâ coronata.

§. 1. *Disco luteo, radio discolore.*

\* *Seminibus nudis.*

1. *A. ALTISSIMA* (L. mant. 474.) caule erecto ramoso, foliis bipinnatifidis, laciniis lanceolatis acutis subdentatis dentibus inferioribus subreflexis, paleis lanceolatis cuspidatis. ① in campis Galliæ australis. — Fl. fr. 4. p. 203. — J. Bauh. hist. 3. p. 120. f. 1. (v. v.)
- β. *tomentosa*, laciniis obtusis. Circâ Prades agri Ruscinonensis reperiit Ph. Thomas. (v. s.)
2. *A. FRERGERIA* (W. sp. 3. p. 2182.) caule erecto ramoso fastigiato, ramis subunifloris, lateralibus apice longioribus, foliis bipinnatis, pinnis linearibus acutis integris, involucri squamis membranaceis. ① in arenosis Druentiæ propè Avenionem. Corsicâ (cl. Soleirol.) — Fl. fr. 5. p. 482. (v. v. DC.)
3. *A. COTULA* (L. sp. 1261.) foliis bipinnatis glabris puberulisve, pinnis subulatis tripartitis, paleis lineari-setaceis acutis, seminibus tuberculosiis. ① in incultis et arvis. — Fl. fr. 4. p. 206. Maruta fetida Cass. (v. v.)
4. *A. MIXTA* (L. sp. 1260.) caule ramoso, foliis sessilibus semipinnatifidis, laciniis dentatis, paleis lanceolatis duris acutis, seminibus levibus. ① in campis Galliæ occidentalis, australis, orientalisque. — Fl. fr. 4. p. 204. Ormenis bicolor Cass. — Moris. 3. s. 6. t. 12. u. 15. (v. v.)
5. *A. INCRASSATA* (Lois. not. 129.) caule ramoso patulo, foliis pinnatifidis pubescentibus, laciniis inferioribus integris acutis, superioribus longioribus pinnatim incis, pedunculis demùm incrassatis, paleis acutis, seminibus 4-gonis apice umbilicatis. ①? in arenosis et lapidosis Narbonæ, Arelatis, Nicææ. — Fl. fr. 5. p. 482. *A. nicæensis* W. sp. 3. p. 2182. (v. v.)

6. *A. VISCATA* (Brot. phyt. 1. v. 15.) caule diffuso, foliis bipinnatis lineari-fili-  
formibus tripartitis, squamis oblongis paleisque obovatis obtusis margine  
fascis. ① circa Telonem et Niczam. — Fl. fr. 5. p. 482. (v. s.)
7. *A. NOBILIS* (L. sp. 1260.) caule basi ramoso decumbente, foliis bipinnatis,  
pinnis tripartitis lineari-subulatis subvillosis, paleis mollibus lanceolatis ob-  
tusis. ½ in pascuis siccis. — Fl. fr. 4. p. 205. — Lob. ic. t. 770. f. 2. (v. v.)
- \*\* *Seminibus membranâ brevi coronatis.*
8. *A. ARVENsis* (L. sp. 1261.) foliis bipinnatis pinnis lineari-lanceolatis pubes-  
centibus acutis, paleis lanceolatis acuminatis, seminibus membranâ trun-  
catâ coronatis. ② in arvis. — Fl. fr. 4. p. 206. (v. v.)
9. *A. MONTANA* (L. sp. 1261.) caule ascendente, foliis pinnatis pubescentibus,  
pinnis linearibus, terminali trifida, pedunculis elongatis nudis tomentosis,  
paleis lanceolatis acutis, seminibus membranâ inæqualiter truncatâ coro-  
natis. ½ in Pyrenæis, Cebennis, montibus Arvernæ, Galloprovinciæ. —  
Fl. fr. 5. p. 483. *A. saxatilis* DC. syn. fl. gall. 291. (v. v.)
10. *A. MARITIMA* (L. sp. 1259.) caule herbaceo, foliis pinnatifidis carnosis sub-  
tus punctatis, lacinii in foliis superioribus lanceolatis, in foliis inferioribus  
incisus apice dilatatis, seminibus membranâ brevissimâ denticulatâ coro-  
natis. ½ in arenosis secus Mediterraneum. — Fl. fr. 4. p. 204. (v. v.)

§. 2. *Radio concolore luteo.*

11. *A. TINGITORIA* (L. sp. 1263.) foliis subtus pubescentibus triplinatifidis ser-  
ratâ, caule erecto ramoso, seminibus membranâ integerrimâ coronatis. ½ in  
Galliâ mediâ et orientali. — Fl. fr. 4. p. 208. — Barr. ic. t. 465. (v. s.)

XXVI. *ANACYCLUS Pers. ench. 2. p. 464. DC. fl. fr. 5. p. 480.*

Involucrum hemisphæricum imbricatum squamis ovato-lanceola-  
tis lanceolatisve; flosculi omnes tubulosi aut radiati in disco herma-  
phroditi, in radio feminei fertiles; receptaculum conicum paleaceum;  
semina compressa margine membranacea nuda.

1. *A. CLAVATUS* (Pers. ench. 2. p. 465.) caule erectiusculo, foliis subglabris, pedi-  
culis subflore incrassatis, floribus radiatis, invol. squamis acutis, paleis gla-  
berrimis spathulatis mucronatis. ① Narbonæ. — Fl. fr. 5. p. 481. *Anthemis*  
*clavata* Desf. atl. 2. p. 287. Ligulæ albescentes. (v. s.)
2. *A. TOMENTOSUS* (DC. fl. fr. 5. p. 481.) caule demisso, foliis villosis, pediculis  
cylindraceis, floribus radiatis, invol. squamis subacutis, paleis cuneatis tra-  
pezoideis acutis pilis coronatis. ① circa Monspelium, Narbonam. — *Anthe-  
mis* *biaristata* Fl. fr. 4. p. 214. Ligulæ albescentes. (v. v.)
3. *A. VALENTINUS* (L. sp. 1258.) foliis villosis, floribus flosculosis radiatisve,  
flosculorum lacinii inæqualibus, duabus aristatis erectis, invol. squamis  
subacutis, paleis lanceolato-spathulatis subacutis pilis coronatis. ① circa Car-  
cassonam, Perpignanum, Niczam. — Fl. fr. 4. p. 202. Lam. ill. t. 700. f. 1. (v. v.)
4. *A. RADIATUS* (Lois. gall. 583.) foliis villosis, floribus radiatis, flosculorum la-  
ciniis inæqualibus, 2 aristatis, invol. squamis obtusis, paleis spathulatis mu-  
cronatis glaberrimis. ① in provinciis mediterraneis. — Fl. fr. 5. p. 481. *An-  
themis* *Valentina* L. sp. 1262. Fl. fr. 4. p. 208. Ligulæ latæ flavæ. (v. v.)
5. *A. PURPURASCENS* (Pers. ench. 2. p. 465.) caule folisque subglabris, floribus  
radiatis, flosculorum lacinii inæqualibus, invol. squamis obtusis, paleis gla-  
berrimis. ① in provinciis mediterraneis. — Fl. fr. 5. p. 481. Ligulæ flavæ  
subtus purpureæ. Præced. var. ? (v. s.)

XXVII. *ACHILLEA L. gen. 971.*

Involucrum ovatum imbricatum; flores radiati, flosculis hermaphro-  
ditis, ligulis brevibus femineis paucioribus; receptaculum planum an-  
gustum paleaceum; semina nuda.

§. 1. *Floribus luteis.*

1. *A. TOMENTOSA* (L. sp. 1264.) foliis bipinnatifidis villosis, pinnis confertissimis, laciniis linearis-lanceolatis integerrimis, corymbo composito.  $\frac{1}{2}$  in sterilibus Delphinatûs, Galloprov. — Fl. fr. 4. p. 210. — Clus. hist. 1. p. 330. f. 2. (v. v.)
2. *A. AGRATUM* (L. sp. 1264.) foliis oblongis obtusis serratis in petiolum attenuatis fasciculatis, corymbo composito coarctato.  $\frac{1}{2}$  in petreis subhumidis Gallie australis. — Fl. fr. 4. p. 209. — Lob. ic. 1. 489. f. 2. (v. v.)

§. 2. *Florum radiis albis.*

3. *A. HERBA-ROTA* (All. ped. t. 9. f. 3.) foliis cuneiformibus basi attenuatis serratis cauleque glabris.  $\frac{1}{2}$  in editissimis Alpibus Delphinatûs. — Fl. fr. 4. p. 210. (v. v. in Helv.)
4. *A. PYRÆNICA* (L. sp. 1266.) foliis linearibus acuminatis æqualiter argute serratis glabris.  $\frac{1}{2}$  in pratis subalidis. — Fl. fr. 4. p. 211. — Clus. 2. p. 12. f. 1. (v. v.)  
*β. pubescens*, foliis glanduloso-pubescentibus. In Pyrenæis. (v. v.)
5. *A. CLAVENNE* (L. sp. 1266.) foliis tomentoso-incanis subpinnatifidis laciniis lanceolatis obtusis terminali 3-dentatâ, inferioribus petiolatis, corymbo simplici subumbellato.  $\frac{1}{2}$  in Cebennis ex Lois. — Jacq. austr. t. 76. (v. v. D. in Alp. Italiâ.)
6. *A. MOSCHATA* (Jacq. austr. app. t. 33.) foliis pectinato-pinnatis glabris, pinnis linearibus obtusiusculis integerrimis punctatis.  $\frac{1}{2}$  inter lapides et saxa Alpium Delphinatûs. — Fl. fr. 4. p. 213. (v. v. D. in Helv.)
7. *A. ATRATA* (L. sp. 1267.) foliis pectinato-pinnatis glabris, pinnis linearibus acuminatis subtripartitis, cyma simplici, pedunculis pubescentibus, caule simplicissimo.  $\frac{1}{2}$  in Pyrenæis ex Lapeyr. et alpinis Delphinatûs? ex Vill. — Fl. fr. 4. p. 214. — Clus. hist. 1. p. 336. f. 2. (v. v. in Helv.)
8. *A. NANA* (L. sp. 1267.) foliis pinnatis villosis, pinnis dentatis linearibus, radicalibus petiolatis bipinnatis, caule simplicissimo, corymbo simplici compacto.  $\frac{1}{2}$  in saxosis editiorum Alpium Delphinatûs, Cebennarum, Pyren. Orientalium. — Fl. fr. 4. p. 213. All. ped. t. 9. f. 2. (v. v.)
9. *A. CHAMÆMELIOLIA* (Pourr. act. toul. 3. p. 305.) foliis pectinato-pinnatifidis glabris, pinnis linearibus integerrimis, corymbo composito subundo coarctato.  $\frac{1}{2}$  in Pyrenæis oriental. propè Villefranche et Olette. — Fl. fr. 4. p. 212. (v. v. DC.)  
*β. capillata*, corymbo subsimplici. — A. capillata Lapeyr.
10. *A. MACROPHYLLA* (L. sp. 1265.) foliis pinnatis glabris, pinnis lanceolatis inciso-serratis extimis confluentibus, corymbo composito.  $\frac{1}{2}$  in sylvis montosis et lapidosis Alpium Delphinatûs ex Ray et Vill. — Fl. fr. 4. p. 212. — Barr. ic. t. 991. (v. v. in Sabaudiâ.)
11. *A. DENTIFERA* (DC. fl. fr. 5. p. 285.) caule piloso, foliis pinnatifidis rachi alatâ serrato-incisâ, laciniis inciso-serratis oblongis, superioribus pilosis, floribus fastigiatis corymbosis, invol. cylindrici ovati squamis hirtis.  $\frac{1}{2}$  in Alpibus Galloprovinciæ. — Moris. s. 6. t. 11 n. 5 et 14. (v. s.)
12. *A. COMPACTA* (Lam. dict. 1. p. 27.) foliis bipinnatifidis laciniis lanceolatis inciso-serratis, rachi foliaceâ integrâ, corymbis fastigiato-compositis, involucri ovato-globosi squamis hirtis.  $\frac{1}{2}$  in sylvis montosis Delphinatûs? Jurassii. — Fl. fr. 4. p. 215, excl. syn. All. et Wild. An satis à sequente distincta? (v. s.)
13. *A. TANACETIFOLIA* (All. ped. n. 666.) foliis bipinnatifidis sparse pilosis, rachi integrâ, laciniis linearibus integris serratisve, corymbo composito patulo, invol. globosi squamis lucidis hirtis.  $\frac{1}{2}$  in sylvis pratensque montosis Alpium Delph., Galloprov., Pyrenæorum. — Fl. fr. 4. p. 214. (v. v.)
14. *A. LIGUSTICA* (All. ped. t. 53. f. 2.) foliis bipinnatifidis breviter pubescenti-

- bus, laciniis linearibus inciso-serratis, rachi alata integerrima, corymbo composito fastigiato, invol. ovati squamis breviter pubescentibus.  $\frac{1}{2}$  in collibus Nicææ, Corsicæ. — Fl. fr. 4. p. 215. (v. s.)
15. *A. ODORATA* (L. sp. 1268.) foliis radicalibus petiolatis subtus pilosis bipinnatis, pinnulis linearibus obtusis integris, caulinis pinuatifidis, corymbo simplici.  $\frac{1}{2}$  in asperis apricis secus agros in Pyren. Orient. et Alpibus Delphinatûs. — Fl. fr. 5. p. 486. — Barr. ic. t. 992. (v. v. DC.)
16. *A. MILLEFOLIUM* (L. sp. 1125.) foliis subtriplicato-pinnatifidis, rachi integerrima, laciniis linearibus subacuminatis minimis, summis confluentibus, involucrio glabro.  $\frac{1}{2}$  ad vias et in arvis frequens. — Fl. fr. 4. p. 215. Fl. dan. t. 737. (v. v.)
17. *A. SETACEA* (Waldst et Kit. hung. t. 80.) foliis subtriplicato-pinnatifidis, laciniis lineari-setaceis mucronatis confertissimis pilosis integris, rachi integrâ angustissima, involucrio subhirsuto.  $\frac{1}{2}$  in Pyren. Orient. et Galloprovincia propè Avenionem. — Fl. fr. 5. p. 485. (v. v.)
18. *A. NOBILIS* (L. sp. 1268.) foliis caulinis bipinnatifidis, laciniis linearibus subdentatis, rachi alata dentata, radicalibus triplicato-pinnatifidis, corymbo composito fastigiato densissimo, involucrio hirsuto.  $\frac{1}{2}$  in campis siccis et collibus Gallie orientalis, medie australisque. — Fl. fr. 4. p. 216. (v. v.)

## XXVIII. ARTEMISIA L. gen. 945.

Involucrum ovatum aut rotundatum imbricatum; flosculi omnes tubulosi, in disco hermaphroditi 5-dentati, in radio graciles pauciores integri feminei fertiles; semina nuda; receptaculum nudum aut pilosum.

## SECT. I. ABSINTHIUM Hall. helv.

Receptaculum hirsutum.

1. *A. ABSINTHIUM* (L. sp. 1188.) foliis incanis, radicalibus triplicato-pinnatifidis obtusiusculis, superioribus pinnatifidis acutis, laciniis lanceolatis, floralibus indivisis, floribus globosis pedunculatis nutantibus.  $\frac{1}{2}$  in lapidosis incultis et montosis. — Fl. fr. 4. p. 189. — Cam. epit. 452. ic. (v. v.)
2. *A. ARBORASCENS* (L. sp. 1188.) foliis tripinnatifidis sericeis cinereis, foliolis linearibus, floribus globosis, ramulis floriferis simplicibus.  $\frac{1}{2}$  Olbiæ et in Corsicâ (Req.). — Fl. fr. 4. p. 190. — Lob. ic. t. 753. f. 1. (v. s.)
3. *A. CORYMBOSA* (Lam. dict. 1. p. 265.) suffruticosa, foliis palmato-multifidis, lobis linearibus 3-vel-4-partitis, floribus globosis pedunculatis nutantibus, involucrio pubescente.  $\frac{1}{2}$  in apricis demissiorum montium Delphin., Galloprov., Nicææ, et agri Ligericini. — Fl. fr. 4. p. 190. — Lob. ic. t. 769. f. 1. Flores paniculati nec minime corymbosi. (v. s.)
4. *A. GLACIALIS* (L. sp. 1187.) caule herbaceo simplicissimo, foliis omnibus palmato-multifidis candidis sericeis, capitulis terminalibus glomeratis sessilibus globosis 40-floris, invol. squamis lanceolato-ovatis margine atratis.  $\frac{1}{2}$  in editissimis Alpibus Delphinatûs. — Fl. fr. 4. p. 191. All. ped. t. 8. f. 3. (v. v.)
5. *A. MUTELLINA* (Vill. dauph. 3. p. 244. t. 35.) caule herbaceo simplicissimo, foliis omnibus palmato-multifidis albido-sericeis, capitulis subovatis 14-floris, inferioribus pedunculatis, summis sessilibus, invol. squamis lineari-lanceolatis margine fuscis. ① in saxosis editioribus Alpium, montium Arvernæ et Pyrenæorum. — Fl. fr. 5. p. 478. *A. rupestris* Fl. fr. 4. p. 191. non L. (v. v. D. in Helv.)

## SECT. II. DRACUNCULUS.

Receptaculum nudum.

\* *Floribus subglobosis.*

6. *A. SPICATA* (L. syst. 744.) caule herbaceo simplicissimo, foliis cano-sericeis, radicalibus palmato-multifidis, caulinis pinnatifidis, summis linearibus inte-



- gerrimis obtusis, floribus axillaribus, infimis pedunculatis.  $\frac{1}{2}$  in summis Alpibus et Pyrenæis. — Fl. fr. 4. p. 192. — Barr. ic. t. 462. Magna cum præcedente affinitas. (v. v.)
7. *A. CHAMÆMELIFOLIA* (Vill. daph. 3. p. 250. t. 35.) caule erecto basi ramoso glabro, foliis subglabris, inferioribus tripinnatis, superioribus bipinnatis, foliolis linearibus acutis, floribus paniculatis pedunculatis nutantibus.  $\frac{1}{2}$  in apricis humilioribus montium Delphinatû. — Fl. fr. 4. p. 193. (v. s.)
8. *A. MARITIMA* (L. sp. 1186.) caule herbaceo ramoso ascendente, ramulis cernuis, foliis niveo-tomentosis multifidis, laciniiis planis linearibus, rameis simplicibus linearibus, floribus oblongis tomentosis sessilibus.  $\frac{1}{2}$  in maritimis. — Fl. fr. 4. p. 196. Wood. med. bot. t. 122. (v. v.)
9. *A. CAMPESTRIS* (L. sp. 1185.) caule procumbente ramoso virgato, foliis cauliniis setaceis pinnatis glabris, radicalibus pinnatis laciniiis trifidis, floribus subovatis pedunculatis.  $\frac{1}{2}$  in campestribus lapidosis siccis. — Fl. fr. 4. p. 194. A. procerâ Lapeyr. ex cl. Benth. Oligosporus campestris Cass. — Cam. epit. 939. ic. (v. v.)
10. *A. CRITHMIFOLIA* (L. sp. 1186.) caule ascendente, foliis carnosis obtusis, cauliniis pinnatis pinnis simplicibus bifidiave linearibus, floralibus indivisis, radicalibus lobis linearilanceolatis sæpè sericeis, floribus oblongis pedunculatis erectis.  $\frac{1}{2}$  in arenosis maritimis circâ Telonem et ex Bayonnâ ad Nannetes. — Fl. fr. 4. p. 478. An præcedentis var.? (v. v.)
11. *A. TANACETIFOLIA* (L. sp. 1188.) caule simplicissimo herbaceo, foliis bipinnatifidis subpubescentibus, laciniiis linearilanceolatis acuminatis integerrimis, racemo nudo terminali, floribus subrotundis nutantibus, squamis ovatis obtusis margine scariosis.  $\frac{1}{2}$  in Alpibus Delphinatû. — Fl. fr. 4. p. 193. All. ped. t. 10. f. 3 et t. 70. f. 2. (v. s.)
12. *A. DRACUNCULUS* (L. sp. 1189.) foliis glabris lanceolatis utrinquè attenuatis, floribus subrotundis pedunculatis erectis.  $\frac{1}{2}$  E. Sibiria orta, in hortis culta, in Lotharingia spontanea? — Fl. fr. 4. p. 195. Oligosporus condimentarius Cass. — Gm. sib. 1. t. 59 et t. 60. f. 1. (v. s. c.)
- \* \* Floribus ovatis vel cylindricis.
13. *A. CÆRULESCENS* (L. sp. 1189.) caule suffruticoso ramoso, foliis incanis lanceolatis, superioribus integerrimis, inferioribus 2-aut 3-partitis, floribus oblongis paniculatis nutantibus.  $\frac{1}{2}$  in maritimis circâ Nicæam ex All. — Fl. fr. 4. p. 195. — Lob. ic. t. 765. f. 2. (v. s.)
- $\beta$ . Corsica, foliis præsertim inferioribus subglabris. — In Corsicâ. (v. s.)
14. *A. VULGARIS* (L. sp. 1188.) caule herbaceo, foliis pinnatifido-incanis subtus tomentosis suprâ atro-viridibus, floralibus indivisis linearilanceolatis, floribus subsessilibus oblongis erectis, involucri tomentosis.  $\frac{1}{2}$  ad vias in incultis. — Fl. fr. 4. p. 195. — Lob. ic. t. 764. f. 2. (v. v.)
15. *A. ARAGONEENSIS* (Lam. dict. 1. p. 269.) caule erecto basi fruticoso lamato, foliis sericeis bipinnatifido-palmatis, floralibus simplicibus, floribus ovatis paniculatis sessilibus tomentosis.  $\frac{1}{2}$  in Navarrâ ad limites Hispaniæ in declivibus apricis. — Fl. fr. 5. p. 499. — Barr. ic. t. 447. (v. s. ex Hisp.)
16. *A. GALLICA* (W. sp. 3. p. 1834.) caule erecto ramoso tomentoso, ramis erectis, foliis incanis bipinnatisve laciniiis linearibus obtusis, cauliniis pinnatis, summis linearibus, capitulis oblongis subsessilibus erectis sub 3-floris.  $\frac{1}{2}$  in maritimis Nicææ, Occitanicæ, agri Ruscinonensis, Vindeensis (Vendée) et circâ Nannetes. — Fl. fr. 4. p. 197. (v. v.)
- $\beta$ . congesta, ramis abbreviatis, floribus congestis. — Circâ Massiliam reperiit amic. Solier. An species? (v. s.)
17. *A. ABROTANUM* (L. sp. 1185.) frutescens, caule stricto, foliis inferioribus bipinnatis, superioribus pinnatis papillaceis, involucri pubescentibus hemisphaericis.  $\frac{1}{2}$  in Galliâ australi? in hortis odoris gratiâ culta. — Fl. fr. 4. p. 197. — Moris. s. 6. t. 2. n. 2. (v. v. c.)

18. *A. PANICULATA* (Lam. dict. 1, p. 265.) frutescens, caule ramoso patente, foliis omnibus bipinnatis capillaceis, involucriis glabris hemisphaericis. ♂ in Galliâ australi Lam., ex Occitaniâ et agro Ruscinonensi, Pyren. editioribus? ex Lapeyr. — Fl. fr. 4. p. 198. — Lob. ic. 768. f. 2. (v. s. c.)

XXIX. *COTULA* L. gen. n. 968.

Involucrum pauciseriatum squamis imbricatis obtusis margine scariosis; flosculi omnes tubulosi, in disco hermaphroditi 4-dentati, in radio feminei fertiles; receptaculum conicum nudum; semina ovato-compressa nuda.

1. *C. AUREA* (L. sp. 1257.) foliis bipinnatifidis, laciniis simplicibus incisive capillaribus. ① in provinciis Australioribus. — *Anacyclus aureus* L. mant. 287. Fl. fr. 4. p. 202. *Marcelia aurea* Cass. (v. s.)

XXX. *LONAS* Gært. carp. 2. t. 165. Juss. ann. mus. 8. p. 173.

Involucrum ovato-rotundum imbricatum squamis ovatis; receptaculum conico-subulatum paleaceum; flosculi omnes fertiles tubulosi androgyni; semina pappo membranaceo oblique truncato denticulato coronata.

1. *L. IXODORA* (Gært. 2. t. 165. f. 5.) ① in Galliâ australi ex Pers., in agro Ruscinonensi ex Lapeyr. Fl. fr. 5. p. 480. — *Athanasia annua* L. (v. v. c.)

XXXI. *TANACETUM* Tourn. t. 261. L. gen. 944.

Involucrum hemisphaericum imbricatum; flores flosculosi; flosculi in centro hermaphroditi 5-lobi, in radio feminei fertiles 3-lobi; receptaculum nudum; semina pappo membranaceo integro coronata.

1. *T. VULGARE* (L. sp. 1148.) foliis bipinnatifidis incisive serratis. ② in ruderatis ad muros. — Fl. fr. 4. p. 189. — Fuchs. hist. 46. ic. (v. v.)

XXXII. *SANTOLINA* Tourn. t. 260. L. gen. 942.

Involucrum hemisphaericum imbricatum; flosculi omnes tubulosi hermaphroditi; receptaculum paleaceum; semina nuda.

1. *S. ROSMARINIFOLIA* (L. sp. 1180. a.) pedunculis unifloris, foliis linearibus, inferioribus subpubescentibus margine tuberculatis, superioribus planis integerrimis glabris, ramis involucriisque glabris. ♂ in Galliâ australi. — Fl. fr. 4. p. 201. — Moris. s. 6. t. 3. n. 22. (v. v.)
2. *S. VIARIS* (W. sp. 3. p. 1798.) pedunculis unifloris, foliis subcylindricis glabris quadrifariam dentatis, dentibus subulatis rectis, ramis involucriisque glabris. ♀ in Galliâ australi circa Narbonam. — Fl. fr. 4. p. 200. (v. s.)
3. *S. squarrosa* (W. sp. 3. p. 1798.) pedunculis unifloris, foliis glabris quadrifariam dentatis dentibus subulatis patentibus, ramis tomentosis, involucriis glabris. ♀ in asperis Occitanis Narbonensis, et agri Ruscinonensis. — Fl. fr. 5. p. 479. — Moris. s. 6. t. 3. n. 17. *Sequentis varietas?* (v. v. DC.)
4. *S. incana* (Lam. ill. t. 671. f. 3.) pedunculis unifloris, foliis sub 4-gonis imcanis quadrifariam dentatis dentibus obtusis, ramis tomentosis, involucriis pubescentibus. ♂ in collibus agri Ruscinonensis, Occitanis circa Narbonam, Nemausim, et Galloprovinciâ circa Massiliam (Solier). — Fl. fr. 4. p. 200. — Moris. l. c. n. 12. (v. s.)

XXXIII. *DIOTIS* Desf. ail. DC. fl. fr. 4. p. 201.

Involucrum hemisphaericum imbricatum; flosculi omnes tubulosi hermaphroditi 5-dentati medio coarctati, basi dilatati in appendices 2 ovarium ciungentes; receptaculum convexum paleaceum; semina nuda.

2. *D. CANDIDISSIMA* (Desf. atl. 2. p. 251.) pedunculis corymbosis, foliis oblongis obtusis crenatis densè lanatis.  $\mathcal{L}$  in arenosis maritimis à Niceâ ad Nannetes. — Fl. fr. l. c. — Gærtn. fruct. 2. t. 165. f. 2. (v. v.)

XXXIV. BALSAMITA *Paill. DC. fl. fr. 4. p. 187.*

Involucrum patens imbricatum; flosculi omnes tubulosi hermaphroditi 5-dentati; receptaculum nudum; semina incompletâ membranâ coronata.

1. *B. MAJOR* (Desf. act. soc. hist. nat. 1. p. 3.) caule subherbaceo, foliis ellipticis crenato-dentatis, inferioribus petiolatis, superioribus sessilibus basi auriculatis, floribus corymbosis.  $\mathcal{L}$  in Gallia australi. — Fl. fr. 4. p. 187. — Dalech. hist. 678. ic. (v. s.)
2. *B. VIRGATA* (Desf. l. c. 1. p. 2.) caule herbaceo basi ramoso ramis unifloris, foliis caulinis sessilibus linearibus acutis dentato-serratis, radicalibus spathulatis obtusis crenato-dentatis.  $\mathcal{L}$  in aridis agri Nicæensis. — Fl. fr. 4. p. 188. Chrysanth. discoidæum All. ped. t. 11. f. 1. (v. v. DC.)
3. *B. AGERATIFOLIA* (Desf. l. c. 1. p. 2.) caule suffruticoso basi simplici superne ramoso, foliis oblongo-spathulatis argutè profundeque serratis dentibus mucronatis.  $\mathcal{L}$  in Corsicâ. — Moris. s. 6. t. 9. n. 16. (v. s.)
4. *B. ANNUA* (DC. fl. fr. 4. p. 187.) caule herbaceo superne ramoso corymboso, foliis radicalibus bipinnatifidis, caulinis confertis pinnatifidis pubescentibus, pinnis linearibus acutis mucronatis.  $\textcircled{1}$  in sabulosis incultis circâ Avenionem, Vindomagum, Arelatem, Nicæam. — Tanacetum annuum L. sp. 1183. — Clus. hist. 1. p. 336. f. 1. (v. s.)
5. *B?* *AUDREATHI* (Req. in ann. sc. nat. t. 5. p. 382.) foliis bipinnatifidis pubescentibus lacinii linearilanceolatis incisè acutis, floribus paucis subcorymbosis.  $\mathcal{L}$  in editiori sylvâ Valdonicellâ Corsicæ (Req. Soleir.). (v. s. nondum florentem.)

XXXV. AMBROSIA *Tourn. t. 252. L. gen. 1057.*

Monoica. *Masc.* Involucrum monophyllum; receptaculum nudum; flosculi tubulosi. *Fem.* Perigonium nullum; involucrum 1-phyllum 1-florum extus muricatum, intus 1-loculare; semen involucre indurato obtectum.

1. *A. MARITIMA* (L. sp. 1401.) foliis bipinnatis obtusis subtus canis, racemis terminalibus solitariis, caule divaricato, ramis villosis.  $\textcircled{1}$  in arenosis maritimis circâ Nicæam. — Fl. fr. 3. p. 326. — Barr. ic. f. 1144. (v. s.)

XXXVI. XANTHIUM *L. gen. n. 1056.*

Monoicum. *Masc.* Involucrum polyphyllum multiflorum; receptaculum paleaceum; flosculi tubulosi. *Fem.* Invol. 1-phyllum flores omnino involvens extus muricatum, intus in loculos 2 unifloros divisum; perigonium 0; semina involucre indurato clauso obtecta.

1. *X. MACROCARPUM* (DC. fl. fr. 5. p. 356.) caule inermi, foliis basi cuneatis subcordatis trinerviis, fructibus oblongis, aculeis conicis incurvis uncinatis, terminalibus corniformibus intus uncinatis.  $\textcircled{1}$  in vineis Oecitaniz inferioris. — *X. orientale* L. sp. 1400. excl. syn. Gærtn. 2. t. 164. f. 9. (v. v.)
2. *X. STRAMARIUM* (L. sp. 1400.) caule inermi, foliis cordatis 3-nerviis, fructibus ovatis, aculeis linearibus rectis apice breviter hamatis, terminalibus corniformibus rectis.  $\textcircled{1}$  ad sepes et vias Galliz mediæ, occidentalis australisque. — Fl. fr. 3. p. 326. — Lob. ic. t. 588. t. 2. (v. v.)
3. *X. SPINOSUM* (L. sp. 1400.) spinis ternatis, foliis trilobis.  $\textcircled{1}$  ad agros et vias Galliz mediterraneæ, circâ Aginnum ex St.-Aman, Burdigalam ex Laterr., Parisios ex Mér. — Fl. fr. 3. p. 327. Lam. ill. t. 765. f. 4. (v. v.)

XXXVII. HELIANTHUS *L. gen.* 979.

Involucrum imbricatum, foliolis patulis vel reflexis; receptaculum latum paleaceum; flores radiati, flosculi hermaphroditis in medio ventricosi, ligulis sterilibus; semina aristis 2 mollibus et caducis coronata.

1. *H. annuus* (*L. sp.* 1276.) foliis omnibus cordatis trinerviatis hirtis, pedunculis incrassatis, floribus cernuis. (1) è Peruvia ortus, in hortis cultus. — *Fl. fr.* 4. p. 220. *Mill. ill. ic.* (v. v.)
2. *H. tuberosus* (*L. sp.* 1277.) foliis triplinerviis scabris acuminatis, inferioribus cordato-ovatis, superioribus ovatis in petiolum decurrentibus, squamis involucri ciliatis. 2/ è Brasiliâ ortus, in hortis cultus. — *Fl. fr.* 4. p. 220. *Jacq. hort. vind. t.* 161. (v. v.)
3. *H. multiflorus* (*L. sp.* 1277.) foliis triplinerviis scabris, inferioribus cordatis, superioribus ovatis acutis, involucri squamis lanceolatis vix ciliatis. 2/ è Virginia ortus, in hortis cultus. — *Fl. fr.* 4. p. 220. — *Pluck. t.* 159. f. 2. (v. v.)

XXXVIII. BIDENS *Tourn.* 262. *L. gen.* 932.

Involucrum calyculatum foliolis exterioribus longioribus patulis; receptaculum planum paleaceum; flores plerumque flosculosi, flosculi omnibus hermaphroditis raro radiatis, ligulis nunc femineis, nunc hermaphroditis; semina 2-5 aristis persistentibus coronata.

1. *B. tripartita* (*L. sp.* 1165.) floribus flosculosis bracteatis, bracteis flore majoribus, foliis tripartitis, foliolis lanceolatis dentatis. (1) in fossis et aquaticis frequens. — *Fl. fr.* 4. p. 219. — *Blackw. t.* 519. (v. v.)
2. *B. cernua* (*L. sp.* 1165.) floribus terminalibus subnutantibus, bracteis lanceolatis integris, foliis subconnatis ovato-lanceolatis serratis. (1) in fossis humidis, ad fontes. — *Fl. fr.* 4. p. 219. — *Tab. ic.* 117. Variet radiata et flosculosa. (v. v.)

XXXIX. TAGETES *Tourn.* t. 278. *L. gen.* 964.

Involucrum polyphyllum simplex squamis coalitis; receptaculum nudum; flores radiati; semina aristis 5 coronata.

1. *T. patula* (*L. sp.* 1249.) caule patulo, foliis pinnato-multifidis, laciniis lanceolatis, ciliato-serratis pedunculis unifloris subincrassatis, involucri laevibus. (1) ex Mexico orta, in hortis culta. — *Fl. fr.* 4. p. 172. — *Dill. elth. t.* 279. f. 361. (v. v.)
2. *T. erecta* (*L. sp.* 1249.) caule erecto, foliis pinnato-multifidis, foliolis lanceolatis ciliato-serratis, pedunculis unifloris ventricosso-incrassatis, involucri angulatis. (1) cum priori. — *DC. syn. gall. p.* 285. — *Col. ecphr. 2. t.* 46. f. 1. (v. v.)

XL. CALENDULA *L. gen.* 990.

Involucrum polyphyllum simplex æquale; flores radiati; flosculi in disco masculi, in radio hermaphroditi; ligulæ femineæ fertiles; semina nuda membranacea irregularia curvata.

1. *C. arvensis* (*L. sp.* 1303.) seminibus cymbiformibus muricatis incurvatis, exterioribus lanceolato-subulatis dorso muricatis. (1) in vineis et arvis. — *Fl. fr.* 4. p. 177. — *Moris. 2. 6. t.* 4. n. 6. (v. v.)
2. *C. officinalis* (*L. sp.* 1304.) seminibus omnibus cymbiformibus muricatis incurvatis. (1) in arvis Galliarum australis. — *Fl. fr.* 4. p. 177. — *Blackw. t.* 106. (v. v.)

## Trib. II. CYNAROCEPHALÆ.

Corollæ omnes tubulosæ, receptaculum paleaceum; stigma

cum apice styli articulatam. — Folia alterna sæpius spinosa; genitalia ex irritatione sæpius contractilia.

† PAPPO NULLO.

**XLII. ECHINOPS** *L. gen. 999. DC. mem. comp. p. 19.*

Involucra uniflora numerosa polyphylla squamis linearibus acutissimis, basi setis oblecta, aggregata super receptaculum nudum globosum in capitulum sphericum basi squamis reflexis minimis cinctum; corollulæ 5-dentatæ; semina pentagona pappo piloso brevissimo coronata.

1. *E. SPHÆROCEPHALUS* (L. sp. 1314.) caule ramoso, foliis pinnatifidis superne pubescentibus subtus incano-lanuginosis, setis involucri dimidium superantibus.  $\mathcal{U}$  in sterilibus et incultis provinciis australiorum. — Fl. fr. 4. p. 71. — Lob. ic. 2. t. 8. f. 2. (v. v.)
2. *E. RITAO* (L. sp. 1314.) caule subsimplici corymboso, foliis pinnatifidis supra glabris subtus tomentosis, setis involucri subtriplo brevioribus.  $\mathcal{U}$  in collibus sterilibus Gallie australis. — Fl. fr. 4. p. 71. — Lob. ic. 2. t. 8. f. 1. — Gouan ill. 74. (v. v.)

**XLIII. CARTHAMUS** *DC. mem. comp. p. 20. — Carthami sp. L.*

Involucrum foliaceum imbricatum subspinosum; flosculi numerosi omnes hermaphroditi tubulosi 5-dentati; staminum filamenta glabra; receptaculi palæ longitudinaliter dissectæ et setacæ; semina glabra lævia.

1. *C. TINCTORIUS* (L. sp. 1262.) caule glaberrimo, foliis ovatis integris spinoso-dentatis, pappo nullo.  $\mathcal{U}$  in Gallia australi culta, circa Nicæam disseminata ex All. — Fl. fr. 4. p. 72. Lam. ill. t. 661. f. 3. (v. v.)

†† PAPPO PILOSO AUT PLUMOSO.

**XLIII. CARDUNCELLUS** *DC. mem. comp. p. 20.*

Involucrum foliaceum imbricatum subspinosum; flosculi numerosi, omnes hermaphroditi tubulosi 5-dentati; staminum filamenta pilis hispida; receptaculi palæ longitudinaliter dissectæ et setacæ; semina lævia, pappus pilosus. — Flores cærulei.

1. *C. MITIASIMUS* (DC. fl. fr. 4. p. 73.) foliis inermibus, radicalibus lanceolatis serratis, caulinis pinnatifidis, involucri subcylindracei squamis subaveniis integris inermibus, caule brevi 1-floro.  $\mathcal{U}$  in Gallia mediâ, occidentali et australi. — *Carthamus mitiasimus* L. sp. 1164. (v. v.)
2. *C. MONSPELIENSIS* (All. ped. n. 563.) foliis spinosis, radicalibus obovatis serratis, caulinis pinnatifidis, invol. ovati squamis exterioribus trinerviis margine et apice spinosis, caule brevi 1-floro.  $\mathcal{U}$  in montosis aridis Gallie australis. — Fl. fr. 4. p. 73. — Lob. ic. 2. p. 20. f. 1. (v. v.)
3. *C. CÆVULUS* (DC. mas.) caule elongato subunifloro, foliis spinosis lanceolatis inciso-dentatis sessilibus amplexicaulibus.  $\mathcal{U}$  in Corsicâ reperiit Ph. Thomas. — *Carthamus cævulus* L. sp. 1163. — Moris. s. 7. t. 34. n. 19. (v. s.)
4. *C. TINGITANUS* (DC. mas.) caule elongato subunifloro, foliis spinosis, radicalibus pinnatifidis laciniis lineari-lanceolatis dentatis, superioribus amplexicaulibus basi subauriculatis lineari-lanceolatis inciso-dentatis.  $\mathcal{U}$  in collibus circa Bonifacio reperiit Soleirol. — *Carthamus tingitanus* L. sp. 1163. An notæ characteristicæ hanc speciem à præcedente discernentes, satis constantes? (v. s.)

XLIV. *ARCTIUM* Dalech. Lugd. 1307. f. 1. Lam. DC. mem. comp. p. 21.

Involucrum imbricatum squamis acutis vix spinosis; receptaculum favosum alveolorum marginibus dentatis; semina lævia prismatica; pappus persistens rigidus pilosus pilis spiraliter contortis.

1. *A. LAMUGINOSUM* (Lam. ill. t. 664.) foliis petiolatis subrotundo-ovatis subcordatis crenato-dentatis inermibus tomentosis. 2. in locis schistosis et apricis Alpium Delphin. et Galloprovinciæ. — Fl. fr. 4. p. 75. — Moris. s. 7. t. 32. n. 4. (v. v. DC.)

XLV. *LAPPA* Tourn. inst. 450. t. 156. DC. mem. comp. p. 20.

Involucrum imbricatum squamis desinentibus in spinam mollem apice hamatam; receptaculum paleaceum; pappus brevis persistens pilis rigidis inæqualibus.

1. *L. TOMENTOSA* (Lam. dict. 1. p. 377.) involucri arachnoideo-tomentosis. ② ad vias et in rudertis. — Fl. fr. 4. p. 77. — Mill. ic. t. 139. (v. v.)
2. *L. GLABRA* (Lam. dict. 1. p. 377.) involucri glabris. ② ad vias et in nemoribus. — J. Bauh. hist. 3. p. 570. f. 1.
  - a. *minor* (DC. mass.) floribus minoribus racemosis. — L. minor Fl. fr. 4. p. 77. eng. bot. t. 1228. (v. v.)
  - β. *major* (DC. mass.) floribus majoribus solitariis. — L. major Fl. fr. 4. p. 77. — Dod. perempt. 58. f. 1. (v. v.)

XLVI. *ONOPORDUM* L. gen. 927. DC. mem. comp. p. 20.

Involucrum imbricatum squamis pungentibus; receptaculum favosum; semina compressa tetragona transverse sulcata; pappus caducus pilosus, pilis basi annulatis connexis.

#### §. 1. *Caulescentia, foliis decurrentibus.*

1. *O. ARABICUM* (L. sp. 1159. excl. Barr. syn.) involucri squamis adpressis ovato-lanceolatis breviter spinosis glabris, foliis pinnatifido-sinuatis dentato-spinosis tomentosis. ② Monspelii, in aridis. — Fl. fr. 5. p. 456. — Pluck. alm. t. 154. f. 5. (v. v.)
2. *O. ACANTHIUM* (L. sp. 1158.) involucri squamis patentissimis lineari-subulatis flosculis brevioribus, foliis tomentosis sinuatis dentato-spinosis, ramis multifloris. ② ad margines viarum. — Fl. fr. 4. p. 74. — Lob. ic. 2. p. 1. f. 1. (v. v.)
3. *O. VIRENS* (DC. fl. fr. 5. p. 456.) involucri squamis viscidulis arachnoideo-pubescentibus plano-concavis rectis patentissimis, foliis vix pubescentibus sinuatis dentato-spinosis, summis elongatis apice subintegris, caule pubescente ramoso. ② Monspelii in aridis. — Barr. ic. t. 501. (v. v.)
4. *O. ILLYRICUM* (L. sp. 1148.) invol. squamis basi latis apice angustis exterioribus recurvo-reflexis, interioribus patentibus, foliis tomentosis sinuatis dentato-spinosis, summis brevissimis, caule tomentoso subsimplici. 2. in siccis et sterilibus Gallie australis. — Fl. fr. 4. p. 74. Gærtn. fruct. t. 181. f. 1. (v. v.)

#### §. 2. *Acaulia, foliis subradicalibus.*

5. *O. PYRÆNAICUM* (DC. fl. fr. 5. p. 457.) involucri plurimis ovatis subsessilibus, squamis lanceolatis spinosis suberectis, foliis petiolatis pinnatifidis dentato-spinosis tomentosis subtus candidissimis. 2. in Pyrenæis. — O. acaule. Fl. fr. 4. p. 75. non L. (v. v.)

XLVII. *ALYBUM* *Vaill. acad. 1718. DC. mem. compos. p. 21.*

Involucrum imbricatum squamis basi foliaceis arete adpressis apice appendiculatis patulis spinescentibus; receptaculum paleaceum; pappus piloso-paleaceus deciduus, pilis basi in anulum connatis.

1. *S. MARIANUM* (Gært. fruct. 2. p. 378. t. 162. f. 2.) foliis amplexicaulibus hastato-pinnatifidis albo-maculatis glabris. ① ad margines viarum et in locis incultis. — *Carduus Marianus*. Fl. fr. 4. p. 78. — Lob. ic. p. 7. f. 2. (v. v.)

XLVIII. *CARDUUS* *Gært. 2. p. 377. t. 162. DC. mem. comp. p. 21. — Cardui sp. L.*

Involucrum imbricatum, squamis simplicibus apice spinosis; receptaculi paleæ longitudinaliter dissectæ in setas lineares; pappus caducus pilosus pilis basi in anulum connatis.

1. *C. FUTANS* (L. sp. 1150.) foliis decurrentibus lanceolatis sinuatis spinosis, floribus solitariis cernuis, involucri squamis patentibus lanceolatis spinosis. ② in rudertis. — Fl. fr. 4. p. 80. — Moris. s. 7. t. 31. n. 6. (v. v.)
2. *C. NIGERSCENS* (Vill. dauph. 3. t. 20.) foliis decurrentibus lanceolatis sinuato-pinnatifidis spinosis, floribus solitariis cernuis, involucri squamis patentibus linearibus breviter spinosis. ② in asperis Delph., Galloprov., Occitanis. — Fl. fr. 5. p. 458. (v. v. DC.)
3. *C. ANTHURUS* (Desf. atl. 2. t. 222.) foliis decurrentibus oblongis sinuato-pinnatifidis margine spinosis utrinquæ tomentosis, capitulis paucis aggregatis breviter pedunculatis terminalibus, involucri squamis linearibus apice subulatis vix spinosis, caule simplici. ② in asperis Galloprovinciæ superioris. — Fl. fr. 5. p. 457. — Herr. ic. 417. (v. s.)
4. *C. CARLINODES* (Gou. ill. p. 62. t. 23.) foliis decurrentibus pinnatifidis spinosis utrinquæ tomentosis, floribus glomeratis, invol. squamis linearilanceolatis exterioribus spinosis, interioribus mollibus apice subscariosis. ② in Pyrenæorum glareosis. — Fl. fr. 4. p. 83. (v. v.)
5. *C. PODACANTHUS* (DC. fl. fr. 4. p. 80.) foliis decurrentibus lanceolatis pinnatifidis spinosis, floribus congestis, pedunculis brevibus alato-spinosissimis, invol. squamis linearilanceolatis subulatis subpatentibus pungentibus. ② in Delphinatûs montosis. — *C. aurescens* Chaix in Vill. dauph. 3. t. 20. (v. s.)
6. *C. CRISPUS* (L. sp. 1150.) foliis decurrentibus oblongis sinuatis margine breviter spinosis, floribus aggregatis terminalibus, invol. squamis linearilanceolatis patulis exterioribus subulato-pungentibus, interioribus mollibus apice scariosis. ② in arvis, cultis, ad sepes provinc. orientaliæ. — Fl. fr. 4. p. 79. — Lœs. pruss. t. 5. (v. v.)
7. *C. ACANTHOIDES* (L. sp. 1150.) foliis decurrentibus sinuato-pinnatifidis margine spinosis, floribus solitariis congestive erectis, involucri squamis linearibus laxius subulatis, exterioribus subpungentibus, interioribus mollibus. ① in arvis incultis. — Fl. fr. 4. p. 79. Vix à precedente distinctus. (v. v.)
8. *C. LEUCOGRAPHUS* (L. sp. 1149.) foliis decurrentibus sinuato-dentatis breviter spinosis subtus arachnoideo-tomentosis, pedunculis longis unifloris tomentosis nudis, involucri squamis linearilanceolatis subspinosis arachnoideis erectis. ① in sterilibus Corsicæ et Galliæ australis. — Fl. fr. 4. p. 78. All. ped. t. 73. *Tyrinnus leucographus* Cass. (v. s.)
9. *C. TENUIFLORUS* (Sm. Fl. brit. 849.) foliis decurrentibus sinuatis subtus tomentosis, capitulis ovato-cylindraceis subaggregatis, involucri squamis linearilanceolatis laxiusculis, exterioribus spinosis, interioribus mollibus acutis. ① ② in locis incultis, ad muros.  
*a. acanthifolius* (DC. mss.) alis caulinis latis, capitulis aggregatis sessilibus, foliis immaculatis. In Galliâ australi. — *C. tenuiflorus* eng. bot. t. 412. Fl. fr. 4. p. 79. — Moris. s. 7. t. 31. n. 13. (v. v.)

- β. *Arabicus* (DC. mss.) alis caulinis latis, floribus aggregatis, foliis albis venis notatis. — In Galliâ australi. — C. *Arabicus* Jacq. ic. rar. t. 166. (v. s.)
- γ. *pycnocephalus* (DC. mss.) alis caulinis angustis, floribus aggregatis subsessilibus. — In Corsicâ et Galliâ australi. — C. *pycnocephalus* Jacq. vind. t. 44. non L. — Moris. s. 7. t. 30. f. 9. n. 17. (v. v. DC.)
- δ. *elongatus* (DC. mss.) alis caulinis angustis valde interruptis, pedunculis elongatis nudis uni-aut paucifloris. — In Galliâ meridionali. — C. *pycnocephalus* Fl. fr. 4. p. 79. (v. v.)
10. C. *distortus* (L. sp. 1152.) foliis decurrentibus oblongis sinuatis spinoso-serratis glabris subtus glaucis, pedunculis longissimis unifloris tomentosis, squamis erectis linearibus acutis. ½ in lapideis herbosis montium Galliæ australis orientalisque. — Fl. fr. 4. p. 81. Jacq. aust. t. 89. — Hall. helv. t. 4. f. 2. (v. v.)
11. C. *argemone* (Lam. dict. 1. p. 700.) foliis semi-decurrentibus oblongo-lanceolatis semi-pinnatifidis spinoso-serratis glabris subtus glaucis, pedunculis breviusculis 1-floris tomentosis, invol. squamis lineari-subulatis strachnoideo-tomentosis, exterioribus subspinosis, interioribus mollibus vix acutis. ½ in Pyrenæis. — Fl. fr. 4. p. 83. Nymium præcedenti affinis. (v. s.)
12. C. *medius* (Gouan ill. p. 62. t. 24.) foliis semi-decurrentibus oblongo-lanceolatis semi-pinnatifidis spinosis serratis subtus glaucis laciniis trilobis, pedunculis paucis longissimis tomentosis unifloris, invol. squamis lineari-subulatis subglabris, exterioribus subspinosis, interioribus mollibus acutis. ½ in Pyrenæorum pratis montanis et humidiusculis. — Fl. fr. 4. p. 82. A præcedente non satis distinctus. (v. v.)
13. C. *carlinifolius* (Lam. dict. 1. p. 700.) foliis decurrentibus lanceolatis pinnatifidis glabris subtus glaucis rigido-spinosis, pedunculis tomentosis unifloris erectis, invol. squamis glaucis lineari-lanceolatis exterioribus subulato-pungentibus, interioribus molliusculis. ½ in Pyrenæis et Alpibus Galloprovinciæ. — Fl. fr. 4. p. 82. (v. v.)
14. C. *personata* (Jacq. aust. t. 348.) foliis caulinis semidecurrentibus oblongis indivisis spinoso-dentatis subtus villosis, radicalibus basi pinnatifidis, floribus subsessilibus congestis, invol. squamis lineari-subulatis subinermibus. ☉ in pascuis humidis montosis. — Fl. fr. 4. p. 84. — Gærtn. fruct. 2. t. 160. f. 1. (v. v.)

#### XLIX. SERRATULA DC. mem. comp. p. 21.

Involucrum imbricatum squamis inermibus acutis; receptaculi palæ longitudinaliter dissectæ in setas lineares; pappus persistens pilis rigidis inæqualibus.

\* *Foliis subtus glabris.*

1. S. *tingitoria* (L. sp. 1144.) foliis subpinnatifidis argutè serratis, laciniis inferioribus minoribus terminali maximâ, flosculis omnibus æqualibus. ½ in nemoribus et pratis umbrosis. — Fl. fr. 4. p. 85. Fl. dan. t. 281. (v. v.)
- β. *indivisa*, foliis omnibus indivisis. (v. v.)
- γ. *incisa*, foliis omnibus incisissimis laciniis subæqualibus. — Fl. fr. 5. p. 458. S. *coronata* Fl. fr. 4. p. 85. excl. desc. — Bocc. mus. t. 37. (v. s.)
2. S. *heterophylla* (Desf. cat. p. 93.) foliis inferioribus longè petiolatis ovatis grossè serratis, superioribus basi pinnatifido-incisissimis, caule simplici unifloro spice nudo. ½ in montibus Delphinatûs ex Vill. — Fl. fr. 4. p. 86. — Vill. daph. 3. p. 23. t. 19. Klasea *heterophylla* Cass. (v. s. c.)
3. S. *nudicaulis* (DC. fl. fr. 4. p. 86.) foliis subradicalibus integris ovalibus aut oblongis, caulinis paucis linearibus sessilibus integris aut vix inciso-dentatis, caule 1-floro subnudo. ½ in montibus Galloprovinciæ, Vapinci. — Centaurea *nudicaulis* L. — Ger. galloprov. t. 5. (v. s.)



\* \* *Foliis subtus albo-tomentosis.*

4. *S. HUMILIS* (Desf. atl. t. 220.) foliis pinnatifidis laciniis oblongis acutis costam non attingentibus, caule 1-floro foliis brevioribus subnudo, involucri squamis laxis glabrisculis.  $\mathcal{U}$  in Cebennis inter Campestre et Salboux, in Pyrenæis ad collum Benaque. — Fl. fr. 5. p. 458. *Carduus mollis* Lapeyr. non L. (v. s.)
5. *S. CYTHAROIDES* (DC. fl. fr. 4. p. 87.) foliis amplius ovatis, inferioribus ad costam fere pinnatifidis laciniis dentatis, mediis ad medium usque incis, summis dentatis, involucri squamis margine scariosis ovato-lanceolatis.  $\mathcal{U}$  in Pyrenæis. — *Cnicus centauroides* L. *Stemmacantha centauroides* Cass. — Moris. s. 7. t. 25. n. 2 bis. (v. v. DC.)

L. SAUSSUREA DC. *mem. comp.* p. 22.

Involucrum imbricatum squamis inermibus; flosculi omnes hermaphroditi; receptaculum paleaceum; pappus duplicis ordinis, pili externi breves denticulati persistentes, interni longi plumosi.

1. *S. ALPINA* (DC. l. c. p. 42.) foliis subtus villosis subdentatis, radicalibus ovato-lanceolatis, floribus terminalibus subumbellatis.  $\mathcal{U}$  in editioribus Alpibus et Pyrenæis. — Fl. fr. 5. p. 466. — Pluck. t. 154. f. 3. (v. v. in Helv.)
2. *S. DISCOLOR* (DC. l. c. p. 43.) foliis subtus niveo-tomentosis dentatis, radicalibus ovatis subcordatis, caulinis ovato-lanceolatis, floribus terminalibus subumbellatis.  $\mathcal{U}$  rarior, in Alpibus Delphinatûs. — Fl. fr. 5. p. 466. — Hall. helv. t. 6. (v. s.)

LI. *CIRSIIUM* *Turn.* t. 255. DC. *mem. comp.* p. 23. — *Cirsium*, *Onotrophe*, *Notobasis*, *Eriolepis*, *Picnomon*, *Lamyra*, *Cass.*

Involucrum obovatum imbricatum squamis apice spinosis; flosculi omnes hermaphroditi æquales; receptaculi paleæ in laciniis setiformes partitæ; stylus simplex, pappi pili plumosi æquales basi in annulum conferruminati.

#### Sect. I. ERYSTHALES DC. *mss.*

Flores ochroleuci, folia sessilia.

1. *C. GLUTINOSUM* (Lam. fl. fr. 2. p. 27.) foliis amplexicaulibus pinnatifidis glabris ciliatis, pedunculis apice cernuis, involucri glutinosi squamis lanceolatis inermibus breviter mucronatis patulis.  $\mathcal{U}$  in Alpium, Jurassi, montium Arvernæ nemoribus udis. — Fl. fr. 5. p. 464. *C. erysthales* Jacq. obs. t. 17. An satis à sequente distinctum? (v. v. DC.)
2. *C. OCHROLEUCUM* (All. ped. n. 546.) foliis amplexicaulibus pinnatifidis glabris ciliatis, pedunculis erectis brevibus albo-lanatis, involucri (non glutinosi) squamis lanceolatis suberectis mucronatis, exterioribus subspinosis.  $\mathcal{U}$  in Alpibus. — Fl. fr. 5. p. 464. — Dalech. lugd. 1094. ic. (v. v.)
3. *C. ZYUGAGNEUM* (DC. fl. fr. 4. p. 115.) foliis amplexicaulibus pinnatifidis apice elongatis subtus villosis, laciniis incis, apice spinescentibus, floribus congestis terminalibus sessilibus, involucri glabri squamis lanceolatis acuminatis inermibus adpressis.  $\mathcal{U}$  in pratis frigidis Delphinatûs et Jurassi. — Fl. fr. l. c. excl. syn. Vill. (v. s.)
4. *C. TATARICUM* (All. ped. n. 550.) foliis amplexicaulibus glabris ciliato-spinulosis inferioribus pinnatifidis laciniis inciso-bifidis dentatis, superioribus indivisis lanceolatis, floribus solitariis foliis oblongo-linearibus stipatis, invol. glabri squamis exterioribus lanceolatis mucronato-spinosis, interioribus linearilanceolatis acuminatis inermibus.  $\mathcal{U}$  in vallibus locisque humidis Alpium Delphinatûs. — Fl. fr. 4. p. 114. *Carduus tataricus* Jacq. austr. t. 90. (v. s.)
5. *C. OLERACEUM* (All. ped. n. 544.) foliis amplexicaulibus cordatis ciliato-spi-

- nosis, caulinis pinnatifidis, superioribus ovatis acutis, summis flores sessiles aggregatos cingentibus et superantibus, involucri squamis lanceolato-acuminatis exterioribus spinosis.  $\mathcal{U}$  in pratis paludosis. — Fl. fr. 4. p. 114. *Cnicus oleraceus* L. *Onotrophe oleracea* Cass. — Lob. ic. 2. t. 11. f. 1. (v. v.)
6. *C. AUSTRIACUS* (Ram. in DC. fl. fr. 4. p. 114.) foliis amplexicaulibus ovato-oblongis subcordatis sinuato-dentatis ciliato-spinosis, floralibus sublinearibus, floribus congestis subracemosis canleque pilos rufescentes gerentibus, involucri squamis exterioribus ciliatis, internis linearibus subinermibus.  $\mathcal{U}$  in Pyrenæis editoribus inter rupes. — *C. cerniolicum* Scop. carn. t. 54. (v. s.)
7. *C. SPINOSISSIMUS* (Scop. carn. n. 1006.) foliis amplexicaulibus pinnatifidis dentatis spinosis pubescentibus, caule simplici, floribus terminalibus sessilibus congestis bracteatis, involucri squamis lanceolatis spinosissimis.  $\mathcal{U}$  in Alpium et Jurassi pratis ndis propè nives. — Fl. fr. 4. p. 113. — Hall. helv. t. 5. (v. v. in Helv.)
8. *C. GLABRUM* (DC. fl. fr. 5. p. 463.) foliis pinnatifidis dentatis spinosis glabris, floribus terminalibus pedicellatis congestis, bracteis angustis spinosissimis florem vix superantibus, invol. squamis acuminatis, exterioribus spinosis, interioribus subinermibus.  $\mathcal{U}$  in Pyrenæis editoribus secus torrentes. — *Cnicus spinosissimus* Lapeyr. (v. v. DC.)

Sect. II. *ACARNA DC. mss.*

Folia decurrentia supernè non setoso-scabra, flores purpurei aut variantes albi.

9. *C. PALUSTRE* (Scop. carn. n. 1004.) foliis decurrentibus lanceolatis sinuato-pinnatifidis marginè spinosis, floribus racemoso-glomeratis, involucri squamis adpressis, exterioribus ovato-lanceolatis mucronato-spinosis, interioribus linearibus acuminatis.  $\mathcal{U}$  in stagnis pratiquè umbrosis. — Fl. fr. 4. p. 111. eng. bot. t. 974. *Onotrophe palustris* Cass. (v. v.)
10. *C. PRATENAE* (DC. fl. fr. 4. p. 113.) foliis breviter decurrentibus sinuatis dentatis inæqualiter setoso-ciliatis utrinquè glabris, pedunculis foliolosis subtomentosis, involucri squamis exterioribus ovato-lanceolatis mucronato-spinosis, interioribus lanceolatis acuminatis inermibus apice patulis.  $\mathcal{U}$  in humidis Galliæ australis ex Lam. et Pyrenæorum. — Moris. s. 7. t. 29. n. 8. (v. s.)
11. *C. MONSPESULANUM* (All. ped. n. 556.) foliis longè decurrentibus oblongo-lanceolatis inæqualiter setoso-ciliatis utrinquè glabris, inferioribus sinuato-crenatis, superioribus integris, pedunculis subnudis tomentosis multifloris, involucri squamis exterioribus ovato-lanceolatis mucronatis, interioribus linearibus lanceolatis acuminatis inermibus erectis.  $\mathcal{U}$  in pratis humidis Delph., Galliæ australis, Pyrenæorum. — Fl. fr. 4. p. 112. — Lob. ic. t. 581. f. 2. (v. v.)
12. *C. PYRÆNÆICUM* (DC. fl. fr. 4. p. 112.) foliis longè decurrentibus oblongo-lanceolatis subtus arachnoideo-villosis canescentibus sinuatis ciliato-spinosis, invol. squamis exterioribus ovato-lanceolatis mucronatis, interioribus linearibus lanceolatis acuminatis inermibus erectis.  $\mathcal{U}$  in Pyren. orientalium pratis. — An varietas præcedentis ut vult cl. Benthani? (v. s.)
13. *C. ACARNA* (DC. fl. fr. p. 111.) foliis decurrentibus lanceolatis lanato-canis dentatis spinosis, superioribus flores aggregatos subsessiles circumdantibus, involucri squamis exterioribus spinâ pinnatâ terminatis.  $\mathcal{U}$  in petrosis et sterilibus Galliæ mediterraneæ. — *Cnicus Acarna* L. sp. 1158. *Piconomon Acarna* Cass. — Clus. hist. 2. p. 155. f. 1. (v. v.)

Sect. III. *EPITRACHYS DC. mss.*

Folia supernè scabra subtus tomentosa pinnatifida sessilia aut decurrentia, flores purpurei aut variantes albi.

14. *C. ITALICUM* (DC. mss.) foliis decurrentibus in alam brevem obtusam spinosis subtus arachnoideo tomentosis supernè setoso-scabris, floralibus flores

subsessiles circumdantibus, involucri ovati glabrique squamis lanceolatis adpressis in spinam productis. ② in Corsicâ (Soleirol.). — Dod. pempt. 738. f. 2. (v. s.)

15. *C. LANCEOLATUM* (Scop. carn. n. 1007.) foliis spinosis decurrentibus in alam lobato-spinosam subtus arachnoideo-tomentosis superne setoso-acabris, involucri ovati subarachnoides squamis lanceolatis setaceo-acuminatis, exterioribus spinosis patentibus. ② ad vias et pagos. — Fl. fr. 4. p. 111. *Eriolepis lanceolata* Cass. — J. Bauh. hist. 3. p. 58. f. 1. (v. v.)
16. *C. PRÆOX* (DC. fl. fr. 4. p. 120.) caule arachnoideo, foliis subdecurrentibus spinosis subtus cano-tomentosis superne spinoso-acabris, floralibus flores cingentibus, involucri subglabri ovati crassi squamis lanceolatis subulato-spinosis subpatulis. ② in montosis et collibus Gallie australis. — *Cnicus ferox* All. ped. t. 50. — Lob. ic. 2. t. 10. f. 1. (v. v.)
17. *C. ERIOPHORUM* (Scop. carn. n. 1008.) foliis sessilibus superne ciliato-acabris subtus niveo-tomentosis, laciniiis geminatis spinosis, involucri crassi globosi lanuginosissimi squamis lanceolatis apice linearibus spinosis patentibus. ② in montosis Pyren., Gallie australis orientalisque. — Fl. fr. 4. p. 120. *Eriolepis lanigera* Cass. — Clus. hist. 2. p. 154. ic. (v. v.)
18. *C. ECHINATUM* (DC. fl. fr. 5. p. 465.) foliis sessilibus hispidis subtus lanatis, laciniiis bilobis divaricatis spinosis, involucri ovati crassi globosi lanati bracteati squamis lanceolatis subulato-spinosis basi adpressis. ② in insulâ Sanctæ Lucie prope Nerbonam (Requien). — *Carduus echinatus* Desf. atl. 2. p. 247. (v. s.)

#### Sect. IV. CHAMELÆON DC. *ms.*

Folia sessilia superne non setoso-acabra, flores purpurei aut variantes albi.

19. *C. CASABONÆ* (DC. fl. fr. 4. p. 121.) foliis sessilibus lanceolatis subtus tomentosis superne glaberrimis margine spinas ternatas gerentibus, floribus axillaribus sessilibus, involucri squamis exterioribus in spinam longam productis. ② in Corsicâ et insulâ Stæchadum. — *Carduus Casabonæ* L. *Lamyra triacantha* Cass. — Barr. ic. t. 1211. (v. s.)
20. *C. SYRIACUM* (Gærtn. fruct. 2. p. 383 t. 163. f. 2.) foliis amplexicaulibus oblongis inciso-dentatis spinosis albo-venosis, floribus subsessilibus terminalibus axillaribusve bracteatis bracteis pinnaefido-spinosis, involucri squamis oblongo-lanceolatis adpressis acuminato-spinosis. ① in Corsicâ. *Notobasis Syriaca* Cass. — Moris. s. 7. t. 30. n. 5. (v. s.)
21. *C. ARVENSE* (Lam. fl. fr. 2. p. 26.) foliis sessilibus oblongis sinnato-pinnaefidis undulatis spinosis subtus glabriusculis tomentosive, caule subglabro multifloro, involucri squamis adpressis exterioribus ovato-lanceolatis mucronatis vix spinosis, interioribus lanceolatis inermibus. ② in vineis et arvis frequens. — Fl. fr. 4. p. 119. *Serratula arvensis* L. sp. 1149. — Moris. s. 7. t. 82. n. 14. (v. v.)
22. *C. ACAULE* (All. ped. n. 558.) caulibus brevissimis subnudis subuni-floris, foliis subradicalibus pinnaefidis dentatis ciliato spinosis, involucri squamis lanceolatis glabris mucronatis subinermibus. ② in herbosis et siccis. — Fl. fr. 4. p. 119. *Onotrophe acaulis* Cass. — Lob. ic. 2. t. 5. f. 1. (v. v.)
23. *C. SULCOSUM* (DC. fl. fr. 4. p. 118.) foliis amplexicaulibus pinnaefidis ciliato-spinosis, laciniiis bilobis basi sursum dentatis, caule subbifloro, involucri squamis mucronatis subinermibus apice patulis, radice tuberosâ. ② in humidis herbosis. — Lob. ic. 2. t. 10. f. 2. (v. v.)
24. *C. ANGELICUM* (Lob. ic. t. 583. f. 1.) foliis lanceolatis sinnato-dentatis ciliato-spinosis subtus lanuginosis, inferioribus subincisis subpetiolatis, caulibus sessilibus semi-amplexicaulibus, flore solitario terminali, involucri squamis lineari-lanceolatis acuminatis subinermibus. ② in Gallie occidentalis, mediæ et australis pratis paludosis. — Fl. fr. 4. p. 119. — Dalech. Ingd. 594. f. 1. (v. v. DC.)

- β. multiflorum*. Fl. fr. 5. p. 465. In agro Aurelianensi et Andegaviensi. (v. s.)
25. *C. TRICEPHALODES* (DC. fl. fr. 4. p. 116.) foliis amplexicanlibus nudis glabris setoso-ciliatis pinnatifidis, caulino- rum lacinii lanceolatis oblongis, pedunculis longis tomentosis, floribus 3-5 aggregatis, involucri squamis adpressis ovato-lanceolatis acuminatis inermibus.  $\mathcal{Z}$  in pascuis Jurassi, Alpium, montis Anrei, Pyrenæorum. — *Carduus rivularis* Jacq. aust. t. gr. — All. ped. t. 35 et 36. (v. v.)
26. *C. SALISBURGENSIS* (Duby mss.) foliis amplexicanlibus scabris setoso-ciliatis glabris integris sinuato-dentatis, inferioribus ovalibus, superioribus linearilanceolatis, pedunculis longis tomentosis 1-2-floris, involucri squamis adpressis lanceolatis inermibus.  $\mathcal{Z}$  in Pyrenæorum et Jurassi pratis udis. — *Cnicus Salisburgensis* W. sp. 2. p. 1675. *Cirsium tricephalodes*  $\gamma$  Fl. fr. 5. p. 464. Precedentis forsitan varietas? (v. s.)
27. *C. AMBROSUM* (All. auct. p. 10.) foliis amplexicanlibus subtus albedo-tomentosis supernè glabris subscabris margine setoso-ciliatis, radicalibus petiolatis integris dentatis, caulinis sessilibus pinnatifidis lacinii dentatis aut bifidis, flore longè pedunculato subsolitario, involucri squamis lanceolatis adpressis subinermibus.  $\mathcal{Z}$  in pratis Alpium sæpius commixtum cum *C. heterophyllo* et *tricephalode* quorum videtur mera proles hybrida. — Fl. fr. 4. p. 116. (v. s.)
- β. Antarcticum*, floribus 2-3 aggregatis. — *Carduus antarcticus* Vill. dauph. 3. t. 19.
28. *C. HETEROPHYLLUM* (DC. fl. fr. 4. p. 117.) foliis amplexicanlibus lanceolatis integris aut apice incisitis subtus niveo-tomentosis supernè glabris margine setoso-ciliatis, pedunculis unifloris tomentosis, involucri squamis glaberrimis adpressis inermibus, exterioribus ovato-lanceolatis mucronatis, interioribus linearibus acuminatis.  $\mathcal{Z}$  in pratis humidis secus vias montanas, in Alpibus, Pyrenæis. — *Carduus polymorphus* Lapeyr. act. tol. 1. t. 19 et 20.
- α. helenioides*, foliis omnibus integris. — *C. helenioides* All. ped. t. 13. (v. v.)
- β. incisum*, foliis inferioribus incisitis. — *C. heterophyllum* All. ped. t. 34. (v. s.)
29. *C. STELLATUM* (All. ped. n. 560.) foliis sessilibus lanceolatis integerrimis subtus tomentosis spinis 2-3 axillaribus basi cum folii margine confluentibus instructis, floribus subsessilibus, involucri squamis lanceolatis subulato-spinosis patentibus. ① circa Nicæam ex All. — Fl. fr. 4. p. 121. *Lamyra stipulacea* Cass. — Barr. ic. t. 418. (v. s.)

## LII. CYNARA Juss. gen. 173. DC. mem. comp. p. 23.

Involucrum maximum ventricosum, squamis numerosis basi carnosius apice spinescentibus integris; receptaculi magni carnosius paleæ longitudinaliter dissectæ in lacinias setiformes; pappus longus plumosus.

1. *C. CARDUNCULUS* (L. sp. 1159.) foliis spinosis omnibus pinnatifidis, involucri squamis ovatis.  $\mathcal{Z}$  *α* sylvestris Monspelii et in Corsicâ ex Viv.; *β* hortensis, in hortis culta. — Fl. fr. 4. p. 108. — Tab. ic. 696. (v. v.)
2. *C. SCOLYMUS* (L. sp. 1159.) foliis subspinosis pinnatifidis indivisiisque, involucri squamis ovatis.  $\mathcal{Z}$  in hortis culta. — Fl. fr. 4. p. 108. — Clus. hist. 2. p. 153. f. 3. Prioris forsitan varietas? (v. v.)
3. *C. HUMILIS* (L. sp. 1159.) foliis pinnatifidis spinosis subtus tomentosis, involucri squamis subulatis.  $\mathcal{Z}$  in Corsicâ ex cl. Viviani. — Pluck. alm. 85. t. 181. f. 2. (v. s. ex Hispaniâ.)

## LIII. LEUZEA DC. fl. fr. 4. p. 109. mem. comp. p. 22.

Involucrum magnum imbricatum squamis scariosis inermibus, exte-

rioribus rotundatis laceris, internis longioribus acutis integris; flosculi omnes hermaphroditi; stigma 1; receptaculi paleæ longitudinaliter dissectæ in lacinias setiformes; semina tuberculosa; pappus longus pilis plumosis æqualibus.

1. *L. CONIFERA* (DC. l. c.) foliis tomentosis radicalibus lanceolatis, caulinis pinnatifidis. ½ in sterilibus Corsicæ et Galliæ mediterranæ. — *Centaurea conifera* L. — Barr. ic. t. 138. (v. v.)

LIV. GALACTITES *Mench. meth. 558. DC. fl. fr. 4. p. 170.*

Involucrum obovatum squamis numerosis lanceolato-subulatis spinosis; flosculi disci hermaphroditi, marginales neutri majores; receptaculi paleæ paucæ caducæ; stylus simplex; pappus longus plumosus.

1. *G. TOMENTOSA* (Mench. meth. 558.) foliis decurrentibus sinuatis spinosis subtus tomentosis. ½ in Corsicâ et in sterilibus Galliæ mediterranæ, circa Tolosam ex Lapeyr., Aginnum ex St.-Am. Burdigalam ex Des Moul. — Fl. fr. l. c. DC. mem. comp. t. 5. (v. v.)

LV. RHAPONTICUM *Lam. fl. fr. DC. mem. comp. p. 21.*

Involucrum magnum imbricatum squamis scariosis rotundatis inermibus; receptaculi paleæ longitudinaliter in setas lineares dissectæ; pappus persistens pilis rigidis inæqualibus.

1. *R. SCARIOSUM* (Lam. fl. fr. 2. p. 38.) foliis ovato-oblongis basi subcordatis denticulatis subtus subtomentosis. ½ in Alpibus Delphinatûs, Galloprovinciæ. — DC. mem. comp. p. 32. Serratula Rhaponticum Fl. fr. 4. p. 87. — Moris. s. 7. t. 25. u. 7. (v. s.)

LVI. CENTAUREA *L. gen. 984. DC. fl. fr. 4. p. 88.*

Involucrum imbricatum squamis foliaceis scariosisve; flosculi disci hermaphroditi, radii neutri majores; receptaculi paleæ laciniatæ; semina hylo laterali receptaculo adfixa; pappus pilosus.

Sect. I. CAUPINA *DC. mem. comp. p. 23.*

Invol. squamæ integræ lanceolato-acuminatæ, pappus setoso-paleaceus.

1. *C. CAUPINA* (L. sp. 1285.) involucri inermibus, foliis pinnatis scabris pinnulis serrulatis linearibus acutis. (1) in sterilibus et collibus Galloprov., Occitaniæ, circa Mimatem, Burdigalam ex Later. — Fl. fr. 4. p. 89. — Barr. ic. t. 1136. (v. v.)

Sect. II. CENTAURIUM *DC. l. c. p. 24.*

Involucri squamæ inermes sæpius marcescentes saltem in margine.

2. *C. AMARA* (L. sp. 1292.) involucri glaberrimi lucidi squamis scariosis obtusis integris fimbriatisve, exterioribus ovatis, interioribus lanceolatis apice spathulatis, foliis tomentosis pubescentibusve lanceolatis, caulibus deumbentibus. ½ in collibus aridis Galliæ australis. — Fl. fr. 4. p. 90. — Lob. ic. t. 548. f. 2. (v. v.)

β. *linearifolia* DC. foliis linearibus utrinquæ tomentosis. — Fl. fr. 5. p. 459. (v. v.)

γ. *nana*, vix pollicaris, foliis linearibus pubescentibus. — Circâ Monspe-  
lium reperiit de Chesnel. (v. s.)

3. *C. ALBA* (L. sp. 1293.) involucri glabri lucidi squamis scariosis mucronatis serrulatis, exterioribus ovatis, interioribus lanceolatis apice spathulatis, foliis infimis pinnatifidis lobis linearibus distantibus, mediis basi dentatis, summis integris, caule erecto. (2) in Occitaniâ ex Gon. et Pyrenæis orienta-

libus ex Lapeyr. ? — Fl. fr. 5. p. 460. — Tab. ic. 153. f. 1. (v. s. ex Hispaniâ ?)

Sect. III. CYANUS DC. l. c.

Involucri squamæ inermes pinnatifido-ciliatæ, pappus pilosus.

4. *C. JACEA* (L. sp. 1293.) involucri basi pubescentis squamis ovatis apice spathulatis scarioso-ciliatis, foliis lanceolatis integerrimis radicalibus subdentatis, ramis angulatis, pappo subnullo.  $\mathcal{L}$  ad oras sylvarum et in pratis ubique. — Fl. fr. 4. p. 91. (v. v.)
5. *C. MUTABILIS* (St.-Amans mem. mns. 1. p. 477. t. 24.) involucri globosi squamis pubescentibus exterioribus ovatis apice lineari-spathulatis longè ciliatis acuminatis, internis lanceolatis apice globoso-spathulatis subintegris, foliis tomentosis, radicalibus obsolete lyratis vel indivisis, canlinis integris.  $\mathcal{L}$  circà Aginnum. (v. s.)
6. *C. NIGRASCENS* (W. sp. 3. p. 2288.) involucri squamis exterioribus ovatis scarioso-ciliatis, internis lanceolatis apice spathulatis integris lacerisve, foliis radicalibus obsolete pinnatifidis, mediis basi subdentatis, superioribus integris.  $\mathcal{L}$  circà Parisiis, Bellocvacum ex D. Graves. et in Pyrenæis. — Fl. fr. 5. p. 460. (v. v. c.)
7. *C. NIGRA* (L. sp. 1288.) involucri squamis exterioribus erecto-plumosis, flosculis equalibus, foliis lanceolatis radicalibus subpinnatifidis, canlinis indivisis serratis.  $\mathcal{L}$  in Galliâ boreali, mediâ et occidentali. — Fl. fr. 4. p. 91. Fl. dan. t. 996. — Gærtn. 2. t. 161. f. 4. Squamæ nigræ. (v. v.)
8. *C. PECTINATA* (L. sp. 1287.) involucri squamis recurvato-plumosis, foliis mucronato-serratis, canlinis inferioribus sinuato-pinnatifidis, superioribus et ramis indivisis.  $\mathcal{L}$  in saxosis montosis Galliæ australis. — Fl. fr. 4. p. 93. (v. v.)
9. *C. PHRYGIA* (L. sp. 1287.) involucri squamis recurvato-plumosis, foliis oblongis pubescentibus indivisis scabris mucronato-serratis.  $\mathcal{L}$  in pascuis editiorum montium. — Fl. fr. 4. p. 92. Fl. dan. t. 520. (v. v.)
10. *C. UNIFLORA* (L. mant. 148.) involucri squamis recurvato-plumosis, foliis lanceolatis subdentatis tomentosis, caule unifloro.  $\mathcal{L}$  in pratis montosis Delph., Galloprovinc., Occitaniz? — Fl. fr. 4. p. 93. — Bocc. mus. p. 20. t. 2. (v. v. DC.)
11. *C. CYANUS* (L. sp. 1289.) involucri squamis ciliatis exterioribus ovatis interioribus lanceolatis, foliis linearibus sessilibus integerrimis subtus lanato-tomentosis, infimis basi pinnatifidis. ① inter segetes frequens. — Fl. fr. 4. p. 95. Bull. herb. t. 221. (v. v.)
12. *C. MONTANA* (L. sp. 1289.) involucri squamis ovatis margine ciliatis, foliis lanceolato-acuminatis decurrentibus tomentosis integerrimis aut basi sinuato-dentatis.  $\mathcal{L}$  in pascuis montium. — Fl. fr. 4. p. 94. C. sensana Vill. — Lob. ic. 542. f. 1. (v. v.)
13. *C. FULLATA* (L. sp. 1288.) involucri squamis lanceolatis apice ciliatis, bracteis lanceolatis involucreis, foliis petiolatis basi amplexicaulibus lyratis breviter hispidis. ① ad sepes in Galliâ australi. — Fl. fr. 4. p. 94. — Lob. ic. t. 542. f. 2. (v. v. DC.)
14. *C. INTYBRICA* (Lam. dict. 1. p. 671.) involucri glaberrimi globosi squamis striatis apice ciliatis ovatis, foliis subrigidis lanceolato-linearibus basi laciniatis pinnatifidisve, laciniis acutis simplicibus, terminali dentatâ, caule 1-floro.  $\mathcal{L}$  circà Massiliam et Narbonam. — Fl. fr. 4. p. 98. C. leucantha Pourr. ex cl. Benth. — Barr. ic. 1229. (v. s.)
15. *C. MACULOSA* (Lam. dict. 1. p. 669.) involucri globosi squamis striatis exterioribus ovatis laxiusculis longè ciliatis acutis, interioribus lanceolatis obtusis, foliis inferioribus bipinnatifidis, superioribus pinnatifidis, caule subcorymboso. ② in agro Ruscinonensi, Cebennensi, Ruthenico, Arvernico et circà Tarones. — Fl. fr. 4. p. 96. C. cœrulestems Lapeyr. C. corymbosa Pourr. ex cl. Benth. — Gm. sib. 2. t. 44. f. 1. 2. (v. v.)

- β. nana*, caule 3. floro vix pollicari. — Circà Massiliam reperit cl. Solier. (v. s.)
16. *C. PANICULATA* (L. sp. 1289.) involucri ovoidei squamis ciliatis ovatis adpressis striatis acutis, foliis inferioribus bipinnatifidis, superioribus pinnatifidis, caule paniculato. ② in arvis et sterilibus Gallis australis, Pyrenæorum, Arvernæ, Delphinatûs. — Fl. fr. 4. p. 97. — Moris. s. 7. t. 28. n. 15. (v. v.)
17. *C. SCABIOSA* (L. sp. 1291.) involucri pubescentis squamis ovatis ciliatis, foliis pinnatifidis scabriusculis, lacinis lanceolatis acutis basi inciso-dentatis pinnatifidiæve. ½ ad marginem agrorum et sylvarum. — Fl. fr. 4. p. 97. — Moris. s. 7. t. 28. n. 10. (v. v.)

Sect. IV. *CROCODYLIUM* DC. l. c.

Involucri squamæ in spinam simplicem desinentes, pappus pilosus.

18. *C. SALMANTICA* (L. sp. 1299.) involucri globosi glabri squamis ovatis in setulam subspinosam desinentibus, foliis lanceolatis serratis radicalibus lyratis, caule divaricato. ½ in arvis et pascuis maritimis Galloprovinciæ, Monapellii, Carcassoni, agri Rusciniensis. — Fl. fr. 4. p. 106. — Clus. hist. 2. p. 9. f. 1. (v. v.)
- β. inermis*, squamis inermibus. (v. s.)

Sect. V. *CALCITRAPA* DC. l. c.

Involucri squamæ nunc pinnato-spinosæ nunc palmato-spinosæ, pappus pilosus.

19. *C. COLLINA* (L. sp. 1298.) involucri glabri squamis exterioribus ovatis acuminatis ciliato-spinosis, interioribus ciliatis lanceolatis obtusis, foliis caulinis pinnatifidis, radicalibus bipinnatifidis, lacinis lanceolatis acutissimis. ½ in provinciis australibus. — Fl. fr. 4. p. 115. — Clus. hist. 2. p. 8. f. 2. (v. v.)
20. *C. CENTAUREOIDES* (L. sp. 1298.) involucri glabri squamis exterioribus ovatis acuminatis, interioribus obtusis, foliis lyrato-pinnatis subintegrisve, lacinis lanceolato-ovatis mucronatis, terminali maximâ dentatâ. ½ Monapellii secus rivos. — Fl. fr. 4. p. 105. — Col. œphr. 1, t. 35. Præcedentis var? (v. s.)
21. *C. NAFIFOLIA* (L. sp. 1295.) involucri globosi glabri vix basi pubescentis squamis adpressis ovatis apice transversè appendiculo lineari palmato-spinoso erecto spathulatis, foliis radicalibus lyratis obtusis, caulinis oblongis decurrentibus, caule erecto ramoso. ① in Corsicâ circà Ajaccio. — Fl. fr. 5. p. 461. — Pluck. t. 94. f. 2. (v. s.)
22. *C. ASPERA* (L. sp. 1296.) involucri glabri squamis exterioribus palmato-spinosis, spinâ 3-5-cuspidatâ, foliis lanceolatis sessilibus sinuato-dentatis scabriusculis. ½ in arvis et sterilibus à Nicæâ ad Nigri Monasterii insulam. — Fl. fr. 4. p. 98. (v. v.)
23. *C. SPHEROCEPHALA* (L. sp. 1295.) involucri globosi arachnoidæi squamis ovatis apice palmato-spinosis, spinâ 5-cuspidatâ reflexâ, foliis lanceolatis pinnatifidis lacinis dentato-serratis, radicalibus petiolatis, superioribus amplexicaulibus, caule procumbente. ½ in Corsicâ. — *C. cæspitosa* Vahl. symb. 2. p. 93. — Moris. s. 7. t. 27. n. 9. (v. s.)
24. *C. SKRIBIS* (L. sp. 1294.) involucri glaberrimi globosi squamis palmato-spinosis, spinâ 5-7-cuspidatâ reflexâ, foliis oblongis tomentosis amplexicaulibus decurrentibus basi subincisis dentatis dentibus acuminatis subspinosis. ½ in arvis circà Nicæam ex All., Aquas Sextias ex Gér., Monapellium ex Gouan et in Delphinatû ex Vill. — Fl. fr. 4. p. 99. — Pluck. t. 38. f. 1. (v. s. c.)
25. *C. SONCHIFOLIA* (L. sp. 1294.) involucri glaberrimi globosi squamis palmato-spinosis, foliis oblongis tomentosis amplexicaulibus semi-decurrentibus repando-dentatis dentibus subspinosis, superioribus integerrimis. ①? secus stagna et rivos in Galloprovinciâ ex Gér., agro Nicæensi ex All. — Fl.

- fr. 4. p. 99. — Pluck. t. 39. f. 1, 94. f. 1. An à præced. satis distincta? (v. s. c.)
26. *C. MYACANTHA* (DC. fl. fr. 4. p. 101.) involucri glabri squamis apice corneis spinoso-dentatis, foliis sessilibus lineari-oblongis subtomentosis dentato-serratis vel basi sublaciniatis, caule ramoso glabro, pappi subnulla. ② Parisiis circa Vincennes et Versailles. — Ic. rar. t. 23. (v. s.)
27. *C. ARULA* (Lam. dict. 1. p. 674.) involucri globosi basi tomentosi squamis ovatis apice palmato-spinosis, spinis patulis 5, mediâ duplicato-spinosâ, capitulis axillaribus terminalibusque, foliis lanceolatis integerrimis semi-decurrentibus radicalibus lyratis obtusis. ① Circâ Narbonam, Monspelium, Arelatem, Nicæam et in Corsicâ. — Fl. fr. 4. p. 104. — Col. œschr. 1. p. 31. ic. (v. s.)
28. *C. MELITENSIS* (L. sp. 1297.) involucri squamis subpalmato-spinosis spinis rectis, capitulis subgeminatis sessilibus, foliis oblongis subincanis profundè dentatis pinnatifidisve, laciniis distantibus acutiusculis. ① in Galliâ australiori, Monspelii. — Fl. fr. 4. p. 104. (v. s.)
29. *C. SOLSTITIALIS* (L. sp. 1297.) involucri globosi tomentosi squamis exterioribus subpalmato-spinosis spinis rectis, foliis canescentibus lineari-lanceolatis decurrentibus integerrimis, radicalibus lyratis. ① in siccis Galliæ australis; sæpè cum seminibus Medicaginis in provinciis borealibus seritur, sed non permanet. — Fl. fr. 4. p. 103. — Moris. s. 7. t. 34. n. 9. (v. v.)
30. *C. DIFFUSA* (Lam. dict. 1. p. 676.) involucri squamis lanceolatis ciliato-spinosis spinâ terminali longâ robustiore floribus breviorè, foliis radicalibus bipinnatifidis, caulinis linearibus integris, caule ramosissimo. ② Monspelii, ex lanis barbaris seta? — Fl. fr. 5. p. 462. (v. v.)
31. *C. POUZIN* (DC. cat. monsp. 91.) involucri squamis rotundo-ovatis adpressis apice patente corneo pinnatim 5-7-spinoso, floeculis fertilibus breviter papposis, sterilibus nudis, foliis oblongis inferioribus pinnatifidis, superioribus rariter serratis, caule erecto ramosissimo. ② secûs vias Monspelii, Narbonæ, agri Ruscin. ex Benth. — Fl. fr. 5. p. 462. (v. v.)
32. *C. CALCITRAPA* (L. sp. 1297.) involucri glaberrimi squamis exterioribus ovato-rotundis apice spinosis spinâ mediâ robustâ longissimâ flores superante intimis spatulatis scariosis integris, foliis pinnatifidis linearibus dentatis, radicalibus lyratis, caule ramosissimo stristo piloso, pappo nullo. ① in sterilibus petrosis ad vias. — Fl. fr. 4. p. 100. — Clus. hist. 2. p. 7. f. 3. (v. v.)  
*β. autumnalis* (DC.) foliis inferioribus 2-pinnatifidis, floralibus pinnatifidis, laciniis linearibus dentatis, caule hirsuto. ① secûs vias Monspelii, Lugduni. — Fl. fr. 5. p. 462. (v. v. DC.)
33. *C. CALCITRAPOIDES* (L. sp. 1297.) involucri glabri squamis exterioribus ovato-rotundis apice spinosis spinâ mediâ longissimâ, intimis spatulatis scariosis integris, foliis lineari-oblongis integris vel subdentatis, caule altissimo, seminibus papposis. ① in Delph., Galloprov. et circâ Nemausim. — Fl. fr. 4. p. 100. (v. s.)

## Sect. VI. CNICUS DC. l. c. non L.

Involucrum squamis apice spinosis, involucreto foliolis magnis, apice spinosis, pappus duplex, semina apice margine crenato onusta. — An genus distinctum?

34. *C. BENEDICTA* (L. sp. 1296.) involucri squamis lanceolatis linearibusve sublanatis apice duplicato-spinosis, foliis semi-decurrentibus dentato-spinosis. ① in cultis circâ Montem-Albanum ex Monspelium ex Benth. Gat. Aquas-Sextias ex Ger. et Massiliam (Solier). — Fl. fr. 4. p. 100. — Gærtn. fruct. t. 162. f. 5. (v. s.)

## LVII. KENTROPHYLLUM Neck. DC. mem. comp. p. 24.

Involucrum ventricosum imbricatum squamis interioribus cartilagineis apice ciliato-spinosis, exterioribus foliaceis pinnatifidis bracteas



æmulantibus; flosculi disci hermaphroditi, radii neutri majores; semina tetragona hylo laterali receptaculo adfixa; pappus paleaceo-pilosus.

1. *K. LANATUM* (DC. mss.) foliis inferioribus pinnatifidis dentatis, summis amplexicaulibus pinnatifido-dentatis spinosis, caule lanuginoso. ① in sterilibus ad vias. — *Centaurea lanata* Fl. fr. p. 4. p. 102. — Lob. ic. 2. t. 13. f. 1. (v. v.)

LVIII. *CARLINA* Tourn. t. 285. L. gen. 410. DC. mem. comp. 23.

Involucrum squamis exterioribus sinuato-spinosis basi conniventibus apice divaricatis, interioribus sæpè simplicibus acutis, intimis multò longioribus scariosis ligulæformibus coloratis et radium mentientibus; receptaculi paleæ in lacinias setiformes fissæ; pappus plumosus.

1. *C. ACANTHIFOLIA* (All. ped. t. 51.) acaulis, foliis longè petiolatis pinnatifidis tomentosis, laciniiis dentato-angulosis spinosis, involucri squamis spinosis spinis ramosis. ② in collibus et submontanis siccis Delphin., Galloprov., Occitanie superiori, Arvernæ. — Fl. fr. 4. p. 123 excl. var. (v. s.)
2. *C. CYNARA* (Pouret ex DC. fl. fr. l. c.) acaulis, foliis pinnatifidis sessilibus subtus tomentosissimè subglabris, involucri squamis spinosis spinis simplicibus. ② in Pyrenæis. — *C. acanthifolia* ß Fl. fr. 4. p. 123. Radii intensius flavi quàm in specie præcedente. (v. v. DC.)
3. *C. CHAMELSON* (Vill. dauph. 3. p. 31.) caule simplici unifloro, foliis pinnatifidis glaberrimis, laciniiis inciso-dentatis spinosis, involucri squamis spinosis spinis ramosis. ② in pascuis submontanis siccis et sterilibus. — *C. acaulis* L. sp. 1160. *C. subacaulis* DC. fl. fr. 4. p. 122.  
*a. acaulis*, Lob. ic. 2. t. 4. f. 1. (v. v.)  
*ß. caulescens*, Lob. ic. 2. t. 4. f. 2. (v. v.)
4. *C. LANATA* (L. sp. 1160.) caule uni-aut multifloro lanato, foliis lanceolatis sinuato-spinosis, involucri radio intimo purpurascente. ① in siccis Gallie mediterraneæ. — Fl. fr. 4. p. 124. *Mitina lanata* Cass. — Garid. aix. p. 86. t. 21. (v. v.)
5. *C. VULGARIS* (L. sp. 1161.) caule uni-aut multifloro corymboso pubescenti-arachnoideo, foliis lanceolatis dentatis spinosis subtus pubescentibus, involucri squamis radiantibus albidis. ② in siccis et petrosis. — Fl. fr. 4. p. 124. Gærtn. fruct. 2. t. 163. f. 1. (v. v.)
6. *C. CORYMBOSA* (L. sp. 1160.) caule multifloro corymboso glabriusculo, foliis lanceolatis sinuato-spinosis glabris, involucri squamis radiantibus luteis. ① in sterilibus Gallie australis. — Fl. fr. 4. p. 124; 5. p. 467. *Mitina corymbosa* Cass. — Col. ceph. t. 27. — Barr. ic. t. 594. (v. v.)

LIX. *STÆHELINA* L. gen. 938. DC. mem. comp. p. 37.

Involucrum cylindricum imbricatum squamis inermibus sæpius apice coloratis; flosculi omnes hermaphroditi; antheræ basi caudatæ; stylus bifidus; receptaculum paleaceum; pappus ramosus.

1. *S. DUMA* (L. sp. 1176.) foliis sessilibus linearibus denticulatis subtus tomentosissimis, involucri squamis interioribus elongatis apice coloratis. ③ in sterilibus et siccis provinciarum australium usque ad Aginum (St.-Am.). — Fl. fr. 4. p. 107. — Ger. galloprov. t. 6. — Barr. ic. t. 406. (v. v.)
2. *S. ARBORESCENS* (L. syst. veg. 615.) foliis petiolatis ellipticis obtusis integerrimis subtus sericeo-tomentosis. ③ in insulis Stæchadum ex Ger. — Fl. fr. 4. p. 107. — Alp. ex. 32. ic. (v. s. c.)

LX. *ATRACTYLIS* L. gen. 930. DC. mem. com. p. 20.

Involucrum ovatum connivens squamis integris acuminatis, cinctum

foliis longioribus pinnatifido-spinosis; flosculi omnes hermaphroditi, sæpius omnes tubulosi, interdum marginales ligulæformes; receptaculi paleæ in lacinias setiformes fissæ; pappus plumosus.

1. *A. CANCELLATA* (L. sp. 1162. excl. syn. Alp. et Moris.) caule tomentoso, foliis arachnoideo-tomentosis amplexicaulibus lanceolatis ciliato-spinosis, floribus bracteatis, bracteis erectis pinnatifido-spinosis. ① in sterilibus petrosis apricis Monspelii, Niceæ. — Fl. fr. 4. p. 125. — Lam. ill. t. 662. f. 1. (v. v.)
2. *A. HUMILIS* (L. sp. 1162.) caule glabro, foliis sinuatis oblongo-lanceolatis dentato-spinosis spinis simplicibus vel geminatis, flore terminali, involucri squamis mucronatis glabris. ② in saxosis circa Narbonam. — Fl. fr. 4. p. 126. Lam. ill. t. 660. (v. s.)

LXI. *XERANTHEMUM* Gærtn. 2. p. 399. t. 165. DC. mem. comp. p. 21.

Involucrum imbricatum squamis inermibus scariosis, interioribus longioribus coloratis radium æmulantibus; flosculi disci hermaphroditi fertiles, radii feminei steriles pauci; receptaculum paleaceum; semina disci pappo 5-aristato coronata, radii nuda.

1. *X. ANNUUM* (L. sp. 1201. v. s.) involucri squamis externis scariosis glabris, interioribus lanceolatis obtusis patentibus. ① in arvis et collibus siccis Gallie medice australis occidentalisque. — Fl. fr. 4. p. 130. — Clus. hist. 2. p. 11. f. 2. (v. v.)
2. *X. IMPERITUM* (W. sp. 3. p. 1902.) involucri squamis subacutis margine scariosis dorso tomentosis, interioribus lanceolatis acutis conniventibus. ① in arvis collibusque aridis Gallie medice, australis occidentalisque. — Fl. fr. 4. p. 130. — Moris. s. 6. t. 12. n. 1. (v. v.)

Subordo III. *CICHORACEÆ* Juss. gen. 168. — *Lactucæ* Cass. dict. sc. nat. 20. p. 355.

Flosculi omnes ligulati et hermaphroditi; receptaculum vix carnosum. — Folia alterna; flores lutei rarius cærulei sæpè meteorici. Succus proprius in plurimis lacteus.

† **FRUCTUS COMPRESSUS AUT TETRAGONUS; PAPPUS ALBUS SQUAMOSO-PILOSUS.**

LXII. *SCOLYMUS* Tourn. t. 273. L. gen. 922.

Involucrum imbricatum spinosum; receptaculum paleaceum; pappus nullus aut squamoso-pilosus.

1. *S. GRANDIFLORUS* (Desf. atl. 2. t. 218.) caule subsimplici villosa erecto, foliis decurrentibus villosis pinnatifidis spinoso-dentatis, inferioribus petiolatis, capitalis solitarii sessilibus, involucri squamis exterioribus obtusis, interioribus acutis spinulosis. ② in agro Ruscinonensi circa Collioure et Port-Vendres ex Lapeyr. Arnott et Benth. (v. s. ex Barbaria.)
2. *S. MACULATUS* (L. sp. 1142.) caule ramoso, foliis decurrentibus glabris margine cartilagineis sinuato-pinnatifidis spinosis, foliis involucralibus pectinato-multifidis, invol. squamis omnibus acuminato-spinosis, seminibus nudis. ① ad marginem agrorum in Gallia australi. — Fl. fr. 4. p. 69. — Clus. hist. 2. p. 153. f. 1. (v. v.)
3. *S. HISPANICUS* (L. sp. 1143.) caule apice ramoso, foliis decurrentibus scabris margine non cartilagineis sinuato-pinnatifidis spinosis, inferioribus petiolatis, foliis involucralibus foliaceis dentato-spinosis, invol. squamis omnibus acuminato-spinulosis, seminibus paleis involutis. ② ad margines agrorum in Gallia à Nicea ad Nannetes. — Fl. fr. 4. p. 69. — Clus. hist. 2. p. 153. f. 2. (v. v.)

LXIII. UROSPERMUM *Juss. gen. 170.*

Involucrum simplex 8-phyllum squamis coalitis; receptaculum nudum; semina transversè sulcata; pappus plumosus stipitatus stipite cavo conico incurvo basi incrassato.

1. *U. DALSCHAMPRI* (Desf. cat. go.) involucris pubescentibus inermibus, foliis runcinato-dentatis.  $\mathcal{L}$  secus vineas et prata provinciarum austral. — Fl. fr. 4. p. 62. — Dalech. ligd. 569. f. 1. (v. v.)
2. *U. FICOIDES* (Desf. cat. go.) involucris hispido-aculeatis, foliis runcinatis-dentatis, caulibus auriculato-sagittatis. (1) ad margines viarum in Gallia australi. — Fl. fr. 4. p. 63. — C. Bauh. prod. p. 60. f. 2. (v. v.)  
 $\beta$ . *asperum*, caule subunifloro, foliis superioribus subintegris. — *U. asperum*. Fl. fr. 4. p. 63. (v. s.)

LXIV. PICRIDIMUM *Desf. adl. 2. p. 221. DC. fl. fr. 4. p. 15.*

Involucrum imbricatum basi tumidum squamis margine membranaceis; receptaculum nudum; semina tetragona subincurvata transversè tuberculata; pappus sessilis pilosus.

1. *P. VULGARE* (Desf. l. c.) pedunculis squamosis incrassatis, squamis cordiformibus, involucri squamis margine glabris. (1) in provinciis mediterraneis. — Fl. fr. 4. p. 16. *Sonchus picroides* Lam. — Lob. ic. t. 236. f. 2. (v. v.)
2. *P. ALBUM* (DC. fl. fr. 4. p. 16.) pedunculis subnudis, involucri squamis puberulis margine ciliatis.  $\mathcal{L}$  in montosis provinciarum australium. — *Crepis albida* Vill. dauph. 3. t. 33. *Paleyia albida* Cass. (v. v.)

LXV. SONCHUS *L. gen. 908.*

Involucrum oblongum imbricatum basi ovatum; receptaculum nudum; semina longitudinaliter striata; pappus brevis sessilis pilosus.

§. 1. *Floribus cæruleis.*

1. *S. ALPINUS* (Fl. dan. t. 182.) pedunculis involucrisque hirsutis, floribus racemosis bracteatis, foliis runcinatis basi sagittatis glabris subtus glaucis.  $\mathcal{L}$  in sylvis umbrosis editiorum montium. — Fl. fr. 4. p. 14. — Cam. epit. 281. ic. (v. v.)
2. *S. PLUMIERI* (L. sp. 1117.) pedunculis bracteis involucrisque glabris, floribus paniculato-corymbosis, foliis runcinatis amplissimis.  $\mathcal{L}$  in umbrosis et saxosis editioribus Vogesorum, Alpium, Pyren., Arvernæ. — Fl. fr. 4. p. 15. (v. v. in Helv.)

§. 2. *Floribus luteis.*

3. *S. PALUSTRIS* (L. sp. 1116.) pedunculis involucrisque subumbellatis glanduloso-hispidis, foliis basi sagittatis lobis acutis, inferioribus runcinatis.  $\mathcal{L}$  in stagnorum et fossarum oris. — Fl. fr. 4. p. 14. — Pet. angl. t. 14. f. 7. (v. v.)
4. *S. ARVENSIS* (L. sp. 1116.) pedunculis involucrisque hispidis subumbellatis, foliis runcinatis basi oordatis dentato-ciliatis.  $\mathcal{L}$  in arvis. — Fl. fr. 4. p. 14. — Lob. ic. t. 237. f. 1. *Nimia* cum sequente affinitas. (v. v.)
5. *S. OLERACEUS* (L. sp. 1116.) pedunculis sub capitulo subtomentosis umbellatis, involucris glabris, foliis oblongo-lanceolatis amplexicaulibus margine ciliatis subinnatis, inferioribus pinnatifidis. (1) — Fl. fr. 4. p. 13. *Lepicaune spinulosa* Lapeyr. abr. 480 ex cl. Arnott.  
 $\alpha$ . *lævis*, in cultis et hortis. — *S. lævis* Vill. — Cam. ep. 279. ic. (v. v.)  
 $\beta$ . *asper*, in sterilibus. — *S. asper* Vill. Pluck. t. 61. f. 5. (v. v.)
6. *S. PECTINATUS* (DC. rapp. 2. p. 78.) pedunculis ramosis pilos glandulosos hinc inde gerentibus, foliis pectinato-pinnatifidis, inferiorum laciniis sub

æqualibus integris aut basi appendiculatis, superiorum laciniiis æqualibus linearibus integerrimis, involucri hispidis. ½ in rupestribus circà Collioure. — Fl. fr. 5. p. 434. (v. v.)

7. *S. TENERRIMUS* (L. sp. 1117.) pedunculis subhispidis umbellatis, foliis inferioribus subbipinnatifidis laciniiis inæqualibus, superioribus pinnatifidis laciniiis lanceolatis serratis basi inciso-dentatis, involucri basi tomentosis. ① in maritimis Gallie mediterraneæ. — Fl. fr. 4. p. 13. — Pluck. t. 93. f. 3. (v. v.)
8. *S. MARITIMUS* (L. sp. 1116.) pedunculis subsolitariis terminalibus nudis, foliis lanceolatis amplexicaulibus indivisis retrorsum argutè dentatis. ½ in maritimis à Nicæa ad Rapellam et in stagnis subæalis Delphinatidæ. — Fl. fr. 4. p. 12. All. ped. t. 16. f. 2. (v. v.)

#### LXVI. LACTUCA Tourn. t. 267. L. gen. 909.

Involucrum oblongum imbricatum squamis margine membranaceis; receptaculum nudum; pappus stipitatus pilosus mollis fugax.

##### §. 1. *Floribus luteis.*

1. *L. SATIVA* (L. sp. 1118.) foliis rotundatis caulinis cordatis, caule corymboso. ① patria ignota, in hortis culta. — Fl. fr. 4. p. 9. — Moris. s. 7. t. 2. n. 2. (v. v.)
2. *L. LACINIATA* (Roth. cat. 1. p. 90.) foliis obtusis glabris, inferioribus pinnatifidis laciniatis, superioribus runcinatis, laciniiis infimis stipularibus elongatis. ① in Cenomaniâ culta. — Fl. fr. 5. p. 433. — Lob. ic. 242. f. 1. (v. v. c.)
3. *L. SYLVESTRIS* (Lam. dict. 3. p. 406.) foliis sinuato-pinnatifidis amplexicaulibus verticalibus carinâ aculeatis apice acutis, floribus laxè paniculatis, caule aculeato. ① ad margines viarum. — Fl. fr. 4. p. 10. Camer. epit. 300. ic. (v. v.)
4. *L. VIROSA* (L. sp. 1119.) foliis oblongis denticulatis amplexicaulibus horizontalibus carinâ subaculeatis, caule inermi. ② ad asepes, muros et agros. — Fl. fr. 4. p. 10. — Moris. s. 7. t. 2. n. 16. Præcedentis var? (v. s.)
5. *L. SALIGNA* (L. sp. 1119.) foliis amplexicaulibus carinâ aculeatis glabrisve, radicalibus pinnatifidis laciniiis linearibus acuminate recurvis, caulinis linearibus integerrimis. ① ad margines agrorum et vinearum. — Fl. fr. 4. p. 11. — Barr. ic. t. 136. (v. v.)

##### §. 2. *Floribus cæruleis.*

6. *L. TENERRIMA* (Pourr. act. toul. 3. p. 322.) caule ramoso divaricato, foliis radicalibus pinnatifidis laciniiis integris aut basi dentatis, caulinis linearibus integerrimis sagittatis, involucri squamis exterioribus linearibus, seminibus ovatis planis. ½ in agro Narbonensi et Ruscinonensi circà Olette et Villefranche. — Fl. fr. 4. p. 11. (v. v.)
7. *L. PERENNIS* (L. sp. 1120.) foliis pinnatifidis laciniiis linearibus sursum dentatis, summis linearibus sagittatis, involucri squamis exterioribus lanceolato-ovatis, seminibus lineari-lanceolatis utrinquè attenuatis sulcatis. ½ ad vineas, in agris lapideis et saxosis apricis. — Fl. fr. 4. p. 11. — Dalech. hist. 566. f. 2. (v. v.)
8. *L. CICHORIIFOLIA* (DC. fl. fr. 5. p. 434.) foliis radicalibus oblongis pectinato-dentatis, dentibus longis subretrois, floralibus integerrimis linearibus sagittatis, floribus corymbosis, pedicellis divaricatis. ½ in Pyren. orientilibus. (v. s.)

#### LXVII. CHONDRILLA Gærtn. fruct. 2. p. 362. t. 158. DC. fl. fr. 4. p. 8.

Involucrum squamis erectis linearibus, calyculatum squamis minimis aut nullis; receptaculum nudum; pappus stipitatus pilosus.

1. *C. MURALIS* (Lam. dict. 2. p. 78.) foliis lyrato-pinnatifidis dentatis, lobo terminali 5-angulari, involucri squamis 5-6, pappi stipite semine brevior. (1) ad vetustos muros. — Fl. fr. 4. p. 8. *Prenanthes muralis* L. *mycelis angulosa* Cass. — Clus. hist. 2. p. 146. f. 2. (v. v.)
2. *C. JUNCEA* (L. sp. 1120.) foliis radicalibus runcinatis, caulinis linearibus integris, involucri squamis 7-10, pappi stipite semine longiore. 2/ ad margines arborum. — Fl. fr. 4. p. 8. — Clus. hist. 2. p. 144. f. 2. (v. v.)

LXVIII. *PRENANTHES* Vaill. L. gen. 911.

Involucrum squamis margine imbricatis, calyculatum squamis inæqualibus ovatis; receptaculum nudum; pappus sessilis pilosus simplici serie; flosculi pauci.

§. 1. *Floribus purpurascensibus.*

1. *P. PURPUREA* (L. sp. 1121.) foliis oblongo-lanceolatis amplexicaulibus cordatis denticulatis subtus glaucis, caule ramoso. (1) in sylvis lapideis montosis. — Fl. fr. 4. p. 6. Gært. fruct. 2. t. 57. f. 1. (v. v.)
2. *P. TENUIFOLIA* (L. sp. 1120.) foliis linearibus integerrimis amplexicaulibus, caule simplici. 2/ in sylvis Delphinatûs, Cebennarum. — Fl. fr. 4. p. 6. All. ped. t. 33. f. 2. (v. s.)

§. 2. *Floribus luteis.*

3. *P. VIMINEA* (L. sp. 1120.) foliis decurrentibus, inferioribus pinnatifidis extrorsum dentatis, summis linearibus, caule ramoso divaricato viscoso, involucri squamis 5-6 lanceolatis. 2/ in lapideis provinciarum austral., occidentaliæque. — Fl. fr. 4. p. 6. All. ped. t. 52. f. 2. *Phanixopas decurrens* Cass. (v. v.)
4. *P. PULCHRA* (DC. fl. fr. 4. p. 7.) foliis subscabris, radicalibus runcinatis, caulinis amplexicaulibus ovato-lanceolatis basi sagittatis, paniculâ corymbosâ divaricatâ nudâ, involucri squamis 15-20 lineari-lanceolatis, calyculi ovatis. (1) in cultis et vineis Galliæ mediæ, occidentalis australisque. — *Crepis pulchra* L. *P. hieracifolia* Wild. *Phæcasium lampsanoides* Cass. — Moris. s. 7. t. 5. n. 13 et 37. (v. v.)
5. *P. BULBOSA* (DC. fl. fr. 4. p. 7.) foliis lanceolato-oblongis subdentatis glabris, scapo unifloro supernè hirsuto, involucri glabri squamis 13-20 lanceolatis. 2/ in maritimis Galloprovinciæ, Occitaniz, Armoraciæ. — *Leontodon bulbosum* L. *Hieracium bulbosum* Wild. *Ætheorhiza bulbosa* Cass. — Lob. ic. t. 230. f. 2. (v. v.)

† † FRUCTUS ELONGATUS AD APICEM PLUS MINUS ATTENUATUS; PAPPUS ALBUS AUT NULLUS. — INVOLUCRUM UNISERIATUM.

## I PAPPUS NULLUS.

LXIX. *LAMPSANA* Tourn. t. 272. DC. fl. fr. 4. p. 3.

Involucrum squamis lineari-lanceolatis, calyculatum; receptaculum nudum; semina pappo destituta caduca involucri foliolis minimè cincta.

1. *L. VETIDA* (Scop. carn. ed. 2. n. 989.) scapo unifloro, foliis runcinato-pinnatifidis, lacinia terminali rhomboideâ. 2/ in Pyrenæis ex Lapeyr. — Fl. fr. 4. p. 3. — Mich. gen. t. 28. (v. s. ex Helv.)
2. *L. COMMUNIS* (Lam. dict. 3. p. 414.) caule erecto striato ramoso, foliis ovatis petiolatis angulato-dentatis. (1) in cultis. — Fl. fr. 4. p. 4. — Lob. ic. t. 209. f. 1. (v. v.)
3. *L. MINIMA* (Lam. dict. 3. p. 414.) scapo 1-3-floro, pedunculis fistulosis clavato-incrassatis, foliis ovato-oblongis denticulatis. (1) in pascuis siccis et arenosis passim. — Fl. fr. 4. p. 3. *Arnoseris pusilla* Gært. 2. t. 157. f. 3. — Clus. hist. 2. p. 142. f. 2. (v. v.)

LXX. RHAGADIOLUS *Tourn. t. 272. DC. fl. fr. 4. p. 4.*

Involucrum calyculatum, squamis linearibus intus concavis, internis semina matura cingentibus; receptaculum nudum; semina non papposa sæpius inflexa vel contorta, matura non caduca.

1. *R. nudus* (Gærtn. fruct. 2. p. 354.) seminibus lævibus patentibus, foliis lyratis lacinia terminali rotundo-ovata maximâ. ① in Galloprovincia, circa Nemausum et Monspelium. — Fl. fr. 4. p. 5. (v. s.)
2. *R. stellatus* (Gærtn. fruct. 2. t. 157. f. 2.) seminibus lævibus patentibus, foliis caulinis lanceolatis subdentatis. ① in arvis Gallie australioris. — Fl. fr. 4. p. 4. — Lob. ic. t. 249. f. 2. (v. v.)

## II PAPPUS PILOSUS.

LXXI. ZACINTHA *Gærtn. fruct. 2. p. 358. DC. fl. fr. 4. p. 48.*

Involucrum 8-phyllum basi calyculatum demum coriaceum sulcatum; receptaculum nudum, pappus pilosus subdentatus sessilis brevissimus.

1. *Z. verrucosa* (Gærtn. fruct. 2. t. 157. f. 7.) ① in sterilibus Corsicæ, Niceæ, Galloprovinciæ. — Fl. fr. 4. p. 48. — Moris. s. 7. t. 1. n. 4. (v. v.)

LXXII. BARKHAUSIA *Mench. meth. 537. DC. fl. fr. 4. p. 41. — Hostia, Barkhausia, Villemetia Cass.*

Involucrum calyculatum, maturum sulcato-costatum, squamis exterioribus laxis; receptaculum nudum; pappus pilosus stipitatus.

1. *B. apargioides* (Duby mss.) scapo subnudo paucifloro involucrique laxiusculo pilis nigris hispidis, foliis radicalibus lanceolatis dentatis glabris. ② in Pyrenæis. — *Crepis apargioides* W. sp. 3. p. 1594. Villemetia apargioides Cass. Semina longè stipitata. (v. s.)
2. *B. alpina* (DC. l. c.) foliis ovatis cordato-sagittatis amplexicaulibus denticulatis, pedunculis elongatis unifloris, involucri hispidis, squamis exterioribus membranaceis scariosis. ① in pratis Alpium Galloprovinciæ ex Ger. — *Crepis Alpina* L. sp. 1134. Gærtn. fruct. 2. t. 158. f. 8. (v. s. c.)
3. *B. bellidifolia* (DC. fl. fr. 5. p. 449.) glabra subcarnosa multicaulis, foliis subintegris, radicalibus oblongo-spathulatis, caulinis oblongis basi auriculatis. ① in Corsicæ locis incultis. — *Crepis bellidifolia* Lois. gall. t. 18. (v. s.)
4. *B. suffreniana* (DC. cat. monsp. 83.) caule erecto subsimplici basi hispidis, foliis radicalibus oblongis basi angustatis sinuatis, caulinis paucis linearibus integris, pedicellis unifloris, involucri hispidulis, capitulis antè anthesin cernuis. ① in aridis lapideis Galloprovinciæ. (v. s.)
5. *B. setosa* (DC. fl. fr. 4. p. 44.) setis in caule pedunculis involucri nervis et oris foliorum sparsis, foliis inferioribus pinnatifido-dentatis lacinia terminali maximâ, caulinis dentatis, summis linearibus, involucri squamis exterioribus patulis. ② in Corsicâ et circâ Avenionem. — Ic. rar. 1. t. 19. (v. s.)
6. *B. rupea* (Mench. meth. 537.) foliis radicalibus runcinato-lyratis lacinia anguloso-dentatis, caulinis amplexicaulibus lanceolatis, inferioribus pinnatifidis, involucri hispidis, squamis exterioribus scariosis. ① secus vias Niceæ, Monspelii, Aginni ex St.-Amans. — Fl. fr. 4. p. 42. — C. Bauh. prod. 68. f. 2. Flores rubelli. (v. s.)
7. *B. pectida* (DC. fl. fr. 4. p. 42.) foliis acuminatis runcinato-pinnatifidis scabris sessilibus, summis lanceolatis basi profundè incisis, involucri ovatis pilosis angulatis. ① in incultis. — Lob. ic. t. 226. f. 1. (v. v.)

8. *B. TARAXACIFOLIA* (DC. fl. fr. 4. p. 43.) foliis scabris, radicalibus lyrato-runcinatis obtusis, caulinis lanceolatis amplexicaulis basi dentatis, involucri subtomentosis, squamis exterioribus margine submembranaceis. (2) in pratis lapidosis. — *Crepis biennis* Lapeyr. abr. p. 84. ex cl. Arnoit. — Lob. ic. t. 239. f. 2. (v. v.)

β. *præcox*, foliis superioribus basi dilatato-auriculatis. — *Crepis præcox* Balb. Misc. t. 9. (v. s.)

γ. *intybacea*, foliis superioribus basi auriculatis, involucrio glabro. — In Occitania. — *B. intybacea* DC. cat. 82. fl. fr. 5. p. 449. (v. s.)

9. *B? LEONTODON* (DC. fl. fr. 4. p. 43.) foliis glabris runcinatis dentatis, radicalibus obtusis aut brevissimè mucronatis, caulinis acuminatis, scapo nudo multifloro ascendente, involucri subtomentosis, squamis exterioribus adpressis. (2) in Corsicâ (Ph. Thomas). — Seminibus immaturi pappus sessilibus, maturi subsessilibus. An hujus generis? (v. s.)

LXXIII. *PTEROTHECA Cassini* in bull. phil. 1821. p. 125. — *Genus novum?* DC. syn. gall. 261.

Involucrum subuniseriatum, squamis adpressis margine membranaceis; receptaculum planum, paleis longissimis filiformibus instructum; semina marginalia plerumque nuda, externâ facie striata internâ 3-5-alata, interna papposa striata apice in collum attenuata; pappus albus pilosus.

1. *P. NEMAUSENSIS* (Cass. l. c.) foliis runcinato-lyratis dentatis obtusis brevissimè mucronatis, scapo nudo multifloro, involucri squamis exterioribus lanceolatis. (1) in Galliâ mediterraneâ. — *Andryala Nemausensis* Vill. dauph. 3. t. 26. Fl. fr. 4. p. 38. *Trichocrepis bifida* Visiani dalm. (v. v.)

LXXIV. *CREPIS Manch. meth.* p. 534. DC. fl. fr. 4. p. 38.

Involucrum calyculatum laxum, fructigerum sæpè costatum; receptaculum nudum; pappus pilosus sessilibus niveus.

1. *C. DIFFUSA* (DC. cat. monsp. 98.) foliis glabris subradicalibus remotè dentatis sinuatis runcinatisve, superioribus subintegris subsagittatis, caule basi ramoso diffuso multifloro, involucri pubescentibus, seminum costis lævibus. (1) in agris, ruderatis. — Fl. fr. 5. p. 448. *C. virens* α. et γ. Fl. fr. 4. p. 39. Sequenti nimium affinis. (v. v.)
2. *C. STRICTA* (DC. cat. monsp. 99.) foliis glabris, radicalibus runcinato-pinnatifidis, caulinis paucissimis, inferioribus basi dentato-pinnatifidis, superioribus subintegris, caule erecto, corymbo laxo, involucri pubescentibus, seminum costis lævibus. (1) in Galliâ mediâ, australi et orientali passim. — Fl. fr. 5. p. 447. Scop. carn. t. 47. (v. v.)
3. *C. VIRENS* (Vill. dauph. 3. p. 142.) foliis glabris lanceolato-runcinatis, superioribus basi dentatis sagittatis, floribus corymboso-paniculatis, involucri pubescentibus, seminum costis lævibus. (1) in pascuis ruderatisque. — Fl. fr. 4. p. 39. non L. (v. v.)
4. *C. SCABRA* (W. sp. 3. p. 1603.) foliis hispidis runcinatis, caulinis sessilibus basi dentato-auriculatis, ramis floralibus paniculato-corymbosis lævibus, caule suberecto striato, involucri squamis carinato-canaliculatis pubescentibus, seminum costis lævibus. (1) Nicæ, Narbonæ, in agro Andegaviensi. — Fl. fr. 5. p. 446. (v. s.)
5. *C. BIENNIS* (L. sp. 1136.) foliis hispidis runcinato-pinnatifidis, summis sessilibus lanceolatis dentatis, ramis floralibus paniculato-corymbosis scabris, caule elato recto sulcato, seminum costis lævibus. (2) in pratis. — Fl. fr. 4. p. 39. 5. p. 446. eng. bot. t. 149. (v. v.)

6. *C. DROSCORIDIS* (L. sp. 1133.) foliis radicalibus lyrato-runcinatis, caulinis hastatis lanceolatis, inferioribus dentatis, caule erecto, involucri tomentosis, fructigeris subrotundo-ovatis angulosis. ① in siccis secus vias Alsatiæ. — Fl. fr. 4. p. 40. Ic. rar. t. 18. *Gatyna globulifera* Cass. (v. s.)
7. *C. TECTORUM* (L. sp. 1135. non fl. fr.) foliis glabris sinuato-pinnatifidis, superioribus sagittato-linearibus integris margine subrevolutis, seminum oblongo-attenuatorum costis acabris apice muricatis. ① in campestribus Alsatiæ et probabiliter Galliæ totius. — Fl. fr. 5. p. 448. (v. s.)
8. *C. LACHENALII* (Gochm. diss. t. 3.) foliis glabris, radicalibus spathulato-lanceolatis dentatis, superioribus linearibus margine revolutis integerrimis, caule gracili erecto paucifloro, seminum costis aspere-muricatis. ① in arvis Alsatiæ. — Fl. fr. 5. p. 449. (v. s.)

## LXXV. TARAXACUM Hall. helv. p. 23. Juss. gen. 169.

Involucrum duplex squamis exterioribus brevibus sæpius patulis; receptaculum nudum; pappus stipitatus pilosus. — Scapi uniflori.

1. *T. PALUSTRE* (DC. fl. fr. 4. p. 45.) foliis lanceolatis sinuato-dentatis glabrinaculis, involucri exterioris adpressi squamis ovatis subacuminatis. ② in paludosis et pratis humidis. — *Hedypnois paludosa* Scop. carn. 2. t. 48. (v. v.)
2. *T. OBOVATUM* (DC. rapp. voy. 2. p. 83.) foliis obovatis obtusis vix retro-dentatis, involucri exterioris patuli squamis lanceolato-ovatis. ② Monsipelli et in Galloprovinciâ. — *Leontodon obovatus* W. hort. berol. t. 47. — J. Bauh. hist. 2. p. 1037. f. 2. (v. v. DC.)
3. *T. LEVIGATUM* (DC. cat. monsp. 49.) foliis runcinato-pinnatifidis, laciniis valde inæqualibus, majoribus reflexis basi subincisis, involucri exterioris patuli squamis lanceolatis. ② in rupestribus apricis passim. — Fl. fr. 5. p. 450. — Barr. ic. t. 237. An à sequente satis distinctum. (v. v.)
4. *T. DENS-LEONIS* (Desf. atl. 2. p. 228.) foliis runcinatis glabris, laciniis lanceolatis dentatis, involucri exterioris reflexi squamis lineari-lanceolatis. ② ubique in pratis. — Fl. fr. 4. p. 45. *Leontodon Taraxacum* L. Bull. herb. t. 217. — Moris. s. 7. t. 8. n. 1. Tres præcedentes species ut hujusce varietates habent cl. St.-Amans et Benth. (v. v.)

## III PAPPUS PLUMOSUS.

## LXXXVI. HELMINTHIA Juss. gen. 170.

Involucrum duplex, interius squamis 8 æqualibus, exterius squamis 5 foliaceis laxis; semina transversim striata; pappus plumosus pedicellatus.

1. *H. XCHIMONES* (Gærtn. fruct. 2. t. 159. f. 2.) involucri exterioris squamis latis ovato-cordatis subspinosis. ① in arvis, ad vias. — Fl. fr. 4. p. 58. — Lob. ic. t. 577. f. 2. (v. v.)
- β. mollis*, involucri squamis subinermibus. — In pratis salis circa Arelatem detexit cl. A. Richard. (v. s.)
2. *H. SPINOSA* (DC. fl. fr. 4. p. 58.) involucri exterioris squamis lanceolatis margine apiceque spinosis. — In Pyrenæis ex herb. Lemonnier. Ic. rar. t. 21. (v. s.)

## LXXXVII. PICRIS Juss. gen. 170.

Involucrum calyculatum, calyculi squamis lineari-lanceolatis; receptaculum nudum; semina transversè striata, pappus plumosus sessilis vel subsessilis.

1. *P. HIERACIOIDES* (L. sp. 1115.) caule scabro ramoso-divaricato, foliis aspersi-



mis lanceolatis dentatis, floribus corymbosis, seminibus lanceolatis striatis.  $\mathcal{L}$  in agris. — Fl. fr. 4. p. 57. Lam. ill. t. 648. f. 2. (v. v.)

2. *P. PYRENAICA* (L. sp. ed. 1. p. 792.) caule scabro ramoso-divaricato, pedunculis 1-floris, foliis asperis lanceolatis dentibus patentibus.  $\mathcal{L}$  in Pyrenæis. Fl. fr. 5. p. 454. *P. tuberosa* Lapeyr. Radix napiformis (Lap.). An satis à præcedente distincta? (v. s. imperf.)
3. *P. PAUCIFLORA* (W. sp. 3. p. 1557.) caule scabro subramoso, foliis asperissimis lanceolatis denticulatis, pedunculis elongatis incrassatis, seminibus lineari-elongatis sulcato-echinatis apice attenuatis. ① in arvis Gallie australis. — Fl. fr. 4. p. 57. Gærtn. 2. t. 159. f. 2. (v. v.)

††† FRUCTUS BREVIS, AD APICEM ATTENUATUS BASI TRUNCATUS;  
PAPPUS NULLUS AUT PILOSUS.

#### LXXVIII. HIERACIUM L. gen. 913.

Involucrum imbricatum, receptaculum nudum aut rarius pilis sparsis semine brevioribus onustum; pappus sessilis pilosus sæpius sordidè rufescens.

##### §. 1. *Leontodontoidea* DC. syn. gall. 258.

Foliis radicalibus subglabris nec glaucis nec coriaceis; scapis nudis uni aut paucifloris.

1. *H. AUREUM* (Vill. dauph. 3. t. 33.) scapo 1-floro subnudo, involucri hispido, foliis lanceolato-spathulatis runcinato-dentatis glabrisculis.  $\mathcal{L}$  in pascuis Alpium, Pyrenæorum. — Fl. fr. 4. p. 17. (v. v.)
2. *H. PRÆMORSUM* (L. sp. 1126.) scapo nudo racemoso, floribus superioribus primoribus, foliis ovatis vix denticulatis puberulis, radice præmorsâ.  $\mathcal{L}$  in pascuis Alsatie, Lotharingie. — Fl. fr. 4. p. 18. 5. p. 434. — Gm. sib. 2. t. 13. f. 2. (v. s.)
3. *H. AURANTIACUM* (L. sp. 1126.) scapo basi folioso hispido, floribus corymbosis, pedunculis glomeratis, foliis oblongis scutinculis piloso-hispidis.  $\mathcal{L}$  in Alpibus, Jurasso, Vogesis. — Fl. fr. 4. p. 18. Jacq. aust. t. 410. Fl. auran-tiaci. (v. v.)
4. *H. ALPINUM* (L. sp. 1124.) scapo 1-floro subnudo villosa, involucri villosissimo, foliis lanceolatis integerrimis acutis villosis.  $\mathcal{L}$  in pascuis Alpium, montium Arvernæ, Cebennarum, Pyren. ex Lapeyr. — Fl. fr. 4. p. 19. eng. bot. t. 1120. — Moris. s. 7. t. 7. n. 5. (v. v.)  
 $\beta$ . *glabratum* (Ser. mss.) foliis glabris. — H. glabratum W. Fl. fr. 5. p. 435. (v. s.)
5. *H. PUMILUM* (W. sp. 3. p. 1562.) scapo unifloro piloso 2-3-folio, foliis villosis lanceolatis obtusis brevissimè mucronatis, superioribus integris, radicalibus remotè dentatis sessilibus.  $\mathcal{L}$  in Alpibus Delphinatûs (Ph. Thomas). — Fl. fr. 5. p. 435. (v. s.)
6. *H. HALLERI* (Vill. dauph. 3. p. 104.) scapo 1-2-floro subnudo involucri piloso, foliis caulinis linearibus brevissimis, radicalibus petiolatis subvillosis ovato-oblongis basi runcinato-dentatis.  $\mathcal{L}$  in Alpibus Delph. — Fl. fr. 4. p. 19. *H. hybridum* Vill. 3. t. 26. An à præcedente satis distinctum? (v. s.)

##### §. 2. *Andryaloidea* DC. l. c.

Plantæ subobtectæ pilis albis longis mollibus sublentè articulatis subplumosis.

7. *H. SAXATILE* (Vill. dauph. 3. t. 29.) scapo subnudo subunifloro villosa, involucri pubescente aut hirta, foliis obovatis villosa-lanatis integrinerviis breviter mucronatis.  $\mathcal{L}$  in rupibus apricis Pyrenæorum et Alpium Delph. — Fl. fr. 4. p. 22. *H. sericeum* Lapeyr. abr. 477. (v. s.)

8. *H. ANDRYALOIDEUS* (Lam. dict. 2. p. 364.) caule erecto subsimplini tomentoso supernè nudo, foliis tomentosis, inferioribus ovali-oblongis basi runcinatis dentatis, supero lineari integro.  $\mathcal{U}$  in rupibus apricis Delphinatùs. — Fl. fr. 4. p. 22. Vill. dauph. 3. f. 29. (v. v.)
9. *H. LAMATUM* (Vill. dauph. 3. p. 120.) caule erecto subramoso basi folioso lanato, floribus corymbosis longè pedunculatis, foliis ovatis lanatis subintegris, receptaculis subpilosis.  $\mathcal{U}$  in locis apricis montium imprimis australiorum. — Fl. fr. 4. p. 21. — Dill. elth. t. 150. f. 281. (v. s.)

$\beta$ . *uniflorum*. (v. s.)

10. *H. ELONGATUM* (Lapeyr. abr. 476.) caule basi longè barbato erecto subsimpli foliisque laxè villosis, radicalibus oblongis basi attenuatis, caulinis basi cordatis amplexicaulibus, involucris villosis patulis.  $\mathcal{U}$  in asperis Pyrenæorum præsertim orientalium: — Fl. fr. 5. p. 436. (v. s.)
11. *H. VILLOSUM* (L. sp. 1130.) caule erecto subramoso foliisque villosis, radicalibus oblongo-lanceolatis dentatis, caulinis ovatis subamplexicaulibus subdentatis, involucris villosissimis patulis.  $\mathcal{U}$  in montanis. — Fl. fr. 4. p. 20. Jacq. aust. t. 89. (v. v.)
12. *H. FLEXUOSUM* (W. sp. 3. p. 1581.) caule erecto subramoso basi glabro, foliis glaucis inferioribus oblongo-lanceolatis glabris basi attenuatis, superioribus sessilibus subpilosis, involucris villosis.  $\mathcal{U}$  in Alpibus Delphinatùs. — Fl. fr. 5. p. 436. Præcedentis var? (v. s.)

### §. 3. *Pilosellæ* DC. l. c.

Plantæ glaucescentes subcoriaceæ hirtæ aut ciliatæ, pilis sparsis longis albis rigidulis.

13. *H. PILOSELLA* (L. sp. 1125.) scapo unifloro nudo, involucris pilosis, stolonibus repentibus, foliis integerrimis ovatis subtus tomentosis.  $\mathcal{U}$  in locis aridis. — Fl. fr. 4. p. 23.
- $\alpha$ . *vulgaris* (DC.) foliis utrinquè viridibus, involucri breviter pilosi pilis atris. — Cam. epit. t. 708 et 709. (v. v.)
- $\beta$ . *incana* (DC.) foliis subtus incanis, involucri tomentoso.
- $\gamma$ . *Peleteriana* (Ser.) involucri pilosissimo. — H. Peleterianum Merat. Fl. fr. 5. p. 437. (v. s.)
14. *H. BREVISCAPUM* (DC. fl. fr. 5. p. 439.) scapo subnudo 2-4-floro foliis vix longiore basi barbato subvelutino, floribus confertis subsessilibus, involucris albo-villosis, foliis oblongo-linearibus subacutis utrinquè rufescenti-pilosis.  $\mathcal{U}$  in summis Pyrenæis orientalibus. — H. pumilum Lapeyr. non W. Præcedentis forsan varietas? (v. v.)
15. *H. AURICULA* (DC. fl. fr. 4. p. 24.) scapo subnudo 1-30-floro, floribus corymbosis, stolonibus nullis aut reptantibus, foliis integerrimis oblongis subtus glabris.  $\mathcal{U}$  in sterilibus. — H. Lactucella Wallr. sch. crit. p. 408.
- $\alpha$ . *pilosa*, foliorum limbis pilosis. — H. Auricula L. sp. 1126. (v. v.)
- $\beta$ . *dubia*, foliorum limbis subglabris. — H. dubium L. sp. 1125. Sm. eng. fl. 3. p. 357. (v. v.)
- $\gamma$ . *angustifolia*, foliis lineari-lanceolatis. — In Pyrenæis. — H. angustifolium W. Vill. voy. t. 3. f. 4. Fl. fr. 5. p. 438. (v. v. in Helv.)
16. *H. CYMOSUM* (L. sp. 1126.) caule scapiformi oligophyllo foliisque hispidis, inferioribus oblongo-lanceolatis obtusis basi attenuatis, reliquis lineari-lanceolatis acutis sessilibus, floribus umbellato-corymbosis confertis, pedicellis brevibus subtrifidis involucrisque rufescenti-hispidis.  $\mathcal{U}$  in siccis montosis Galloprovinciæ et Delphinatùs. — Fl. fr. 5. p. 440. Vill. voy. t. 4. f. 2. (v. DC.)
- $\beta$ . *collinum*, stolonosum, floribus submajoribus. — In Alsatiâ et Jurasso. — H. collinum Gochn. diss. p. 17. t. 1. Fl. fr. 5. p. 440. (v. v.)
17. *H. FALLAX* (W. enum. 822.) caule scapiformi stricto oligophyllo hispido,

floribus corymboso-paniculatis, pedunculis elongatis haxis ramosis, pedicellis subconfertis, involuero pilis nigris hirtis, foliis inferioribus lanceolato-spathulatis subpetiolatis pilosis acutis subintegris.  $\frac{1}{2}$  ab Alsatia ad Delphinatum. — Fl. fr. 5. p. 442. Flores majores quam in specie sequente. (v. s.)

$\beta$ . *hybridum*, scapo paucifloro uni-aut bis bifido. — H. *hybridum* Vill. voy. t. 2. f. 2. Fl. fr. 5. p. 442. (v. s.)

18. H. *FRAXETUM* (Vill. voy. t. 2. f. 1.) caule scapiformi stricto oligophyllo glabro, floribus laxè corymbosis, pedunculis elongatis ramosis centrali brevissimo 1-floro, pedicellis laxis, involucri hirtis, foliis lineari-lanceolatis acutis subglabris subintegris.  $\frac{1}{2}$  in pratis siccis à Galloprovincia ad Alsatiam. — Fl. fr. 5. p. 441. H. *piloselloides* Fl. fr. 4. p. 25 non Vill. (v. v.)

19. H. *PILOSELLOIDES* (Vill. dauph. t. 27. non fl. fr.) caule scapiformi oligophyllo subglabro, floribus paucis laxè corymboso-paniculatis, pedunculis laxis subsimplicibus centrali 1-floro aliis longiore, involucri hirtis, foliis lanceolatis acutis integris subglabris.  $\frac{1}{2}$  in Alpibus Delphinatûs. — Fl. fr. 5. p. 441. — Vill. voy. t. 3. f. 3. Hanc suspicor nil nisi præcedentis varietatem esse. (v. v. DC.)

20. H. *STATICIFOLIUM* (Vill. dauph. t. 27.) scapo oligophyllo glabro ramoso subtrifloro, pedunculis longis squamosis unifloris, foliis lineari-lanceolatis subdentatis glabris.  $\frac{1}{2}$  in Alpibus secus torrentes. — Fl. fr. 4. p. 25. All. ped. t. 81. f. 2. (v. v.)

21. H. *GLAUCUM* (All. ped. t. 28. f. 3 et t. 81. f. 1.) caule erecto ramoso folioso infernè subpiloso, foliis lanceolatis sessilibus dentatis utrinquè attenuatis.  $\frac{1}{2}$  secus torrentes Alpium Delphin., Pyren. — Fl. fr. 4. p. 26. (v. v.)

$\beta$ . *decipiens*, caule basi glabro, foliis lineari-lanceolatis. — H. *porrifolium* Fl. fr. 4. p. 26. non L. (v. s.)

#### §. 4. *Pulmonariæ* DC. l. c.

Caule folioso, involuero pilis nigris sæpius hirtis.

22. H. *ALBIDUM* (Vill. dauph. 3. t. 31.) caule ramoso hispido, foliis lineari-lanceolatis sessilibus semi-amplexicaulibus dentatis utrinquè attenuatis, pedunculis incrassatis, involucri laxis hispidis.  $\frac{1}{2}$  in pascuis Vogesorum, Alpium Delphinatûs. — Fl. fr. 4. p. 32. All. ped. t. 29. f. 3. (v. s.)

23. H. *TUBULOSUM* (Lam. dict. 2. p. 367.) caule ramoso hispido-viscoso, foliis lineari-lanceolatis profundè dentatis sessilibus basi attenuatis, involucri hispidis subviscosis, ligulis apice callosis canaliculato-tubulosis.  $\frac{1}{2}$  in Alpibus Delphinensibus. — Fl. fr. 4. p. 32. (v. s. ex monte Cenisio.)

24. H. *MONTANUM* (Jacq. aust. t. 190.) caule simplici unifloro, foliis ovato-lanceolatis denticulatis sessilibus, receptaculo subpiloso.  $\frac{1}{2}$  in pratis Alpium Delph. et Jurassi. — Fl. fr. 4. p. 29. — Vill. dauph. t. 23. — Boec. mus. 148. t. 113. (v. v.)

25. H. *GRANDIFLORUM* (All. ped. t. 29. f. 2.) caule simplici ascendente sulcato piloso-viscoso, foliis lanceolatis incurvatè dentatis, radicalibus subpetiolatis, caulinis amplexicaulibus sagittato-hastatis subdentatis, involucri hispidis, pappo albissimo.  $\frac{1}{2}$  in pascuis Alpium, montium Arvernæ, Mimatis et Pyrenæorum. — Fl. fr. 4. p. 33. (v. v.)

26. H. *AMPLEXICAULE* (L. sp. 1129.) caule ramoso, foliis ovato-cordatis amplexicaulibus basin versùs dentatis, radicalibus subobtusis breviter mucronatis, pedunculis involucrique hirsutis.  $\frac{1}{2}$  in lapideis Alpium Pyren., et montium Corbariensium. — Fl. fr. 4. p. 31. All. ped. t. 15. f. 1 et t. 30. f. 2. *Lepicaune balsamea* Lapeyr. H. *humile* Lap. ex cl. Arnott. *Catonia amplexicaulis* Cass. (v. v.)

27. H. *OVATUM* (Lapeyr. abr. suppl. 129.) caule simplici 1-4-floro, foliis glaberrimis mucronatis, radicalibus lanceolato-spathulatis petiolatis integris denticulatis, superioribus cordato-lanceolatis amplexicaulibus, peduncu-

lis involucrique hirsutis.  $\frac{1}{2}$  in Pyrenæis demissioribus. — An prioris var? (v. s.)

28. *H. CERINTHOIDES* (L. sp. 1129.) caule simplici villosa, foliis pilosis denticulatis, radicalibus obovatis, caulinis oblongis semi-amplexicaulibus, involucri hirsutis.  $\frac{1}{2}$  in Pyrenæis et Alp. Delphinatûs. — Fl. fr. 4. p. 27. Gon. ill. t. 22. f. 4. An satis distincta sp? (v. s.)
29. *H. BLATTARIOIDES* (L. sp. 1129.) caule simplici sulcato glabriusculo, foliis radicalibus oblongis basi profunde dentatis, caulinis lanceolatis hastato-sagittatis amplexicaulibus dentatis, involucri laxi pilosis.  $\frac{1}{2}$  in pascuis lapidosi Pyrenæorum, Alpium et Jurassi. — Fl. fr. 4. p. 33. *Crepis Austriaca* All. ped. t. 30. f. 1. *Catonia blattarioides* Cass. (v. v.)
30. *H. LAMPSANOIDES* (Gou. ill. t. 21. f. 3.) caule simplici, foliis caulinis lyrato-runcinatis amplexicaulibus hirsutis, floribus paniculatis, involucri hispidis.  $\frac{1}{2}$  in sylvis et pratis Pyrenæorum. — Fl. fr. 4. p. 28. (v. s.)
31. *H. PRÆNANTHOIDES* (Vill. dauph. 3. p. 108.) caule subsimplici, foliis radicalibus ovato-lanceolatis petiolatis, caulinis lanceolato-cordatis amplexicaulibus denticulatis pubescentibus, mediis basi obtusè subappendiculatis, floribus racemoso-corymbosis.  $\frac{1}{2}$  in apricis Alpium, Jurassi et montium Arverniz. — Fl. fr. 4. p. 24. *H. spicatum* All. ped. t. 27. f. 3. (v. v.)
32. *H. COMPOSITUM* (Lapeyr. abr. 476.) caule erecto villosa apice paniculatum ramoso, ramis bifidis ramosis 2- $\infty$ -floris, foliis denticulatis subtus præsertim in nervo villosissimis, radicalibus ovatis, caulinis cordatis amplexicaulibus.  $\frac{1}{2}$  in Pyrenæis orientalibus. — Fl. fr. 5. p. 444. (v. s.)
33. *H. SUCCISÆVOLIVUM* (All. ped. n. 786.) caule simplici scabriusculo, foliis integris, caulinis lanceolatis semi-amplexicaulibus, radicalibus petiolatis oblongis obtusè scabris, floribus corymbosis.  $\frac{1}{2}$  in Jurasso et Pyrenæis orientalibus. — Fl. fr. 4. p. 28. (v. v.)
34. *H. PALUDOSUM* (L. sp. 1129.) caule simplici, foliis glabris oblongis basi attenuatis runcinato-dentatis, caulinis amplexicaulibus, floribus paniculatis, involucri hispidis.  $\frac{1}{2}$  in humidis sylvis montanis. — Fl. fr. 4. p. 34. All. ped. t. 28. f. 2. et t. 31. f. 2. (v. v.)
35. *H. UMBELLATUM* (L. sp. 1131.) caule simplici, foliis linearibus subdentatis, floribus corymboso-umbellatis, involucri subglabro.  $\frac{1}{2}$  in pascuis siccis. — Fl. fr. 4. p. 31. Clus. hist. 2. p. 140. ic. (v. v.)
36. *H. SABAUDUM* (L. sp. 1131.) caule simplici, foliis ovato-oblongis subglabris acutis sessilibus subamplexicaulibus basin versùs dentatis, floribus corymbosis, involucri subglabro.  $\frac{1}{2}$  in nemoribus. — Fl. fr. 4. p. 31. *H. sylvaticum* Lapeyr. abr. 472. non Gon. ex. cl. Arnott. — Moris. s. 7. t. 5. n. 59. (v. v.)
37. *H. XRIOPHORUM* (S.-Am. bull. phil. n. 52. t. 2. f. 1.) radice præmorsâ, caulibus sulcatis erectis foliisque lanato-sericeis, foliis inferioribus lanceolatis dentatis acutis, involucri squamis basi sericeis apice subglabris, floribus confertis.  $\frac{1}{2}$  in arenosis agri Syrtici propè la Teste. — Fl. fr. 4. p. 21. (v. v.)
38. *H. PROSTRATUM* (DC. rapp. voy. 78.) radice præmorsâ, caulibus sulcatis prostratis basi villosa apice pubescentibus, foliis inferioribus lanceolatis villosa-sericeis dentatis subobtusis, floribus laxis.  $\frac{1}{2}$  in arenosis maritimis circâ Bayonam. — Fl. fr. 5. p. 437. An prioris var? (v. v.)
39. *H. SYLVATICUM* (Gou. ill. 56.) caule simplici folioso, foliis mollibus oblongis villosis dentatis, inferioribus petiolatis ovatis, floribus paniculatis, involucri breviter hispido.  $\frac{1}{2}$  in sylvis. — Fl. fr. 4. p. 30. — Lob. ic. 587. f. 1. (v. v.)
40. *H. MURORUM* (Lam. dict. 2. p. 365.) caule simplici subnudo, foliis mollibus subtus villosis ovatis basi profunde dentatis, inferioribus petiolatis subcordatis, floribus paucis paniculatis, involucri hispido.  $\frac{1}{2}$  ad muros et in sterilibus. — Fl. fr. 4. p. 30. — Barr. ic. 342. (v. v.)

*β. lanuginosum*, foliorum radicalium limbis petiolisque longis pilis albis undiquè vestitis. — In Pyren. orient. reperit Ph. Thomss. (v. s.)

41. *H. JACQUIN* (Vill. dauph. 3. t. 28.) caule erecto basi ramoso paucifloro, pedunculis involucriisque pilosis, foliis oblongis basi subpinnatifidis.  $\mathcal{L}$  in rupibus montanis Vogesorum, Delph., Galloprov., Pyrenæorum. — Fl. fr. 4. p. 35. (v. v.)
42. *H. PRUNELLEFOLIUM* (Gou. ill. t. 22. f. 30.) caule procumbente basi ramoso paucifloro, pedunculis involucriisque tomentosis, foliis ovatis basi inæqualibus denticulatis petiolatis, petiolis dentatis.  $\mathcal{L}$  in lapideis Alpium et Pyrenæorum. — Fl. fr. 4. p. 34. All. ped. t. 15. f. 2. *Omalocline prunellæfolia* Cass. (v. v.)

## LXXIX. DREPANIA Juss. gen. 169.

Involucrum polyphyllum squamis interioribus erectis, exterioribus patulis subulatis; receptaculum favosum; semina centralia margine membranaceo coronata 2-10-aristata, marginalia pappo sessili piloso brevissimo squamuloso donata.

1. *D. BARBATA* (Desf. atl. 2. p. 232.) involucri foliolis externis interiora æquantibus, pappo 1-3-setoso. ① in arenosis Gallie australis. — Fl. fr. 4. p. 48. Moris. s. 7. t. 4. n. 32. Variat pedunculis simplicibus et proliferis, floribus flavis aut pallidioribus. (v. v.)
2. *D. AMBIGUA* (DC. mss.) involucri foliolis externis paucis interioribus multò brevioribus, pappo 5-10-setoso. ② in Corsicâ. — *Crepis ambigua* Balb. diss. p. 4. t. 1. Fl. fr. 4. p. 40. *Schmidtia ambigua* Cass. (v. s.)

## LXXX. ANDRYALA L. gen. 915.

Involucrum imbricatum; receptaculum pilis semina superantibus onustum; pappus sessilis pilosus; in plurimis (*Rothia* Gærtn.) semina marginalia nuda.

1. *A. INTEGRIFOLIA* (L. sp. 1136.) caule erecto corymboso, foliis tomentosis sessilibus, inferioribus oblongis runcinatis, superioribus lanceolatis subdentatis, pedunculis glanduloso-villosis. ① in sterilibus provinc. australiorum et occidentaliû. — Fl. fr. 4. p. 36. — *Dalech. lugd.* 1116. f. 2. (v. v.)
2. *A. SINUATA* (L. sp. 1137.) caule erecto corymboso, foliis subtomentosis lanceolatis, inferioribus pinnatifidis, mediis sinuato-dentatis sessilibus, superioribus integerrimis. ② in Galliâ australi. — Fl. fr. 5. p. 444. (v. s.)
3. *A. LYRATA* (Pourr. act. toul. 3. p. 308.) caulibus subdiffusis, foliis adpressè velutinis, radicalibus lyratis, mediis basi subpinnatifidis, summis subintegris.  $\mathcal{L}$  in arenosis circâ Narbonam, Perpinianum. — Fl. fr. 5. p. 445. — *Lob. ic.* 231. f. 2. (v. v.)
4. *A. INCANA* (DC. fl. fr. 5. p. 445.) caulibus erectis laxè paniculatis, foliis oblongo-linearibus subdentatis integrisve, caule involucriisque velutinocanis, flosculis omnibus papposis.  $\mathcal{L}$  secus fluvium in arenosis vallium Plan et Gistan Pyrenæorum. — *Crepis incana* Lapeyr. abr. 483. Præcedentis var? (v. s. incompletam.)

+++ FRUCTUS CYLINDRACEUS. PAPPUS PLUMOSUS AUT SQUAMOSUS.

## LXXXI. ROBERTIA DC. fl. fr. 5. p. 453.

Involucrum simplici serie polyphyllum; receptaculum paleaceum; pappus sessilis plumosus.

1. *R. TARAXACOIDES* (DC. l. c.) glabra, foliis subruncinatis.  $\mathcal{L}$  ? in Corsicâ. — *Seriola taraxacoides* Lois. gall. t. 18. (v. s.)
- β. hirta*, involucrio hirta. — In Corsicâ montibus. (v. s.)

LXXXII. SERIOLA *L. gen.* 917.

Involucrum simplex; receptaculum paleaceum; pappus stipitatus, pilis paucis basi membranaceis apice subplumosis.

1. *S. STERNENSIS* (*L. sp.* 1139.) hispida, foliis obovatis integris dentatisve. ① in Corsicâ, Nicææ. — *Fl. fr.* 4. p. 922. *Lam. ill. t.* 656. f. 1. (v. v.)

LXXXIII. GEROPOGON *L. gen.* 904.

Involucrum simplex; receptaculum paleaceum; pappus stipitatus, disci plumosus, radii 6-aristatus.

1. *G. GLABRUM* (*L. sp.* 1109.) foliis glabris, involucri simplicibus. ① circa Nicæam. — *Fl. fr.* 4. p. 66. *Lam. ill. t.* 646. (v. s.)

LXXXIV. HYPOCHÆRIS *Vaill. L. gen.* 928.

Involucrum oblongum imbricatum; receptaculum paleaceum; pappus plumosus stipitatus aut in disco sessilis.

Sect. I. ACHYROPHORUS *Scop. carn. ed. 2. vol. 2. p.* 116.

Pappi omnes pedicellati.

1. *H. HELVETICA* (*Jacq. ic. rar. t.* 4.) caule simplici folioso subunifloro, foliis oblongo-lanceolatis dentatis, involucri villosissimo. 2℥ in pascuis montosis Jurassi, Delphinatûs, Pyrenæorum. — *H. uniflora* *Vill. dauph. 3. t.* 23. *Fl. fr.* 4. p. 46. *Porcellites Helvetica* *Cass. (v. v.)*
2. *H. MACULATA* (*L. sp.* 1140.) caule subnudo, ramo solitario, foliis ovato-oblongis subdentatis sæpius maculatis, involucri glabro. 2℥ in ericetis et collibus sterilibus. — *Fl. fr.* 4. p. 46. *Fl. dan. t.* 149. *Porcellites maculata* *Cass. (v. v.)*  
*β. simplex*, caule unifloro. — *Fl. fr.* 5. p. 451. (v. v.)
3. *H. RADICATA* (*L. sp.* 1140.) caule ramoso nudo sublævi, foliis runcinatis obtusis scabris, pedunculis squamosis, involucri squamis dorso hispidis. 2℥ ad vias et in pratis frequens. — *Fl. fr.* 4. p. 47. *Porcellites radicata* *Cass. — Lob. ic. t.* 238. f. 1. (v. v.)
4. *H. BALSMI* (*Lois. not. 124.*) subglabra, radice exili, foliis radicalibus repando-sinuatis, scapo simplici aut ramoso, involucri squamis glaberrimis, seminum omnium pappis stipitatis. ① in arenosis et humidiusculis. — *Fl. fr.* 5. p. 452. (v. s.)
5. *H. PINNATIFIDA* (*Ten. ad cat. h. neapol. app. 1. ed. 2. (1819) p.* 60. n. 147.) caule ramoso, foliis radicalibus runcinato-pinnatifidis, lacinii linearilanceolatis obtusis subintegris, caulinis linearibus, involucri squamis linearibus. 2℥ in montosis Corsicæ reperit *Salzmann* et sub nomine *Seriolæ taraxacifoliæ* divulgavit. (v. s.)

## Sect. II. EUNYPOCHÆRIS.

Pappi radii sessiles.

6. *H. GLABRA* (*L. sp.* 1141.) foliis radicalibus subtus glaberrimis, supernè piloso-hispidis, caule ramoso nudo, involucri squamis linearilanceolatis lucidis internis membranaceo-scariosis. ① in sylvis humidiusculis. — *Fl. fr.* 4. p. 47. *Lam. ill. t.* 646. f. 1. (v. v.)  
*β. minor*, caule 1-floro minimo. (v. s.)

LXXXV. TRAGOPOGON *Juss. gen.* 170.

Involucrum simplex 8-10-phyllum squamis coalitis; receptaculum nudum; semina longitudinaliter striata; pappus plumosus stipitatus, stipite tenui.

\* *Floribus luteis.*

1. *T. PRATENSE* (*L. sp.* 1109.) foliis glabris basi canaliculatis, pedunculis apice teretibus, involucri corollæ radio sublongioribus, squamis lanceolatis acutis,

seminibus asperis. ③ in pratis frequens. — Fl. fr. 4. p. 64. — Fuchs. hist. 821. ic. (v. v.)

2. *T. MAJUS* (Jacq. aust. t. 29.) foliis glabris planis basi latioribus, pedunculis apice incrassatis, involucri corollæ radio longioris squamis 10-12 lanceolato-ovatis acuminatis, seminibus echinatis. ½ in apricis Gallie medii, australis occidentalisque. — Fl. fr. 4. p. 64. Lam. ill. t. 646. f. 1. (v. v.)
3. *T. MINUTUM* (Gou. monsp. 342.) caule foliisque basi supernè tomentosis, pedunculis apice incrassatis, involucri corollæ radio non longioribus. ③ in sterilibus apricis Gallie mediterranæ. — Fl. fr. 4. p. 64. Garid. aix 466. t. 95. Præced. varietas? (v. s.)

\*\* *Floribus cæruleis vel violaceis.*

4. *T. FORNIFOLIUM* (L. sp. 1110.) caule stricto glabro, foliis lineari-ensiformibus basi dilatatis, involucri squamis longè acuminatis corollâ planissimâ duplò longioribus. ② in pratis provinciarum australiorum — Fl. fr. 4. p. 65. Jacq. ic. rar. t. 159. (v. v.)
5. *T. CROCIFOLIUM* (L. sp. 1110.) caule stricto tomentoso, foliis lineari-subulatis, involucri squamis lanceolatis acutis corollâ sublongioribus. ② in montibus subherbosis Gallie australis. — Fl. fr. 4. p. 65. — Col. ecphr. t. 230. (v. v. DC.)

LXXXVI. *THRINCIA* Roth. cat. bot. 1. p. 98. DC. fl. fr. 4. p. 51.

Involucrum imbricatum; receptaculum favosum; semina disci pappo sessili plumoso et inæquali donata, radii breviter papposa, pappo abortivo.

1. *T. HIRTA* (Roth. l. c.) foliis lanceolatis sinuato-dentatis hirtis, scapo 1-floro, involucri glabro basi squamulis ancto. ½ in sterilibus arenosis ad vias. — Fl. fr. 4. p. 51. Leontodon hirtum L. sp. 1123. — C. Bauh. prod. p. 63. ic. (v. v.)
2. *T. HIRPIDA* (Roth. cat. 1. p. 99.) foliis lanceolatis obtusis dentatis hispidis pilis furcatis, scapo 1-floro piloso, involucri canescente nudo. ① in lapidosis et arenosis provinciarum australium occidentaliūque. — Fl. fr. 4. p. 52. Hyoseris taraxacoides Vill. dauph. 3. t. 25. (v. s.)
3. *T. TUBEROSA* (DC. fl. fr. 4. p. 52.) radice tuberosâ, foliis lyratis hirtis scabris, scapo unifloro nudo apice hirtio. ½ in pratis circâ Monspelium, Nicæm, Aquas Sextias, Massiliam. — Leontodon tuberosum L. sp. 1123. — Lob. ic. t. 232. f. 1. (v. s.)

LXXXVII. *LEONTODON* Juss. gen. p. 170. DC. fl. fr. 4. p. 53. — *Apargia* Schreb. Wild.

Involucrum imbricatum; receptaculum favosum; pappus plumosus sessilis, pilis aliis squamosis aliis sericeis.

1. *L. INCAVUM* (DC. fl. fr. 4. p. 56.) foliis oblongis subdenticulatis cano-scabris pilis apice radiantibus, scapo unifloro pubescente, involucri squamis linearibus acuminatis pubescentibus. ½ in pascuis excelsis Alpium Delphinensium, circâ Monspelium ex cl. Benth. — Hieracium incavum L. Jacq. austr. t. 287. — Vill. 3. t. 24. (v. s.)
2. *L. squamosum* (Lam. dict. 3. p. 529.) foliis oblongis glabris subdentatis, scapo unifloro squamoso supernè hirtio, involucri hirti squamis linearibus acutis. ½ in pascuis montosis. — Fl. fr. 4. p. 54. *L. pyrenaicum* Gou. ill. t. 22. f. 1. 2. *Apargia Alpina* Wild. — All. ped. t. 14. f. 4. (v. v.)
3. *L. MONTANUM* (Lam. fl. fr. 3. p. 640.) foliis runcinato-dentatis glabris, scapis unifloris nudis supernè incrassatis hirtis, involucri villosis. ½ in lapidosis Alpium Delph., et Pyrenæorum ex Lapeyr. — Fl. fr. 4. p. 54. *Hedypnois taraxaci* Vill. dauph. 3. t. 26. — All. ped. 1. t. 31. f. 1. (v. v.)
4. *L. HASTILE* (L. sp. 1123.) foliis runcinato-pinnatifidis glaberrimis, laciniis lanceolatis basi dilatatis subdentatis axi inclinatis, scapo 1-floro nudo glabro,

involucris squamis interioribus glabris.  $\mathcal{U}$  in pascuis humidis Gallie occidentalis, mediz, orientalisque. — Fl. fr. 4. p. 55. L. danubiale Jacq. anst. t. 164. — All. ped. t. 70. f. 3. Vill. dauph. 3. t. 24. (v. v.)

$\beta$ . *arenarium*, foliis lineari-lanceolatis indivisis subdentatis. — In sabuletis maritimis agri Syrtici propè Teste de Buch reperiit cl. L. Dufour. (v. s.)

5. L. *hirsutum* (L. sp. 1124.) radice præmorsâ, foliis pinnatifido-oblongis laciniis nervo medio perpendicularibus hispidis pilis bifurcatis, scapo unifloro subglabro, involucri squamis lanceolatis acutis subhirsutis.  $\mathcal{U}$  in lapideis apricis. — Fl. fr. 4. p. 55. excl. var.  $\beta$  et  $\gamma$ . 5. p. 453. (v. v.)

6. L. *crispum* (Vill. dauph. 3. p. 84. t. 25.) radice simplici verticali, foliis runcinato-pinnatifidis laciniis linearibus lanceolatis nervo medio inclinatibus hispidis pilis apice ramosis, scapo 1-floro hirtio, involucri squamis lanceolatis acutis basi sublanatis.  $\mathcal{U}$  in siccis Delph., Occitaniz, Pyren. ex Lapeyr., Bellovacii ex cl. Graves. — Fl. fr. 5. p. 454. (v. v.)

7. L. *Villarsii* (Lois. gall. 514.) radice subsimplici verticali, foliis pinnatifidis laciniis linearibus lanceolatis nervo medio perpendicularibus hispidis pilis simplicibus, scapo 1-floro, involucri squamis hispidulis glabrisve, interioribus lanceolatis acuminatis.  $\mathcal{U}$  in siccis apricis Delph. infer., Galloprov., Occitaniz, agri Ruscinonensis. — Fl. fr. 5. p. 454. L. hirtum Vill. dauph. 3. p. 82. t. 25. (v. v.)

8. L. *autumnale* (L. sp. 1123.) foliis lanceolatis pinnatifidis glabris, scapis ramosis declinatis squamosis, pedunculis sub involucrio tumescentibus, involucri squamis puberulis lanceolatis acuminatis.  $\mathcal{U}$  ad margines agrorum et viarum. — Fl. fr. 4. p. 53. — Fuchs. hist. 320. ic. (v. v.)

$\beta$ . *simplex*, caule 1-floro. — eng. bot. t. 830. (v. v.)

LXXXVIII. *PODOSPERMUM* DC. fl. fr. 4. p. 61. — *Scorzonera* sp. L.

Involucrum imbricatum squamis margine membranaceis; semina pedicellata pedicello cavo incrassato; receptaculum seminibus delapsis tuberculato-muricatum; pappus sessilis plumosus.

1. P. *calcitrapæfolium* (DC. fl. fr. 5. p. 455.) caule basi ramoso, foliis petiolatis pinnatifidis, laciniis lanceolatis acutis, terminali longiore lanceolata, superioribus simplicibus, seminibus angulatis glaberrimis.  $\mathcal{U}$  in provinciis australibus et circâ Parisios. — P. resedifolium Fl. fr. 4. p. 61. non L. — Barr. ic. 800. (v. v.)

2. P. *laciniatum* (DC. fl. fr. 4. p. 62.) caule ramoso erecto, foliis petiolatis pinnatifidis laciniis linearibus acuminatis, terminali longissimâ linearibus lanceolata, superioribus linearibus, seminibus subangulatis glaberrimis.  $\mathcal{U}$  ad marginem agrorum. — *Scorzonera laciniata* L. Gærtn. fr. 2. t. 159. f. 1. S. petiolaris Lapeyr. abr. supp. 120. (v. v.)

3. P. *subulatum* (DC. fl. fr. 4. p. 61.) caule subramoso infrâ folioso superne nudo, foliis rigidis linearibus carinato-subulatis.  $\mathcal{U}$  Sorricinii. — An *Scorzonera pinifolia* Gon. ill. 53. var? (v. s. in specim. imperfecto.)

LXXXIX. *SCORZONERA* DC. fl. fr. 4. p. 59. — *Scorzonera* sp. L.

Involucrum imbricatum squamis margine membranaceis; receptaculum nudum; semina sessilia; pappus substipitatus plumosus.

1. S. *hirsuta* (L. mant. 278.) caule 1-floro, foliis linearibus subhirsutis obtusis, involucri squamis lanceolatis acuminatis glabris, seminibus lanatis.  $\mathcal{U}$  in lapideis sterilibus Occitaniz, Galloprovinciz circâ Massiliam et montem Victoriz, Rupellam. — Fl. fr. 4. p. 60. — Hieracium capillaceum All. ped. t. 31. f. 3. (v. v. DC.)

2. S. *angustifolia* (L. sp. 1113.) caule subnudo unifloro apice tomentoso, foliis sublinearibus basi lanatis, involucri squamis subglabris subobtusis, exterioribus lanceolato-ovatis, interioribus linearibus oblongis.  $\mathcal{U}$  in Gallia mediâ, australi occidentalisque. — Fl. fr. 4. p. 60. Gm. sib. 2. p. 2. t. 1. (v. v.)



*β. Provincialis*, caule foliisque glabris. — In paludosis salis Galloprovinciæ. (v. s.)

3. *S. ARISTATA* (Ram. in DC. fl. fr. 4. p. 922.) caule subnudo 1-floro, foliis linearibus acuminatis longissimis glabris, involucri squamis inferioribus foliaceis laxis subulatis florem æquantibus, interioribus obtusis, seminibus tuberculosus.  $\frac{1}{2}$  in herbosis Pyren. editiorum. — *S. grandiflora* Lapeyr. (v. v. DC.)
4. *S. HUMILIS* (L. sp. 1112.) caule subnudo subunifloro, foliis inferioribus oblongo-lanceolatis 5-7-nerviis acuminatis, superioribus linearibus, involucri squamis exterioribus ovatis subacutis, interioribus lanceolatis obtusis.  $\frac{1}{2}$  in pratis rariis. — Fl. fr. 4. p. 59. — Clus. hist. 2. p. 138. f. 2. (v. v.)
5. *S. HISPANICA* (L. sp. 1112.) caule ramoso 5-6-floro, foliis amplexicaulibus integris basi denticulatis.  $\frac{1}{2}$  in oleraceis hortis culta. — Fl. fr. 4. p. 59. — Blak. t. 406. (v. v.)
6. *S. PARVIFLORA* (Murr. syst. veg. ed. 14. p. 710.) caule ascendente folioso basi ramoso, foliis lanceolato-ensiformibus glabris amplexicaulibus acuminatis nervosis, involucri glabri squamis obtusis, exterioribus ovatis interioribus lanceolatis.  $\frac{1}{2}$  in littoribus Mediterraneis ex cl. Benth. — Jacq. aust. 4. t. 395. (v. s. ex Hung.)
7. *S. GRAMINIFOLIA* (L. sp. 1112.) caule folioso sub-1-floro apice tomentoso, foliis lineari-ensiformibus amplexicaulibus acutissimis nervosis, involucri subvillosi squamis exterioribus acuminatis, interioribus lanceolatis acutis.  $\frac{1}{2}$  in Galliâ australi. — Jacq. obs. 4. p. 13. t. 100. sed in nostrâ capitulum foliis non stipatum. (v. s.)
8. *S. PURPUREA* (L. sp. 1113.) caule stricto 1-2-floro basi lanato, foliis lineari-subulatis canaliculato-triquetris basi pubescentibus, involucri squamis obtusis exterioribus ovato-oblongis lanuginosis, interioribus sublinearibus.  $\frac{1}{2}$  in pratis umbrosis circa Mimatem reperiit cl. Prost. — Jacq. aust. t. 35. — Gmel. sib. 2. t. 2. Fl. purpurei. (v. s.)

XC. HYOSERIS Lam. dict. 3. p. 158.

Involucrum calyculatum; receptaculum nudum; seminum centralium pappus pilosus inæqualis, radii squamatus subabortivus.

1. *H. RADIATA* (L. sp. 1137.) foliis glabris runcinato-pinnatifidis laciniis dentato-angulatis reflexis, terminali 3-dentatâ, scapis unifloris nudis, involucri squamis exterioribus minimis lanceolatis.  $\frac{1}{2}$  in collibus maritimis Galliæ australis. — Fl. fr. 4. p. 49. — Pluck. t. 37. f. 2. (v. v.)
2. *H. SCABRA* (L. sp. 1138.) foliis lyrato-pinnatifidis dentatis scabriusculis, laciniis inflexis, scapis unifloris nudis apice incrassatis, involucri exteriori subnullo. ① in olivæ circa Nicæam. — Fl. fr. 4. p. 49. — Bocc. mus. 2. t. 106. (v. s.)
3. *H. HYDYNHOIS* (L. sp. 1138.) caule tereti ramoso, foliis inferioribus oblongis sinuato-dentatis basi attenuatis, superioribus subamplexicaulibus lanceolatis, involucri fructigeris glabris. ① inter segetes et in sterilibus Delph. et provinciæ mediterranearum. — Fl. fr. 4. p. 50. — Lob. ic. t. 239. f. 1. (v. v.)  
*β. nana*, caule nano 1-floro. (v. s.)  
*γ. rhagadioloides* (DC. mss.) involucri hirtellis. — *H. rhagadioloides* L. sp. 1139. Fl. fr. 4. p. 50. (v. s.)
4. *H. CAERTICA* (L. sp. 1139.) caule ramoso, foliis radicalibus sinuato-dentatis apice subretusis, caulibus subdentatis oblongis subcordatis amplexicaulibus, pedunculis tumidis apice fistulosis, involucri hirtis. ① in arvis circa Nicæam. — Fl. fr. 4. p. 51. Gærtn. fruct. 2. t. 160. f. 2. (v. s.)

XCI. CATANANCHE Tourn. t. 271. L. gen. 920.

Involucrum imbricatum scarosum; receptaculum paleaceum; pappus sessilis 5-squamosus squamis mucronatis.

1. *C. CAERULEA* (L. sp. 1142.) involucri squamis medio lineatis, inferioribus ovatis mucronatis, foliis villosis linearibus basi subpinnatifidis. 2/ in sterilibus Gallie australis occidentalisque. — Fl. fr. 4. p. 67. — Barr. ic. t. 1134. (v. v.)

XCII. CICHORIUM *Tourn. t. 272. L. gen. n. 921.*

Involucrum calyculatum, externum 5-phyllum, internum 8-phyllum squamis basi coalitis; receptaculum nudum vel subpilosum; pappus sessilis squamosus semine brevior.

1. *C. INTENSUS* (L. sp. 1142.) floribus sessilibus subaxillaribus geminis, foliis runcinatis nervis subhirsutis. 2/ ad visa. — Fl. fr. 4. p. 68. — Lob. ic. t. 228. f. 2. (v. v.)
2. *C. ERYDRIA* (L. sp. 1142.) pedunculis axillaribus geminis, altero elongato 1-floro, altero brevissimo sub 4-floro, foliis oblongis denticulatis. ① ex India (W.) orta, in hortis culta. — Fl. fr. 4. p. 68. (v. v.)

Ordo LVII. LOBELIACEÆ *Juss. ann. mus. 18. p. 1.*

Calyx ovario adhærens limbo 5-lobato aut integro. Cor. monopetala calyci inserta irregularis 5-loba suprâ profundè longitudinaliter fissa. Stam. 5 calyci inserta. Antheræ oblongæ filamento adnatæ longitudinaliter dehiscentes. Stylus simplex. Stigma urceolo membranaceo seu coronâ ciliatâ obvallatum. Fructus capsularis 2-aut rarè 4-locularis loculis polyspermis apice dehiscentibus, aut drupaceus nucularis 2-aut (abortu) 1-locularis, loculis polyspermis. Embryo rectus cylindricus in albumine tenui carnosus inclusus. Radicula ad hylum versa. — Herbæ aut suffrutices foliis alternis, floribus axillaribus terminalibusve. An potiùs ut Campanulacearum tribus tantùm habendæ?

I. LOBELIA *L. gen. n. 1006.*

Calyx 5-dentatus; cor. inæqualiter 5-loba; stam. 5; antheræ oblongæ in tubum connatæ; stigma obtusum sub-2-lobum; capsula calyce coronata 2-3-locularis seminibus numerosis tenuissimis.

1. *L. DORTMANNIA* (L. sp. 1318.) foliis linearibus 2-locularibus integerrimis, scapo simplici subnudo racemoso. 2/ in stagno *Casaux* propè Burdigalam ex Laterr. — Fl. fr. 3. p. 713. Fl. dau. t. 39. (v. s. ex Belgio.)
2. *L. URRUS* (L. sp. 1321.) caule erectiusculo, foliis inferioribus subrotundis crenatis, superioribus lanceolatis serratis, floribus racemosis. ① in Gallia occidentalis et mediæ sylvis humidis. — Fl. fr. 3. p. 715. — Moris. s. 5. t. 5. n. 56. (v. v.)
3. *L. LAURENTIA* (L. sp. 1321.) caule folioso prostrato, foliis lanceolato-ovalibus subcrenatis, pedunculis solitariis unifloris. ① in Corsicâ. — Fl. fr. 3. p. 731. Bocc. mus. 2. t. 27. fig. maj. (v. s.)
4. *L. MINUTA* (L. mant. 292.) foliis radicalibus ovato-spathulatis obtusis crenatis integrisve, scapâ capillaribus 1-floris foliis 2 minutis linearibus instructis. ① in Corsicâ. — Fl. fr. 3. p. 716. — Bocc. mus. 2. p. 35. t. 27. fig. min. An præcedentis var? (v. s.)

Ordo LVIII. CAMPANULACEÆ. *Juss. gen. p. 163.*

Calyx ovario adhærens 5-lobatus. Corolla 1-petala summo tubo calycis inserta 5-loba regularis aut rarius irregularis, sæpius marcescens. Stamina 5 sub corollâ calyci inserta, corollæ laciniis alterna. Antheræ 2-loculares oblongæ distinctæ rarius coalitæ. Ovarium suprâ glandulosum. Stylus unicus. Stigma 3-5-fidum. Caps. 3-5-locularis sæpius 3-locularis, foraminulis aut valvis 3-5 incompletis lateraliter dehiscens, loculis polyspermis, seminibus angulo loculorum interiori affixis. Embryo rectus in albumine carnoso; radícula infera. — Herbæ foliis alternis, floribus distinctis aut capitatis involucretis.

† *Antheræ connatæ.*

I. JASIONE *L. gen. n. 1005.*

Calyx 5-fidus; corolla rotata, tubo brevissimo, laciniis 5 longis linearibus; stam. 5, antheris in tubum coalitis; stigma 2-fidum; capsula 2-locularis. — Flores aggregati in involucre polyphylo.

1. *J. nummulus* (Pers. ench. 2. t. 215.) caulibus cespitosis tortuosis ascendentibus brevibus sub capitulo hirtis, foliis linearibus subundulatis obtusis glabriusculis ciliatisve, involucri foliolis lanceolato-ovatis mucronatis flores sessiles subæquantibus, calycis laciniis linearis-etaceis. ♀ in summis Pyrenæorum orientalium pascuis siccis. — Fl. fr. 5. p. 433. (v. v.)

2. *J. montana* (L. sp. 1317.) caule ramoso, foliis obtusis undulatis linearilanceolatis, pedunculis elongatis, involucri foliolis ovatis interioribus acuminatis, floribus pedicellatis, calycis laciniis linearibus acuminatis. ♂ in sterilibus siccis. — Fl. fr. 3. p. 717. — Dalech. lugd. 364. f. 1.

α. *lævis*, caule ramosissimo erecto, foliis glabriusculis, calyce glabro. (v. v.)

β. *hirsuta*, caule vix ramoso hirsuto, foliis hirsutis, involucri acutis, calyce subglabro. (v. v.)

γ? *maritima*, incano-pilosissima, caule prostrato subsimplici, foliis brevibus, involucri squamis exterioribus obtusis, calycibus hirsutis. ♂ in maritimis Gallie occidentalis. — An distincta sp? (v. s.)

3. *J. perennis* (Lam. dict. 3. p. 216.) caule simplici, foliis obtusis planis lanceolatis, pedunculis elongatis, involucri foliolis ovatis profunde dentatis acuminatis. ♀ in montosis Delph., Arvernæ, agri Lugd., Ligericini et Pyrenæorum. — Fl. fr. 4. p. 717. Lam. ill. t. 724. f. 2. (v. v.)

†† *Antheræ distinctæ.*

II. PHYTEUMA *L. gen. n. 220.*

Calyx 5-fidus; corolla rotata tubo brevissimo, laciniis 5 longis linearibus; stam. 5; stigma 3-partitum; capsula 3-locularis foraminulis lateralibus dehiscens. — Flores in spicas aut capitula bracteata congesti.

1. *P. pauciflora* (L. sp. 241.) foliis lanceolato-spathulatis obtusis, capitulo subfolioso, bracteis subrotundo-cordatis ciliatis subobtusis floribus subduplo brevioribus. ♀ in editoribus Alpibus Delph. et Pyrenæis (Ph. Thomas). — Fl. fr. 3. p. 710. (v. s.)

2. *P. HEMISPHERICA* (L. sp. 241.) foliis sublinearibus subintegerrimis caule vix brevioribus, capitulo subrotundo, bracteis ovatis inæqualibus acuminatis ciliatis floribus brevioribus.  $\mathcal{L}$  in Alpibus, editoribus montibus Arvernæ, Cebennis, Pyrenæis. — Fl. fr. 3. p. 710. — Moris. s. 5. t. 5. n. 53. (v. v.)
3. *P. SERRATA* (Viv. cors. app. 1.) foliis lineari-lanceolatis denticulatis, radicalibus obtusis petiolatis caule vix brevioribus, capitulo subrotundo, bracteis glaberrimis lanceolato-ovatis basi cordatis dentatis acutis floribus brevioribus.  $\mathcal{L}$  in montibus Corsicæ reperiit Ph. Thomas. (v. s.)
4. *P. CHARMELII* (Vill. dauph. 2. p. 516. t. 11.) foliis radicalibus petiolatis ovato-cordatis subdentatis, caulinis lineari-elongatis acuminatis denticulatis, bracteis linearibus capitulo rotundo longioribus.  $\mathcal{L}$  in Alpibus Delph., Gallopr., Pyrenæis. — Fl. fr. 3. p. 712. (v. v.)
5. *P. SCHREUCHERI* (All. ped. t. 39. f. 2.) foliis radicalibus petiolatis ovato-ellipticis lineari-lanceolatisve denticulatis, caulinis lineari-elongatis acuminatis vix denticulatis, bracteis linearibus angustissimis valdè inæqualibus capitulo rotundo 1-3-plò longioribus.  $\mathcal{L}$  in Pyrenæis ad montem Marboré ex cl. Benth. — Fl. fr. 3. p. 711. (v. s. ex Helv.)
6. *P. ORBICULARIS* (L. sp. 242.) foliis oblongo-lanceolatis crenato-serratis obtusis, radicalibus petiolatis lanceolato-cordatis, bracteis cordato-lanceolatis basi latioribus capitulo subrotundo brevioribus.  $\mathcal{L}$  in montosis. — Fl. fr. 3. p. 711. — Barr. ic. t. 525. (v. v.)
7. *P. RETONICA-FOLIA* (Vill. dauph. 2. p. 518. t. 12. f. 3.) foliis simpliciter crenatis, radicalibus lanceolato-cordatis, caulinis lineari-lanceolatis, inferioribus longè petiolatis, spicà oblongà subnudà.  $\mathcal{L}$  in Alpibus et Pyrenæis. — Fl. fr. 3. p. 713. (v. s.)
8. *P. SCORONERÆ-FOLIA* (Vill. dauph. 2. p. 519. t. 12. f. 2.) foliis subdentatis inferioribus oblongis petiolatis, superioribus linearibus sessilibus utrinquè attenuatis, bracteis paucis lineari-acuminatis spicà oblongà subsextuplò brevioribus.  $\mathcal{L}$  in Delphinatûs montibus. — Fl. fr. 3. p. 713. (v. s.)
9. *P. SPICATA* (L. sp. 242.) foliis radicalibus obtusis cordato-ovatis duplicato-dentatis petiolatis petiolo alato, caulinis lineari-lanceolatis dentatis sessilibus, bracteis paucis lineari-lanceolatis acuminatis spicà elongatà subquadruplò brevioribus.  $\mathcal{L}$  in sylvaticis humidis. — Fl. fr. 3. p. 714. — C. Bauh. prod. t. 32. f. 1. (v. v.)
10. *P. HALLERI* (All. ped. p. 430.) foliis obtusis, radicalibus cordato-ovatis duplicato-dentatis, inferioribus longè petiolatis, caulinis lanceolatis subsessilibus, bracteis numerosis lanceolatis acuminatis inæqualibus spicà ovatà subbrevioribus, stylis hirsutis flore longioribus emarginato-bifidis.  $\mathcal{L}$  in Pyrenæorum montibus Esquierry, Bergons, Erealids. — Fl. fr. 3. p. 714. *P. ovata* Willd. (v. v. DC.)

III. *PRISMATOCARPUS* L'Her. *sert. angl. DC. fl. fr. 3. p. 708.* — Campanulæ *sp. L.*

Corolla rotata limbo plano; capsula prismatica 2-3-locularis versùs apicem dehiscens. Cætera ut in Campanulâ.

1. *P. HYBRIDUS* (L'Her. l. c.) caule stricto basi subramoso, foliis oblongis subcrenatis, floribus solitariis, laciniis calycinis corollâ brevioribus. ① inter segetes. — Fl. fr. 3. p. 709. *Campanula hybrida* L. sp. 239. — Moris. s. 5. t. 2. n. 22. (v. v.)
2. *P. SPECULUM* (L'Her. l. c.) caule ramosissimo diffuso, foliis oblongis crenatis, floribus solitariis, laciniis calycinis corollam patulam æquantibus. ① in agris ad versuras. — Fl. fr. 3. p. 708. *Campanula Speculum* L. — Lob. ic. t. 418. (v. v.)

IV. *CAMPANULA* L'Her. *DC. fl. fr. 3. p. 696.* — Campanulæ *sp. Tourn. L.*

Calyx 5-fidus sinubus interdum reflexis; corolla campanulata 5-fida,

stam. 5 filamentis basi latioribus; stigma 4-rarius 5-partitum; capsula 3-rarius 5-locularis.

§. 1. *Capsulis obtectis sinubus calycis reflexis.*

1. *C. ALLIOWII* (Vill. dauph. 2. p. 512. t. 10.) caule unifloro hispido, foliis lineari-lanceolatis undulatis hispidis, radicalibus obtusis. ② in Alpibus Delph., Galloprov. — Fl. fr. 3. p. 706. C. Alpestris All. ped. t. 6. f. 3. (v. v. DC.)
2. *C. BARBATA* (L. sp. 236.) caule simplici, foliis lanceolatis villosis obtusis, racemo terminali laxo, corollâ barbata cernuis, laciniis calycinis acutis. ② in pascuis montanis Lothar. ex Buch., Belpin. — Fl. fr. 3. p. 706. — C. Bauh. prod. p. 36. f. 1. — Moris. s. 5. t. 3. n. 35. (v. v. in Helv.)
3. *C. MEDIUM* (L. sp. 236.) caule indiviso erecto folioso, foliis lanceolatis lineari-busve crenatis, floribus erectis, calycis laciniis lanceolato-ovatis ciliato-hispidis. ② in Galloprov. sylvis apricis, sæpius in hortis colitur. — Fl. fr. 3. p. 707. — Clus. hist. 2. p. 172. f. 2. Stigmata 5, capsulæ 5-loculares. (v. v. s. DC.)
4. *C. SPICATA* (L. mant. 337.) hispida, caule simplicissimo, foliis linearibus integris, florum sessilium spicâ laxâ, bracteis calycem superantibus. ② in Delphin. et Galloprov. subalpinis. — Fl. fr. 3. p. 708. — All. ped. t. 46. f. 2 et t. 47. f. 1. (v. v. D. in Valesiâ.)
5. *C. SPICIOSA* (Pourr. act. tol. 3. p. 309.) hispida, caule simplici folioso, pedunculis axillaribus unifloris, foliis linearibus subcrenatis longissimis, stigmatibus 3, capsulis 3-locularibus. ② in petrosis montosis Pyren., Gallie Narbonensis, agri Ligericini (Prost.), propè Monspelium loco dicto *Capouladoux* ex Benth. — Fl. fr. 3. p. 706. — C. longifolia Lapeyr. fl. pyr. 1. t. 6. (v. v. DC.)

§. 2. *Calycis sinubus non reflexis.*

6. *C. THYRSOIDEA* (L. sp. 235.) hispida, caule simplicissimo sulcato, foliis lanceolato-linearibus obtusis, floribus spicatis, spicâ densâ terminali oblongâ basi foliosâ. ② in Jurassi, Alpium et Pyrenæorum ex Lapeyr. declivibus asperis. — Fl. fr. 3. p. 704. Jacq. obs. t. 21. (v. v.)
7. *C. CERVICARIA* (L. sp. 235.) hispida, caule erecto simplici, foliis lanceolato-linearibus undulatis obtusis, radicalibus in petiolum attenuatis, floribus sessilibus capitato-glomeratis. ② in sylvaticis asperis montosis passim. — Fl. fr. 3. p. 703. — Bauh. prod. p. 36. f. 2. (v. v. in Helv. D.)
8. *C. PETREA* (L. sp. 236.) caule erecto simplici aut basi ramoso tomentoso, foliis scabris obtusis subtus cano-tomentosis, radicalibus petiolatis oblongo-lanceolatis crenatis basi attenuatis, caulinis sessilibus semialexicaulibus lineari-lanceolatis, florum capitulis terminalibus, seminibus exsertis. ② in Galloprovincia orientali loco dicto *les escalles d'Eglen*. — Clus. hist. p. 333. ic. (in herb. Richard v. cl. Alph. DC.)
9. *C. GLOMERATA* (L. p. 235.) caule obsolete angulato simplici glabresculo, foliis scabris obtusis, radicalibus petiolatis oblongo-lanceolatis dentatis basi cordatis, caulinis sessilibus semialexicaulibus, florum capitulis terminalibus axillaribusque. ② in aridis, collibus calcareis. — Fl. fr. 3. p. 703. — Lob. ic. t. 326. f. 2. (v. v.)
10. *C. TRACHELIUM* (L. sp. 235.) caule erecto obsolete angulato, foliis cordato-lanceolatis scabris inæqualiter serratis acuminatis, inferioribus petiolatis, floribus pedunculatis axillaribus, calycibus ciliatis laciniis lanceolato-ovatis. ② in sylvis. — Fl. fr. 3. p. 703. C. urticifolia Schmidt ex cl. Alph. DC. fl. fr. 3. p. 702. — Lob. ic. t. 326. f. 1. (v. v.)
11. *C. LATIFOEA* (L. sp. 233.) caule erecto simplicissimo tereti, foliis ovato-lanceolatis scabris crenato-dentatis, inferioribus petiolatis, floribus solitariis axillaribus terminalibusque breviter pedunculatis, calycis glabri laciniis nervo medio hispidulo aut glabro, corollis ciliatis. ② in sepius umbrosis montium. — Fl. fr. 3. p. 701. — Lob. ic. 2. t. 278. f. 2. (v. v.)

12. *C. RAPUNCULOIDES* (L. sp. 234.) caule erecto ramoso, foliis scabris cordato-lanceolatis, superioribus linearibus, floribus secundis axillaribus spicatis, calycis patentissimi pubescentis laciniis lineari-lanceolatis.  $\mathcal{L}$  in agris, siccissimis. — Fl. fr. 3. p. 702. — Moris. s. 5. t. 3. n. 32. (v. v.)
13. *C. RHOMBOIDALIS* (L. sp. 233.) caule striato, foliis sessilibus serratis, inferioribus subrhomboidis, caulinis acutis ovatis lanceolatisve, floribus racemosis secundis, calycis patentissimi laciniis subulato-linearibus.  $\mathcal{L}$  in pratis Alpium Delph., Gallopr., Arvernæ, Pyren. — Fl. fr. 3. p. 701. *C. lanceolata* Lapeyr. — Barr. ic. t. 567. (v. v.)
14. *C. RAPUNCULUS* (L. sp. 232.) caule erecto striato, foliis undulatis pilosis, radicalibus lanceolato-ovalibus, caulinis lineari-lanceolatis, panicula coarctata, calycis laciniis glabris subpatulis lineari-subulatis. ② in sylvaticis, arvis. — Fl. fr. 3. p. 699. — Fuchs. hist. 214. ic. (v. v.)
15. *C. PENSICIFOLIA* (L. sp. 232.) glabra, caule erecto simplici, foliis radicalibus lanceolato-obovatis serratis in petiolum attenuatis, caulinis lanceolato-linearibus subserratis sessilibus remotis, laciniis calycinis subpatulis lanceolato-linearibus.  $\mathcal{L}$  in sylvaticis. — Fl. fr. 3. p. 700. — Cus. hist. 2. p. 171. f. 3. (v. v.)
16. *C. PATULA* (L. sp. 232.) foliis radicalibus lanceolato-ovalibus dentato-sinuatis subpubescentibus basi attenuatis, caulimis linearibus glabris, panicula patula, calycis tubo obconico glabro laciniis basi denticulatis lineari-subulatis patentibus ferè longitudine corollæ.  $\mathcal{L}$  in sepibus et versuris passim. — Fl. fr. 3. p. 699. *C. decurrens* L. auct. — Dill. elth. t. 58. f. 68. (v. v.)
17. *C. LINIFOLIA* (Lam. dict. 1. p. 579.) caule decumbente glabro paucifloro, foliis radicalibus oblongo-lanceolatis subdentatis, superioribus linearibus lanceolatis, calycibus glabris laciniis subpatulis subulato-linearibus.  $\mathcal{L}$  in pratis frigidis Alpium, Arvernæ montium et Pycnæorum. — Fl. fr. 3. p. 698. *C. Schenckleri* Vill. dauph. 2. t. 10. (v. v.)
- β. Valdensis* (DC.) pubescens, caule subanifloro. — *C. Valdensis* All. ped. t. 6. Fl. fr. 3. p. 698. (v. v.)
18. *C. ROTUNDIFOLIA* (L. sp. 232.) caule decumbente, foliis radicalibus subrotundo-cordatis crenatis, caulinis lineari-lanceolatis dentatis, superioribus linearibus integerrimis, calycis patentissimi laciniis subulatis.  $\mathcal{L}$  in muris, saxosis, pascuis montosis. — Fl. fr. 3. p. 697. — Lob. ic. t. 328. f. 1. (v. v.)
- β. velutina* (DC.) caule foliisque velutinis. — Propè Massiliam. (v. v.)
19. *C. FUSILLA* (Jacq. coll. 2. p. 79.) caulibus decumbentibus hispidulis, foliis radicalibus longè petiolatis ovatis dentatis, caulinis lanceolatis crenatis, superioribus linearibus, calycis patentissimi laciniis setaceis.  $\mathcal{L}$  in rupetribus sylvaticis. — Fl. fr. 3. p. 697. — C. Bauh. prod. p. 34. ic. (v. v.)
20. *C. CENISIA* (All. ped. t. 6. f. 2.) minuta, caulibus diffusis unifloris foliosis, foliis ovatis glabris integerrimis subciliatis, floribus 5-fidis.  $\mathcal{L}$  in Alpibus Delph. et Galloprov. — Fl. fr. 3. p. 696. (v. s.)
21. *C. HEDERACEA* (L. sp. 240.) glabra, caulibus filiformibus decumbentibus laxis, foliis cordatis quinquelobis petiolatis, floribus longè pedunculatis, calycis laciniis linearibus. ① in umbrosis humidis provinciarum occidentaliū et rariis australium. — Fl. fr. 3. p. 696. — Moris. s. 5. t. 2. n. 18. (v. v.)
22. *C. ERINUS* (L. sp. 240.) pubescens, caule ramoso dichotomo, foliis sessilibus ovatis profundè dentatis, superioribus oppositis, floribus axillaribus subsessilibus.  $\mathcal{L}$  in petrosis provinciarum australium et rariis occidentaliū. — Fl. fr. 3. p. 705. — Moris. s. 5. t. 3. n. 25. (v. v.)

Ordo LIX. VACCINIEÆ. DC. *theor. elem. p.* 216.

Calyx adhærens 1-phyllus persistens, integer aut 4-5-6-lobus. Corolla toro inserta monopetala 4-5-partita, sæpius urceolata. Stamina definita distincta, toro inserta, sæpius decem. Antheræ sæpè bicornes. Ovarium unicum. Stylus 1. Stigma 1. Bacca calyce persistente coronata 4-5-locularis, loculis oligospermis. Embryo rectus in albumine carnosio; cotyledones brevissimæ, radícula longa infera. — Suffrutices foliis alternis coriaceis.

I. VACCINIUM *L. gen.* 483.

Calyx integer aut dentatus; corolla 4-fida; stam. 8; bacca globosa umbilicata oligosperma.

## Sect. I. MYRTILLUS.

Corolla campanulato - globosa 4-dentata, antheræ medio 2-cornes. — Folia decidua.

1. *V. ULIGINOSUM* (*L. sp.* 499.) pedunculis unifloris, foliis obovatis lanceolatis integris obtusis glabris venosis, ramis teretibus.  $\frac{5}{2}$  in uliginosis montosis passim. — *Fl. fr.* 3. p. 687. — *Clus. hist.* 1. p. 62. f. 1. (v. v.)
2. *V. MYRTILLUS* (*L. sp.* 498.) pedunculis unifloris, foliis serratis ovatis mucronulatis, ramis angulatis.  $\frac{5}{2}$  in ericetis, sylvaticis, arenosis. — *Fl. fr.* 3. p. 687. — *Lob. ic.* 2. t. 109. f. 1. (v. v.)

## Sect. II. VITIS-IDÆA.

Corolla campanulata 4-fida, antheræ muticæ. — Folia persistentia.

3. *V. VITIS-IDÆA* (*L. sp.* 500.) foliis obovatis obtusis revolutis integerrimis subtus punctatis, racemis terminalibus nutantibus.  $\frac{5}{2}$  in sylvaticis montium. — *Fl. fr.* 3. p. 687. — *Cam. epit.* 136. ic.

Sect. III. OXYCOCCOS *Pers. ench.* 1. p. 419.

Corolla 4-partita reflexa, antheræ muticæ. — Folia persistentia.

4. *V. OXYCOCCOS* (*L. sp.* 500.) caule filiformi repente, foliis ovatis integerrimis revolutis, pedunculis unifloris.  $\frac{4}{2}$  in uliginosis passim. — *Fl. fr.* 3. p. 688. — *Lob. ic.* t. 109. f. 2. (v. v.)

Ordo LX. ERICINÆ. *Desv. journ. bot.* 1813. p. 28. — Rhodoracæ et Ericacæ *DC. fl. fr.* 3. p. 671 et 675. — Rhododendra et Ericæ *Juss. gen.*

Calyx persistens liber 3-4-sæpius 5-partitus. Corolla basi calycis inserta, 4-5-fida aut partita, marcescens. Stamina corollæ laciniis alterna aut dupla, distincta, basi calycis aut corollæ inserta. Antheræ basi bicornes. Ovarium unicum liberum; stylus 1; stigma 1. Fructus multilocularis baccatus aut sæpius capsularis, polyspermus, multivalvis valvis suturâ aut medio dehiscentibus. Semina minuta placentæ centrali affixa. Embryo rectus in albumine carnosio. Cotyledones semicylindricæ. — Herbæ aut Frutices foliis sæpè persistentibus alternis verticillatisve integris aut dentatis.

Trib. I. *ERICACEÆ DC. fl. fr. l. c.*

Fructus valvis medio dehiscentibus.

† *Fructus baccatus.*

1? *EMPETRUM Tourn.* 421. *L. gen.* 1100.

Dioicum. Calyx 3-partitus; corolla 3-petala. *Masc.* Stam. 3, filamentis longis. *Fem.* Stylus subnullus; stigma radiis 6-9; bacca orbiculato-depressa 1-locularis 6-9-sperma; radícula inferior. — An hujus familiæ? aut ut vult. cl. Nuttall (*gen. am.* 2. p. 233.) novæ familiæ vicinæ Coniferis typus?

1. *E. nigrum* (L. sp. 1450.) procumbens, foliis oblongis glabris, stigmatibus 9-radiato. ♂ in petrosis Vogesorum, Jurassi, Alpium, Montis-Aurei, Pyrenæorum passim. — *Fl. fr.* 3. p. 686. — *Cam. epit.* 77. ic. (v. v.)

II. *ARBUTUS L. gen.* 552.

Calyx minimus 5-partitus; cor. ovata limbo parvo 5-fido revoluta; stamina 10; bacca 5-locularis.

Sect. I. *UNEDO.*

Stamina basi villosa, antheræ apice biperforatæ, bacca tuberculosa, loculi polyspermi.

1. *A. UNEDO* (L. sp. 566.) caule arboreo, foliis glabris obtusè serratis, paniculâ terminali. ♂ in maritimis Galliæ mediterranæ et austro-occidentalis. — *Fl. fr.* 3. p. 682. — *Barr. ic. t.* 673 et 674. (v. v.)

Sect. II. *MAIRANIA Neck. et bot.* 1. p. 219.

Stamina glabra, antheræ apice non biperforatæ, bacca lævis, loculi monospermi.

2. *A. ALPINA* (L. sp. 566.) caulibus procumbentibus, foliis rugosis serratis. ♂ in Alpibus et Pyrenæis. — *Fl. fr.* 3. p. 682. — *Clus. hist.* 1. p. 61. ic. (v. v.)

3. *A. UVA-URSI* (L. sp. 566.) caulibus procumbentibus, foliis integerrimis



lævibus margine revolutis.  $\S$  in Jurasso, Alpibus, Cebennis, Corbariensibus, Pyrenæis. — Fl. fr. 3. p. 683. Fl. dan. t. 33. (v. v.)

†† *Fructus capsularis.*

III. PYROLA *Tourn. inst. t. 132. Gærtn. fruct. 1. t. 63.* — *Pyrolæ sp. L.*

Calyx 5-fidus vel 5-partitus; cor. 5-partita serè 5-petala; stylus filiformis staminibus longior; stamina filiformi-subulata; capsula 5-locularis, basi angulis dehiscens. — An cum sequente *Monotropeis* affinior? (DC. mss.)

\* *Stylo erecto.*

1. *P. uniflora* (L. sp. 568.) foliis suborbiculatis, sæpè uniflora, calyce 5-partito, corollâ apertâ.  $\frac{1}{2}$  in Alpibus Delph. et Pyrenæis. — Fl. fr. 3. p. 685. — *Moris. s. 12. t. 10. n. 2.* (v. v. D. in Helv.)
2. *P. secunda* (L. sp. 567.) foliis elliptico-ovatis dentatis, racemo unilateralis, calyce 5-fido, corollâ campanulatâ, stylo exserto.  $\frac{1}{2}$  in montosis sylvaticis. — Fl. fr. 3. p. 685. — *Daléch. hist. p. 1148. f. 4.* (v. v.)
3. *P. minor* (L. sp. 567.) foliis rotundo-ovatis crenatis, racemo laxo, calyce 5-fido, corollâ campanulatâ stylum subsequente.  $\frac{1}{2}$  in umbrosis montosis. — Fl. fr. 3. p. 684. *Radius mon. pyr. t. 1. P. rosea Sm. eng. bot. p. 2543. Rad. pyr. t. 2.* (v. v.)

\*\* *Stylo declinato.*

4. *P. rotundifolia* (L. sp. 567.) foliis ovalibus subrotundis orbicularibusve obtusis, racemo laxo, calyce 5-partito, corollâ apertâ, stylo corollâ duplò longiore.  $\frac{1}{2}$  in sylvaticis humidis. — Fl. fr. 3. p. 684. — *Lob. ic. t. 294. f. 2.* (v. v.)

IV. CHIMOPHILA *Pursh. fl. am. bor. 1. p. 276.* — *Pyrolæ sp. L.*

Calyx 5-fidus; cor. 5-partita serè 5-petala patens; stylus brevis teres ovario immersus; stam. medio in discum dilatata; capsula 5-locularis dehiscens.

1. *C. umbellata* (Nutt. gen. am. 1. p. 274.) foliis cuneato-lanceolatis sessilibus, pedunculis subumbellatis subquaternis, filamentorum stipite nudo ciliato.  $\S$  in sylvis Vogesorum parte dictâ *ban de la Roche* rarissima ex Oberlin. — Fl. fr. 5. p. 431. — *Moris. s. 12. t. 10. n. 5.* (v. s. ex Germaniâ.)

V. ANDROMEDA *L. gen. n. 549.*

Calyx minimus 5-partitus; corolla 5-fida laciniis reflexis; stamina 10, capsula 5-locularis, 5-valvis.

1. *A. polifolia* (L. sp. 564.) foliis alternis lanceolatis revolutis, pedunculis aggregatis, corollis ovatis.  $\S$  in uliginosis circâ Rothomagum et in turfosis Jurassi, Vogesorum, Ruthenorum (Romergue). — Fl. fr. 3. p. 681. — *Pluck. t. 175. f. 1.* (v. v.)

VI. ERICA *Salisb. soc. lin. 6. p. 369.* — *Ericæ sp. L.*

Calyx 4-partitus; corolla campanulata sæpè ventricosa 4-fida; stam. 8; capsula 4-8-locularis 4-8-valvis.

1. *E. scoparia* (L. sp. 502.) foliis ternis angustis linearibus, calycis laciniis ovatis, antheris aristatis inclusis, corollis campanulatis, stigmatè exserto peltato.  $\S$  in sterilioribus provinc. med., austr. et occidentaliun. — Fl. fr. 3. p. 648. — *Lob. ic. t. 2. f. 2.* (v. v.)
2. *E. arborea* (L. sp. 502.) foliis ternis, ramulis incano-villois, corollâ globoso-campanulatâ glabrâ, antheris aristatis inclusis, stylo exserto peltato,

- calycis laciniis ovatis.  $\frac{3}{5}$  in provinciis australibus et austro-occidentalibus. — Fl. fr. 3. p. 677. — Clus. hist. 1. p. 41. ic. — Lob. t. 214. f. 1. (v. v.)
3. *E. RAMULOSA* (Viv. aen. bot. 1. p. II. p. 169.) foliis quaternis quiniseve, ramulis strictis coerctatis subglabris, floribus capitato-congestis, calycis laciniis lanceolatis, corollis ovoideis stylo aequalibus, antheris aristatis inclusis.  $\frac{3}{5}$  in collibus Corsicæ australis. — Viv. fragm. ital. 4. t. 7. *E. Corsica* DC. fl. fr. 3. p. 677. ic. rar. t. 17. (v. v.)
4. *E. MULTIFLORA* (L. sp. 503.) caule basi decumbente, foliis quaternis quiniseve linearibus angustissimis acutis, floribus pedicellatis, calycis laciniis ovatis, corollis ovatis, antheris muticis longè exsertis, stylo exserto.  $\frac{3}{5}$  in ericetis prov. occidentalium et circà Parisios. — Fl. fr. 5. p. 430. (v. v.)
5. *E. VAGANS* (L. mant. 230.) caule erecto, foliis quaternis quiniseve linearibus latis obtusis, floribus longè pedunculatis, calycis laciniis linearilanceolatis membranaceis, corollis cylindraceis, antheris muticis longè exsertis, stylo exserto.  $\frac{3}{5}$  in Galliâ meridionali. — Fl. fr. 3. p. 679 et 5. p. 430. (v. v.)
6. *E. MEDITERRANEA* (Thunb. diss. n. 41.) caule erecto, foliis ternis quaterniseve linearibus angustis obtusiusculis, floribus pedicellatis, calycis laciniis linearibus triangularibus, corollis cylindricis, antheris muticis styloque exsertis.  $\frac{3}{5}$  circà Bardigalam ex cl. Des Moulins (bull. soc. lin. burd. 1. p. 34.) — Curt. bot. mag. t. 471. (v. s. ex Italiâ.)
7. *E. CINEREA* (L. sp. 501.) foliis ternis, calycis laciniis linearibus, corollis ovatis, antheris aristatis inclusis, stylo exserto, stigmate capitato.  $\frac{3}{5}$  in arenosis aridis, ruentosis, sylvaticis Galliæ medim, australis occidentalisque. — Fl. fr. 3. p. 676. — Clus. hist. 1. p. 43. f. 2. (v. v.)
8. *E. CILIARIS* (L. sp. 503.) foliis ternis quaternisque subovatis ciliatis, racemis secundis, calycis laciniis lanceolatis, corollis ovatis inflatis, antheris muticis inclusis, stylo exserto.  $\frac{3}{5}$  in Galliâ occidentali. — Fl. fr. 3. p. 678. — Clus. hist. 1. p. 46. ic. (v. v.)
9. *E. TETRALIX* (L. sp. 502.) foliis quaternis linearibus ciliatis, floribus capitatis pedicellatis, calycis longè ciliati laciniis pedicellisque tomentosis, corollis ovatis, antheris aristatis inclusis, stylo subincluso.  $\frac{3}{5}$  in humidis turfosis Pyrenæorum, Galliæ occidentalis et medim. — Fl. fr. 3. p. 676. Fl. dan. t. 81. (v. v.)

VII. *CALLUNA* Salisb. trans. linn. l. c. t. 6. — *Erica* sp. L.

Calyx 4-partitus duplicatus; corolla campanulata 4-fida; stam. octo; capsula 4-locularis dissepimentis receptaculo affixis, 4-valvis, valvulis suturâ dehiscentibus.

1. *C. ERICA* (DC. fl. fr. 3. p. 681.) foliis sessilibus minimis 4-fariam imbricatis.  $\frac{3}{5}$  in sterilibus, sylvaticis, arenosis. *Erica vulgaris* L. Cam. epit. 75. ic. (v. v.)

Trib. II. *RHODORACÆE* DC. fl. fr. 3. p. 671.

Fructus capsularis, valvis suturis dehiscentibus.

VIII. *MENZIEZIA* Smith plant. ic. fasc. 3. 56. Juss. ann. mus. 1. p. 55. t. 4.

Calyx 4-partitus; corolla oblongo-ovoidea, limbo patente 4-dentato reflexo; stam. 8 basi corollæ inserta.

1. *M. DABEOCI* (DC. fl. fr. 3. p. 674.) floribus terminalibus racemosis, foliis alternis ovatis revolutis subtus incano-tomentosis.  $\frac{3}{5}$  in Pyrenæis occidentalisibus et Andegaviâ. — *Erica Dabeoci* L. sp. 509. (v. v.)

IX. *AZALEA* L. gen. n. 212.

Corolla infundibuliformis aut campanulata 5-fida inæqualis; stam. 5 sub pistillo inserta.

1. *A. procumbens* (L. sp. 217.) ramis diffusis procumbentibus, foliis ellipticis glabris margine revolutis, calycis laciniis æqualibus, stylo incluso. ♀ in Alpibus et Pyrenæis. — Fl. fr. 3. p. 674. *Loiseleuria procumbens* Deav. Journ. bot. 1813. t. 3. p. 35. — Clus. hist. j. p. 75. f. 3.

X. RHODODENDRON L. gen. n. 548.

Calyx 5-partitus; corolla subinfundibuliformis limbo patente 5-lobo; stam. 10 declinata.

1. *R. vernalinum* (L. sp. 562.) foliis ovato-oblongis glabris subtus leprosis, corollis infundibuliformibus. ♀ in Alpibus et Pyrenæis. — Fl. fr. 3. p. 673. — Lob. ic. t. 366. f. 2. (v. v.)  
 2. *R. nigrum* (L. sp. 562.) foliis ellipticis ciliatis subtus ferrugineo-punctatis, corollis infundibuliformibus. ♀ in summo Jurassi monte *Thoiry* ex Hall., Alp. Delphin. ex Vill., Pyren. monte *Sisay* ex Lapeyr. — Fl. fr. 3. p. 673. — Lob. ic. t. 468. f. 1. (v. v. D. in Ital.)

XI. LEDUM L. gen. n. 546.

Calyx minimus 5-dentatus; cor. 5-partita sub 5-petala; stamina 5-10 inno calyci inserta; capsula 5-locularis 5-valvis.

1. *L. palustre* (L. sp. 561.) foliis linearibus margine revolutis, subtus tomentosis. ♀ in Alsatia turfosis uliginosis ex Mapp. et Will. — Fl. fr. 3. p. 672. — Cam. epit. 546. ic. (v. ex Germ.)

Ordo LXI. MONOTROPEÆ. Nutt. gen. amer. 1. p. 272.

Calyx 5-partitus persistens interdum nullus aut bracteis irregularibus suppletus. Cor. perigyna persistens, petalis 4-5 nunc imâ basi coalitis, nunc connexis in tubum apice 4-5-dentatum. Stam. petalis dupla, corollæ basi inserta, alia petalis opposita, alia alterna. Antheræ excentricè peltatæ, filamentis adnatæ, sæpius 1-loculares; appendices filiformes recurvæ inter stamina sitæ. Ovarium liberum? Stylus 1. Stigma simplex discoideum. Caps. 5-locularis 5-valvis valvis medio septiferis, septis axeos basi coalitis. Semina numerosa minutissima pelliculâ samaræformi inclusa, apice interdum alata. — Herbæ orobanchoideæ carnosæ coloratæ, in radicibus arborum parasiticæ aphyllæ, nempè squamis foliorum loco onustæ.

I. MONOTROPA L. gen. n. 737. Gartn. fl. carp. 3. t. 185. f. 6.

Calyx 4-5-sepalus coloratus; petala totidem sepalis alterna et cum iis decidua. Stam. 8-10; filamenta subulata; ovarium liberum; stylus simplex cylindræus; caps. 8-10 sulcis exarata 4-5-locularis 4-5-valvis; albumen carnosum.

1. *M. hyemalis* (L. sp. 555.) floribus lateralibus octandris, terminali decandro. ♀ parasitica ad radices arborum parasitica. — Fl. fr. 4. p. 291. — Moris. hist. 3. s. 12. t. 16. n. 20. (v. v.)

## Subclassis III. COROLLIFLORÆ.

Calyx liber gamosepalus, sepalis nempè plus minùs inter se coalitis. Petala in corollam hypogynam liberam coalita. Stamina corollæ inserta. Ovarium liberum.

Ordo LXII. EBENACEÆ. *Vent. tabl. p. 443.* — *Guyacanæ Juss. gen. 155.*

Calyx monosepalus apice divisus. Corolla calyci inserta, monopetala lobata lobis corollæ laciniis numero æqualibus. Stamina corollæ inserta, definita aut indefinita sæpè monadelphæ. Ovarium liberum aut cum calycis basi subadhærens multiloculare loculis 1-2-ovulatis; stylus simplex aut divisus; stigma simplex aut divisum. Capsula aut bacca 1-∞-locularis, loculis 1-spermis. Embryo rectus in albumine carnoso. Cotyledones planæ. Radicula ad hylum versa. — Arbores aut Frutices foliis alternis simplicibus integris aut dentatis, floribus axillaribus.

Trib. I. *DIOSPYRÆÆ*. — *Ebenacæ Rich. R. Brown prod. p. 524.*

Ovarium liberum multiloculare, loculis 1-2-spermis, seminibus pendulis.

I. *DIOSPYROS* L. *gen. 1161.*

Calyx urceolatus 4-6-fidus; corolla imo calyci inserta urceolata 4-6-fida; stamina 8-16 imæ corollæ inserta interdum sterilia; stigmata 4 aliquandò bifida; bacca infernè calyce cincta 8-12-locularis.

1. *D. Lotus* (L. sp. 1510.) foliis ovato-oblongis infernè discoloribus punctis viridibus subcallosis notatis. ♂ in Occitaniâ cultus et nunc subspontaneus. — Fl. fr. 3. p. 670 et 5. p. 429. — Cam. epit. p. 156 et 157. ic. (v. v. c.)

Trib. II. *STYRACÆÆ* *Rich. anal. du fr. p. 48.*

Ovarium interdum adhærens 4-spermum, seminibus 2 ascendentibus, 2 inversis.

II. *STYRAX* *Tourn. inst. t. 369. L. gen. 595.*

Calyx urceolatus integer aut 5-fidus; corolla imo calici inserta urceolata, limbo brevi 3-7-lobo; stam. 6-16, filamentis basi coalitis corollæ tubo insertis, stigma 1; drupa exsucca supernè coriacea, nuce osseâ sphericâ.

1. *S. OFFICINALE* (L. sp. 635.) foliis ovatis subtus villosis, racemis simplicibus folio brevioribus. ♂ 5 in Galloprovinciâ et Nicæ sylvis maritimis. — Fl. fr. 3. p. 671. — Cam. epit. 38. ic. (v. v. D.)

Ordo LXIII. JASMINEÆ Juss. gen. 104. *Ach. Rich. mem. soc. hist. nat. t. 2.* — Jasmineæ et Oleineæ *Link et Hoffm. fl. portug. R. Brown prod. 520 et 522.*

Flores hermaphroditi aut in Fraxino polygami. Calyx 1-sepalus, in parte inferiori tubulosus 4-5-8-lobus. Corolla 1-petala regularis 4-5-8-loba aut partita, æstivatione mox contortâ mox valvatâ, aliquandò nulla. Stam. 2 corollæ inserta; filamenta brevissima; antheræ introrsæ 2-loculares longitudinaliter dehiscentes. Ovarium liberum sessile 2-loculare, loculis 2-ovulatis ovulis pendulis; stylus simplex; stigma 2-lobum. Fructus mox siccus dehiscentis indehiscensve 1-2-locularis, loculis 1-2-spermis; mox carnosus 1-2-locularis. Semina integumento membranaceo tenui aut crasso induta. Albumen album carnosum aut subcorneum aliquandò tenuissimum. Radicula cylindrica ad hylum versa. — Arbores aut frutices, foliis sæpius oppositis. Flores axillares aut racemosi terminales.

Trib. I. JASMINEÆ Vent. tabl. 1. p. 311.

Fructus carnosus.

I. OLEA Tourn. inst. t. 370. L. gen. n. 20.

Calyx parvus 4-dentatus; corolla tubo brevi, limbo 4-fido; stamina 2; lacinie stigmatis emarginatæ; drupa nuce 2-loculari 2-spermâ, abortu 1-loculari 1-spermâ.

1. O. EUROPEA (L. sp. 11.) foliis lanceolatis integerrimis paginis discoloribus, racemis paniculatis. ♀ culta in Galliâ mediterraneâ, in Corsicâ sylvestris forsan occurrens. — Fl. fr. 3. p. 497. Lam. ill. t. 8. f. 2. (v. v.)

II. PHYLLIREA Tourn. inst. t. 367. L. gen. n. 19.

Calyx minimus 4-dentatus; corolla brevis 4-fida; stamina 2; bacca 1-locularis 1-sperma.

1. P. ANGUSTIFOLIA (L. sp. 10.) foliis lineari-lanceolatis integerrimis acutis. ♀ in Galliâ australi. — Fl. fr. 3. p. 500. Lam. ill. t. 8. f. 3. (v. v.)

2. P. LATIFOLIA (Lam. dict. 2. p. 502.) foliis lanceolatis oblongisque subobtusis integerrimis serratisque. ♀ in Galliâ australioris collibus. — Fl. fr. 3. p. 499. — P. latifolia et media L. sp. 10. — Lob. ic. 2. p. 132. f. 2. Clus. hist. 1. p. 62. (v. v.)

III. LIGUSTRUM Tourn. inst. t. 367. L. gen. n. 18.

Calyx minimus 4-dentatus; corolla tubo brevi, limbo 4-fido patente; bacca 1-locularis 2-4-sperma.

1. L. VULGARE (L. sp. 10.) foliis elliptico-lanceolatis glabris, racemis compositis coarctatis. ♀ in glareosis, ad sepes. — Fl. fr. 3. p. 501. Lam. ill. t. 7. (v. v.)

IV. JASMINUM Tourn. inst. t. 368. L. gen. n. 27.

Calyx 5-fidus; corolla tubulosa limbo 5-partito plano, laciniis obliquis; bacca bilocularis loculis monospermis; semina arillata.

1. *J. OFFICINALE* (L. sp. 9.) foliis oppositis pinnatis, foliolis acuminatis, lacinii calycinis subulatis. 5 ex Indiâ ortum, in hortis cultum. — Fl. fr. 3. p. 500. Lam. ill. t. 7. f. 1. (v. v. c.)
2. *J. FRUTICANS* (L. sp. 9.) foliis alternis ternatis simplicibusque, foliolis subcuneatis obtusis, lacinii calycinis subulatis. 5 in Galliâ australi ad sepes et vineas. — Fl. fr. 3. p. 500. — Lob. ic. 2. p. 52. f. 1. Flores flavi. (v. v.)
3. *J. HUMILE* (L. sp. 9.) foliis alternis simplicibus ternatis pinnatisve, foliolis ovato-oblongis acutiusculis, ramis angulatis, lacinii calycinis brevissimis. 5 in Galloprovinciâ propè Grasse. — Fl. fr. 5. p. 394. — Lob. ic. 2. t. 106. f. 1. Fl. flavi. (v. s.)

Trib. II. *LILACEÆ* Vent. tabl. 1. p. 306. DC. fl. fr. 3. p. 494.

Fructus siccus.

V. *LILAC* Tourn. inst. t. 372. Fl. fr. l. c. — *Syringa* L. gen. n. 22.

Calyx parvus 4-dentatus; corolla tubulosa limbo 4-partito; stam. 2 intrâ tubum; capsula ovata compressa 2-locularis medio 2-valvis, 2-sperma; valvæ naviculares medio semi-septiferae.

1. *L. VULGARIS* (Lam. fl. fr. 2. p. 305.) foliis cordatis. 5 ex Oriente orta, in hortis culta et passim subspontanea. — Fl. fr. 3. p. 495. *Syringa vulgaris* L. — Duham. arb. 1. t. 138. (v. v. c.)
2. *L. FRAXEA* (Lam. l. c.) foliis lanceolatis integris pinnatifidisque. 5 in hortis colitur. — Fl. fr. 3. p. 495. *Syringa Persica* L. — Mill. dict. t. 164. f. 1 et 2. (v. v. c.)

VI. *FRAXINUS* L. gen. n. 1160.

Calyx o vel 3-4-partitus; corolla o vel 4-partita quasi 4-petala; samara pendula basi monosperma.

Sect. I. *FRAXINUS* Tourn. 343.

Flores calyce et corollâ destituti, nunquam hermaphroditi; antheræ sessiles.

1. *F. EXCELSIOR* (L. sp. 1509.) foliolis subsessilibus lanceolato-oblongis attenuatis serratis, samaris apice obliquè emarginatis. 5 in subhumidis. — Fl. fr. 3. p. 496. (v. v.)

Sect. II. *ORNUS* Tourn. l. c.

Flores calyculati tetrapetali; antheræ pedicellatae.

- 2? *F. ARGENTEA* (Lois. gall. 2. p. 697.) foliis impari-pinnatis 3-jugis, foliolis petiolatis ovato-lanceolatis apice attenuatis crenato-serratis cinereo-argenteis. 5 in fissuris rupium in montibus Corsicæ. (v. s. sine fl. et fructu.)
3. *F. FLORIFERA* (Scop. carn. n. 1250.) foliolis lanceolatis attenuatis petiolatis serratis. 5 in collibus Alsatiæ ex Lam., Galloprovinciæ ex Ger. et Gar. — Fl. fr. t. 3. p. 496. *F. Ornus* L. (v. v. c.)

Ordo LXIV. APOCYNÆÆ *Juss. gen. p. 143.*

Calyx monosepalus 5-divisus persistens. Corolla monopetala hypogyna 5-loba regularis, æstivatione imbricatâ rarissimè valvatâ, decidua. Stamina 5 imæ corollæ inserta laciniis limbi alterna; filamenta connata aut distincta; antheræ 2-loculares. Ovaria 2; styli 2 arcuè approximati sæpè brevissimi. Stigma 1 capitatum. Fructus 2-follicularis, folliculis elongatis 1-locularibus rimâ internâ dehiscentibus. Semina ad marginem folliculi inserta imbricata plana pendula ad umbilicum nuda aut comosa. Albumen carnosum. Embryo rectus. Radicula supera. — Herbæ aut frutices interdum volubiles sæpè lactescentes, foliis oppositis integris.

Trib. I. *ASCLEPIADEÆ* *R. Brown. in mem. soc. wern. 1. p. 12.*

Filamenta sæpius connata. Antheræ 2-loculares septivæ semi-completis nunc sub 4-loculares; pollen ad dehiscentiam antherarum coalescens in massas numero loculos æquantes rariùve per paria confluentes et solitariè vel geminatim vel quaternatim processibus 5 stigmatis affixas.

I. *ASCLEPIAS* *L. gen. 306.*

Corolla 5-partita reflexa; corona summo tubo filamentorum imposita 5-phylla, foliolis cucullatis; antheræ membranâ terminatæ; massæ pollinis compressæ apice attenuato affixæ pendulæ; stigma depressum muticum; semina comosa.

Sect. I. *GOMPHOCARPUS* *Brown. l. c. 1. p. 38.*

Coronæ foliola subcucullata utrinquè uni-dentata absque lacinulâ interiori, folliculi inflati aculeis mollibus echinati.

1. *A. FRUTICOSA* (L. sp. 315.) caule fruticoso, foliis lineari-lanceolatis glabris, pedunculis pedicellisque pubescentibus. ♀ in Corsicâ. — Pluck. alm. t. 138. f. 2. (v. a.)

Sect. II. *ASCLEPIAS* *Brown. l. c.*

Coronæ foliola cucullata è fundo exserentia lacinulam corniformem; folliculi graciles læves.

2. *A. SYRIACA* (L. sp. 313.) foliis planis ovalibus subtus tomentosis, caule simplicissimo, umbellis nutantibus. ♀ in hortis culta, nunc spontanea in Galliâ australi et occidentali. — Fl. fr. 3. p. 669. (v. v. c.)

II. *CYNANCHUM* *Brown l. c. — Cynanchi et Asclepiadis sp. L.*

Corolla subrotata 5-partita; corona 1-phylla 5-10-loba; antheræ membranâ terminatæ; massæ pollinis ventricosæ pendulæ; stigma apiculatum; folliculi læves; semina comosa.

1. *C. MONSPELIACUM* (L. sp. 311.) caule volubili herbaceo, foliis reniformi-cordatis acutis glabris, corollæ laciniis lanceolatis acutiusculis. ♀ in maritimis

Gallie mediterraneæ, Santoniæ (Des Moul.), Rupellæ. — Fl. fr. 3. p. 667. — Cav. ic. t. 60. (v. v.)

2. *C. Vincetoxicum* (Brown. mem. wern. soc. 1. p. 47.) caule erecto, foliis ovatis acuminatis margine tenuissimè ciliatis, racemo terminali, corollis imberbibus, coronâ 5-lobâ. ♀ in sylvaticis et glareosis. — Aocl. Vincetoxicum L. sp. 314. Fl. fr. 3. p. 668. — Lob. ic. t. 630. f. 1. (v. v.)
3. *C. minus* (Brown. l. c. p. 48.) caule supernè subvolubili, foliis ovatis basi barbatis, racemo terminali, corollis barbatis, coronâ semi-decemfidâ. ♀ in collibus sterilibus Gallie mediterraneæ. — Aocl. nigra L. sp. 315. Fl. fr. 3. p. 668. — Lob. ic. t. 630. f. 2. (v. v. DC.)

Trib. II. *VINCEÆ*. — Apocynæ Brown. prod. nov.-holl. p. 465.

Staminum filamenta distincta; antheræ 2 biloculares longitudinaliter dehiscentes; pollen granulosum stigmati immediate applicitum.

### III. *NERIUM* L. gen. 297.

Calyx 5-partitus; corolla infundibuliformis limbo obtusè 5-partito, laciniis basi intus appendiculatis; appendices 5 sæpè latæ faucem coronantes; antheræ apice in filum desinentes; stylus 1; stigma truncatum orbiculo insidens; semina comosa.

1. *N. Oleander* (L. sp. 305.) foliis lanceolatis angustis ternis subtus costatis, calycinis laciniis squarrosis, appendicibus, planis trifidis. ♂ in Galloprovinciâ, agro Nicœnsi et Corsicâ. — Fl. fr. 3. p. 666. — Lob. ic. t. 364. f. 1. (v. v. c.)

### IV. *VINCA* L. gen. 295.

Calyx 5-partitus; corolla hypocrateriformis tubo longo, fauce prominula 5-gonâ, limbo plano obtusè 5-lobato; antheræ approximatæ; stylus 1; stigma capitatum basi annulatum; semina nuda.

1. *V. major* (L. sp. 304.) caulibus erectiusculis, foliis ovato-subcordatis subciliatis, pedunculis unifloris solitariis folio (sæpius) brevioribus. ♀ in sylvis et collibus prov. austral. occidentalinque usque ad Nannetes et Andegavum. — Fl. fr. 3. p. 665. — Lob. ic. t. 636. f. 1. (v. v.)
2. *V. minor* (L. sp. 304.) caulibus procumbentibus, foliis oblongo-lanceolatis margine glabris, pedunculis 1-floris solitariis axillaribus folio longioribus. ♀ in sepibus et sylvis. — Fl. fr. 3. p. 665. Lam. ill. t. 172. f. 2. (v. v.)



Ordo LXV. GENTIANEÆ *Juss. gen.* 141.

Calyx monosepalus divisus persistens. Corolla monopetala regularis sæpè marcescens, limbo partito æquali, lobis æstivatione imbricatis, laciniis calycinis numero æqualibus, sæpiùs 5. Stamina totidem corollæ inserta et ejus laciniis alterna. Ovarium liberum. Stylus unicus aut rariùs scissione 2; stigma simplex aut bilobum. Capsula polysperma bivalvis 1-2-locularis, valvis apice dehiscentibus, margine introflexis et in bilocularibus dissepimentum constituentibus. Semina valvarum margini inserta. Embryo rectus in centro albuminis carnosì; radícula sæpiùs infera. — Herbæ glabræ amaræ, foliis oppositis sæpiùs integris sessilibus.

† *Capsula unilocularis.*I. MENYANTHES *Tourn. inst. t. 15. DC. fl. fr. 3. p. 647.* — *Menyanthis sp. L.*

Calyx 5-partitus; corolla infundibuliformis limbo patente 5-lobò æquali intùs hirtò; stamina 5; stylus 1; stigma capitatum 2-5-sulcum; capsula 1-locularis parietibus seminiferis.

1. *M. TRIPOLIATA* (L. sp. 208.) foliis 3-foliolatis glaberrimis longè petiolatis. 2½ in palustribus et aquosis. — Fl. fr. 3. p. 647. Lam. ill. t. 100. f. 1. — *Acopa Moris. s. 15. t. 2. n. 5. Fl. albescentes aut rosei.* (v. v.)

II. VILLARSIA *Gmel. syst. veg. 1. DC. fl. fr. 3. p. 648.* — *Menyanthis sp. L.*

Calyx 5-partitus; corolla rotata tubo brevi, limbo patente 5-lobò lobis margine ciliatis; stamina 5; stylus 1; stigma bilobum lobis crenulatis; capsula unilocularis suturis parietalibus seminiferis; semina biserialia margine membranacea.

1. *V. NYMPHOIDES* (Vent. choix. n. 9. p. 2.) foliis cordato-orbiculatis natan-  
tibus, floribus umbellatis, corollis ciliatis. 2½ in fluviis et aquis stagnantibus  
passim. — Fl. fr. 3. p. 648. Men. nymphoides L. sp. 207. Lam. ill. t. 100. f. 2.  
Flores flavi. (v. v.)

III. CHLORA *L. mant. 10.*

Calyx 8-partitus; corolla hypocrateriformis, tubo brevi, limbo 8-partito; stamina 8 brevissima fauci inserta; stylus 1; stigma 4-fidum; capsula 1-locularis.

1. *C. PERFOLIATA* (L. mant. 10.) foliis perfoliatis, calycibus 8-partitis, lobis  
subulato-linearibus. ① in collibus aridis. — Fl. fr. 3. p. 649. — Cam. epit. 427.  
ic. (v. v.)
2. *C. SÆSSILIFOLIA* (Desv. soc. am. 1. t. 3. f. 2.) foliis subamplexicaulibus, ca-  
lycibus 6-7-fidis, laciniis lanceolatis corollâ sublongioribus basi plus minùsve  
coactis. ① in maritimis Rupellæ, Narbonæ, Arelatis, Massiliæ (Solier). — Fl.  
fr. 5. p. 426. (v. v. DC.)

IV. SWERTIA *L. gen. n. 321.*

Calyx sub-5-partitus; corolla rotata tubo brevissimo, limbo phano 5-partito laciniis lanceolatis singulis basi intùs geminâ glandulâ ciliatâ instructis; stamina 5 corollâ breviora.

1. *S. PRÆNENS* (L. sp. 328.) corollis ecalcaratis 5-fidis, pedunculis tetragonis subulatis, caule indiviso, foliis radicalibus ovalibus.  $\frac{1}{2}$  in turfosis montosis Jurassæ, Alpium, Arvernæ, Pyrenæorum. — Fl. fr. 3. p. 650. — Clus. hist. 1. p. 316. f. 2. (v. v.)

V. *GENTIANA* Froel. gent. 9. Willd. sp. n. 512.

Calyx 5-fidus; corolla basi tubulosa, campanulata aut infundibuliformis, limbo 4-5-6-fido, laciniis integris aut ciliatis interdum laciniulis interjectis distinctis; stamina 5 tubo corollæ inserta; antheræ interdum connatæ; stylus 2-partitus; stigmata duo; capsula 1-locularis.

Sect. I. *CORLANTHE* Froel. gent. 15.

Corolla subcampanulata 4-9-fida fauce et limbo nuda.

1. *G. LUTEA* (L. sp. 329.) foliis latis ovatis nervosis, floribus verticillatis pedicellatis, calycibus membranaceis unilaterialibus, corollis rotatis 5-8-fidis acutis.  $\frac{1}{2}$  in montium pratis. — Fl. fr. 3. p. 650. — Lob. ic. t. 308. f. 2. Fl. lutei. (v. v.)
2. *G. BONSERI* (Lapeyr. abr. 132.) foliis ovato-lanceolatis obtusis, floribus verticillatis sessilibus, calycibus membranaceis uni-lateralibus, corollis campanulatis 6-fidis laciniis ovatis, sinibus dentatis.  $\frac{1}{2}$  in Pyrenæis frequens, in Alpihus rarior. — Fl. fr. 5. p. 426. Variat flore flavescente et nigro-punctato. (v. v.)
3. *G. PURPUREA* (L. sp. 227.) foliis ovato-lanceolatis subacutis, caulinis angustioribus, floribus verticillatis sessilibus, corollis campanulatis 6-fidis, laciniis subrotundo-spathulatis, calycibus membranaceis unilaterialibus.  $\frac{1}{2}$  in Alpihus. — Fl. fr. 3. p. 652. Fl. purpurei aut albi. (v. v.)
4. *G. BELOBA* (DC. fl. fr. 3. p. 653.) corollis subsexfidis verticillatis, calycibus membranaceis bilobis, lobis obtusis integris æqualibus.  $\frac{1}{2}$  in Galloprovinciæ montibus Sedenensibus. — Ic. rar. t. 15. (v. s.)
5. *G. PUNCTATA* (L. sp. 329.) foliis ovato-lanceolatis acutis, floribus verticillatis sessilibus, calycibus membranaceis truncatis lobis inæqualibus tubo calycino brevioribus, corollis subsexfidis campanulatis punctatissimis, laciniis ovato-acutis.  $\frac{1}{2}$  in Alpihus et Pyrenæis. — Fl. fr. 3. p. 653. — Barr. ic. t. 69. Fl. purpureo-flavescens. (v. v.)
6. *G. CRUCIATA* (L. sp. 334.) foliis lanceolatis obtusis, caulinis sessilibus basi coadunatis vaginantibus, floribus verticillatis sessilibus, corollis 4-fidis imberbibus.  $\frac{1}{2}$  in montosis siccis. — Fl. fr. 3. p. 653. — Clus. hist. 1. p. 313. f. 1. Fl. cærulei. (v. v.)
7. *G. ASCLEPIADEA* (L. sp. 329.) foliis amplexicaulibus ovato-lanceolatis acuminatis, floribus oppositis axillaribus solitariis subsessilibus, corollis 5-fidis campanulatis.  $\frac{1}{2}$  in Alpihus et montibus Galliæ mediteranææ. — Fl. fr. 3. p. 654. — Clus. hist. 1. p. 312. f. 2. Fl. cærulei. (v. v. D. in Helv.)
8. *G. PNEUMONANTHE* (L. sp. 330.) foliis linearilanceolatis linearibusve obtusiusculis, floribus terminalibus axillaribusve sessilibus, calycis laciniis linearibus, corollis 5-fidis campanulatis.  $\frac{1}{2}$  in pratis humidis. — Fl. fr. 3. p. 654. — Clus. hist. 1. p. 313. f. 2. Fl. cærulei. (v. v.)
9. *G. ACALYPS* (L. sp. 330.) foliis patulis oblongo-lanceolatis acutis, corollæ 5-fidæ caulem quadrangulum subæquantis laciniis acutis.  $\frac{1}{2}$  in Alpihus et Pyrenæis. — Fl. fr. 3. p. 654. excl. var.  $\gamma$ . — Barr. ic. t. 106. Fl. cærulei. (v. v.)
10. *G. ALPINA* (Vill. dauph. 2. p. 526. t. 10.) foliis obovatis latitudine vix longioribus obtusiusculis, corollæ campanulatæ 5-fidæ caule brevissimo longioris laciniis subobtusis.  $\frac{1}{2}$  in Alpihus et Pyrenæis. — Fl. fr. 5. p. 427. — Barr. ic. t. 105. Fl. cærulei. (v. v.)

Sect. II. CALATHIA *Froel. gent. 57.*

Corolla infundibuliformis, fauce nudâ, limbo non ciliato 5-10-fido.

11. *G. VERNA* (L. sp. 331.) caule simplici pumilo, foliis lanceolato-ovatis acutis, radicalibus confertis majoribus, corollæ 5-fidæ laciniis acutis margine erosis.  $\mathcal{Z}$  in pratis Jurassi, Alpium, Pyrenæorum. — Fl. fr. 3. p. 656. *G. utriculosa*  $\beta$ . Lapeyr. abr. 135. ex cl. Boileau. — Barr. t. 98 et 109. f. 1 et 2. Fl. cærulei. (v. v.)
12. *G. BAVARICA* (L. sp. 331.) caule simplici, foliis obovato-spathulatis obtusis, radicalibus confertis imbricatis, caulinis minoribus, corollæ 5-fidæ laciniis obtusis serratis.  $\mathcal{Z}$  in Alpibus. — Fl. fr. 3. p. 656. — Barr. ic. t. 10. fig. super. (v. v.)
13. *G. PYRÆNICA* (L. mant. 55.) caule infernè subramoso, foliis lanceolato-linearibus, corollâ 10-fidâ laciniis alternis minoribus obtusis crenatis.  $\mathcal{Z}$  in Pyrenæis orientalibus. — Fl. fr. 3. p. 657. Gon. ill. 7. t. 2. f. 2. (v. v. DC.)
14. *G. UTRICULOSA* (L. sp. 332.) caule ramoso, foliis radicalibus ovatis confertis, calycibus inflatis plicato-carinatis, corollæ 5-fidæ hypocrateriformis laciniis ovato-lanceolatis angustis.  $\textcircled{1}$  in pratis humidis Alpium, Jurassi, Alsatiæ, Nestler. — Fl. fr. 3. p. 657. — Barr. ic. t. 48 et 122. f. 2. Fl. azurei. (v. v. D. in Italiâ.)
15. *G. NIVALIS* (L. sp. 332.) caule submultifloro, ramis alternis unifloris, foliis caulinis lanceolatis, calycibus 5 lineis fuscis notatis, corollâ 5-fidæ laciniis ovato-lanceolatis.  $\textcircled{1}$  in pratis editiorum Alpium et Pyren., Montis-Aurei ex Delarb., Burgundiae? ex Dur. Fontisbleaudi? ex Thuil. — Fl. fr. 3. p. 656. — Hall. helv. t. 17. f. 5. — Barr. t. 103. f. 2. (v. v.)

Sect. III. ENDOTRICHIA *Froel. gent. 86.*

Corolla 4-5-fida, fauce squamis multifidis acutis instructâ.

16. *G. CAMPESTRIS* (L. sp. 334.) caulibus ramosissimis, foliis ovato-lanceolatis, laciniis calycinis 2 maximis obovatis, 2 minoribus linearibus, corollis 4-fidis hypocrateriformibus obtusis fauce barbatis.  $\textcircled{1}$  in pratis montosis. — Fl. fr. 3. p. 659. Fl. dan. t. 367. Fl. purpureo-violacei. (v. v.)
17. *G. GERMANICA* (Willd. sp. 1. p. 1346.) caule ramosissimo, foliis ovato-lanceolatis, pedunculis internodiis longioribus brevioribusve, calycibus æqualiter 5-fidis, corollis 5-fidis hypocrateriformibus fauce barbatis.  $\textcircled{1}$  in pratis montosis. — Fl. fr. 3. p. 658. *G. Amarella* W. sp. 1. p. 1347. — Barr.-ic. t. 112 et 510. f. 2. Fl. purpureo-violacei. (v. v.)
18. *G. GLACIALIS* (Vill. dauph. 2. p. 532.) caule infernè ramoso folioso, foliis ovatis, laciniis calycinis æqualibus, floribus longè pedicellatis, corollis 4-fidis hypocrateriformibus fauce barbatis.  $\textcircled{1}$  in Alpibus Galloprovinciæ et Pyren. — Fl. fr. 3. p. 659. Fl. cærulei, tubo pallidiore. (v. v.)

Sect. IV. CROSSOPETALUM *Froel. gent. 109.*

Corolla 4-fida infundibuliformis, laciniis ciliatis.

19. *G. CILIATA* (L. sp. 334.) caule flexuoso angulato, foliis lanceolatis linearibusque, corollis 4-fidis, laciniis serratis medio inciso-ciliatis.  $\mathcal{Z}$  in humidis montosis passim. — Fl. fr. 3. p. 659. — Bar. ic. t. 121. Fl. azurei. (v. v.)

†† *Capsula 2-locularis.*

- VI. *CHIRONIA* *Schmidt bohem. Wild. sp. n. 394.* — *Erythræa* *Ren. Rich. in Pers. ench.* — *Gentianæ* *sp. L.*

Calyx 5-fidus; corolla infundibuliformis limbo 5-loba; stylus declinatus; stamina 5 tubo corollæ inserta; antheræ post deflorationem spiritaliter contortæ; capsula 2-locularis.

1. *C. LINEARIFOLIA* (Lois. not. 155.) ramoso-dichotoma, foliis omnibus linea-

ribus obtusiusculis, calyce 5-partito lobis linearibus obtusiusculis tubum subæquantibus. ② in arenosis Druentia propè Avenionem. — Fl. fr. 5. p. 428. — Barr. ic. 423. — Fl. rosei. (v. s.)

2. *C. SPICATA* (W. sp. 1. p. 1069.) ramis dichotomo-paniculatis, foliis lanceolato-oblongis, floribus sessilibus subsecundis, calyce bracteisque corollæ tubo subæqualibus. ① in Gallia australis humidis. — Fl. fr. 3. p. 662. — Barr. ic. t. 1242. — Fl. rosei. (v. v.)

3. *C. CENTAURIUM* (Smith. brit. 1. p. 257.) caule erecto, foliis ovatis aut ovato-oblongis, floribus corymbosis aut paniculato-corymbosis, calycis 5-fidi laciniis subulatis tubo corollæ brevioribus. ① in sylvis, arvis, ad ripas frequens. — Fl. rosei. Mirum in modum varians adeò ut vix in tres sequentes varietates distingui possit.

a. *fascicularis*, caule subsimplici, floribus fasciculato-corymbosis subsessilibus, calycibus plerumque tubo corollæ dimidiò brevioribus. — *C. Centaurium* Fl. fr. 3. p. 660. (v. v.)

β. *pulchella* (DC. fl. fr. 5. p. 428.) caule ramosissimo, floribus paniculato-corymbosis plerumque pedicellatis, calycibus plerumque tubo corollæ subæquantibus. — *C. pulchella* Swartz act. holm. 1783. t. 3. f. 8. g. Fl. fr. 3. p. 661. — Vaill. bot. t. 6. f. 1. (v. v.)

γ. *nana*, caule pumilo simplicissimo, floribus terminalibus axillaribusve subsolitariis, calyce tubum plerumque subæquante. (v. v.)

4. *C. OCCIDENTALIS* (DC. fl. fr. 5. p. 428.) caule nano è collo subramoso floribus brevioribus, foliis ovato-oblongis, calyce ultrà medium 5-fido laciniis linearibus subulatis. ① in arenosis maritimis à Bayonà ad Armoraciam passim. — *Flos flavus*. Sequentis forsàn var? (v. s.)

5. *C. MARITIMA* (Wild. sp. 1. p. 1069.) caule elongato simplici paucifloro dichotomo-corymboso, foliis ovato-lanceolatis, laciniis calycinis linearibus subulatis. ① in Gallia mediterraneæ maritimis. — Fl. fr. 3. p. 662. *Gentiana maritima* L. mant. 55. Cav. ic. t. 269. f. 1. — Barr. ic. t. 468. *Flores flavi*. (v. v.)

VII. *EXACUM* Wild. sp. 2. p. 634. DC. fl. fr. 3. p. 663. — *Gentianæ* sp. L.

Calyx 4-partitus; corollæ tubo globoso, 4-loba; stamina 4; antheræ post fecundationem non tortæ; stylus unicus; stigma 2-fidum; caps. 2-locularis.

1. *E. CANDOLLI* (Bast. supp. 22.) glaucum, caule ramoso subdichotomo gracilimo, foliis linearibus lanceolatis trinerviis, pedicellis elongatissimis, calycis laciniis linearibus subulatis rectis. ① in herbis inundatis circà Domfront ex cl. de Brébisson, Andegaviam, Nannetes, Monspelium. — Fl. fr. 5. p. 429. *E. pusillum* β DC. ic. rar. t. 16. Fl. rosei. (v. v. DC.)

2. *E. PUSILLUM* (DC. fl. fr. 3. p. 663.) caule ramosissimo dichotomo, foliis oblongo-lanceolatis 3-nerviis, calycis laciniis linearibus subulatis, corollæ limbo connivente. ① in locis hyeme inundatis Gallia mediæ et occidentalis passim. — *Gent. pusilla* Lam. dict. 2. p. 645. — Vaill. bot. 32. t. 6. f. 2. Fl. flavescens. (v. v. D.)

3. *E. FILIFORME* (Wild. sp. 1. p. 638.) caule filiformi subramoso, foliis radicalibus subrotundis, caulinis subulatis, corollæ limbo patente. ① in pratis humidiusculis sylvaticis passim. — Fl. fr. 3. p. 663. *Gent. filiformis* L. — Vaill. bot. t. 6. f. 3. Fl. flavescens. (v. v.)

Ordo LXVI. POLEMONIDEÆ *Juss. gen.* 136.

**Calyx** 1-sepalus divisus. **Corolla** regularis 5-loba. **Stamina** 5 medio corollæ tubo inserta, ejusdem laciniis alterna. **Ovarium** liberum. **Stylus** simplex. **Capsula** calyce persistente tecta 3-locularis 3-valvis polysperma, valvis medio intus septiferis seu costâ prominente instructis, axi centrali trigono valvularum septis angulis applicito. **Embryo** rectus in albumine corneo; **radicula** infera; **cotyledones** ellipticæ foliaceæ. — **Herbæ** foliis oppositis aut rariùs alternis.

I. POLEMONIUM *Tourn. inst. t. 61. L. gen. n. 217.*

**Calyx** 5-fidus; **corolla** rotata tubo brevi, limbo 5-lobo; **staminum** filamenta basi latiora; **antheræ** incumbentes.

1. *P. CÆRULEUM* (L. sp. 162.) foliis pinnatis, calycibus corollæ tubo longioribus. ♀ in Jürasso et Pyrenæis ex Lapeyr., frequenter in hortis cultum. — Fl. fr. 3. p. 646. Lam. ill. t. 106. f. 1. Variat floribus albis et cæruleis. (v. v. c.)

Ordo LXVII. CONVULVACEÆ *Juss. gen.* 132. (1)

**Calyx** 5-lobatus, persistens. **Corolla** monopetala hypogyna, limbo regulari 5-lobo. **Stam.** 5 imæ corollæ inserta, ejusdem laciniis alterna. **Ov.** simplex liberum, basi glandulâ annulari cinctum, 2-4-locularis, loc. sæpiùs 1-2-spermis. **Stylus** unicus apice quandoquæ ad basin usque partitus. **Stigma** acutum vel capitatum. **Caps.** 1-4-locularis, dissepimento 1-4 valvularum marginibus affixo, in centro basi seminifero. **Semina** subangulosa basi hilo notata. **Embryo** convolutus. **Rad.** infera; **cotyl.** corrigatæ; **albumen** mucilaginosum.

**Herbæ** suffrutices frutices et arbores; **plures** volubiles lactescentesque. **Folia** alterna, formâ varia. **Inflorescentiæ** variæ.

I. CONVULVULUS *L. gen. n. 215.*

**Calyx** 5-partitus nudus vel bracteis 2 imbricatus; **cor.** campanulata 5-plicata; **stamina** inclusa, sæpè inæqualia; **ov.** 2-loculare, loculis 2-spermis, rarò 3-loculare; **stylus** unicus; **stigma** filiforme aut capitatum, **capsula** 1-2-locularis.

§. 1. *Floribus subtus 2 bracteis magnis cinctis.*

1. *C. SOLDANELLA* (L. sp. 226.) caule lævi prostrato, foliis reniformibus obtusis-

(1) Hujus familiæ specierum Callicarum monographiam mihi communicavit cl. amic. Choisy.

simis apice rarè emarginatis, pedunculis 1-floris, calyce membranaceo, corollâ purpureâ, stigm. ovatis complanatis.  $\mathcal{Y}$  frequens in maritimis sabulosis. — Fl. fr. 3. p. 641. — Lob. ic. 602. f. 2. (v. v.)

2. *C. SEPTUM* (L. sp. 218.) caule glabro volubili angulato, foliis sagittato-acuminatis, pedunculis 1-floris quadrangulis, calyce membranaceo, corolla alba rarè rosea spectabili, stigmatibus ovatis complanatis.  $\mathcal{Y}$  ubique in sepibus. — Fl. fr. 3. p. 630. Fl. dan. t. 648. (v. v.)

$\beta$ . *maritimus* (Gou. fl. monsp. 27.) omni parte gracilior, internodiis longioribus foliis crassiusculis. — In maritimis circâ Mompelunum. (v. s.)

§. 2. *Floribus nudis, caule erecto aut diffuso.*

3. *C. SAXATILIS* (Vahl. symb. 3. p. 33.) caule suffruticoso ramoso, ramis foliis calycibusque lanatis, foliis lineari-lanceolatis acutis sessilibus, pedunculis longis capitatis, capitulis multibracteatis, calyce acuminato, stigm. linearibus.  $\mathcal{Y}$  In rupestribus circâ Perpignanum. — Fl. fr. 5. p. 424. C. capitatus Cav. ic. 2. t. 89. — Barr. ic. t. 470. (v. s.)

$\beta$ . *argenteus* (Ch.) pilis argenteo-sericeis potiùs compressis, calyce minùs elongato. — Circâ Cujes propè Telonem. — Conv. linearis DC. fl. fr. 5. p. 424. (v. s.)

4. *C. CANTABRICA* (L. sp. 225.) caule ascendente ramosissimo villosa, foliis lineari-lanceolatis acutis sessilibus villosis, ped. elongatis 1-3-floris, calyce lineari inæquali, stigm. linearibus.  $\mathcal{Y}$  in totius regionis Mediterranei maris montosis et rupestribus, circâ Burdigalam ex Des Moulins. — Fl. fr. 3. p. 642. Jacq. aust. t. 296. (v. v.)

5. *C. LINEATUS* (L. sp. 224.) caule minimo suffruticoso sericeo-viloso, foliis oblongis apice dilatatis acutis in petiolum attenuatis sericeis, ped. 3-6-floris paniculatis, calyce cuneato-lanceolato.  $\mathcal{Y}$  in rupestribus maritimis Galloprovinciæ. — Fl. fr. 3. p. 642. — Triumf. obs. gr. ic. (v. v.)

$\beta$ . *intermedius* pube magis argenteâ, caule longiore recto. — Circâ Avenionem. — C. lineatus  $\beta$  DC. fl. fr. 5. p. 413.

6. *C. TRICOLOR* (L. sp. 225.) caule herbaceo tereti villosa, foliis lanceolato-obovatis subspathulatis basi ciliatis, ped. 1-floris folio sæpè longioribus, calyce ovato-lanceolato acuto, corollâ tricolore, stigm. linearibus. (1) in hortis vulgaribus, hab. Siciliam, Hispaniam, Italiam, etc. — Fl. fr. 3. p. 641. — Moris. s. 1. t. 4. n. 4. (v. v. c.)

7. *C. SICULUS* (L. sp. 223.) caule herbaceo prostrato tereti villosa, foliis ovatis acutis subpubescentibus basi truncatis, ped. 1-floris post anthesin nutantibus, calyce lanceolato acuto ciliato, corollâ minimâ cærulescente, stigm. filiformibus. (1) in arenosis agri Syrtici, Ruscinonensis. — Fl. fr. 3. p. 640. — Boeck. sic. 89. t. 48. (v. s.)

§. 3. *Caule volubili, floribus nudis.*

8. *C. ARVENSIS* (L. sp. 218.) caule angulato glabro, foliis sagittatis subauriculatis, ped. sæpiùs unifloris angulatis glabris, calyce submembranaceo obtuso, corollâ campanulatâ albo-rosâ, stigm. linearibus.  $\mathcal{Y}$  ubique in agris et cultis. — Fl. fr. 3. p. 640. Fl. dan. t. 459. eng. bot. t. 312. (v. v.)

9. *C. ALTRAMOIDES* (L. sp. 222.) caule hirsuto tereti elongato, foliis inferioribus ovato-cordatis sinuatis aut sublobatis, superioribus lobatis laciniosisque, ped. rectis 1-2-floris longis, calyce ovato-lanceolato inæquali, cor. speciosâ.  $\mathcal{Y}$  in Galliâ mediterraneâ ad rupes et muros. — Fl. fr. 3. p. 641. — Barr. ic. t. 312. — Lob. ic. 623. f. 2. (v. v.)

$\beta$ . *ferrugineus* (Chois.) caule tomentoso-ferrugineo. — In Galliâ meridionali. (v. s.)

II. *CRESSA* L. gen. n. 313.

Calyx 5-partitus; cor. tubulosa 5-partita; stam. exserta; ov. 2-loculare, loc. 2-spermis; styli 2; stigm. capitatum; caps. monosperma.

1. *C. CRETICA* (L. sp. 325.) caule herbaceo humili non lactescente pubescente, foliis integris adpressis, floribus apice congestis. ① in humidis montibus Gallie meridionalis. — Fl. fr. 3. p. 643. — Pluck. t. 43. f. 6. (v. v.)

III. CUSCUTA L. *gen. n.* 170.

Calyx 4-5-fidus; cor. globoso-urceolata, limbo 4-5-fido; stam. subtus squamosa; ov. 2-loculare, loc. 2-spermis; stigm. 2; capsula circumscissa; embryo acotyledoneus.

1. *C. MAJOR* (Bauh. pin. 219.) floribus subsessilibus, corollæ lobis sæpius reflexis, stylis è basi arcuatis divergentibus, stigm. acutis. ① parasitica in multis plantis, vulgaris. — Fl. fr. 3. p. 644. (v. v.)
2. *C. MINOR* (Bauh. pin. 219.) floribus sessilibus, corollæ lobis rectis, stylis apice divergentibus, stigm. acutis. ① parasitica in fruticibus permultis vulgarissima. — Fl. fr. 3. p. 644. Lam. ill. t. 88. (v. v.)
3. *C. MONOCTYMA* (Vahl. symb. 2. p. 32.) caulibus tuberculis crassis, floribus in brevi pedunculo glomerato-sessilibus, corollâ urceolatâ, lobis rectis, stylis coadunatis, stigm. globosis. ① in vinetis Occitanie inferioris. — Fl. fr. 5. p. 425. — Buxb. cent. 1. p. 15. t. 23. (v. v.)

Ordo LXVIII. BORRAGINÆ Juss. *gen.* 128.

Calyx 1-sepalus 5-lobus persistens. Corolla 5-loba sæpius regularis, fauce nudâ aut squamis 5 obtectâ, æstivatione imbricatâ. Stamina 5 corollæ adfixa et cum ejusdem laciniis alterna. Ovarium liberum 2-4-lobum disco hypogyno appositum; stylus simplex persistens inter ovarii lobos nascens; stigma integrum aut bilobum. Nuces aut cariopsides 2-4 uniloculares 1-spermæ latere interno stylo adnatæ. Albumen nullum. Embryo rectus; radícula infera, cotyledones foliaceæ. — Herbæ aut suffrutices, folia alterna sæpius aspera, sicca maculis parvis conspersa; flores in racemos unilaterales incurvos sæpè digesti.

Sect. I. NUCE 2 BILOCULARES.

I. CERINTHE L. *gen. n.* 186.

Calyx 5-partitus; corolla tubulosa ventricosa 5-fida; antheræ erectæ vix exsertæ; nuces 2 osseæ 2-loculares 2-spermæ.

1. *C. ASPERA* (Roth. cat. 1. p. 33.) foliis subciliatis asperis, corollis obtusis patulis calyce longioribus. ① in arvis provinciarum australium. — Fl. fr. 3. p. 618. (v. v.)
2. *C. MINOR* (L. sp. 196.) foliis glabris, corollis acutis clausis calyce longioribus. ② in montibus Delphin. ex Vill., Galloprovinciæ superioris. — Fl. fr. 3. p. 619. Jacq. aust. t. 124. (v. v. DC.)

Sect. II. NUCE 4 UNILOCULARES.

† *Nuces cohærentes.*

II. HELIOTROPIMUM Tourn. *inst. t.* 57. L. *gen. n.* 179.

Calyx tubulosus 5-fidus aut 5-dentatus; corolla hypocrateriformis 5-fida dentibus 5 interjectis; stigma subconicum.

1. *H. PERUVIANUM* (L. sp. 187.) caule fruticoso, foliis lanceolato-ovatis, spicis nummosis aggregato-corymbosis. ♂ ex Peruvia ortum, in hortis colitur. — Fl. fr. 3. p. 620. Mill. ic. t. 143. (v. v. c.)
2. *H. EUROPEUM* (L. sp. 187.) caule erecto herbaceo, foliis ovatis integerrimis tomentoso-scabris rugosis, spicis conjugatis, calycibus 5-fidis. ① in glareosis aridis. — Fl. fr. 3. p. 620. Jacq. aust. 3. t. 207. (v. v.)
3. *H. SUPINUM* (L. sp. 187.) caule prostrato herbaceo, foliis ovatis obsolete crenulatis plicatis subtus incanis, spicis conjugatis, calycibus 5-dentatis. ① in Galliae mediterraneae humidis rariis. — Fl. fr. 3. p. 620. Gon. monsp. p. 17. t. 1. (v. v. DC.)

†† *Nuces distinctæ, corollæ faux nuda.*

III. *ECHIUM* Tourn. inst. t. 54. L. gen. n. 191.

Calyx 5-partitus; corolla tubo brevi, limbo ampliore campanulato, obliquè 5-lobo, laciniis inæqualibus, 2 superioribus longioribus, infimâ minore acutâ reflexâ; nuces tuberculatæ.

1. *E. VULGARE* (L. sp. 200.) caule erecto tuberculato hispido herbaceo, foliis lanceolatis hispidis, floribus spicatis lateralibus, spicis axillaribus. ② ad vias et agros. — Fl. fr. 3. p. 621. Lam. ill. t. 94. f. 1. (v. v.)
2. *E. AUSTRALE* (Lam. ill. n. 1860.) caule herbaceo hispido erecto, foliis ovatis utrinquè attenuatis, floribus spicatis unilateralibus, spicis terminalibus, bracteis basi ovato-cordatis acuminatis, corollis stamina apice villosa æquantibus? ♀ propè Vindomagus (Beaucaire) reperit cl. A. Richard. — Fl. fr. 3. p. 622. (v. s.)
3. *E. PLANTAGINEUM* (L. mant. 202.) caule herbaceo villosa basi prostrato, foliis radicalibus rotundo-ovatis petiolatis, floralibus subcordatis, staminibus glabris corollâ sublongioribus. ② in agro Nicænsi et Narbonensi. — Fl. fr. 3. p. 622. Jacq. vind. x. t. 45. (v. s.)
4. *E. VIOLACEUM* (L. mant. 42.) caule hispido erecto, foliis oblongo-lanceolatis obtusis, floribus spicatis unilateralibus, corollis stamina glabra æquantibus calyce duplò longioribus. ① in Gallia australi. — Fl. fr. 3. p. 622. (v. v.)
5. *E. PYRENAICUM* (L. mant. 334.) caule pilosissimo herbaceo ramosissimo, foliis lineari-lanceolatis hispido-scabris, capitulis pedunculatis axillaribus, staminibus glabris corollâ pilosâ longioribus. ♀ in siccis Galliae australis et austro-occidentalis in Medulco (Médoc) et Santoniâ (Des Moulins). — Fl. fr. 3. p. 621. *E. asperum* Lam. *E. italicum* L? (v. v.)
6. *E. CALYCINUM* (Viv. fragm. t. 4.) caulibus herbaceis basi prostratis, foliis ovato-oblongis, floribus subsparsis, corollâ subregulari calyce vix longiore, calycibus fructiferis productioribus basi inflatis. ② in maritimis Nicæ, Massiliæ (cl. Solier). — Fl. fr. 5. p. 419. *E. prostratum* Tenor. nap. 1. t. 12. (v. s.)

IV. *LITHOSPERMUM* Tourn. inst. t. 55. L. gen. n. 181.

Calyx 5-partitus; corolla infundibuliformis parva 5-loba; antheræ oblongæ in fauce corollæ inclusæ; stigma obtusum bifidum; nuces nitidæ aut glabræ, læves aut rugosæ.

1. *L. FRUTICOSUM* (L. sp. 190.) caule fruticoso, foliis linearibus hispidis margine revolutis, calyce corollâ multò brevior, staminibus corollam æquantibus, nucibus nitidis lævibus. ♂ in Galliae australioris aridis. — Fl. fr. 3. p. 625. — Garid. six t. 15. (v. s.)
2. *L. PROSTRATUM* (Lois. gall. 105. t. 4.) suffruticosum procumbens, foliis lanceolato-linearibus pilosis margine subrevolutis, corollis puberulis calyce



- duplò longioribus, staminibus sauci villosæ æqualibus.  $\mathcal{L}$   $\S$  circa Bayonam, Aquas Tarbellicas et in Armoraciâ extremâ. — Fl. fr. 5. p. 419. (v. v. D.)
3. *L. OLEAFOLIUM* (Lapeyr. abr. supp. p. 28.) suffrutescens, caulibus diffusis, foliis elliptico-oblongis obtusis suprâ viridibus pilis sparsis brevibus adpressis subtus sericeo-argenteis, corollis calyce duplò longioribus extus villosis stamina superantibus.  $\S$  in rupestribus Pyren. orientalium. Rami incani. (v. s. D.)
4. *L. TINCTORIUM* (L. sp. 1. p. 132.) caulibus decumbentibus, foliis lanceolatis obtusis, floralibus subcordatis, floribus axillaribus, corollis calyce longioribus, staminibus inclusis, seminibus tuberculosi.  $\mathcal{L}$  in Galliæ australis arenosis sterilioribus. — Fl. fr. 3. p. 624. — J. Bauh. hist. 3. p. 584. ic. (v. v.)
5. *L. PURPUREO-CERULEUM* (L. sp. 190.) caulibus floralibus erectis, sterilibus prostrato-repentibus, foliis lanceolatis acutis, floribus axillaribus, nucibus lævissimis nitidis.  $\mathcal{L}$  in nemoribus et agris passim. — Fl. fr. 3. p. 624. Jacq. aust. t. 14. (v. v.)
6. *L. OFFICINALE* (L. sp. 189.) caule erecto, foliis lanceolatis acuminatis, corollis calycem vix superantibus, nucibus lapideis nitidis subtuberculatis.  $\mathcal{L}$  in ruderatis. — Fl. fr. 3. p. 623. Lam. ill. t. 91. (v. v.)
7. *L. AUREUM* (L. sp. 190.) caulibus erectis, foliis lanceolatis margine revolutis, corollis vix calycem superantibus, nucibus rugosis. (I) in agris et arvis. — Fl. fr. 3. p. 623. — Moris. s. 11. t. 31. n. 3. (v. v.)
8. *L. APULUM* (Vahl. symb. 2. p. 33.) caulibus erectis, foliis acutis, caulinis linearibus, radicalibus lineari-lanceolatis, spicis terminalibus secundis, bracteis lanceolatis, nucibus nitidis in costis tuberculatis. (I) in Galliæ mediterraneæ maritimis. — Fl. fr. 3. p. 624. — Lob. ic. 578. f. 1. (v. v.)
- V. *NONEA* Medik. philos. bot. 1. p. 31. DC. fl. fr. 3. p. 626. — *Lycopsidis* sp. L.

Calyx 5-fidus fructifer inflatus; corolla tubo erecto limbo 5-lobo æquali; stamina intra tubum; nukes sulcatæ.

1. *N. VIOLACEA* (DC. l. c.) canle prostrato hispido, foliis oblongo-lanceolatis integerrimis. (I) circa Monspelium ex Gouan, in agro Ruscionensi ex Lapeyr. — *Lycopsis vesicaria* L. sp. 198. — *Buglossum* II. Moris. s. 11. t. 26. f. 10. (v. s. c.)
2. *N. ALBA* (DC. fl. fr. 5. p. 420.) canle erecto basi ramoso, foliis integris oblongo-linearibus, pilis rigidis eglandulosis. (Y) inter segetes secus Rhodanum infrâ Avenionem, propè Agde ex cl. Benth. — Fl. albi. (v. s.)
3. *N. LUTEA* (DC. fl. fr. 3. p. 626. in adn.) pilis aliis rigidis longis acutis, aliis brevioribus apice glandulosis, calycibus ultra medium divisis. (I) in insulis Stæchadum et agro Ruscionensi propè Prato de Mollo ex cl. Benth. — Fl. fr. 5. p. 420. Fl. lutei. (v. s. spec. imperf.)

VI. *PULMONARIA* Tourn. t. 55. L. gen. n. 184.

Calyx campanulatus 5-fidus 5-gonus; corolla infundibuliformis sub 5-loba erecto-patens; stigma obtusum emarginatum.

1. *P. OFFICINALIS* (L. sp. 194.) foliis radicalibus ovato-cordatis maculatis, calyce vix 5-fido, nucibus lucidis lævissimis lenticulari-convexis submarginatis.  $\mathcal{L}$  in nemoribus. — Fl. fr. 3. p. 627. — Moris. s. 11. t. 29. n. 4. (v. v.)
- $\beta$ . *mollis*, foliis immaculatis. — *P. mollis* Schrad., et Fl. fr. 5. p. 420. (v. v.)
2. *P. ANGSTIFOLIA* (L. sp. 194.) foliis radicalibus lanceolatis immaculatis in petiolum attenuatis, calyce vix 5-fido.  $\mathcal{L}$  in sylvis montosis. — Fl. fr. 3. p. 627. Fl. dan. t. 483. (v. v.)

## VII. ONOSMA Tourn. L. gen. n. 187.

Calyx 5-fidus; corolla tubulosa limbo erecto sub 5-lobo; antheræ sagittatæ basi lobis connexæ; stigma simplex; nuces ovatæ lucidæ.

1. *O. xchioides* (L. sp. 196.) foliis lanceolatis hispidis, corollis cylindricis obtusis, fructibus erectis.  $\frac{1}{2}$  in Galliæ australis aridis glareosis. — Fl. fr. 3. p. 627. Lam. ill. t. 93. (v. v.)

+++ *Nuces distinctæ, corollæ faux squamis instructa.*

•

## VIII. SYMPHITUM Tourn. t. 56. L. gen. n. 185.

Calyx 5-fidus; corolla cylindrico-campanulata tubo brevissimo, limbo tubulato-ventricoso 5-lobo lobis brevibus, squamæ subulatæ conniventes.

1. *S. tumerosum* (L. sp. 195.) foliis semidecurrentibus, inferioribus in petiolum angustatis, limbi laciniis brevissimis obtusis.  $\frac{1}{2}$  in humidis prov. austral., agri Mimatensis (Prost), Pyrenæorum ex Lapeyr. — Fl. fr. 3. p. 628. Jacq. aust. t. 225. — Moris. s. 11. t. 29. n. 2. (v. s.)
2. *S. officinale* (L. sp. 195.) foliis ovato-lanceolatis omnibus sessilibus decurrentibus.  $\frac{1}{2}$  in pratibus, incultis, et ad sepes. — Fl. fr. 3. p. 628. Fl. dan. t. 664. — Moris. s. 11. t. 29. n. 1. (v. v.)

IX. LYCOPSIS Desf. cat. par. 73. DC. fl. fr. 3. p. 634. — *Lycopsidis* sp. L.

Calyx 5-fidus, corolla infundibuliformis 5-loba erecto-patens tubo incurvo; squamulæ ad faucem ovatæ prominulæ conniventes; stigma emarginatum, nuces basi insculptæ.

1. *L. arvensis* (L. sp. 199.) foliis lanceolatis hispidis, calycibus florentibus erectis.  $\frac{1}{2}$  in arvis, ruderatis. — Fl. fr. 3. p. 634. Lam. ill. t. 92. *Anchusa arvensis* Lehms. (v. v.)

## X. ANCHUSA L. gen. 182.

Calyx 5-fidus; corolla infundibuliformis 5-loba erecto-patens, lobis integris; cætera *Lycopsidis*.

1. *A. italica* (Retz obs. 1. p. 12.) caule exaltato, foliis lucidis strigosis oblongis utrinque attenuatis hispidis, calyce hispido sub-5-partito, bracteis linearilanceolatis, floribus fauce barbatis.  $\frac{1}{2}$  in ruderatis, cultis. — Fl. fr. 3. p. 631. Trew. dec. 2. p. 14. t. 13. *A. officinalis* Gon. an L? (v. v.)
2. *A. angustifolia* (L. sp. 191.) foliis oblongo-lanceolatis integris, floribus spicatis, calycibus 5-fidis, bracteis linearilanceolatis.  $\frac{1}{2}$  in Galliæ australi et occidentali. — Fl. fr. 3. p. 632. — Lob. ic. 676. f. 2. (v. v.)
3. *A. undulata* (L. sp. 191.) strigosa, foliis lanceolato-linearibus sinuatis, pedicellis bracteis lanceolato-ovatis brevioribus, corollæ laciniis angustis, calycibus fructiferis inflatis.  $\frac{1}{2}$  in arvis Monspelii. — Fl. fr. 3. p. 633. Mill. dict. t. 59. (v. v. DC.)
4. *A. sempervirens* (L. sp. 192.) foliis hispidis, inferioribus ovatis subsinuatis petiolatis, superioribus sessilibus lanceolatis, pedunculis diphyllis capitatis, calyce hispido.  $\frac{1}{2}$  in arvis cultis propè Dax ex Thore, Bardigalam ex Latterade, Nannetes ex Bon., Maclovii ex cl. Petit-Thouars. — Fl. fr. 3. p. 633. — Lob. ic. 755. f. 2. (v. v. c.)

## XI. BORRAGO Tourn. t. 53. L. gen. n. 188.

Calyx 5-partitus; corolla rotata 5-fida plerumque patens; squamulæ obtusæ emarginatæ; fructus rugosi, calyce connivente tecti.

1. *B. OFFICINALIS* (L. sp. 197.) hispida, foliis radicalibus petiolatis, superioribus sessilibus, pedunculis ramosis. ① in hortis et cultis. — Fl. fr. 3. p. 638. — Black. t. 36. (v. v.)
2. *B. LAXIFLORA* (Desf. cat. par. DC. fl. fr. 5. p. 422.) hispidula, caulibus procumbentibus, foliis radicalibus oblongis basi in petiolum angustatis, caulinis sessilibus, pedicellis distantibus unifloris. ½ in Corsicâ. — Anch. laxiflora Fl. fr. 3. p. 631. (v. s.)  
*β. nana*, caule brevissimo unifloro. — Campanula pygmaea fl. fr. 3. p. 705. ips. obs. DC. in herb. Lam. nunc Rœper.

XII. ASPERUGO Tourn: t. 54. L. gen. n. 189.

Calyx 5-fidus inæqualis dentibus interjectis; corolla tubo brevi 5-lobo; squamulæ convexæ conniventes; semina calyce conduplicato et compresso tecta.

1. *A. PROCUMBENS* (L. sp. 198.) floribus axillaribus, calycibus post maturationem dilatatis. ① in ruderatis pinguibus. — Fl. fr. 3. p. 634. — Lam. ill. t. 54. — Buglossum 13. Moris. s. 11. t. 26. f. 11. (v. v.)

XIII. MYOSOTIS L. gen. n. 180.

Calyx 5-fidus aut 5-partitus; corolla hypocrateriformis tubo brevi, limbo plano 5-lobo lobis emarginatis; squamulæ convexæ conniventes; nuces læves aut scabræ.

Sect. I. LAPPULA Mœnch. met. 417.

Nuces echinatæ.

1. *M. LAPPULA* (L. sp. 189.) caulibus erectis, foliis lanceolatis pilosis, calycibus 5-partitis, lobis per maturationem patulis. ① in argillosis, ruderatis. — Fl. fr. 3. p. 630. Lam. ill. t. 91. (v. v.)

Sect. II. ECHIOIDES Mœnch. met. 416.

Nuces læves.

2. *M. LUTEA* (Pers. ench. 1. p. 156.) caule herbaceo erecto simplici, foliis oblongo-lanceolatis alternis pilosis, inferioribus spathulatis, calycibus campanulatis 5-partitis hirsutis tubo corollæ subbrevioribus, nucibus triquetris pellucidis lævissimis. ① in pinetis circâ *S.-Bonet-le-Froid* propè Lugdunum. (D. Foudras ex cl. Balbis.) — Anchusa lutea Cav. ic. 1. t. 69. f. 1. Fl. Intei. (v. s.)
3. *M. ARVENSIS* (Mœnch. hass. n. 153.) caule ramoso foliisque oblongis pilosis, calycibus campanulatis 5-partitis hirsutis tubo corollæ longioribus, nucibus lævibus. ① in pratis et arvis. — Fl. fr. 3. p. 629. *M. arvensis* Bull. herb. t. 355. (v. v.)
4. *M. FUSILLA* (Lois. not. 36. t. 1 f. 2.) radice exili, caulibus ramosis brevissimis pilosis, ramis decumbentibus, foliis inferioribus ovato-spathulatis superioribus oblongo-lanceolatis, calycibus hirsutis 5-fidis, corollis minimis, nucibus lævibus. ① in agris Corsicæ, Massiliæ (Solier). — Fl. fr. 5. p. 421. (v. s.)
5. *M. PERENNIS* (Mœnch. hass. n. 154.) caule foliisque lanceolatis hispidis, calycibus campanulatis 5-fidis pilosis tubo corollæ brevioribus, nucibus lævibus. ½ in rivulis, humidis inque sylvaticis. — Fl. fr. 3. p. 629. (v. v.)
6. *M. NANA* (Vill. dauph. 2. t. 13.) cespitosa, caulibus simplicibus minimis paucifloris, foliis lanceolato-oblongis villosis, calycibus hirsutissimis 5-partitis, laciniis obtusiusculis, nucibus margine crenulatis. ½ in editioribus Alpibus Delph. et Galloprov. — Fl. fr. 3. p. 630. (v. v. in Helv.)

XIV. CYNOGLOSSUM *L. gen. n. 183.*

Calyx 5-partitus; corolla brevis infundibuliformis 5-loba; squamulae convexae conniventes; stigma emarginatum; nuces depressae interiore latere stylo affixae.

## Sect. I. EUGLOSSUM.

Nuces planae echinatae.

1. *C. OFFICINALE* (L. sp. 192.) caule erecto, foliis lanceolatis basi attenuatis sessilibus tomentosis, laciniiis calycinis ovato-lanceolatis obtusis, staminibus corollâ brevioribus. ① in sylvaticis, sterilibus passim. — Fl. fr. 3. p. 635. Lam. ill. t. 92. f. 1. (v. v.)
2. *C. MONTANUM* (Lam. fl. fr. 2. p. 277.) foliis lanceolatis lucidis glabriusculis subtus scabris, inferioribus petiolatis, calycis hispiduli laciniiis lanceolatis obtusis corollam subaequantibus, staminibus corollâ brevioribus. ② in Alpibus et Pyrenæis. — Fl. fr. 3. p. 635. (v. s.)
3. *C. PICTUM* (Ait. kew. 1. p. 179.) caulibus erectis apice ramosis, foliis lanceolatis tomentosis, superioribus basi cordatis, corollis calyci subaequalibus venosis, fructibus convexis. ② in aridis provinc. austral. et occidentalium. — Fl. fr. 3. p. 636. — Clus. hist. 2. p. 162. f. 2. (v. v.)
4. *C. CARIIVOLIUM* (L. sp. 193.) caulibus à basi ramosis, foliis incano-sericeis lanceolatis obtusis, radicalibus apathulatis, staminibus corollâ brevioribus, calycis lanuginosi laciniiis lanceolatis corollâ duplò brevioribus. ② in provinciis australibus. — Fl. fr. 3. p. 636. (v. v.)

Sect. II. OMPHALODES *Tourn. t. 58.*

Nuces laeves calathiformes, margine membranaceo internè dentato.

5. *C. LINIFOLIUM* (L. sp. 193.) foliis lineari-lanceolatis glabris margine denticulato-scabris. ① in sterilibus Galliae occidentalis, Galloprovinciae propè Carpentras et Lotharingiae propè Nancy (Monnier). — Fl. fr. 3. p. 638. — Barr. ic. t. 1234. (v. s.)
6. *C. OMPHALODES* (L. sp. 193.) foliis subglabris petiolatis, radicalibus ovato-cordatis, caulinis ovatis. ② Pedemontii nemorosis indigena, frequenter in hortis culta. — Fl. fr. 3. p. 637. — Moris. s. 11. t. 26. n. 3. (v. v. c.)

## Ordo LXIX. SOLANEÆ Juss. gen. 124.

Calyx monosepalus æqualis sæpius persistens, 5-fidus aut partitus. Corolla 1-petala sæpius regularis 5-fida decidua, æstivatione plicatâ. Stamina 5 basi corollæ inserta, ejusdem lobis alterna. Ovarium liberum simplex; stylus 1; stigma simplex aut bisulcum. Fructus bilocularis polyspermus, nunc capsularis bivalvis, dissepimento valvis parallelo, nunc baccatus receptaculis seminiferis centralibus. Semina numerosa; albumen carnosum; embryo annularis vel spiralis; cotyledones semiteretes elongatæ. — Herbæ aut rarius frutices, foliis alternis simplicibus lobatisve, floribus sæpè extrâ-axillaribus; interdum quinta pars floris demitur.

† *Fructus baccatus.*

## I. LYCIUM L. gen. n. 262.

Calyx brevis tubulosus; corolla infundibuliformis tubo brevis, limbo 5-fido; filamenta 5 basi villosa; stigma sulcatum; bacca subrotunda; embryo uncinatus subcentralis.

1. *L. BARBARUM* (L. sp. 192.) foliis oblongis glabris, ramis flexuosis nutantibus, calycibus 2-labiatis. ♀ in hortis et in sepibus aliquandò quasi spontaneum, in agro Ruscimonensi ex Benth. — Fl. fr. 3. p. 616. (v. v. c.)
2. *L. EUNOMÆUM* (L. mant. 47.) foliis oblongis glabris, ramis flexuosis, calycibus quinquedentatis. ♀ in Gallia australi. — Fl. fr. 3. p. 616. — Mich. gen. t. 106. f. 1. (v. v. c.)

## II. CAPSICUM Tourn. t. 66. L. gen. n. 252.

Calyx 5-fidus; corolla rotata tubo brevissimo, limbo 5-fido patente; antheræ 5 erectæ conniventes longitudinaliter dehiscentes; bacca exsucca; embryo semicircularis.

1. *C. ANNUUM* (L. sp. 270.) caule herbaceo, pedunculis solitariis. (1) ex America meridionali ortum, in hortis cultum. — Fl. fr. 3. p. 615. Lam. ill. t. 116. f. 1. (v. v. c.)

## III. SOLANUM Dunal. sol. p. 115. — Solani sp. L.

Calyx monosepalus persistens 5-10-divisus; corolla monopetala rotata, tubo brevissimo, limbo 4-5-6-diviso patente; antheræ 4-5-6 oblongæ apice poris duobus dehiscentes; bacca subrotunda 2-3-4-6-locularis; embryo spiralis.

§. 1. *Caulibus foliisque aculeatis.*

1. *S. ACULENTUM* (Dun. sol. p. 208. t. 3. f. E.) caule aculeato subherbaceo, foliis ovatis subsinnatis tomentosis aculeatis, floribus multipartitis, baccis 2-3-locularibus, seminibus nudis. (1) ex India ortum, in hortis Galliarum australis cultum. — *S. Melongena* L. sp. 266. Fl. fr. 3. p. 615. — Pluck. phyt. t. 226. f. 2. — Moris. s. 13. t. 2. n. 2. (v. v. c.)

§. 2. *Caule foliisque inermibus.*

\* *Foliis angulato-repandis integrisve.*

2. *S. VILLOSUM* (Lam. dict. 4. p. 289.) caule herbaceo tereti villoso, foliis ovatis angulato-dentatis villosis canescentibus, floribus subumbellatis. (1) in

*caltia*. — Fl. fr. 3. p. 613. — Dill. elth. t. 274. f. 353. *Bacca croceo-rubra*. (v. v.)

3. *S. OCHROLEUCUM* (Bast. journ. bot. 1814. 3. p. 20.) caule herbaceo erecto, ramis angulatis dentatis pubescentibus, foliis ovato-oblongis angulato-sinuatibus subhirsutis, floribus subumbellatis. ① circa Andegavum. — Fl. fr. 5. p. 418. *Baccæ* è flavo et viridi variz. (v. s.)
4. *S. HUMILE* (Wild. enum. p. 236.) caule herbaceo patulo, ramis subangulatis subdentatis subpubescentibus, foliis ovatis repandis superioribus subintegris, floribus subumbellatis. ① in ruderatis agri Aurelianensis et Andegavensis. — Fl. fr. 5. p. 417. *Bacca* luteo-virescens. (v. s.)
5. *S. MINIATUM* (Wild. enum. 236.) caule herbaceo, ramis pubescentibus angulato-alatis alia dentatis, foliis ovatis repandis glabriusculis, floribus subumbellatis. ① in ruderatis Parisiorum ex Tourn., agri Aurelianensis et Andegavensis, in sylvis Vasconiz prope *la Teste* (Des Moul.), circa Monsapellum ex Benth. — Fl. fr. 5. p. 417. *Baccæ* pallide rubræ. (v. s.)
6. *S. NIGRUM* (Wild. en. p. 236.) caule herbaceo angulato, foliis ovatis dentatis nudiusculis, floribus subumbellatis. ① in hortis et ruderatis. — Fl. fr. 3. p. 613. *S. nigrum* æ L. sp. 266. *Baccæ* nigræ. — An tres præcedentes species ritè ab hac distinguuntur? (v. v.)
7. *S. PSEUDO-CAPSICUM* (L. sp. 263.) caule fruticoso, foliis oblongo-lanceolatis subrepandis glabris, pedunculis subunifloris extrafoliaceis. § in hortis cultum, nunc in Bearniæ pago *Arette* subsponsaneum. — Fl. fr. 5. p. 417. — Sabb. h. rom. t. 59. (v. v. DC.)

\* \* *Foliis lobatis integrisve.*

8. *S. DULCAMARA* (L. sp. 264.) caule fruticoso flexuoso, foliis ovato-cordatis glabris, superioribus laciniatis, corymbis suboppositifoliis. ① in sepibus. — Fl. fr. 3. p. 612. — Lob. ic. t. 266. f. 1. (v. v.)

\* \* \* *Foliis impari-pinnatisectis.*

9. *S. TURKOSUM* (L. sp. 265.) caule herbaceo, foliis impari-pinnatisectis segmentis inæqualibus, alternis minutis, pedicellis articulatis, corollis 5-angulatis. ② ex Amer. austr. ortum, ubique colitur. — Fl. fr. 3. p. 613. — Clus. hist. 2. p. 79. ic. *Varietates innumeræ*. (v. v. e.)

#### IV. LYCOPERSICUM *Tourn. inst. 150. t. 63. Dun. mon. sol. p. 109. — Solani sp. L.*

Calyx monosepalus persistens 5-6-partitus; corolla monopetala 5-6-fida; antheræ conicæ membranâ apice elongatâ connatæ, intus longitudinaliter dehiscentes; semina villosa.

1. *L. ESCULENTUM* (Dun. sol. p. 113. t. 3. f. 3.) pilosum pilis diversis, foliis inæqualiter pinnatisectis, segmentis incisus subtus glaucescentibus, floribus pluribus coalitis, baccois torulosis. ② ex America? ortum, colitur in hortis. — Sol. Lycopersicum L. sp. 265. Fl. fr. 3. p. 615. — Cam. epit. 821. ic. (v. v. c.)

V. *PHYSALIS* L. *gen. n. 250.*

Calyx 5-fidus, corolla rotata 5-fida; antheræ 5 oblongæ erectæ conniventes; bacca globosa, calyce ampliato vesicario tecta.

1. *P. ALKERRINGI* (L. sp. 262.) foliis geminis integris acutis, caule herbaceo. ② in humidis, umbrosis, scrobibus passim. — Fl. fr. 3. p. 612. — Lob. ic. t. 262. f. 2. (v. v.)

#### VI. ATROPA *Certh. 2. p. 240. t. 131. DC. fl. fr. 3. p. 611. — Atropæ sp. L.*

Calyx campanulatus 5-fidus; corolla campanulata calyce duplò longior 5-loba æqualis; filamenta 5 filiformia; bacca globosa calyci insidens.

1. A. *BELLADONNA* (L. sp. 260.) foliis ovalibus integerrimis, floribus axillaribus breviter pedunculatis. 2/ in sylvaticis montanis. — Fl. fr. 3. p. 611. — Lob. ic. t. 263. f. 1. (v. v.)

++ *Fructus capsularis.*

VII. *DATURA* L. gen. n. 246.

Calyx magnus tubulosus ventricosus 5-angularis apice 5-fidus caducus, basi orbiculatâ peltatâ persistente; corolla maxima infundibuliformis tubo longo limbo 5-angulari 5-plicato 5-acuminato; stamina 5; stigma bilamellatum; capsula echinata aut lævis ovata 2-locularis loculis dissepimento prominente bi aut multipartitis.

1. D. *STRAMONIUM* (L. sp. 255.) foliis ovatis sinuatis glabris basi in petiolum attenuatis, pericarpis ovatis spinosis. ① in ruderatis. — Fl. fr. 3. p. 609. — Gard. aix. t. 88. (v. v.)  
2. D. *TATULA* (L. sp. 255.) foliis cordatis glabris dentatis, pericarpis ovatis spinosis. ① prope Burdigalam ex cl. Bory, Perpignanum ex Lapeyr. et Benth., inter Avenionem et Nemausim ex cl. Requier. (v. v. c.)

VIII. *NICOTIANA* Tourn. t. 41. L. gen. n. 248.

Calyx urceolatus 5-fidus; corolla multò longior infundibuliformis 5-fida regularis; stamina 5; stigma emarginatum; capsula 2-valvis.

1. N. *TASACUM* (L. sp. 258.) foliis sessilibus oblongo-lanceolatis acuminatis, inferioribus decurrentibus, corollæ fauce inflato-ventricosâ, limbi laciniis acuminatis, ① ex Americâ anno 1560 allata, in prov. variis præsertim in Galliâ australi et Alatiâ colitur. — Fl. fr. 3. p. 608. — Lob. ic. t. 584. f. 2. (v. v. c.)  
2. N. *RUSTICA* (L. sp. 258.) caule tereti, foliis petiolatis ovatis integerrimis, corollæ tubo cylindrico calyce longiore, limbi laciniis rotundatis obtusis. ① culta in Galliâ australi. — Fl. fr. 3. p. 609. Bull. herb. t. 289. (v. v. c.)

IX. *HYOSCYAMUS* Tourn. t. 42. L. gen. n. 247.

Calyx tubulosus 5-fidus; corolla infundibuliformis limbo patente obliquè 5-lobo inæquali; stamina 5; stigma capitatum; capsula ovata utrinquè compressa et sulco exarata, apice circumscissa seu operculata.

1. H. *NIGRA* (L. sp. 257.) foliis amplexicaulibus acute sinuatis, floribus sessilibus. ① in ruderatis. — Fl. fr. 3. p. 607. — Lob. ic. t. 268. f. 2. (v. v.)  
2. H. *ALBUS* (L. sp. 257.) foliis caulinis petiolatis cordatis obtusè sinuatis acutis, floralibus integerrimis, floribus subsessilibus, corollis ventricosis. ① in Galliâ mediterraneâ. — Fl. fr. 3. p. 607. Lam. ill. t. 117. f. 2. — Clus. hist. 2. p. 84. f. 1. (v. v.)  
3. H. *AUREUS* (L. sp. 257.) foliis petiolatis dentatis acutis, floribus pedunculatis, fructibus pendulis. ② in ruderatis Nicæ ex All., Mompeliis ex Gouan., agri Narbonensis ex cl. Benth. — Fl. fr. 3. p. 608. — C. Bauh. prod. p. 92. ic. (v. s. c.)

X. *RAMONDIA* Rich. in Pers. ench. 1. p. 216. DC. fl. fr. 3. p. 606.

Calyx 5-partitus; corolla 5-loba subinæqualis basi ad loborum sinus hirsuto-maculata; stamina 5 approximata, antheris apice perforatis; caps. 1-locularis 2-valvis, valvulis margine introflexis septiferis.

1. R. *PYRENEICA* (Rich. 1. c.) foliis radicalibus supernè villosis, subtus laugine fuscâ obtectis. 2/ in Pyrenæis. — Fl. fr. 3. p. 606. Verbascum Myconi L. Mill. ic. t. 277. Chaixia Myconi Lapeyr. abr. supp. 39. (v. v.)

XI. *VERBASCUM* L. gen. n. 245.

Calyx 5-partitus; corolla rotata 5-loba inæqualis; stamina 5 inæqua-

lia, filamentis inclinatis ut plurimum basi villosis; capsula 2-valvis ovata aut globosa.

§. 1. *Foliis decurrentibus.*

1. *V. THAPSUS* (L. sp. 252.) caule simplici, foliis lanceolato-ovatis subintegris utrinque lanato-tomentosis decurrentibus, floribus spicatis subsessilibus spica densa, bracteolis calycem lanatum superantibus, laciniis calycinis lanceolatis capsulam subtomentosam subæquantibus. ② in glareosis. — Fl. fr. 3. p. 600. — Lob. ic. t. 561. f. 2. — Fuchs. t. 346. ic. (v. v.)
2. *V. THAPSIFORME* (Schröd. verb. 1. p. 21.) caule simplici, foliis lanceolato-ovatis decurrentibus crenulatis tomentosis superioribus acuminatis, racemo spicato denso, bracteis calycem lanatum superantibus, corollæ laciniis obovatis rotundatis, antheris 2 oblongis. ② in sterilibus Parisiorum, Andegaviæ, Monspeli. — Fl. fr. 5. p. 412. An satis distincta species? (v. s.)
3. *V. AUSTRALE* (Schröd. mon. 1. p. 28. t. 2.) caule subsimplici, foliis lanuginosis crenatis, radicalibus oblongo-lanceolatis in petiolum attenuatis, caulinis oblongis acutis decurrentibus, superioribus lato-ovatis cuspidatis semidecurrentibus, floribus breviter pedicellatis fasciculatis, fasciculis remotiusculis in spicam elongatam laxam dispositis, calycis lanati laciniis lanceolato-ovatis capsulam tomentosam subæquantibus. ② Monspeli in sterilibus. — Fl. fr. 5. p. 413. (v. v.)
4. *V. PHLOMORDES* (L. sp. 253.) foliis subdecurrentibus ovatis acutis utrinque lanuginoso-tomentosis crenatis, inferioribus in petiolum attenuatis, floribus subsessilibus fasciculatis, fasciculis subremotis in spicam aliam basi laxam dispositis, calycis lanati laciniis lanceolato-ovatis, filamentis staminum barbatis flavescens. ② in ruderatis et ad margines viarum. — Fl. fr. 3. p. 601. — Lob. ic. t. 561. f. 1. (v. v.)
5. *V. CRASSIFOLIUM* (DC. fl. fr. 3. p. 601 non Schröd.) caule subsimplici, foliis subintegris lanuginoso-tomentosis inferioribus petiolatis, floribus sessilibus subfasciculatis in spicam densam dispositis, calycis lanati laciniis lanceolatis capsulam superantibus, filamentis staminum glabris. ② in Galliâ Narbonensi et circa Suessones. — Dalech. hist. 1361. ic. — J. Bauh. hist. 3. p. 872. ic. (v. v.)
6. *V. CANDIDISSIMUM* (DC. fl. fr. 5. p. 413.) caule ramoso, foliis decurrentibus lanuginoso-tomentosis candidissimis crenulatis, inferioribus oblongo-lanceolatis acutis, superioribus ovato-acuminatis, racemo paniculato, floribus breviter pedicellatis fasciculatis fasciculis remotis, calycis lanati laciniis linearilanceolatis capsulâ brevioribus. ② in sterilibus Monspeli. (v. v.)
7. *V. SINUATUM* (L. sp. 254.) caule ramoso, foliis radicalibus oblongis pinnatifido-repandis tomentosis, caulinis subdecurrentibus amplexicaulis rameisque cordatis nudiusculis, racemo paniculato, floribus sessilibus fasciculatis, fasciculis remotis, laciniis calycinis lanceolato-ovatis. ② in aridis ad margines viarum in Galliâ australi et austro-occidentali usque ad Burdigalam ex cl. Laterr. — Fl. fr. 3. p. 605. — Cam. epit. p. 882. ic. (v. v.)

§. 2. *Foliis non decurrentibus.*

8. *V. LEPTOSTACHYON* (DC. fl. fr. 5. p. 415.) caule simplici, foliis oblongis acutis integris tomento crasso adhærente albidis, inferioribus in petiolum attenuatis, superioribus semiaplexicaulis, floribus subsessilibus fasciculatis, fasciculis in racemum simplicem elongatum dispositis, laciniis calycinis lanceolato-ovatis. ② in agris lapidosis Monspeli. (v. s.)
9. *V. MAYALE* (DC. fl. fr. 5. p. 415.) foliis supernè virescentibus subrè pulverulento-tomentosis albidis crenulatis oblongo-lanceolatis acutis inferioribus petiolatis, floribus sessilibus subsolitariis, racemo laxo spicato simplici, calycis hirsuti laciniis linearibus capsulâ brevioribus, staminibus 2 glabris longioribus. ② in sterilibus Pyren. orient., Occitanæ, Galloprovinciæ, Niceæ, Corsicæ. (v. v.)



10. *V. MIXTUM* (Ram. in DC. fl. fr. 3. p. 603.) foliis oblongis acutis obsolete crenatis sessilibus pulverulento-tomentosis, inferioribus subpetiolatis, ramis paniculæ villosis, filamentis barbatis violaceis. — In Pyrenæis. An à *V. mayali* distinctum? †
11. *V. BLATTARIA* (L. sp. 254.) caule ramoso, foliis amplexicaulibus crenatis oblongis glabris, radicalibus sinuatis, superioribus acuminatis, floribus longè pedicellatis solitariis remotis in racemum elongatum ramosum dispositis, calycis glandulosi laciniis lanceolato-linearibus capsulâ glabriusculâ brevioribus. ② in argillosis ad ripas. — Fl. fr. 3. p. 604. — Lob. ic. t. 564. f. 2. (v. v.)
12. *V. BLATTARIOIDES* (Lam. dict. 4. p. 225.) foliis oblongis amplexicaulibus crenatis glabriusculis, radicalibus sinuato-crenatis, floribus fasciculatis fasciculis 2-4-floris remotis, calycis glanduloso-villosi laciniis lanceolato-linearibus capsulâ puberulâ brevioribus. ② ad vias passim. — Fl. fr. 3. p. 605. — Lob. ic. 564. f. 1. An præcedentis var? (v. s.)
13. *V. CHAIXI* (Vill. dauph. 2. t. 13.) foliis oblongis cœrdatis petiolatis dentatis superne glabriusculis subtus villosis, inferioribus basi inciso-dentatis aut pinnatifido-incisis, floribus pedicellatis fasciculatis, fasciculis remotis in racemum ramosum paniculatum dispositis, calycis parvi lanuginosi laciniis linearibus apice interdum subglabris. ② in Delph. ex Villars, Galloprovinciâ propè Massiliam (Solier), Occitaniâ propè Monspelium et Pyrenæis orientalibus. — Fl. fr. 3. p. 605. — *V. dentatum* Lapeyr. Fl. fr. 5. p. 416. (v. s.)
14. *V. NIGRUM* (L. sp. 253.) foliis oblongo-cordatis petiolatis crenatis superne pubescentibus glabriusculisve, subtus tomentoso-candiantibus, floribus pedicellatis fasciculatis, racemo simplici, calycis minimi lanuginosi laciniis lineari-lanceolatis capsulâ tomentosâ brevioribus, filamentis barbatis purpurascensibus. ② ad vias. — Fl. fr. 3. p. 603. — Fuchs. hist. p. 849. ic. (v. v.)
15. *V. RAMOSISSIMUM* (Bast. supp. 42.) foliis oblongis acutis crenatis utrinquè villosis semiamplexicaulibus, caule tenuiter velutino, floribus pedicellatis fasciculatis in paniculam ramosam dispositis, calycibus parvis pubescentibus. ② in collibus et agris Andegaviæ. — Fl. fr. 5. p. 416. An distincta species? †
16. *V. ALOPHURUS* (Thuil. par. II. 1. p. 110.) foliis ovatis subincanis crenatis inferioribus petiolatis, floribus pedicellatis fasciculatis, fasciculis remotis, racemo simplici, calycis parvi subglabri laciniis linearibus capsulâ tomentosâ brevioribus. ② in aridis circa Lutetiam, Mimatem (Prost), et in Andegaviâ ex Bast. — Fl. fr. 3. p. 603. An ut vult cl. Merat *V. nigri* var? (v. s.)
17. *V. LYCHNITIS* (L. sp. 253.) caule angulato paniculato, foliis ovatis crenatis superne subglabris subtus tomentosis, floribus pedicellatis fasciculatis, fasciculis remotiusculis, calycis parvi laciniis apice glabris linearibus capsulâ tomentosâ brevioribus. ② ad vias in montosis sterilibus. — Fl. fr. 3. p. 602. Fl. dan. t. 586. (v. v.)
18. *V. FLOCCOSUM* (Walds. et Kit. 1. t. 79.) foliis superne pubescentibus subtus tomento caduco floccoso albis acutis, inferioribus oblongis subpetiolatis superioribus ovatis, racemo paniculato, calycis laciniis linearibus acutis pedicellisq; vix tomentosis. ② in sterilibus Lutetiæ ex Merat. — Fl. fr. 5. p. 414. An satis à sequente distinctum? (v. v. DC. in Italiâ.)
19. *V. PULVERULENTUM* (Vill. dauph. 2. p. 490.) caule tereti paniculato, foliis ovato-oblongis subserratis utrinquè pulverulento-tomentosis, floribus pedicellatis fasciculatis, fasciculis subremotis, calycis parvi pulverulenti laciniis linearibus capsulâ brevioribus. ② in ruderalis ad margines viarum passim. — Fl. fr. 3. p. 602. (v. v.)
20. *V. LONGIFOLIUM* (DC. fl. fr. 5. p. 414.) foliis oblongis acutis angustis integris tomento subflavicante onustis, inferioribus in petiolum attenuatis, racemo paniculato, floribus pedicellatis fasciculatis, fasciculis remotis, calycis parvi lanati laciniis lanceolatis ② in sterilibus Monspelii. (v. s.)

Ordo LXX. ANTIRRHINÆ *Juss. gen.* 118. — Personatæ  
*DC. fl. fr.* 3. p. 573.

Calyx 1-sepalus divisus sæpè persistens. Corolla irregularis, limbo diviso plerumquè bilabiato personato; stamina 4 didynama corollæ inserta. Ovarium liberum; stylus 1; stigma simplex aut 2-lobum. Capsula uni-aut 2-locularis 2-valvis valvis concavis plus minùsve segregatis, interdum 2-partitis; dissepimento centrali valvis parallelo nunc libero simplici, nunc duplici cum valvis introflexis coalito. Semina numerosa medio dissepimento adfixa. Albumen carnosum; embryo rectus; cotyledones semicylindricæ. — Herbæ foliis oppositis aut alternis, floribus foliis superioribus minoribus bracteatis.

† *Stamina 4 didynama; capsulæ 2-loculares.*

I. GRATIOLA *L. gen.* 29.

Calyx 5-partitus, basi 2-bracteatus; corolla tubulosa subbilabiata, labio superiore emarginato, inferiore 3-lobo; stamina 4, 2 sterilia; capsula ovata, bilocularis dissepimento simplici.

1. *G. OFFICINALIS* (*L. sp.* 24.) foliis lanceolatis serratis subtrinerviis, floribus pedunculatis. 2ℓ in humidiusculis. — *Fl. fr.* 3. p. 598. — *Lob. ic. t.* 435. *f.* 2. (v. v.)

II. DIGITALIS *L. gen.* 758.

Calyx 5-partitus inæqualis; corolla campanulata, limbo obliquè 4-lobo inæquali; stigma simplex aut 2-lamellatum; capsula ovata acuminata.

1. *D. PARVIFLORA* (*Lam. fl. fr.* 2. p. 333.) foliis ovato-lanceolatis denticulatis glaberrimis, lacinii calycinis lanceolato-linearibus acutis, corollis acutis, labio superiore bifido. 2ℓ in saxosis rupestribus montosis. — *Fl. fr.* 3. p. 597. *D. lutea* *L. sp.* 867. non *Poll.* — *Lob. ic. t.* 573. *f.* 2. (v. v.)
2. *D. GRANDIFLORA* (*Lam. fl. fr.* 2. p. 332.) foliis lanceolatis serrulatis subtus pubescentibus, calycis lacinii lanceolatis pubescentibus, corollis inflatis galeâ emarginatis. 2ℓ in montosis umbrosis Galliæ orientalis. — *Fl. fr.* 3. p. 596. *D. ochroleuca* *Jacq. vind. r. t.* 57. (v. v.)
3. *D. PURPUREA* (*L. sp.* 866.) foliis lanceolato-ovatis obsoletè crenatis utrinquè pubescentibus, inferioribus petiolatis, lacinii calycinis ovatis acutis, corollis obtusis, labio superiore integro. (2) in montibus sylvaticis Galliæ mediæ, australis occidentalisque. — *Fl. fr.* 3. p. 595. *Lam. ill. t.* 525. *f.* 1. (v. v.)
4. *D. PURPURASCENS* (*Elmig. diss.* 45.) foliis lanceolatis serratis basi amplexicaulibus supernè glaberrimis subtus pubescentibus, calycis lacinii lanceolato-ovatis subacutis, corollâ cylindricâ subventricosâ ad oras villosulâ galeâ obtusâ emarginatâ, labii inferioris lobo medio oblongo obtuso. 2ℓ in Alsatiæ, Arverniæ, Pyrenæorum dumetis montanis. — *Fl. fr.* 5. p. 411. *D. hybrida* *Kœler. in Journ. phys.* 1782. t. 1. *f.* 1. 2. Hybrida videtur inter 2 precedentes. (v. s.)

III. ANARRHINUM *Desf. ad.* 2. p. 52. *DC. fl. fr.* 3. p. 594.

Calyx 5-partitus; corolla tubulosa interdum calcarata vel basi deorsum prominens, fauce patente, palato prominulo destituta; capsula subrotunda apice foraminulis 2 dehiscens.

1. *A. bellidifolium* (Desf. l. c.) foliis radicalibus obovato-lanceolatis obtusis serratis, caulinis partitis integerrimis. ② in sterilibus Gallie australis et austro-orientalis et rivata occidentalis. — Fl. fr. 3. p. 595. — Bauh. prod. 186. ic. (v. v.)

IV. ANTIRRHINUM Tourn. t. 75. Juss. gen. p. 120.

Calyx 5-partitus; corolla non calcarata basi gibba, tubo inflato, limbo 2-labiato, suprā 2-fido reflexo, infrā 3-lobo, palato ad faucem prominulo; capsula basi obliqua, apice foraminulis tribus dehiscens.

1. *A. OROSTIUM* (L. sp. 860.) foliis lanceolatis angustis, floribus subsessilibus subspicatis, lacinii calycinis linearibus inæqualibus corollam capsulamque superantibus. ① in agris et arvis. — Fl. fr. 3. p. 593. — Cam. epit. 923. ic. (v. v.)

2. *A. MAJUS* (L. sp. 859.) caulibus basi glaberrimis apice pubescentibus, foliis lanceolatis alternis rameis oppositis, floribus racemosis, calycis glanduloso-pubescentis lacinii ovatis obtusis corollā multō brevioribus, capsulā calycem superante. ② in maceris saxosis provinc. austr. et in hortis persæpè cultum undè in muros vicinos migrat. — Fl. fr. 3. p. 593. — Lob. ic. t. 404. f. 2. (v. v.)

β. *latifolium*, foliis ovatis aut ovato-lanceolatis glabris ant pubescentibus. — In asperis apricis Galloprovinciæ, agri Ruscinonensis. — *A. latifolium* Mill. dict. n. 4. Fl. fr. 5. p. 411. — Boec. mus. t. 41. (v. v.)

3. *A. MOLLE* (L. sp. 860.) foliis ovatis obtusissimis oppositis tomentosis, lacinii calycinis ovalibus, caulibus procumbentibus. ② in Alpibus, Pyrenæis Hispaniæ conterminis ex cl. Arnott. — Fl. fr. 3. p. 594. †

4. *A. SEMPERVIRENS* (Lapeyr. fl. pyr. 1. p. 7. t. 4.) foliis ellipticis sempervirentibus oppositis cauleque frutescente divaricato-pubescentibus, calycis pubescentis lacinii brevibus lanceolatis corollā multō brevioribus, capsulā calycem superante. ③ in Pyrenæis centralibus ad rupes calcarias. — Fl. fr. 3. p. 593. (v. v.)

5. *A. ASARINA* (L. sp. 860.) caulibus procumbentibus diffusis villosis, foliis oppositis cordatis crenatis, calycis hirsutis lacinii linearibus corollā multō brevioribus capsulamque superantibus. ② in Pyrenæis orientalibus, Cebennis, montibus Ligericinis et Velaunorum ex Arnaud. — Fl. fr. 3. p. 594. — Lob. ic. t. 601. f. 2. (v. v. DC.)

V. LINARIA Tourn. t. 76. Juss. gen. p. 120.

Calyx 5-partitus, lacinii 2 inferioribus remotis; corolla ringens basi calcarata, tubo inflato, limbo 2-labiato, suprā 2-fido reflexo, infrā 3-lobo; capsula ovata aut globosa apice biforis multivalvis; semina marginata.

Sect. I. CHÆNORRHINUM DC. fl. fr. 5. p. 410.

Faux aperta palato prominulo destituta, capsula subrotunda.

1. *L. ORIGANIFOLIA* (DC. fl. fr. 3. p. 591.) foliis obovatis oppositis, inferioribus glabris, floralibus alternis cauleque ascendente ramoso pubescentibus, calycis pubescentis lacinii spathulato-linearibus corollā multō brevioribus capsulam subsuperantibus, calcare recto obtuso. ② ③ in rupestribus Pyrenæorum, Cebennarum, Galloprov., Delph., Montis-Aurei ex L. — *Antirrhinum origanifolium* L. sp. 852. *A. villosum* Lapeyr. non L. — Barr. ic. 598. — Fl. purpureo-cærulescentes. (v. v.)

2. *L. RUBRAIFOLIA* (DC. fl. fr. 5. p. 410.) foliis inferioribus oppositis subrotundis glabris carnosius subtus rubris, superioribus alternis, caule suberecto apice pubescente, calycis pubescentis lacinii spathulato-linearibus corollā brevioribus capsulam superantibus, calcare subdivergente acuto. ① in collibus rupestribus Massiliæ, Monspelii. — Magu. bot. 25. ic. Fl. cærulescentes. (v. s.)

3. *L. minor* (Desf. atl. 2. p. 46.) caule ramosissimo diffuso, foliis lanceolatis obtusis ellipticisque alternis, inferioribus oppositis, calycis laciniis inaequalibus spatulato-linearibus corollâ subbrevioribus capsulam subsuperantibus. ① in cultis, ruderatis, glareosis. — Fl. fr. 3. p. 591. Ant. minus L. sp. 852. — Lob. ic. t. 406. f. 1. Fl. purpureo-violacei. (v. v.)

## Sect. II. LINARIASTRUM.

Corolla ringens, palatum ad faucem prominulum.

\* *Folii angulatis.*

4. *L. CYMBALARIA* (Mill. dict. n. 17.) caulibus glabris procumbentibus, foliis reniformibus cordatis alternis glabris longè petiolatis 5-7-lobis mucronulatis, pedunculis folio brevioribus, laciniis calycinis lanceolatis capsulâ brevioribus, calcare obtuso recurvo. ② in rupibus et muris antiquis passim. — Fl. fr. 3. p. 583. Ant. *Cymbalaria* L. — Cam. epit. 860. ic. Fl. caeruleascentes palato flavo. (v. v.)
5. *L. HEPATICIFOLIA* (Dub. mss.) caulibus procumbentibus diffusis glabris, foliis longè petiolatis oppositis reniformi-cordatis 3-5-lobis lobo medio maximo, calycis glaberrimi laciniis linearibus subaequantibus capsulam subsuperantibus, calcare recto obtuso. ② in Corsicâ. — Ant. *hepaticifolium* Poir. supp. 4. p. 29. Fl. prioris sed majores. (v. s.)
6. *L. AQUETRILLOSA* (Dub. mss.) caulibus filiformibus procumbentibus diffusis pilosis, foliis reniformibus alternis longè petiolatis pilosis obsoletè 3-lobis lobis rotundatis mucronulatis subaequalibus, pedunculis filiformibus, laciniis calycinis linearibus, calcare brevi recto obtuso. ① in Corsicâ humantibus. — Ant. *aquetrilobum* Viv. oors. p. 10. Flores L. *cymbalariae* sed minores. An satis à L. pilosâ distincta? (v. s.)
7. *L. SPURIA* (Mill. dict. n. 15.) caule procumbente, foliis ovato-reniformibus cordatis pilosis alternis, inferioribus oppositis obsoletè dentatis, calycis pilosi laciniis ovato-lanceolatis capsulam subsuperantibus, calcare incurvo acuto. ① in arvis. — Fl. fr. 3. p. 584. Ant. *sparium* L. sp. 851. — Fuchs. hist. 167. ic. Fl. flavi, labio superiore violaceo. (v. v.)
8. *L. ELATINE* (Desf. atl. 2. p. 37.) pilosa, caulibus procumbentibus ramosissimis hirsutis, foliis breviter petiolatis alternis ovato-hastatis, pedunculis filiformibus folia superantibus, calycis laciniis lineari-lanceolatis acutis capsulam subsuperantibus, calcare incurvo acuto. ① in arvis. — Fl. fr. 3. p. 584. Ant. *Elatine* L. sp. 851. Fl. præcedentis palato caeruleascente. (v. v.)
9. *L. CYRHOSEA* (Willd. enum. 639.) caulibus elongato-filiformibus procumbentibus hirsutis, foliis breviter petiolatis subglabris lanceolato-hastatis, pedunculis capillaceis passim cyrrhescentibus folia superantibus, laciniis calycinis lanceolatis margine scariosis acuminatis capsulâ subbrevioribus, calcare subrecto acuto. ① in arenosis Corsicæ, Stœchadum, Telonis (cl. d'Urville). — Fl. fr. 5. p. 407. Ant. cyrrhosum L. Jacq. vind. t. 82. Fl. parvi caeruleascentes. (v. s.)
- \*\* *Folii integerrimis inferioribus verticillatis.*
10. *L. REFLEXA* (Desf. atl. 2. p. 42.) glabra, caulibus patulis, foliis ovato-lanceolatis sessilibus glabris ternis, floralibus alternis, pedunculis axillaribus, fructiferis recurvatis, laciniis calycinis lanceolatis acuminatis capsulam subsuperantibus, calcare longissimo recto acuminato. ① in Corsicâ ex Vallo. All. — Fl. fr. 3. p. 585. — All. misc. taur. 2. p. 205. t. 1. Fl. pallidè caeruleascentes flavascentesve. (v. s. ex Neap.)
11. *L. TRIPHYLLOA* (Mill. dict. n. 2.) glabra, caulibus erectis, foliis ternis ovatis obtusis trinerviis, apicè terminali, floribus subsessilibus, laciniis calycinis lanceolatis ovatisque obtusis capsulam subaequantibus, calcare acuto incurvo. ① in Corsicâ (Soleirol), Santoniâ? ex Lam. — Fl. fr. 3. p. 585. Ant. *triphyllum* L. — Booc. sic. p. 45. t. 22. Cor. pallidè purpurascens palato flavo. (v. s.)
12. *L. THYMIFOLIA* (DC. fl. fr. 3. p. 587.) glaberrima, caulibus procumbentibus,

foliis ovalibus obtusis oppositis ternatisque glabris, floribus capitatis, laciniis calycinis spathulato-ovatis obtusis capsulâ brevioribus, calcare subrecto subacuto.  $\frac{1}{2}$  in arenosis maritimis Gallie austro-occidentalis usque ad ostium Garumne. — Ant. thymifolium Vahl. Lois. gall. t. 10. Fl. flavi. (v. v.)

13. *L. PELISSERIANA* (DC. fl. fr. 3. p. 589.) glabra, caule erecto subsimplici, foliis surculorum sterilibus ternis lanceolatis ovatisve, caulinis linearibus, superioribus alternis, floribus pedunculatis racemoso-capitatis, pedunculis bracteis subæquantibus, calycis laciniis linearibus acuminatis capsulam superantibus, calcare longissimo recto acuto.  $\frac{1}{2}$  in saxosis rupestribus Gallie occidentalis, mediæ australisque. — Ant. Pelisserianum L. sp. 855. Ant. gracile Pers. ench. 2. p. 156. — Barr. ic. t. 1162. Fl. parvi purpurei. (v. v.)
14. *L. ARVENSI* (DC. fl. fr. 3. p. 588.) caule erecto ramoso, foliis linearibus inferioribus quaternis, floribus racemosis, calycis glanduloso-viscosi laciniis linearibus acutis capsulâ brevioribus, calcare incurvo acuto. ① in arvis cultis præsertim in prov. australibus. — *L. arvensis* & Desf. — Dill. elth. t. 163. f. 198. Fl. minimi cærulescentes. (v. v.)
15. *L. SIMPLEX* (DC. fl. fr. 3. p. 588.) caule erecto simplici basi glabro apice glanduloso-pubescente, foliis sublinearibus inferioribus quaternis, floribus racemosis, calycis glanduloso-viscosi laciniis spathulato-linearibus obtusis capsulâ brevioribus, calcare recto subobtus. ① in Galliâ australi passim. — *L. arvensis* & Desf. Ant. simplex Wild. A. parviflorum Jacq. ic. rar. 3. t. 499. non Desf. Fl. minimi flavi. An à priori satis distincta sp? (v. s.)
16. *L. ARENARIA* (DC. ic. rar. t. 14.) pubescenti-viscosa, caulibus ramosis numerosissimis diffusis, foliis lanceolato-linearibus acutis sparsis, inferioribus verticillatis oblongis obtusis, floribus subpedunculatis, bracteis pedunculum multo superantibus, spicâ elongatâ, calycis subhirsuti laciniis lanceolato-linearibus acutis capsulam subæquantibus, calcare recto acuto. ① in Armoricæ et Neustriæ usque ad Cherbourg arenosis maritimis. — Fl. fr. 5. p. 409. Fl. lutei minimi. (v. v.)
17. *L. SAXATILIS* (DC. fl. fr. 3. p. 590.) pubescenti-viscosa, caule decumbente, foliis lanceolato-linearibus sparsis villosis-viscosis, inferioribus verticillatis ovatis obtusis subpetiolatis, floribus spicatis, calycis laciniis lanceolato-ovatis acutis,  $\frac{1}{2}$  in Armoricæ maritimis. — Ic. rar. t. 13. Fl. fr. 5. p. 409. Fl. parvi flavi. (v. s.)
18. *L. SUPINA* (Desf. atl. 2. p. 44.) caule diffuso apice subpiloso, foliis glaucis subquaternis linearibus superioribus sparsis, floribus racemoso-capitatis, calycis laciniis sublinearibus obtusis capsulâ brevioribus, calcare recto acuto. ① in arenosis Gallie mediæ, australis occidentalisque. — Fl. fr. 3. p. 588. Ant. supinum Lin. L. Thuillierii. Mer. fl. par. 240. — Clus. hist. 1. p. 321. ic. flavi. (v. v.)
  - β. *pyrenaica* (Lapeyr.) foliis inferioribus lanceolato-linearibus, caulibus apice calycibusque glanduloso-pilosis. — In Pyrenæis et Alpibus. — *L. Pyrenaica* Fl. fr. 3. p. 587 et 5. p. 408. Ic. rar. t. 11. A. glaucum Lapeyr. abr. 352, non L. ex cl. Arnott. (v. v.)
  - γ. *maritima*, foliis omnibus verticillatis, laciniis calycinis ovato-oblongis. — In maritimis Gallie occidentalis. — *L. maritima* DC. syn. p. 232. ic. rar. t. 12. Fl. fr. 5. p. 408. (v. v.)
19. *L. FLAVA* (Desf. atl. 2. t. 136.) caulibus subdiffusis, foliis lanceolatis alternis, inferioribus ovatis ternis oppositisque, floribus capitatis, calycis glabri laciniis linearibus acutis, calcare recto acuto. ① in Corsicâ. — Fl. fr. 3. p. 729. Fl. flavi. An *L. supina* var? (y. s. specimen imperf.)
20. *L. JUNCEA* (Desf. atl. 2. p. 43.) caulibus gracilibus ramosis virgatis, foliis linearibus carnosis alternis, infimis ternis, floribus pedunculatis racemosis, calycis glabri laciniis linearibus subobtusis capsulam subæquantibus, calcare recto acuto. ① in arenosis et inter segetes circa Aquas-Tarbellicas et Burdigalam (Des Moulins). — Fl. fr. 3. p. 729. Ant. junceum L. sp. 858. Ant. spar-teum L. sp. 854. Fl. flavi. (v. v. D.)

21. *L. ALPINA* (DC. fl. fr. 3. p. 590.) glabra, caule diffuso petalo, foliis quaternis linearibus glaucis, inferioribus ovalibus, floribus racemosis, calycis laciniis lineari-lanceolatis linearibusve acutis capsulam subaequantibus, calcare recto acuto. ② in Jurasco, Alpihus et Pyrenæis. — Ant. alpinum L. sp. 856. — Clus. hist. 1. p. 222. f. 2. Fl. purpurei palato flavescente. (v. v.)
22. *L. PURPUREA* (Mill. dict. n. 5.) glabra, caule erecto ramoso, foliis linearibus quaternatim verticillatis, superioribus alternis, floribus pedunculatis racemoso-spicatis, pedunculis bractea brevioribus, calycis laciniis linearibus capsula brevioribus, calcare incurvo acuto. ½ in collibus circa Fontembellaqueum ex Merat, Beauvais ex Graves et propè S.-Omer ex cl. Desmazières. — Fl. fr. 5. p. 407. — Dod. pempt. 183. f. 2. Fl. purpurei. (v. s. ex Italiâ.)
23. *L. STRIATA* (DC. fl. fr. 3. p. 580.) radice repente, caulibus paniculatis suberectis, foliis linearibus confertis alternis sparsisque infimis verticillatis, calycis glabri laciniis lanceolatis acutis capsula brevioribus, calcare brevi recto obtuso. ½ in campis sterilibus, ad muros Gallie medice, australis occidentalisque. — Ant. monspessulanum et A. repens L. sp. 854. — Dill. elth. t. 163. f. 197. Flores striati cærulescentes albescentesve. (v. v.)
24. *L. VERSICOLOR* (Moench. meth. 523.) caule erecto spicato apice glanduloso-viscoso, foliis lineari-lanceolatis linearibusve, inferioribus ternis quaternisque, pedunculis calycibusque villosis-viscidis, laciniis lineari-spathulatis capsulam superantibus, calcare recto acuto. ① in Galliâ australi? — Fl. fr. 3. p. 586. Ant. versicolor Jacq. ic. rar. 1. t. 116. (v. s. c.)
25. *L. CHALEPENSIS* (Mill. dict. n. 12.) glabra, caule erecto, foliis lineari-lanceolatis linearibusve alternis inferioribus verticillatis, floribus racemosis, calycis corollâ longioris laciniis linearibus acutis, calcare longissimo gracili acuto deflexo. ① in cultis circa Monspelium? — Fl. fr. 3. p. 589. — Triumf. obs. t. 87. f. 2. Fl. albescentes. (v. s. ex Orient.)

\*\*\* *Foliis omnibus alternis integerrimis.*

26. *L. VULGARIS* (Moench. meth. 524.) caule erecto, foliis lanceolatis linearibus acutis confertis, spicis terminalibus, calycis laciniis linearibus acutis capsula brevioribus, calcare subrecto acuto. ½ in rudertis. — Fl. fr. 3. p. 592. Ant. Linaria L. sp. 858. — Cam. epit. 930. ic. Fl. flavi. (v. v.)
27. *L. GENISTIFOLIA* (Mill. dict. n. 14.) caule supra paniculato virgato flexuoso, foliis lanceolatis acuminatis, calycis laciniis linearibus acutis capsulam subaequantibus, calcare recto acuto. ½ in Pyrenæis, Delphicis, Alsatia ex Mapp. — Fl. fr. 3. p. 591. — Clus. hist. 1. p. 322. f. 1. (v. s.)

#### VI. SCROPHULARIA Tourn. t. 74. L. gen. n. 756.

Calyx 5-lobus; corolla globosa, limbo coarctato bilabiato, suprâ biloba squamâ intus sæpè interjectâ, infra brevior 3-loba; stigma unicum; capsula subrotunda acuminata, valvis integris basi introflexis, dissepimento duplici.

1. *S. PERFORATA* (L. sp. 866.) caule 4-6-angulo, foliis cordatis dentatis lucidis inferioribus petiolatis, pedunculis axillaribus 2-4-floris, calycis laciniis lanceolatis acutis. ① in umbrosis et rupestribus Corsicæ, Niceæ ex All., Galloprov., Occitanie, circa Aquas Tarbellicas, Dianæ Armoracæ (cl. de Brébisson). — Fl. fr. 3. p. 580. — Cam. hort. t. 43. (v. v.)
2. *S. VERNALIS* (L. sp. 864.) foliis cordatis pubescentibus duplicato-serratis petiolatis, paniculis axillaribus dichotomis, calycis laciniis lanceolatis obtusis, capsula ovata. ② in Galliâ Narbonensi ex Lam., in Pyrenæis propè St.-Beat ex cl. Benth., circa Bellocvacum ex cl. Graves, Cadomum ex cl. de Bréb., Rothomagus (cl. Le Prévost). — Fl. fr. 3. p. 579. — Barr. ic. t. 273. (v. s.)
3. *S. AQUATICA* (L. sp. 864.) foliis cordatis lanceolatis glabris crenatis petiolatis decurrentibus obtusis, caule membranis angulato, racemis terminalibus,

- pedunculis 6-10-floris, calycis glabri laciniis rotundato-ovatis margine scariosis. ② ad ripas et in aquaticis. — Fl. fr. 3. p. 579. Fl. dan. t. 507. (v. v.)
4. *S. SCORONOMA* (L. sp. 864.) foliis cordatis triangularibus duplicato-serratis pubescentibus acutis petiolatis, pedunculis axillaribus 3-chotomis 5-12-floris, calycis pubescentis laciniis rotundis obtusissimis margine scariosis. ④ in humidis Nicæ ex All., circa Cherbourg (cl. Le Prévoist). — Fl. fr. 3. p. 580. — Moris. a. 5. t. 35. n. 6. (v. s.)
5. *S. MONOSA* (L. sp. 863.) foliis glabris cordatis lanceolato-ovatis acutis dentatis petiolatis, caule obtusangulo, calycis glabri laciniis rotundato-ovatis non scariosis, pedunculis sub 5-floris. ④ in umbrosis succulentis. — Fl. fr. 3. p. 579. — Cam. epit. 866. ic. (v. v.)
6. *S. SCOROLII* (Hop. exs. Pers. ench. 2. p. 160.) caulibus obtusè 4-gonis, foliis pubescentibus petiolatis inferioribus ternatis cordatis crenato-dentatis, superioribus lanceolatis argutè dentatis, floribus racemoso-paniculatis, pedunculis sub 10-floris, pedicellis elongatis, calycis glabri laciniis rotundato ovatis obtusis margine scariosis. ④ in umbrosis Pyrenæorum. — Fl. fr. 5. p. 406. *S. auriculata* Scop. carn. t. 32. (v. v.)
7. *S. TRIFOLIATA* (L. sp. 865.) foliis glabris, inferioribus ternato-pinnatis obtusis, superioribus simplicibus inciso-crenatis, laciniis calycinis rotundatis obtusissimis, pedunculis axillaribus ramosis 3-6-floris. ② in Corsicâ ex Linneo. — Fl. fr. 3. p. 581. — Pluck. t. 313. f. 6. (v. s. c.)
8. *S. SAMBUICIFOLIA* (L. sp. 865.) caule glabro obsoletè tetragono, foliis petiolatis, inferioribus interruptè pinnatis foliolis 5-7 inæqualibus ovato-cordatis glabris crenatis, petiolis appendiculatis, superioribus ovatis crenatis indivisis aut basi incisis appendiculatisve, pedunculis axillaribus 3-5-floris, laciniis calycinis rotundis obtusis margine scariosis, capsulâ ovato-acuminatâ. ④ in Corsicâ. — Moris. a. 5. t. 8. n. 6. Fl. magni. (v. s.)
9. *S. AURICULATA* (L. sp. 864.) caule 4-gono, foliis oblongis cordatis subtus hirsutis duplicato-dentatis basi appendiculatis, paniculâ racemosâ terminali, pedunculis multifloris, calycis laciniis rotundis margine scariosis. ④ in agro Nicænsi ex All. — Fl. fr. 3. p. 580. — Lob. ic. t. 533. f. 1. (v. s. c.)
10. *S. LUCIDA* (L. sp. 865.) foliis inferioribus petiolatis subcarnosis glaberrimis bipinnatis laciniis ovato-rotundis crenatis, superioribus sessilibus, foliolis pinnatifidis lobatisve, floribus racemoso-paniculatis, pedunculis sub 5-floris, laciniis calycinis rotundato-ovatis margine pellucido-scariosis, corollæ palato subtus appendiculato. ② ④ ? in arenosis Nicæ ex All. — Fl. fr. 3. p. 582. — Bocc. mus. 2. p. 166. t. 117. (v. s. ex littore Genuensi.)
11. *S. CANINA* (L. sp. 865.) foliis pinnatis, pinnis indivisis lobatisve lanceolatis dentatis, summis confluentibus, racemo terminali subnudo, pedunculis 4-8-floris, bracteis minimis, calycis laciniis ovato-rotundis obtusis margine pellucido-scariosis. ④ in lapideis et sterilibus. — Fl. fr. 3. p. 582. Clus. hist. 2. p. 209. f. 1. (v. v.)
12. *S. RAMOSISSIMA* (Lois. 2. p. 381.) caule suffruticoso ramosissimo ramis patulis, foliis subcarnosis glabris lanceolatis, inferioribus pinnatifidis inciso-dentatisve superioribus denticulatis sessilibus, floribus racemosis, pedunculis simplicibus 1-floris medio 2-bracteolatis, calycis laciniis rotundis obtusis margine scariosis. ⑤ in Corsicæ, Nicæ et Galloprovinciæ orientalis arenosis maritimis. — Fl. fr. 5. p. 406. *S. frutescens* Fl. fr. 3. p. 729. excl. ayu. (v. v. D.)

VII. *ERINUS* L. gen. n. 771.

Calyx 5-fidus; corolla tubulosa limbo plano 5-lobo subæquali, laciniis apice cordato-emarginatis; capsula ovata 2-valvis valvis apice 2-fidis, marginibus valvarum inflexis 2-locularis.

1. *E. ALPINUS* (L. sp. 878.) floribus racemosis, foliis spathulatis. ④ in montium locis lapideis apricis. — Fl. fr. 3. p. 578. et 5. p. 405. Lam. ill. t. 521. — Barr. ic. t. 1123. (v. v.)

†† *Stamina 4 didynama; capsula unilocularis.*

VIII. *LINDERNIA* All. misc. 3. p. 178. L. mant. p. 252.

Calyx 5-partitus; cor. tubulosa ringens suprâ brevissima emarginata, infrâ 3-fida inæqualis; stamina 4 didynama: filamenta breviora apice 2-dentata, altero dente nudo, altero antherifero; stigma emarginatum.

1. *L. PYRIDARIA* (All. l. c.) foliis oblongo-ovatis integerrimis sessilibus, pedunculis axillaribus solitariis unifloris. ① in paludibus spongiosis inundatis Gallie occidentalis, mediz, borealis orientalisque passim. — Fl. fr. 3. p. 577. *Cephraria gratioloides* L. sp. 876. — Lind. als. 1. p. 152. t. 1. (v. v. DC.)

IX. *LIMOSELLA* L. gen. 776.

Calyx 5-fidus irregularis; corolla campanulata minima 5-loba subæqualis; stamina 4 aut abortu 2; stigma globosum, ovarium basi 2-loculare apice 1-loculare; capsula ovoidea.

2. *L. AQUATICA* (L. sp. 881.) foliis lanceolato-spathulatis, scapis foliis brevioribus. ½ in inundatis ad fluviorum ripas Gall. occid., mediz, borealia orientalisque passim. — Fl. fr. 3. p. 576. — Lind. als. 156. t. 5. (v. v.)

## Ordo LXXI. OROBANCHEÆ Juss. ann. mus. 12. p. 445.

Calyx monosepalus bracteatus, aliquandò profundè in laciniis bracteiformes divisus. Corolla hypogyna monopetala irregularis limbo bilabiato. Stam. 4 corollæ inserta; filamenta 2 longiora. Ovarium simplex liberum 1-loculare polyspermum; stylus unicus. Stigma simplex aut bilobum. Capsula 1-locularis 2-valvis, valvulis liberis in nervo longitudinali seminiferis. Albumen carnosum aut subcorneum in parte superiori lateraliter embryonem parvum fovens. — Herbæ luridæ subcarnosæ sæpiùs parasiticæ, caulibus mox nudis mox squamosis, foliis mox radicalibus, mox caulinis squamiformibus, floribus terminalibus solitariis spicatisve bracteatis.

I. *OROBANCHE* L. gen. 779. Wallr. *diaskeus*. p. 28. *Vauch. monog.* p. 1.

Calyx 1-2-partitus, bracteis 1-3-suffultus; corolla monopetala tubulosa ringens 4-5-fida; stamina 4 didynama; stylus 1; stigma capitatum emarginato-bilobum; ovarium basi glandulosum.

Sect. 1. *OSPROLEON* Wallr. l. c. p. 21. *Vauch. l. c.* p. 37.

Calyx 1-2-partitus, bractea 1 suffultus; corolla 4-fida, capsula lateraliter dehiscens. — Scapus simplicissimus.

1. *O. RAPUM* (Thuil. par. ed. 2. p. 317.) scapo in bulbum carnosum imbricatim-squamosum incrassato, squamis bracteisque superioribus ovato-oblongis hirsuto-glutinoso, laciniis calycinis duobus bifidis, lobis linearibus æqualibus, corollâ inflato-ventricosâ bilabiata, labio superiore 2-lobo, inferiore 3-lobo, stigmate bilobato luteo. ½ in sterilibus et argillosis Genistæ tineto-



rim parasitica. — Wall. l. c. p. 3a. O. major Lam. ill. t. 55t. DC. fl. fr. 3. p. 488 non L. O. Genistæ tinctoriæ Vaucl. l. c. t. 1. — Moris. s. 12. t. 16. f. 2. n. 1. Planta flavo-rubens. (v. v.)

β. *affinis*, duplō triplōve grācilior, squamis basi dilatatis, floribus paucis abbreviatis, filamentis basi dilatatis. — Genistæ sagittali parasitica. — O. Genistæ-sagittalis Vaucl. l. c. O. vulgaris Lam. dict. 4. p. 6at? DC. fl. fr. 3. p. 489? An species?

2. O. *fatida* (Desf. atl. 2. t. 144?) scapo ultrā basin incrassato, squamis bracteisque ovato-oblongis remotis furfuraceo-hirsutissimis, laciniis calycinis in dentes lineares divisis, corollā inflato-ventricosā extus glanduloso-asperā, labio superiore brevi subintegro, inferno trilobo, stigmate profundē diviso. ½ in lapidosis Galloprovinciæ propē Mées, Pyren. Orientalium et Tolosæ ex Lapeyr., Vasconiz (Des Moul.). — Fl. fr. 5. p. 3ga. O variegata Wallr. p. 40. O. Cytisi-scoparii Vaucl. l. c. Omnium maxima; flores fulvi. (v. v. DC.)
3. O. *cavita* (Viv. cors. 11.) scapo basi squamis elongatis linearibus onusto, bracteis lineari-acuminatis externē hirsutis florem superantibus, corollā 4-fidā incurvā subcylindricā lobis rotundatis crenato-ciliatis, filamentis capsulāque glaberrimis, stigmate 2-lobo recurvo. ½ in Corsicā propē Bonifacio ex Viv. — An huic potius quam præcedenti referendum synonym. Desf. (v. s.)
4. O. *major* (L. sp. 882. excl. syn. non DC.) bulbo mediocri squamoso, squamis lanceolatis non adpressis, laciniis calycinis bifidis, corollā inflatā bilabiātā, stigmate bifido exserto, staminibus albidis subexsertis. ½ in arenosis Burdigalæ Ulici Europæo parasiticam reperiit cl. Des Moulins. — Eng. bot. t. 421. O. Ulicis-Europæi Vaucl. l. c. Planta albo-fusca. (v. a.)
5. O. *medicaginis*, scapo ad basin vix inflato, squamis inferioribus numerosis lanceolatis nigrescentibus, superioribus abbreviatis paucioribus, laciniis calycinis 2-lobis, corollā elongatā, fauce constrictā, labio inferiore trilobo, superiore galeato subemarginato, stigmate bifido reflexo flavo-rubescente. ½ Medicagini sativæ parasiticam reperiit cl. prof. Vaucher et nomine O. Medicaginis-sativæ designat l. c. t. 2. Pl. flavescens. •
6. O. *minor* (Sutt. trans. linn. 4. p. 178.) scapo ad basin incrassato modicē squamoso, bracteis lanceolatis pilosis, laciniis calycinis nunc simplicibus nunc bifidis, corollā arcuato-cylindricā, labio superiore rotundato-crenulatō integro, inferiore trifido lobis æqualibus rotundato-crenulatis, staminibus ultrā basin corollæ insertis, antheris nigrescentibus, stigmate bilobo purpurascente transversē rimoso. ½ in agris et pascuis Trifolio pratensi parasitica. — Fl. fr. 3. p. 489. eng. bot. t. 422. O. Trifolii-pratensis Vaucl. l. c. t. 4. Flores pallidē violacei aut sordidē albescentes. (v. v.)
7. O. *ruinosa* (Lapeyr. abr. supp. 87.) scapo fistuloso piloso pruinoso-cæsiō in bulbum mediocrem dilatato, squamis lanceolatis, bracteis angustis corollā brevioribus, laciniis calycinis profundē et angustē bifidis, corollā labiatā, labiis ore plicatis, superiore bifido, inferiore trilobo lobis denticulatis, stigmate profundē diviso dilutē purpurascente. ① in Pyrenæis orientilibus Fabæ parasitica. — O. Fabæ-sativæ Vaucl. l. c. t. 5. Pl. albi. (v. v. c.)
8. O. *Erythrum* (DC. fl. fr. 3. p. 490.) scapo subæquali, squamis bracteisque lanceolatis, floribus patentibus sparsis glanduloso-viscidulis remotis, laciniis calycinis lanceolatis rarē bifidis, corollæ labio superiore rotundato-crenulatō, inferiore trilobo, lobis obtusē crenatis, stigmate bilobo rubescente. ½ in aridis montosis, parasitica in Thymum serpillum. — O. Thymi-Serpilli Vaucl. l. c. t. 6. Planta rubescens. (v. v. DC.)  
β. *epithymoides*, scapo subinflato, corollā venosā, labio superiore bilobe lobis superincumbentibus. — In sylvia Galloprov. propē Brignolles reperiit cl. Vaucher parasiticam Thymo vulgari et hanc habet ut speciem propriam quam nomine O. thymi-vulgaris designat mon. p. 53.
9. O. *galii*, scapo mediocriter incrassato ad basin squamoso, squamis bracteisque lanceolatis nigrescentibus, laciniis calycinis lanceolatis vel irregula-

riter bifidis, corollâ bilabiata inflatâ, labio superiore fornicato subemarginato, inferiore trilobo lobis rotundatis, filamentis inflexis, antheris nigrescentibus, stigmate bilobo atro-rubescente.  $\mathcal{L}$  in glareosis, Galio Mollugini, erecto etc. parasitica. — *O. Galii-Molluginis* Vauch. l. c. t. 7. Scapi rubelli; flores purpurascens. (v. v.)

10. *O. HEDERÆ* (Vauch. orob. p. 56. t. 8.) scapo in bulbam parçè squamosum incrassato, squamis lanceolatis paniculatis, bracteis apice attenuatis reflexis, corollâ bilabiata, labio superiore integro, inferiore trilobo lobis rotundatis, laciniis calycinis subintegris, staminibus inferioribus approximatis, superioribus divaricatis, stigmate emarginato lutescente.  $\mathcal{L}$  *Hederæ* parasiticam in Vasconiâ reperiit cl. Des Moul. (v. s. ex Genuâ.)

11. *O. RUBI*, scapo in bulbam squamosum incrassato, pilis albidis non glandulosis, onusto, squamis numerosis nigrescentibus, laciniis calycinis bifidis, corollâ labio superiore bifido, inferiore trilobo, lobis rotundatis erosus, staminibus suprâ basin adnatis, stigmate bilobo lutescente.  $\mathcal{L}$  in Galloprovinciâ arvis propè Luc, Rubo fruticoso parasiticam reperiit cl. Vauch. — *O. Rubi fruticosi* Vauch. l. c. t. 9. Scapi rubelli; flores dilutè lutei.

12. *O. ERYNGII*, scapo ad basin incurvato squamoso, squamis numerosis lanceolatis semi-adherentibus, laciniis calycinis profundè bifidis, lobis linearibus, corollâ labio superiore fornicato vix bifido, inferiore trilobo subplicato, stigmate bilobo rubescente.  $\mathcal{L}$  in provinciis australibus detexit cl. Vaucher *Eryngio campestri* parasiticam et *O. Eryngii-campestris* nominat l. c. p. 58. t. 10. Scapi rubelli hirsuti; flores rubello-albidi.

13. *O. CONCOLOR* (Dub. mss.) scapo ad basin dilatato squamoso gracili, squamis numerosis lanceolatis luteo-nigrescentibus, floribus numerosis, laciniis calycinis hirsutis sæpè bifidis, corollâ bilabiata luteâ, labio superiore subbilobo, inferiore 3-lobo undulato, staminibus ovario stigmateque bilobo luteis.  $\mathcal{L}$  — *Scabiosæ columbariæ*, *Cherophyllo sylvestri* et *Menthæ arvensi* parasiticam in Galloprov. ex Foro-Julii usque ad Cannas reperiit cl. Vaucher et nomine *O. Scabiosæ-columbariæ* designat mon. p. 59. t. 11. Planta omnis lutea.

14. *O. ELATIOR* (Sutt. trans. linn. 4. p. 178. t. 17.) scapo subæquali gracili piloso-glutinoso, squamis lanceolatis, floribus paucis remotis, laciniis calycinis bifidis, lobis linearibus, corollâ tubulosâ cylindricâ venosâ bilabiata, limbo plicato et fimbriato, labio superiore emarginato, inferiore trilobo, stigmate bifido subfusco.  $\mathcal{L}$  in agris et sylvaticis *Centaureæ Scabiosæ* parasitica. — Fl. fr. 3. p. 490. *O. Cent. scabiosæ* Vauch. l. c. p. 61. ic. (v. v.)

15. *O. RIGENS* (Lois. gall. 384.) scapo glabro, squamis lanceolatis rigentibus imbricatis, staminibus infernè nudis, stigmate bilobo, lobis distantibus, stylo glaberrimo.  $\mathcal{L}$  in Corsicâ ex Lois. — Fl. fr. 5. p. 392. *O. nudiflora* Wallr. sched. crit. 1. p. 310? †

Sect. II. *TRIONYCHON* Wallr. diask. p. 58. Vauch. monog. p. 65.

Calyx 4-5-fidus, bracteis 3 suffultus, corolla 5-fida; capsula apice dehiscens. — Scapus simplex aut ramosus.

16. *O. CÆRULEA* (Vill. dauph. 2. p. 406.) scapo simplicissimo, squamis oblongis, calyce 4-fido lobis oblongo-linearibus capsulam superantibus, corollâ tubulosâ subæquali inclinatâ, labio superiore bilobo, inferiore 3-lobo, stigmate bilobo flavescens.  $\mathcal{L}$  ? ① in agris et collibus *Artemisiæ vulgaris* parasitica. — Fl. fr. 3. p. 490. eng. bot. t. 423. *O. Artemisiæ-vulgaris* Vauch. l. c. t. 14. Corolla violaceo-cærulescens. (v. v.)

17. *O. COMOSA* (Wallr. sched. crit. 1. p. 314.) scapo simpliciusculo, squamis bracteisque hirsutis, calyce lobis 4 linearibus, corollâ infundibuliformi tubulosâ in faucem palato stris villosis exarata ampliatâ, 5-fidâ lobis rotundatis inferiore minore, antheris demum pilosis, stigmate bilobo griseo-rufescente. ①? in provinciis australibus plurimis plantis parasitica. — *O. vagabunda* Vauch. l. c. t. 15. Scapus cærulescens, corolla cærulea.

18. *O. RAMOSA* (L. sp. 882.) scapo ramoso parçè squamoso, calyce suburceolato 4-fido, lobis ovatis acuminatis, corollâ infundibuliformi-tubulosâ, labio inferiore trilobo lobis æqualibus obtusis rotundatis, staminibus basi pubescentibus, stigmate emarginato albido. (1) in agris parasitica Cannabi sativæ. — Fl. fr. 3. p. 490. Lam. ill. t. 551. f. 2. eng. bot. t. 184. *O. Cannabis* Vauch. L. c. t. 16. — Moris. a. 12. t. 16. f. 7. n. 8. Corolla infernè flavescens apice cærulescens. (v. v.)

II. *LATHRÆA* Lin. gen. 743.

Calyx campanulatus 4-fidus; corolla tubulosa 2-labiata labio superiore galeato; stamina 4 didynama; ovarium basi glandulosum; stylus 1; stigma 2-lobum.

1. *L. CLANDESTINA* (L. sp. 843.) caule ramoso subterrestri, floribus erectis solitariis. 2 in umbris Gallie præsertim occidentalis. — Fl. fr. 3. p. 491. Lam. ill. t. 551. f. 1. (v. v.)  
2. *L. SQUAMARIA* (L. sp. 844.) caule simplicissimo, corollis pendulis, labio inferiore 3-fido. 2 in frigidis umbrosis. — Fl. fr. 3. p. 492. Fl. dan. t. 136. (v. v.)

Ordo LXXII. RHINANTHACEÆ DC. fl. fr. 3. p. 454. —  
Pediculares Juss. gen. 99.

Calyx divisus persistens sæpè tubulosus. Corolla hypogyna irregularis sæpè ringens, æstivatione imbricatâ. Stamina numero definita 2-4, corollæ inserta, bina interdum breviora; antheræ basi setulæ aristatæ. Ovarium liberum; stylus simplex. Capsula 2-locularis 2-valvis valvulis coalitis dissepimento medio seminifero. Albumen carnosum; embryo rectus; cotyledones semiteretes. — Herbæ sæpè siccitate nigrescentes; folia opposita aut alterna; flores singuli bracteati axillares aut spicati. — Ordo nimis Antirrhineis affinis et probabiliter ut vult cl. Brown cum illis conjungendus.

Trib. I. *PEDICULARINÆÆ*.

Corolla irregularis plerumquè bilabiata; stam. 4 didynama.

I. *TOZZIA* Mich. gen. t. 16. L. gen. 745. St-Hil. mém. mus. ann. 7. p. 98.

Calyx campanul. subbilabiatus 4-dentatus; corolla tubulosa limbo 2-labiato subæquali, suprâ 2-lobo infrâ 3-lobo; stam. 4 didynama; antheræ basi setulæ aristatæ; ovarium 2-loculare loculis 2-spermis, ovulis oblongis per totam serè longitudinem affixis inferuè liberis; capsula sphaerica 2-valvis abortu 1-sperma.

1. *T. ALPINA* (L. sp. 844.) caule debili, floribus axillaribus in spicas foliosas interruptas dispositis. 2 in asperis humidis Jurassi, Alpium et Pyrenæorum rariis. — Fl. fr. 3. p. 487. Lam. ill. t. 522. — Col. cephr. 2. p. 49. t. 50. (v. v.)

II. *MELAMPYRUM* L. gen. 742.

Calyx tubulosus 4-fidus; corolla tubulosa 2-labiata compressa labio superiore galeato marginè replicato, inferiore sulcato 3-fido; stamina 4 didynama; capsula oblonga obliquè acuminata compressa, loculis monospermis; semina magna subovoidea.

1. *M. ARVENSE* (L. sp. 842.) foliis linearibus, bracteis ovatis dentato- setaceis coloratis, spicis conicis laxis, calycis pubescentis laciniis seabris lanceolato-linearibus aristato-acuminatis capsulâ longioribus. ① in agris. — Fl. fr. 3. p. 485. Fl. dan. t. 911. (v. v.)
2. *M. CRISTATUM* (L. sp. 842.) foliis linearibus, bracteis cordatis compactis denticulatis imbricatis, spicis quadrangularibus, calycis laciniis linearibus acutis. ① in sylvaticis. — Fl. fr. 3. p. 485. Fl. dan. t. 1104. (v. v.)
3. *M. MEMOROSUM* (L. sp. 843.) foliis lanceolato-ovatis, bracteis dentatis cordato-lanceolatis, summis coloratis sterilibus, floribus secundis lateralibus, calycis lanati laciniis lanceolatis ovatisve acuminatis capsulâ subbrevioribus. ① in Delphinatûs, Jurassi, agri Ligericini (Prost) montibus. — Fl. fr. 3. p. 486. Fl. dan. t. 305. (v. s.)
4. *M. PRATENSE* (L. sp. 843.) foliis elongato-lanceolatis superioribus basi pinnatifido-incisis, floribus axillaribus secundis, calycis laciniis linearibus acuminatis capsulâ brevioribus, corollis clausis. ① in pratis usque ad limitem inferiorem abietis. — Fl. fr. 3. p. 486. (v. v.)
5. *M. SYLVATICUM* (L. sp. 843.) foliis elongato-lanceolatis superioribus basi inciso-dentatis inferioribus integerrimis, calycis laciniis lanceolatis linearilanceolatisve acutis capsulâ subæquantibus, corollis hiantibus. ① in sylvis montosis supra limitem inferiorem abietis. — Fl. fr. 3. p. 488. — Fl. dan. t. 145. (v. v.)

### III. PEDICULARIS L. gen. 746.

Calyx ventricosus, plerumque 5-fidus vel inæqualiter 2-3-lobus; corollâ tubulosa 2-labiata labio superiore compresso galeato elongato sæpius emarginato, inferiore plano patulo 3-lobo; stam. 4 didynama; capsula compressa acuminata sæpè obliqua et calyce longior.

#### \* Foliis quaternis.

1. *P. VERTICILLATA* (L. sp. 846.) foliis canlinis pinnatifidis quaternis, pinnis oblongis dentatis, floribus verticillato-capitatis breviter pedicellatis, calycibus 5-dentatis, corollæ galeâ rectâ obtusâ labium inferius subæquante. ② in Alpibus, montibus Arvernæ, Cebennis, Pyrenæis. — Fl. fr. 3. p. 481. — Hall. helv. t. 9. f. 1. (v. v.)

#### \*\* Corollæ labio superiore utrinquè infrâ medium dentato.

2. *P. PALUSTRIS* (L. sp. 845.) caule ramoso, foliis pinnatis pinnis lanceolatis pinnatifidis dentatis, calycibus 2-partitis cristatis demum ovatis inflatis, corollæ labio superiore infrâ medium denticulato, galeâ dilatâ gibbâ obtusâ truncatâ. ① in paludibus. — Fl. fr. 3. p. 479. Lam. ill. t. 517. f. 1. — Riv. t. 92. (v. v.)

#### \*\*\* Corollæ labio superiore basi edentulo rostrato.

3. *P. ROSTRATA* (L. sp. 845.) caule declinato, foliis pinnatis, pinnis oblongis pinnatifidis dentatis, racemo paucifloro, calyce cylindrico 5-fido dentibus dilatatis serratis, corollæ galeâ falcato-rostratâ. ② in Alpibus et Pyrenæis. — Fl. fr. 3. p. 481. — Hall. helv. t. 8. f. 1. (v. v.)
4. *P. INCARNATA* (Jacq. aust. t. 140.) foliis profundè pinnatifidis, pinnis lineari-lanceolatis subpinnatifidis laciniis inæqualibus tenuissimè serrulatis, spicâ laxâ, calycibus villosis 5-fidis, corollæ galeâ falcato-rostratâ. ① in Alpibus Delph. et Galloprov. — Fl. fr. 3. p. 480. All. ped. t. 3. f. 2. (v. v. DC.)
5. *P. TUBEROSA* (L. sp. 847.) caulibus ascendentibus, foliis pinnatis, pinnis profundè pinnatifidis laciniis denticulatis, spicâ laxiusculâ, calyce campanulato 5-fido glabrinusculo, corollæ galeâ falcato-rostratâ. ② in Alpibus et Pyrenæis. — Fl. fr. 3. p. 483. — Hall. helv. t. 10. (v. v.)
6. *P. GYROFLEXA* (Vill. dauph. 2. p. 426. t. 9.) caule hirsuto, foliis pinnatis pinnis profundè pinnatifidis, laciniis denticulatis, calyce pubescente 5-fido

dentibus serratis, corollæ galeâ arcuatâ, rostro brevi conico. 2½ in Alpibus Delphin., Galloprovinciæ. — Fl. fr. 3. p. 481. — Hall. helv. t. 11. (v. v. DC.)

\*\*\*\* Corollæ labio superiore basi edentulo, galeâ brevi apice bidentatâ.

7. *P. sylvatica* (L. sp. 845.) caule basi ramoso, foliis pinnatis, pinnis ovatis acutè dentatis, spicâ pauciflorâ, calyce glabro 5-dentato fructifero inflato, corollæ galeâ obtusissimâ 2-dentatâ. (1) in sylvis paludosis. — Fl. fr. 3. p. 479. — Lob. ic. t. 748. f. 2 (v. v.)

8. *P. comosa* (L. sp. 847.) radice fasciculatâ, foliis pinnatis, pinnis pinnatifidis lacinii linearibus integris aut subincisis, calycibus 5-dentatis, corollæ galeâ obtusâ 2-dentatâ. 2½ in Alpibus, montibus Arvernæ, Ligeris (Lozère), Pyrenæorum. — Fl. fr. 3. p. 484. All. ped. t. 4. f. 1. Stev. mon. ped. t. 14. *P. asparagoides* Lapeyr. — Fl. flavi. (v. v.)

β. *rubens* (Stev.) corollâ purpureâ. — *P. fasciculata* Wild. sp. 3. p. 218. — Fl. fr. 3. p. 482. (v. v. DC.)

\*\*\*\*\* Corollæ labio superiore omninò edentulo, galeâ brevi.

9. *P. rosæ* (Jacq. ic. rar. t. 1. t. 115.) foliis pinnatis, pinnis linearibus integris aut pinnatifidis acutis, calyce hirsuto tubuloso inæqualiter 5-dentato, corollæ galeâ obtusâ. 2½ in Alpibus editioribus Delph. et Galloprov. — Fl. fr. 3. p. 482. — *P. hirsuta* All. ped. t. 3. f. 1. (v. s.)

10. *P. villosa* (L. mant. 86.) foliis pinnatis, pinnis pinnatifidis lacinii lanceolatis inæqualiter denticulatis, spicâ foliosâ, calyce obliquo 5-dentato, corollæ galeâ obtusissimâ. (2) in summis Vogesis (Mougeot), Alp. Delph., Arvernæ montibus ex Delarh, Pyrenæis. — Fl. fr. 3. p. 482. — Hall. helv. t. 9. f. 3. (v. v. DC.)

#### IV. RHINANTHUS Sm. brit. 649. — Rhinanthi sp. L.

Calyx 4-fidus ventricosus; corolla tubulosa 2-labiata, labio superiore compresso, inferiore plano 3-lobo; stamina 4 didynama; capsula compressa obtusa calyce tecta; semina compressa marginata.

1. *R. olabra* (Lam. fl. fr. 2. p. 352.) calycibus glabris. (1) in pratibus. — Fl. fr. 3. p. 478. *R. cristæ-galli* α et β. L. sp. 840. — Riv. t. 92. (v. v.)

2. *R. hirsuta* (Lam. fl. fr. 2. p. 353.) calycibus villosis. (1) in pratibus siccis. — Fl. fr. 3. p. 478. *R. cristæ-galli* γ. L. sp. 840. Prioris var? (v. v.)

#### V. BARTSIA L. gen. 739.

Calyx non ventricosus 4-fidus; corolla tubulosa 2-labiata; labium superius galeatum erectum integrum, inferius reflexum 3-fidum minimum; stamina galeâ breviora, antheræ tomentosæ; capsula ovata compressa acuminata.

##### Sect. 1. BARTSIA Stev. mon. ped. p. 4.

Corolla brevissimè 2-labiata, semina membranacea striata.

1. *B. alpina* (L. sp. 839.) foliis oppositis ovato-cordatis obtusè serratis, dentibus crebris, florum spicâ simplici subinterruptâ foliosâ, bracteis calycem superantibus, antheris hirsutis, capsulâ pilosâ calycem superante. 2½ in editioribus Vogesi, Jurassi, Alpium, montium Arvernæ, Pyrenæorum. — Fl. fr. 3. p. 476. — Fl. dan. t. 43. (v. v.)

2. *B. spicata* (Ran. Lull. phil. n. 42. p. 141. t. 10. f. 4.) foliis oppositis cordiformibus obtusè dentatis superioribus lanceolatis, dentibus remotis, floribus imbricato-apicatis, spicâ ramosâ, bracteis calyce sub brevioribus, staminibus galeam subæquantibus, antheris hirsutis, capsulâ hirsutâ calycem subæquante. 2½ in Pyrenæis. — Fl. fr. 3. p. 476. *B. Fagonii* Lapeyr. (v. s.)

##### Sect. 11. TRIXAGO Stev. l. c.

Corollæ galea semicylindrica brevis, semina minuta non alata.

3. *B. PURPUREA* (Dub. mes.) caule subsimplici aut à basi ramoso, foliis ovatis dentato-palmatis, floribus spicatis, spicâ basi interruptâ foliaceâ, calycis villosi laciniis lanceolatis obtusis, antheris subglabris, capsulâ glabrâ calycem subæquante. (1) in Galliæ australis pascuis et ad ripas Ataris (cl. L. Dufour). — *Trixago purpurea* Stev. mon. ped. p. 4. *Euphrasia latifolia* L. sp. 841. Fl. fr. 3. p. 473. — Magn. monsp. p. 94. ic. (v. v.)
4. *B. VISCOSA* (L. sp. 839.) caule subsimplici, foliis lanceolatis acuminatis grosse serratis, superioribus alternis, floribus distantibus lateralibus, calycis laciniis linearis lanceolatis elongatis subacutis, staminibus galeâ brevioribus, antheris hirsutis, capsulâ hirsutâ calyce duplò brevior. (1) in paludibus, ad rivulos à Nicæâ usque ad Nemetriam. — Fl. fr. 3. p. 477. — Barr. ic. t. 665. (v. v.)
5. *B. BICOLOA* (DC. ic. rar. t. 10.) pubescens, caule simplici aut subramoso, foliis oppositis lanceolato-linearibus hinc inde serratis, floribus densè spicatis, bracteis ovatis aut ovato-rotundis, calycis lanuginosi laciniis ovatis obtusis, antheris hirsutis, capsulâ villosâ calycis brevior. (1) in arenosis insularum Vindiliis Armoricæ (DC.), Reræ (Des Moul.). — Fl. fr. 5. p. 391. Semina striata non alata. (v. v. DC.)
6. *B. TRIxAGO* (L. sp. ed. 1. p. 602.) tomentosa, caule simplici, foliis oppositis lanceolatis obtusè serratis, floribus densè spicatis, bracteis ovato-rotundis, calycis lanuginosi dentibus minimis obtusis, antheris hirsutis, capsulâ villosâ calycem subæquante. (1) in Galliæ australioris maritimis humidis. — Fl. fr. 3. p. 476. — Moris. s. 11. t. 24. n. 8. Sem. striata. (v. v. DC.)
7. *B. VERSICOLOR* (DC. fl. fr. 3. p. 477.) hirsuta, caule simplicissimo, foliis oppositis, inferioribus linearis lanceolatis crenatis, superioribus linearibus inciso-serratis, spicâ densâ abbreviatâ, bracteis ovato-cordatis acuminatis, calycis lanuginosi dentibus obtusis, staminibus galeâ extus hirsutâ brevioribus, antheris hirsutis. (1) circa Antipolim (Stire). — *Rhinanthus versicolor* Wild. sp. 3. p. 190. — Barr. ic. t. 666. Fl. albido-purpurascens. (v. s.)
8. *B. MAXIMA* (DC. fl. fr. 3. p. 728.) pubescens, caule ramoso, foliis superioribus alternis oblongis obtusè dentatis, floribus densè spicatis, bracteis rotundo-ovatis, calycis lanuginosi dentibus obtusis, stam. galeâ brevioribus, antheris hirsutis, capsulâ villosâ calycem subæquante. (1) in pascuis macilentis circa Antipolim (Balbis), Nicæam ex All? et Corsicæ ex Lois. *Rhinanthus maximus* Wild. sp. 3. p. 189. — Barr. ic. t. 774. n. 2. Semina striata. Fl. maximi. (v. s.)

## VI. EUPHRASIA L. gen. 741.

Calyx 4-fidus; corolla tubulosa 2-labiata, labium superius galeatum emarginatum, inferius trilobum æquale; antheræ 2 aut 4 basi acuminatæ; capsula ovata compressa obtusa emarginata.

Sect. 1. EUPHRASIUM. — *Euphrasia* Stev. mon. ped. p. 4.

Antheræ 2 calcaratæ basi ciliatæ.

1. *E. OFFICINALIS* (L. sp. 841.) caule erecto ramoso, foliis ovatis dentatis dentibus acuminatis crenatisve, laciniis calycinis lanceolatis acuminatis, staminibus galeâ brevioribus, lobis labii corollæ inferioris emarginatis. (1) in pascuis aridis. — Fl. fr. 3. p. 472. et 5. p. 390. Lam. ill. t. 518. f. 1. *Bartsia imbricata* Lapeyr. excl. L. syn. (v. v.)

β. *Alpina*, foliis setaceo-dentatis, calycis glabri laciniis linearis-setaceis. — in Alpium et Pyren. pascuis aridis. — E. Alpina Lam. ill. t. 518. f. 2. Fl. fr. 3. p. 473. E. Salisburgensis Hop. (v. v.)

γ. *minima*, caule subsimplici minimo, foliis obtusis, laciniis calycinis lanceolatis. — In pascuis aridis Alpium et Pyren. — E. minima Jacq. Fl. fr. 3. p. 473. *Bartsia humilis* Lapeyr. ex cl. Benth. (v. v.)

Sect. II. ODONTITIS. — *Odontites* Dill. gen. 117. t. 6. Stev. l. c.

Antheræ omnes mucronatæ.

- a. *E. viscosa* (L. mant. 86.) caulibus erectis ramosis, foliis lineari-lanceolatis triuerviis subintegerrimis, calycibus viscoso-pubescentibus, stam. exsertis galeâ brevioribus, lobis lateralibus labii corollæ inferioris emarginatis, capsulâ hirsutâ calycem æquante. ① in Galliæ australis sterilibus. — Fl. fr. 3. p. 475. — Garid. aix. t. 80. (v. v.)
3. *E. LUTEA* (L. sp. 842.) caule erecto ramoso, foliis lanceolato-linearibus serratis, superioribus integerrimis, calycibus pubescentibus, staminibus exsertis corollâ longioribus, lobis lateralibus labii corollæ inferioris denticulatis, capsulâ hirsutâ calyce brevior. ① in aridis montosis Galliæ orientalis, mediæ australisque. — Fl. fr. 3. p. 474. Jacq. austr. t. 398. (v. v.)
4. *E. LINEIFOLIA* (L. sp. 842.) caule erecto ramoso, foliis linearibus omnibus integerrimis, calyce glabro aut adpressè pubescente, staminibus exsertis corollâ longioribus, capsulâ hirsutâ calycem subsuperante. ① in aridis Galliæ australis et vallium apricarum Alpium. — Fl. fr. 3. p. 475. — Col. cephr. 2. p. 68. t. 69. (v. v.)
5. *E. CORSICA* (Lois. gall. 2. p. 367.) caule basi prostrato ramoso, foliis linearibus integerrimis, floribus axillaribus subspicatis, calycibus corollam subæquantibus. ① in montibus Corsicæ. — Fl. fr. 3. p. 391. — Flores ut in 3 præced. lutei (v. s.)
6. *E. VIRENA* (Bell. app. 33.) caule ramoso, foliis elongato-lanceolatis laxèque dentatis, bracteis flore longioribus, calyce subhirsuto, lobis labii corollæ inferioris obtusis. ① in arvis Narbonæ. — Fl. fr. 5. p. 390. — Fl. vere. (v. s.)
7. *E. ODONTITES* (L. sp. 841.) caule erecto ramoso, foliis lineari-lanceolatis serratis, bracteis flore subbrevioribus, calyce hirsuto, lobis labii corollæ inferioris obtusis. ① in arvis pascuisque sterilibus. — Fl. fr. 3. p. 474. excl. var. β. — Fl. dan. t. 625. (v. v.)

## VII. SIBTHORPIA L. gen. 775.

Calyx 5-partitus; corolla subrotata, tubo brevi, limbo regulari 5-lobo; stamina 4 didynama; stigma capitatum; capsula compressa orbicularis apice dehiscens.

1. *S. EUROPEA* (L. sp. 880.) caulibus gracilibus diffusis procumbentibus, foliis reniformi-cordatis longè petiolatis. ½ in udibus aggeribus, ad margines rivulorum Galliæ occidentalis. — Fl. fr. 3. p. 472. — Pluck. t. 7. f. 6. (v. v. DC.)

## Trib. II. VERONICEÆ.

- Stam. 2; cor. rotata inæqualis.

## VIII. VERONICA Tourn. t. 60. L. gen. n. 25.

Calyx 4-rarò 5-partitus; corolla rotata 4-loba inæqualis; capsula compressa, ovata aut obcordata.

§. 1. *Annua*, pedunculis solitariis 1-floris.

1. *V. CYMBALARIA* (Bod. diss. 1798.) caule ramoso diffuso, foliis rotundis subcordatis sub 5-7-lobis petiolatis florum pedunculis brevioribus, laciniis calycinis ovalibus obtusis ciliatis capsulam obcordatam lateraliter hirsutam subsuperantibus, seminibus in quoque loculo binis magnis umbilicato-concavis margine corrugatis dorso læviusculis. ① in cultis et muris vetustis Temonis, Mompelii. — Fl. fr. 5. p. 389. Viv. fragm. 1. t. 16. f. 1. — Buxb. cent. 1. p. 25. f. 2. (v. s.)
2. *V. HADEREFOLIA* (L. sp. 19.) caule ramoso patulo, foliis rotundato-cordatis petiolatis sub 5-lobis florum pedunculos subsuperantibus, laciniis calycinis ovato-cordatis ciliatis capsulam glabram cordatam superantibus, seminibus in quoque loculo binis umbilicato-concavis extus corrugatis. ① in ruderatis, agris, hortis. — Fl. fr. 3. p. 467. — Fl. dan. t. 428. (v. v.)
3. *V. FILIFORMIS* (Savi. bot. et. 1. p. 15.) caulibus diffusis prostratis, foliis

- ovatis suboordatis grossè dentatis pedunculis duplò triplòve brevioribus, lacinii calycinis lanceolatis divaricatis capsulam obcordatam ciliatam superantibus, seminibus plurimis umbilicato-concavis extùs corrugatis. ① in cultis Niceæ, Telonis, Mompeli et Tolosæ ex Benth. Burdigalæ (Des Monl.) — Fl. fr. 5. p. 388. — Benth. cent. 1. t. 10. f. 1 et 2. V. Buxbaumii Ten. (v. s.)
4. V. *ACASTIS* (L. sp. 18.) caule patulo diffuso, foliis cordato-ovatis inciso-crenatis pedunculis brevioribus, calycis lacinii ovatis acutis capsulâ pubescente longioribus, seminibus plurimis umbilicato-concavis extùs corrugatis. ① in agris, hortis, ad maros, etc. — Fl. fr. 3. p. 467. V. pulchella Bast. es. 414. Fl. fr. 5. p. 388. — Fuchs. p. 22. ic. Variat floribus albis et cæruleis. (v. v.)
5. V. *ARVENsis* (L. sp. 18.) caule ramoso, ramis diffusis, foliis ovato-cordatis crenato-dentatis, floralibus alternis oblongis integerrimis calycem superantibus, floribus subsessilibus, lacinii calycinis lanceolatis obtusis inæqualibus capsulam ciliatam obcordatam superantibus, seminibus plurimis compressis rugulosis. ① in arvis, cultis. — Fl. fr. 3. p. 466. — Fl. dan. t. 515. (v. v.)
6. V. *DIGITATA* (Vahl. symb. 1. p. 2.) caule stricto, foliis flores sessiles superantibus omnibus digitato-partitis lacinii linearibus obtusis, lacinii calycinis linearibus capsulam truncatam superantibus, seminibus plurimis leviusculis subangulatis. ① in agris Mompeli ex Vahl, agri Ruscionensis ex Lapeyr. — Fl. fr. 3. p. 466. (v. s. ex Hisp.)
7. V. *VERNA* (L. sp. 19.) caule stricto subsimplici aut à basi ramoso, foliis inferioribus digitato-partitis pinnatifidisve lacinii inæqualibus obtusis, summis indivisis linearibus calyces subsessiles superantibus, lacinii calycinis linearibus aut lineari-lanceolatis acutis capsulam obcordatam glanduloso-ciliatam superantibus, seminibus plurimis lenticulari-compressis leviusculis. ① in aridis passim. — Fl. fr. 3. p. 465. V. succulenta et V. Bellardi All. ped. t. 22. f. 4. et t. 85. f. 1. (v. v.)
8. V. *TRIPHYLLOS* (L. sp. 19.) pubescens, caule ramoso patulo, foliis infimis oordatis dentatis, superioribus florum pedunculis subbrevioribus 3-5-digitato-partitis lacinii obtusis, lacinii calycinis lanceolato-ovatis obtusis, capsulâ obcordatâ glanduloso-ciliatâ, seminibus umbilicato-concavis extùs rugulosis. ① in arvis, cultis et humidis. — Fl. fr. 3. p. 467. — Lob. ic. 464. f. 1. (v. v.)
9. V. *ACUTIFOLIA* (L. sp. 19.) subpubescens, caule erecto ramoso, foliis ovato-oblongis obtusis crenatis, superioribus integris florum pedunculos subæquantibus, lacinii calycinis ovatis obtusis capsulâ obcordatâ profundè emarginatâ glanduloso-ciliatâ brevioribus, seminibus plurimis lævigatis intùs planis extùs convexis. ① in humidis cultis, etc. — Fl. fr. 3. p. 464. Vaill. bot. t. 33. f. 3. V. Burdigalensis Laterr. fl. bord. ed. 2<sup>a</sup>. p. 87. (v. v.)
10. V. *PÆCOX* (All. suet. 5. t. 1. f. 1.) villosa-pubescens, caule ramoso erecto, foliis ovatis cordatis petiolatis profundè dentatis, superioribus incisive integrisve floris pedunculum superantibus, lacinii calycinis ovato-lanceolatis obtusis capsulam obcordatam glanduloso-ciliatam subsuperantibus, seminibus plurimis umbilicato-concavis extùs lævibus. ① in cultis passim. — Fl. fr. 3. p. 465. (v. v.)
11. V. *PEREGRINA* (L. sp. 20.) glabra, caulibus ramosissimis ramis ascendentibus, foliis lanceolato-oblongis obtusis integerrimis dentatisque, superioribus lineari-lanceolatis flores subsessiles superantibus, calycis lacinii lineari-lanceolatis acutis capsulam glabram obcordatam superantibus, seminibus plurimis lævibus intùs planis extùs convexis. ① in Galliæ australis arvis cultis. — Fl. fr. 3. p. 464. Fl. dan. t. 407. (v. v.)
12. V. *REPENS* (Clar. in DC. fl. fr. 3. p. 727.) caule repente, foliis ovato-rotundis integris pedunculo æqualibus, lacinii calycinis ovato-lanceolatis obtusis capsulâ obcordatâ ciliatâ brevioribus, seminibus plurimis lævibus subcompressis. ① in Corsicæ montibus. — Lois. gall. t. 1. V. tenella Viv. cora. an All. ped. t. 22. f. 1? (v. s.)



§. 2. *Perennes, racemis terminalibus.*

13. *V. NUMMULARIA* (Gou. ill. 1. p. 1. f. 2.) caulibus prostratis fruticosis, foliis confertis subrotundo-ovatis obtusis glabris, racemo brevi conferto, laciniis calycinis lanceolatis obtusis ciliatis capsulâ glabrâ obcordatâ subbrevioribus, semin. plurimis lævibus.  $\frac{1}{2}$   $\frac{3}{4}$  in editioribus Pyrenæorum. — Fl. fr. 3. p. 470. *V. irregularis* Lapeyr. (v. v.)
14. *V. SERRYLLOPOLIA* (L. sp. 15.) caule basi repente, foliis ovatis glabris obtusis, inferioribus obsolete crenatis oppositis, superioribus pedicellos subæquantibus, floribus solitariis racemosis, laciniis calycinis glabris lanceolato-ovatis obtusis capsulâ obcordatâ glabrâ brevioribus, semin. plurimis lævibus minutis lenticulari-compressis.  $\frac{1}{2}$  in sylvaticis et pratis subhumidis. — Fl. fr. 3. p. 471. Fl. dan. t. 492. (v. v.)
15. *V. ALPINA* (L. sp. 15.) caule simplici erecto folioso, foliis oblongo-ovatis obtusis integris aut dentatis, racemo conferto abbreviato, calycis hispidi laciniis lineari-lanceolatis obtusis capsulâ hirtâ obovato-cordatâ brevioribus.  $\frac{1}{2}$  in Alpihus, Monte-Anreo, Pyrenæis. — Fl. fr. 3. p. 471. — Hall. helv. t. 15. f. 2. (v. v.)
16. *V. PAUTICULOSA* (L. sp. 15.) caulibus fruticulosisscendentibus diffusisque, foliis lanceolatis subdentatis obtusis, racemo corymbiformi laxo, calycis pubescentis laciniis lanceolatis obtusis capsulam ovatam pubescentem subæquantibus, seminibus plurimis lævibus lenticulari-compressis.  $\frac{1}{2}$  in Alpihus, Pyrenæis et Corsicæ montibus (Soleirol). — Fl. fr. 3. p. 469. — Hall. helv. t. 16. f. 1. (v. v.)
- β. saxatilis* (Lin.) foliis ovatis obtusissimis. — *V. saxatilis* L. f. supp. 83. Fl. fr. 3. p. 469. (v. v.)
17. *V. BELLIDIOIDES* (L. sp. 15.) caule simplici ascendente sub 4-phyllo, foliis villosis pubescentibusque ovato-oblongis obtusis, radiculis confertis apice crenatis, racemo conferto abbreviato, calycis villosi laciniis lanceolatis obtusis capsulâ villosâ ovatâ subcompressâ brevioribus, semin. plurimis lævibus lenticulari-compressis.  $\frac{1}{2}$  in Alpihus, Pyrenæis. — Fl. fr. 3. p. 470. — Hall. helv. t. 15. f. 1. (v. v.)
18. *V. PORA* (Gou. ill. p. 1. t. 1. f. 1.) caule simplicissimmo, foliis oppositis cordato-ovatis dentatis sessilibus, floribus racemosis, bracteis linearibus pedicella multo brevioribus, laciniis calycinis lineari-lanceolatis obtusis capsulâ multo brevioribus, seminibus plurimis lævibus lenticulari-compressis.  $\frac{1}{2}$  in Pyrenæis. — Fl. fr. 3. p. 469. (v. v.)

§. 3. *Perennes, racemis terminalibus spicatis.*

19. *V. LONGIFOLIA* (L. sp. 13.) caule erecto apice polystachyo, foliis oppositis verticillatisque lanceolatis argutè serratis acuminatis, racemo spiciformi, calycis hirsuti laciniis lanceolato-linearibus acutis capsulâ globosâ compressiusculâ submarginatâ brevioribus, seminibus plurimis minutis lævibus.  $\frac{1}{2}$  in Alsatia ex Mapp.; propè Fontembellaqueum ex Thuill. et in Pyrenæis ex Lapeyr. — Fl. fr. 3. p. 468. — Schrad. veron. spic. t. 2. f. 1. (v. v. c.)
20. *V. SPICATA* (L. sp. 14.) caule ascendente pubescente, foliis oppositis oblongis serratis apice integerrimis, inferioribus crenatis acutis aut subobtusis, racemo spiciformi elongato, calycis hirsuti laciniis lanceolatis acutis capsulâ globoso-compressiusculâ vix emarginatâ brevioribus, semin. plurimis minimis lævibus lenticularibus.  $\frac{1}{2}$  in sylvaticis. — Fl. fr. 3. p. 468. — Vaill. bot. t. 33. f. 4. (v. v.)

§. 4. *Perennes, racemis axillaribus.*

21. *V. ALLIOWII* (Vill. dauph. 2. p. 8.) caule subrepente fruticuloso prostrato glaberrimo, foliis rigidis nitidis glaberrimis ovatis obovatisve acutis serrulatis, racemo spiciformi denso multifloro, laciniis calycinis lineari-lanceolatis ciliatis subacutis capsulâ pubescente obcordatâ brevioribus.  $\frac{1}{2}$  in Alpihus.

- Delph., Gallopr., Pyrenæorum ex Lapeyr. — Fl. fr. 3. p. 463. V. Pyrenaica All. ped. t. 46. f. 3. (v. v. DC.)
22. V. OFFICINALIS (L. sp. 14. æ.) caulibus procumbentibus hirtis ramosis ascendentibus, foliis pilosis rugosis ovatis crenato-dentatis subacutis, racemo laxiusculo, lacinii calycinis lineari-lanceolatis acutis capsulâ ciliatâ obcordatâ brevioribus, seminibus plurimis lenticulari-compressis. ½ in sylvestribus sterilibus. — Fl. fr. 3. p. 463. Lam. ill. t. 13. f. 2. (v. v.)
23. V. PROSTRATA (L. sp. 22.) caulibus prostratis pubescentibus, foliis lineari-lanceolatis inciso-serratis integerrimisque, racemis elongatis post florescentiam laxis, calycis glabri lacinii inæqualibus linearibus subacutis, majoribus capsulam obcordatam glaberrimam subæquantibus, seminibus plurimis lævibus compressis. ½ in collibus, pascuis, siccis passim. — Fl. fr. 3. p. 461. (v. s.)
24. V. CHAMÆDRYS (L. sp. 17.) caulibus prostratis ascendentibus bifariam pilosis, foliis ovatis sessilibus rugosis grossè dentatis inferioribus glabris, racemis laxis elongatis, calycis lacinii lineari-lanceolatis subacutis ciliatis capsulam obcordatam subciliatam superantibus. ½ in pratis, ad sepes. — Fl. fr. 3. p. 460. — Fuchs. 872. ic. (v. v.)
25. V. LATIFOLIA (L. sp. 18.) caule erecto stricto pubescente, foliis ovatis sessilibus subcordatis grossè dentatis incisive, racemis elongatis sublaxis, lacinii calycinis lineari-lanceolatis subacutis ciliatis inæqualibus, majoribus capsulam obcordatam subciliatam subæquantibus. ½ in pascuis Alsatie. — Fl. fr. 5. p. 387. V. pseudo-chamædris Jacq. austr. 1. t. 60. (v. s.)
26. V. TRUCRUM (L. sp. 16.) caulibus ascendentibus villosis, foliis lanceolatis acutis rugosis pilosis inciso-dentatis dentatisque sessilibus, racemis laxis elongatis, calycis hirsuti lacinii linearibus subacutis inæqualibus, majoribus capsulam obcordatam hirsutam superantibus. ½ in pascuis siccis. — Fl. fr. 3. p. 460. — Dalech. hist. 1165. f. 2. ic. (v. v.)
27. V. URTICÆFOLIA (L. f. supp. p. 83.) caule stricto, foliis ovatis sessilibus cordatis acutis grossè serratis, racemis laxis elongatis, calycis pubescentis lacinii lanceolatis subobtusis capsulâ pubescente obcordatâ multo brevioribus. ½ in montosis nemorosis. — Fl. fr. 3. p. 459. — Dalech. hist. 1165. f. 1. (v. v.)
28. V. DUBIA (DC. fl. fr. 3. p. 462.) caule basi radicante ascendente piloso, foliis ovatis subsessilibus grossè crenato-dentatis obtusis basi angustatis, racemis laxis elongatis, lacinii calycinis ovato-lanceolatis acutis capsulam cuneato-obcordatam pubescentem duplò superantibus, loculis monospermis, seminibus lævibus lenticulari-compressis. ½ in collibus Alsatie ex Vill. (v. s. c.)
29. V. MONTANA (L. sp. 17.) caule prostrato petiolisque hirtis, foliis petiolatis ovatis obtusis grossè serratis, racemis laxissimis panicifloris, lacinii calycinis lanceolato-ovatis obtusis capsulâ glabrâ ciliatâ margine undulatâ latissimâ brevioribus, seminibus plurimis magnis lævibus compressis. ½ in umbrosis humidis. — Fl. fr. 3. p. 459. Jacq. austr. t. 109. (v. v.)
30. V. SCUTELLATA (L. sp. 16.) caule debili basi decumbente, foliis sessilibus linearibus subdentatis, racemis laxissimis, pedicellis fructiferis pendulis, lacinii calycinis lanceolatis subobtusis capsulâ glaberrimâ compressâ obcordatâ latissimâ brevioribus, seminibus plurimis lævibus compressis. ½ in inundatis. — Fl. fr. 3. p. 461. Fl. dan. t. 209. V. parvularia Poit. et Turp. fl. paris. t. 14. (v. v.)
31. V. ANAGALLIS (L. sp. 16.) caule erecto, foliis sessilibus ovato-lanceolatis lanceolatisve obsolete serratis, racemis laxis, lacinii calycinis lanceolatis capsulam glaberrimam vix emarginatam superantibus, seminibus plurimis minutis lævibus. ½ ad fossas inque inundatis. — Fl. fr. 3. p. 461. Fl. dan. t. 903. (v. v.)
32. V. BECCABUNGA (L. sp. 16.) caule repente ascendente, foliis glaberrimis ovatis obtusissimis obsolete serratis in petiolum subattenuatis, racemis laxis,

lacinii calycinis lanceolato-ovatis acutis capsulam glaberrimam vix emergentem subquantibus, seminibus plurimis minutis lævibus subovoideis. ♀ ad scaturigines. — Fl. fr. 3. p. 462. — Riv. t. 100. (v. v.)

33. *V. APETALA* (L. sp. 14.) surculis serpentibus apice foliosis, foliis confertis obovatis pubescentibus dentatis, pedunculo nudo axillari paucifloro, floribus corymboso-peniculatis, calycis lacinii linearilanceolatis obtusis capsulâ ovato-cordatâ multo brevioribus. ♀ in Alpium frigidia umbrosis. — Fl. fr. 3. p. 463. — Pluck. t. 114. f. 3. (v. v.)

### Ordo LXXIII. LABIATÆ Juss. gen. p. 110.

Calyx 1-sepalus tubulosus 5-fidus aut 2-labiatus. Corolla hypogyna tubulosa irregularis sæpius 2-labiata, labio superiore sæpius bifido, inferiore 3-fido. Stamina rarius 2, sæpissimè 4 didynama sub labio corollæ superiore inserta. Anth. 2-loculares. Ovarium liberum 4-lobum disco hypogyno crasso appciturum; stylus unicus inter ovarii lobos enatus; stigma bifidum. Semina 4 nuda (cariopsides) basi styli dilatata adnata, imo calyce recondita, interdum pilis obsecta. Albumen in Europæanis speciebus nullum; embryo rectus; radícula infera; cotyledones planæ. — Caulis herbaceus aut suffrutescens tetragonus, folia opposita; flores oppositi aut verticillati, axillares aut spicati, bracteis sæpè stipati.

#### † STAMINA 2 FERTILIA.

##### I. LYCOPUS Tourn. inst. t. 89. L. gen. 33.

Calyx tubulosus 5-fidus ore nudo; corolla tubulosa 4-loba subæqualis, lobo superiore latiore emarginato.

1. *L. EUROPEUS* (L. sp. 30.) foliis sinuato-serratis appendiculatis. ♀ in paludosis et humidis. — Fl. fr. 3. p. 505. — Moris. s. 11. t. 9. f. 5. n. 20. (v. v.)
2. *L. EXALTATUS* (L. f. supp. 87.) foliis pinnatifidis, caule altissimo. ♀ in Gallie australis humidis. — Fl. fr. 3. p. 505. — Pluck. t. 45. f. 1. (v. v. c.)

##### II. CUNILA L. gen. 35.

Calyx cylindricus 10-striatus 5-dentatus, ore villis clauso; corolla 2-labiata, labium superum erectum planum emarginatum, inferum 3-lobum.

1. *C. HYMORBUS* (L. sp. 31.) foliis ovalibus integerrimis, floribus verticillatis. (1) Mompeli ex Gouan. Dubia civis. — Fl. fr. 3. p. 506. — Moris. s. 11. t. 19. f. 9. n. 6. (v. s. c.)

##### III. ROSMARINUS Tourn. inst. t. 92. L. gen. 38.

Calyx compressus 2-labiatus suprâ integer infrâ 2-fidus, ore nudo; corolla 2-labiata, labio superiore 2-fido, inferiore 3-fido.

1. *R. OFFICINALIS* (L. sp. 33.) foliis sessilibus, 3 in Gallie mediterraneæ montosis apricis. — Fl. fr. 3. p. 506. — Moris. s. 11. t. 16. f. 2 et 3. n. 1 et 2. (v. v.)

IV. SALVIA L. *gen. n. 39.*

Calyx subcampanulatus 2-labiatus labio superiore 3-dentato, infero 2-fido, ore nudo; corolla ringens superius fornicata emarginata; antheræ loculis 2, uno fertili, altero abortivo, connectivo filiformi transverse filamento adnato dissitis.

Sect. I. HORMINUM *Tourn. t. 82.*

Labium corollæ superius fornicatum non compressum.

1. *S. OFFICINALIS* (L. sp. 34.) foliis lanceolato-ovatis crenulatis, verticillis paucifloris, calycis bractæas superantis lacinii acuminatis. ♂ in Galliâ australi passim. — Fl. fr. 3. p. 507. — Moris. s. 11. t. 15. f. et n. 1. (v. v.)
2. *S. HORMINUM* (L. sp. 34.) foliis obtusis crenatis, bracteis summis sterilibus majoribus coloratis, stylo labio superiore longiore, calycis bracteam superantis labio superiore subemarginato dentibus mucronatis, labio inferiore lacinii acuminatis. ① secus agros Nicæ ex All., agri Ruseignonnensis ex Lapeyr. — Fl. fr. 3. p. 510. — Lob. ic. t. 555. f. 2. (v. v. c.)
3. *S. VERBENACA* (L. sp. 35.) caule hirtio, foliis lanceolatis serratis sinuatis aut grosse dentato-crenatis læviusculis obtusis, bracteis rotundato-ovatis mucronatis, calycis hirsuti dentibus minimis conniventibus, lacinii mucronatis, stylo labio superiore brevior. ♀ in pascuis siccis montanis Galliæ australis et occidentalis. — Fl. fr. 3. p. 511. — Barr. ic. t. 207. (v. v.)
- ℓ. incisa* (DC.) foliis inferioribus subincisis, corollis calycem vix excedentibus. — In Delphinatû et in muris Aquarum Tarbellicarum. — Barr. ic. t. 208. (v. v.)
4. *S. CLAUDESTINA* (L. sp. 36.) caule subsimplici, foliis oblongo-lanceolatis sinuatis pinnatifidiave dentatis nervosis, bracteis rotundo-ovatis acuminatis, calycis substriati hirsuti labii superioris dentibus minimis conniventibus, inferioris lacinii acutis, corollis calyce duplò longioribus. ♀ in Galliâ australi secus vias primo vers. — Fl. fr. 5. p. 395. *S. præcox* Lois. not. p. 6. — Barr. ic. t. 220. (v. v.)
5. *S. VERTICILLATA* (L. sp. 37.) foliis ovato-cordatis crenato-dentatis, verticillis subnudis multifloris, floribus pedicellatis, calycis hirsuti dentibus distinctis lacinii acuminatis, stylo corollæ labio inferiori incumbente. ① ad agrorum et viarum margines rariùs. — Fl. fr. 3. p. 511. — Riv. monop. t. 60. (v. v. c.)

Sect. II. SCLAREA *Tourn. t. 82.*

Labium superius corollæ compressum.

6. *S. PRÆTENSIS* (L. sp. 35.) foliis nervosis rugosis cordato-oblongis crenatis summis amplexicaulibus, verticillis subnudis, calycis labii superioris dentibus minimis, inferioris lacinii mucronatis, stylo galeâ falcatâ subduplò longiore. ♀ in pratia. — Fl. fr. 3. p. 508. — Cam. epit. 629. ic. (v. v.)
7. *S. PYRÆNAICA* (L. syst. 71.) villosa-viscosa, foliis dentato-sinuatis obtusis, staminibus corollâ duplò longioribus, labio calycis superiore 2-dentato, inferiore 3-lobo lobis lateralibus ovalibus, medio subulato. ♀ in Pyrenæis? — Fl. fr. 5. p. 394. — Herm. parad. 187. ic. Dubia civis. (in herb. cl. Jussieu v. DC.) †
8. *S. SYLVESTRIS* (L. sp. 34.) caule pubescente, foliis oblongo-lanceolatis subcordatis rugosis acutis aut subacutis bicrenatis, bracteis coloratis calyce brevioribus, labii superioris dentibus minimis, inferioris lacinii acuminatis, stylo recurvo galeam superante. ♀ in Galliâ australi, in vineis et ad margines agrorum rariùs. — Fl. fr. 3. p. 508. Jacq. austr. t. 212. (v. v. c.)
9. *S. SCLAREA* (L. sp. 38.) foliis rugosis ovato-cordatis villosis serratis, bracteis coloratis calyce longioribus, calycis dentibus lacinii setaceo-acuminatis.

- ② in aridis montosis Galliæ mediæ, occidentalis australisque. — Fl. fr. 3. p. 508. — Lob. ic. 556. f. 2. (v. v.)
10. *S. GLUTINOSA* (L. sp. 37.) villosa-viscosa, foliis cordato-sagittatis grossè serratis acuminatis, calyce bilabiato, labio superiore dentibus obsoletis 1-dentato, inferiore subbidentato dentibus acutis. 2/ in sylvis montanis. — Fl. fr. 3. p. 509. — Lob. ic. 557. f. 1 et 2. (v. v.)
11. *S. ETRUSCÆ* (L. sp. 39.) caule foliis calycibusque arachnoideo-lanuginosis, foliis oblongo-ovatis incisim dentatis, bracteis recurvatis subspinosis calyce brevioribus, calycis laciniis dentibusque subspinosis. ② in aridis Galliæ australioris, agri Ligericini (Prost.) Arvernæ ex Del. — Fl. fr. 3. p. 509. — Moris. s. 11. t. 13. f. 1. n. 4. (v. v. DC.)

†† STAMINA 4 DIDYNAMA FERTILIA.

V. *AJUGA* Schreb. unilab. 24. — *Ajuga* et *Teucrii* sp. L.

Calyx 5-fidus subæqualis; corolla tubulosa labiata, labio superiore minimo bidentato, inferiore 3-lobo lobo medio magno obcordato; cariopsides reticulatæ.

Sect. I. *CHAMÆPITYS* Tourn. t. 98.

Flores axillares solitarii.

1. *A. CHAMÆPITYS* (Schreb. unilab. 24.) caulibus diffusis hirtis, foliis trifidis laciniis linearibus, floribus axillaribus solitariis folio brevioribus. ① in agris, arenosis, lapidosis. — Fl. fr. 3. p. 514. — Lob. ic. 582. f. 2. (v. v.)
2. *A. PSEUDO-IVA* (Rob. et Cast. in DC. fl. fr. 5. p. 395.) caulibus diffusis lanatis, foliis linearibus parè dentatis margine subrevolutis, floribus axillaribus solitariis flavis. ① secus vias Massiliæ et ad littus occidentale insulæ S.-Luciæ propè Narbonam ex cl. Benth. Inodora. An potiùs ut sequentis varietas habenda? (v. s.)
3. *A. IVA* (Schreb. unilab. 25.) caulibus diffusis lanatis, foliis linearibus antice dentatis, floribus axillaribus solitariis purpureis. ① in Galliâ mediterraneâ. — Fl. fr. 3. p. 514. Moris. s. 11. t. 22. f. 11. n. 3. (v. v.)

Sect. II. *BUGULA* Tourn. t. 98.

Flores axillares spicati.

4. *A. ALPINA* (L. mant. 80.) caule erecto non stolonoso, foliis obovato-spathulatis, radicalibus crenulatis, caulinis dentatis flores sessiles superantibus, calycis hirsuti laciniis linearibus acuminatis. 2/ in sylvis montosis. — Fl. fr. 3. p. 513. (v. v.)
5. *A. REPTANS* (L. sp. 785.) stolonibus reptantibus, foliis obovato-spathulatis obsolete crenatis aut undulatis, floribus subsessilibus, calycis hirsuti bractea brevioris laciniis linearibus acutis. 2/ in sylvaticis humidis. — Fl. fr. 3. p. 512. Lam. ill. t. 501. f. 2. (v. v.)
6. *A. PYRAMIDALIS* (L. sp. 785.) caulibus tetragonis villosis hirsutis non stolonosis, foliis obovatis margine sinuatis obtusis, radicalibus in petiolum attenuatis, superioribus flores sessiles superantibus, calycis hirsuti laciniis acutis, labio corollæ inferiore majore trilobo, intermedio emarginato. ② in sylvaticis. — Fl. fr. 3. p. 513. An satis à priore distincta? (v. s.)
7. *A. GENEVENSIS* (L. sp. 785.) villosa, non stolonosa, foliis ovato-lanceolatis margine sinuatis obtusis, caulinis dentatis radicalibus longioribus, calycis laciniis acutis, lobo intermedio labii corollæ inferioris brevioris. 2/ in aridis sylvaticis passim. — Fl. fr. 3. p. 513. Fl. sæpiùs purpurei. Vix à prioribus sejungenda. (v. v.)

VI. TEUCRIUM *Schreb. similab. p. 17. — Teucrii sp. L.*

*Calyx* tubulosus rursù campanulatus 5-fidus; corolla tubo brevi labiata, labio superiore 2-partito lacinii ad latera deflexis, inferiore 3-lobo lobo medio majore; stamina intrâ fissuram labii superioris exserta.

Sect. I. SCORODONIA *Adans. fam. p. 188.*

*Calyx* 2-labiatus, labio inferiore 4-dentato, superiore lacinia 1 ovata.

1. *T. SCORODONIA* (L. sp. 789.) caulibus erectis tetragonia villosis, foliis ovato-cordatis pubescentibus aut glabresculis, floribus racemosis, racemis axillaribus secundis, calyce 2-labiato, labii inferioris dentibus mucronatis brevibus, labii superioris lacinia ovata mucronata, seminibus lævibus. ♀ in sylvis arenosis. — Fl. fr. 3. p. 516. — Lob. ic. 497. f. 2. Variat calycibus hirtis et glabris. (v. v.)
2. *T. MASSILIENSE* (L. sp. 789 excl. Barr. syn.) caulibus erectis suffruticosis, foliis oblongo-ovatis rugosis crenatis incanis, floribus elongato-racemosis, calyce 2-labiato, seminibus lævibus. ♂ in Corsicâ, Stachadibus. — Fl. fr. 3. p. 519. — Ger. gallop. t. 11. (v. a.)

Sect. II. TEUCRIUM *Tourn. t. 93.*

*Calyx* campanulatus 5-fidus; semina magna pubescentia subreticulata.

3. *T. FRUTICANS* (L. sp. 787.) caule frutescente erecto, foliis ovato-lanceolatis obtusis integerrimis subtus niveis, floribus binis solitariisve axillaribus, calycibus tomentosis campanulatis lacinii ovato-lanceolatis acutis. ♂ in agro Ruscinonensi propè Banyuls. — Fl. fr. 3. p. 515. — Dill. elth. 379. t. 284. (v. v.)
4. *T. PSEUDO-CHAMÆDREYS* (L. sp. 787.) caule erecto hirtio, foliis 3-partitis lacinii linearibus acutis margine subrevolutis, floribus pedicellatis solitariis laxè racemosis, calycis campanulati lacinii linearilanceolatis acuminatis. ♀ in sterilibus maritimis Massiliæ. — Fl. fr. 3. p. 516. — Lob. ic. 385. f. 1. (v. a.)

Sect. III. CHAMÆDREYS *Dill. gen. t. 3.*

*Calyx* tubuloso-campanulatus 5-dentatus; semina parva glabra.

5. *T. BOTRYS* (L. sp. 786.) caulibus ramosis sub 4-gonis hirsutis, foliis pubescentibus mollibus multifidis lacinii linearilanceolatis, verticillis dimidiatis paucifloris, calycis tubulosi basi dilatati dentibus acuminatis, seminibus punctato-reticulatis. ① in apricis cultis. — Fl. fr. 3. p. 515. — Lob. ic. 385. f. 2. (v. v.)
6. *T. LUCIDUM* (L. sp. 790.) caulibus erectis lævibus, foliis ovatis glabris incisodentatis, floralibus dentatis aut integris calycem superantibus, verticillis dimidiatis in racemosis digestis, calycis campanulati dentibus lanceolatis acutis æqualibus, seminibus sublævibus. ♀ in sylvis Alpium austral. Galloprovinciæ. — Fl. fr. 3. p. 518. — Magn. hort. t. 5a. (v. a.)
7. *T. CHAMÆDREYS* (L. sp. 790.) caulibus subprocumbentibus villosis, foliis cuneiformi-ovatis incisis crenatis petiolatis, floribus ternis, calycis campanulati hirti dentibus lanceolato-acuminatis, seminibus lævibus. ♀ in apricis sylvaticis. — Fl. fr. 3. p. 518. *T. pseudo-chamædrys* Salzm. cors. exsicc. — Lob. ic. 491. Variat foliis hirsutis et subglabris. (v. v.)
8. *T. FLAVUM* (L. sp. 791.) caulibus erectis fruticosis pubescentibus, foliis rotundato-ovatis crenatis petiolatis, floralibus lanceolatis integerrimis, verticillis dimidiatis in racemosis digestis, floribus pedunculatis, calycis tomentosi campanulati dentibus ovatis acutis, seminibus sublævibus. ♂ in collibus asperis Galliæ mediterraneæ. — Fl. fr. 3. p. 519. — Lob. ic. 490. f. 1. (v. v.)
9. *T. SCORDIUM* (L. sp. 790.) caulibus diffusis hirtis, foliis oblongis sessilibus dentatis pubescentibus, floribus axillaribus geminis brevissimè pedunculatis, calycis parvi subcampanulati dentibus acutis, seminibus punctato-reticulatis.

- ¶ in paludosis. — Fl. fr. 3. p. 517. — Moris. s. 11. t. 22. f. 8. n. 14. (v. v.)
10. *T. PYRENAICUM* (L. sp. 791.) caulibus procumbentibus hirtis, foliis cuneiformi-orbiculatis crenatis pilosis, floribus pedicellatis terminalibus corymboso-capitatis, calycis campanulati dentibus subæqualibus ciliatis acuminatis, seminibus reticulatis. ¶ in rupestribus et vallibus Pyrenæorum, Arvernæ ex Del. — Fl. fr. 3. p. 520. (v. v.)
11. *T. MONTANUM* (Schreb. unil. 50.) procumbens, caulibus pubescentibus, foliis lanceolatis integerrimis subtus tomentosis superne glaberrimis, floribus terminalibus pedicellatis corymboso-capitatis, calycis campanulati dentibus subæqualibus ovato-acuminatis, seminibus reticulatis. ¶ in saxosis rupestribus siccis. — Fl. fr. 3. p. 521. *T. montanum* et *T. supinum* L. sp. 791. — Lob. ic. 488. f. 1 et 2. (v. v.)
12. *T. POLIUM* (Lam. dict. 2. p. 699.) caulibus suffruticosis ramosis decumbentibus tomentosis, foliis lanceolato-oblongis obtusis crenatis tomentosis, capitulis subrotundis pedunculatis, calycis tomentosi subcampanulati dentibus æqualibus. § in Gallia australis montosis et maritimis. — Fl. fr. 3. p. 521. — Barr. ic. t. 1074 et 1078. Variat floribus albescentibus et purpureis. (v. v.)
13. *T. CAPITATUM* (L. sp. 792.) caulibus suffruticosis ramosis, ramis niveis erectis, foliis lanceolatis linearibusque crenatis tomentosis niveis, capitulis subrotundis breviter pedunculatis, calycis tomentosi subcampanulati dentibus acutis. § in Gallia australis saxosis aridis. — Fl. fr. 3. p. 522. Cav. ic. t. 119. — Barr. ic. t. 1047. Species à præcedente vix distincta et cum hæc probabiliter conjungenda. (v. v.)
14. *T. FLAVIGANS* (Lam. dict. 2. p. 700.) tomentosum, caulibus suffruticosis decumbentibus ramosissimis, foliis linearibus anticè crenatis vel ovatis serratis, summitatibus flavescentibus, capitulis sessilibus subrotundis, calycis aureo-tomentosi subcampanulati dentibus acutis. § in Gallia australis rupestribus. — Fl. fr. 3. p. 521. *T. aureum* Cav. ic. 2. t. 117. — Barr. ic. t. 1073. (v. v.)
15. *T. PSEUDO-HYSSOPUS* (Schreb. unil. p. 45.) caulibus erectis incanis ramosis, foliis linearibus aut lineari-lanceolatis crenatis obtusis, floribus capitatis, capitulis numerosis rotundis in racemum dispositis, calycis tubulosi adpressè cano-tomentosi dentibus parvis subobtusis. § in aridis Corsicæ. — Col. ceph. 1. t. 67. (v. s.)
16. *T. MARUM* (L. sp. 788.) caule suffruticoso erecto, ramis canescentibus, foliis integerrimis ovatis acutis petiolatis subtus tomentosis, floribus racemosis secundis, calycis cano-tomentosi dentibus acutis. ¶ in Corsicæ et Stæchadum montosis. — Fl. fr. 3. p. 516. — Black. t. 47. (v. v. D.)

VII. *HYSSOPUS* Tourn. inst. t. 95. L. gen. n. 709.

Calyx substriatus, fauce nudâ; corollæ labium superius breviter emarginatum, inferius 3-lobum lobo intermedio majore obcordato crenulato.

1. *H. OFFICINALIS* (L. sp. 796.) foliis lanceolatis, floribus verticillatis ramosis secundis, lacinia corollæ intermediâ bilobâ integerrimâ. ¶ in collibus provinciarum austro-orientalium orientaliumque et circa Meduntam propè Parisios ex Merat. — Fl. fr. 3. p. 525. Lam. ill. t. 502. f. 1. — Moris. s. 11. t. 1. f. 2 et 3. n. 1 et 7. (v. v.)

β. *canescens* (DC. fl. fr. 5. p. 396.) caulibus foliisque canescentibus. — in Galloprovincia. (v. s.)

VIII. *MOLUCCELLA* L. gen. 724.

Calyx turbinatus campanulatus dentato-spinosus, dentē superiore remotiore; corollæ calyce angustior.

1. *M. PRUTICOSA* (L. sp. 821.) foliis petiolatis ellipticis obtusis sub-5-dentatis, calycibus infundibuliformibus 8-10-dentatis spinosis, galeâ pilosissimâ.  $\zeta$  inter rupes Galloprovinciæ propè Entrevaux et S.-Arnoux (Solier). — Fl. fr. 3. p. 556. All. ped. t. 2. f. 2. (v. s.)

#### IX. PHLOMIS L. gen. 723.

Calyx angulatus 5-dentatus fauce nudâ dentibus patulis; corolla oblonga labiata superius villosa fornicata compressa incumbens subbifida, inferius triloba, lobo medio majore emarginato.

1. *P. FRUTICOSA* (L. sp. 818.) caule fruticoso, ramis densè tomentosis, foliis ovatis obtusis subtus reticulatis tomentosis, bracteis extus tomentosis ovalibus, verticillis 15-20-floris, calycis dentibus minimis retusis mucronato-subulatis.  $\zeta$  circâ Narbonam et in Pyren. orientilibus ex Lapeyr. — Fl. fr. 5. p. 401. — Dod. pempt. 146. ic. (v. v. c.)
2. *P. LYCHNITIS* (L. sp. 819.) foliis lineari-lanceolatis infernè candicantibus, floralibus sessilibus ovatis, bracteis setaceis lanatis calycis hirsutissimi dentes setaceo-acuminatas æquantibus.  $\zeta$  in Galliâ mediterraneâ. — Fl. fr. 3. p. 555. — Lob. ic. t. 558. f. 1. 2. (v. v.)
3. *P. HERBA-VENTRI* (L. sp. 819.) foliis ovato-lanceolatis serratis subtus hirtis, bracteis subulatis cauleque hirtis, calycis dentibus lanceolato-subulatis erectis.  $\mathcal{U}$  in Galliâ mediterraneâ. — Fl. fr. 3. p. 556. — Lob. ic. 532. f. 1. (v. v.)

#### X. GALEOBDOLOM Huds. ang. 258. DC. fl. fr. 3. p. 554.

Calyx campanulatus 5-dentatus dentibus inæqualibus acutis; corolla calyce longior 2-labiata suprâ fornicata integra, infrâ 3-fida laciniis acutis intermediâ longiore.

1. *G. LUTRUM* (Huds. ang. 258.)  $\mathcal{U}$  in nemoribus montosis. — Fl. fr. 3. p. 554. Galeopsis Galeobdolon L. sp. 810. — Moris. s. 11. t. 11. f. 4. n. 5. (v. v.)

#### XI. LEONURUS L. gen. 722.

Calyx cylindricus pentagonus 5-dentatus, fauce nudâ; corolla calyce vix longior 2-labiata superius villosa integra concava, inferius reflexa 3-partita subæqualis; antheræ punctis nitidis adpersæ.

##### Sect. 1. CHAETURUS Mæench. meth. 402.

Stamina ovariaque glabra.

1. *L. MARRUBIASTRUM* (L. sp. 817.) foliis oblongis dentatis, corollis calyce sub-pungente vix longioribus.  $\mathcal{U}$  in cultis siccis. — Fl. fr. 3. p. 554. Jacq. aust. t. 405. (v. v.)

##### Sect. II. CARDIACA Tourn. inst. t. 87.

Stamina villosa, ovaria apice tomentosa.

2. *L. CARDIACA* (L. sp. 817.) foliis petiolatis, inferioribus cuneiformi-ovatis trilobis dentatis, summis indivisis, laciniâ mediâ labij corollæ inferioris acutâ, calyce pungente.  $\mathcal{U}$  in dumetis, ruderatis. — Fl. fr. 3. p. 553. Bull. herb. t. 273. (v. v.)

#### XII. MARRUBIUM L. gen. 721.

Calyx cylindricus 10-striatus 5-aut-10-dentatus; corolla calyce paulò longior 2-labiata, superius angusta 2-fida, inferius 3-fida, laciniâ mediâ latiore emarginatâ.

1. *M. VULGARIS* (L. sp. 816.) caule erecto, foliis subrotundo-ovatis dentatis rugoso-venosis, calycibus 10-dentatis, dentibus setaceis uncinatis.  $\mathcal{U}$  in ruderatis incultis. — Fl. fr. 3. p. 552. — Moris. s. 11. t. 9. f. 8. n. 1. (v. v.)



2. *M. SUPINUM* (L. sp. 816.) caule decumbente, foliis subrotundis crenatis rugoso-venosis, calycibus 5-dentatis, dentibus setaceis rectis villosis.  $\mathcal{L}$  in Occitania ex Sauv., Linnæo et Benth. — Fl. fr. 3. p. 553. — Boec. mus. t. 69. (v. v. c.)

XIII. BALLOTA L. gen. 720.

Calyx campanulatus 5-gonus 10-striatus 5-dentatus; corolla bilabiata superius concava crenata, inferius 3-loba, lobo medio majore emarginato; semina triangularia.

1. *B. FORTIDA* (Lam. fl. fr. 2. p. 381.) foliis cordatis indivisis serratis.  $\mathcal{L}$  in rudertis. — Fl. fr. 3. p. 552. Lam. ill. t. 508. f. 1. *B. nigra* L. sp. ed. 1. p. 582. non ed. 2. — Cam. epit. 572. ic. (v. v.)

XIV. BETONICA Tourn. inst. t. 96. L. gen. n. 718.

Calyx cylindricus 5-dentatus aristatus fauce nudâ; corollæ tubus gracilis, limbus 2-labiatus suprâ erectus subrotundus integer vel emarginatus, infrâ 3-fidus.

1. *B. OFFICINALIS* (L. sp. 810.) caule 4-gono hirtio, foliis cordatis ovatis aut lanceolatis grossè crenatis obtusis, inferioribus petiolatis, spicâ oblongâ interruptâ, calyce basi glabriusculo in parte superiore hispido aut glabro tubo duplò triplòve breviorè, corollæ galeâ integrâ laciniâ intermediâ labii inferioris crenato-undulatâ.  $\mathcal{L}$  in pascuis, sylvaticis. — Fl. fr. 3. p. 545. *B. stricta* Ait. kew. 2. p. 299. Fl. fr. 3. p. 545. — Fuchs. hist. 350. ic. Variat plus minus pilosa. (v. v.)
2. *B. INCANA* (Mill. dict. n. 3.) caule 4-gono hirtio, foliis ovali-oblongis cordatis obtusis, inferioribus petiolatis, spicâ oblongâ interruptâ, calycis hispido tubo multò brevioris dentibus subulatis brevibus, corollarum galeâ bifidâ, laciniâ intermediâ labii inferioris crenatâ, tubo pubescente subincurvo.  $\mathcal{L}$  in sylvis circa Nannetes. — Fl. fr. 5. p. 400. (v. s. c.)
3. *B. HIRSAUTA* (L. mant. 248.) foliis oblongo-lanceolatis cordatis grossè crenatis obtusis petiolatis, spicâ ovatâ obtusissimâ foliosâ, calyce tubo breviorè glabriusculo, corollæ galeâ integrâ, laciniis lateralibus labii inferioris angustioribus acuminatis.  $\mathcal{L}$  in Alpibus, Monte-Aureo, Pyrenæis. — Fl. fr. 3. 546. — Zanon. p. 46. t. 40. (v. v. in Italiâ D.)
4. *B. ALOPECUROS* (L. sp. 811.) villosa, foliis ovato-cordatis grossè dentatis, inferioribus petiolatis, spicâ oblongâ basi foliosâ, calyce tubum æquante, corollæ galeâ emarginatâ labio inferiore longiore.  $\mathcal{L}$  in Pyrenæis, Alp. Delph. ex Vill., Galloprov. ex Ger. — Fl. fr. 3. p. 547. — Scop. carn. t. 28. (v. v.)

XV. GALEOPSIS Huds. ang. 1. p. 256. DC. fl. fr. 3. p. 542.

Calyx campanulatus 5-dentatus spinosus; corolla tubo brevi, fauce dilatâ 2-dentatâ, limbo 2-labiato suprâ fornicato crenato, infrâ 3-lobo lobis inæqualibus; antheræ intus subhirsutæ.

1. *G. TETRAHIT* (L. sp. 810. var.  $\alpha$ .) caule hispido, internodiis caulinis incrassatis, foliis ovatis hispidis glabrisve grossè serratis acuminatis, calycis hispidi laciniis subulato-pungentibus corollâ subduplò brevioribus, seminibus magnis obsolete 3-quetris leproso-punctulatis. ① in arvis. — Fl. fr. 3. p. 544. — Riv. t. 31. (v. v.)

$\beta$ . *versicolor*, foliis oblongo-ovatis, floribus majoribus ochroleucis versicoloribus. — *G. versicolor* Cart. lond. 6. t. 38. Fl. fr. 3. p. 544. (v. v.)

2. *G. OCHROLEUCA* (Lam. dict. 2. p. 600.) caulibus erectis ramosis, foliis ovato-lanceolatis petiolatis, verticillis distantibus multifloris, calycibus hirsutis, corollis longè tubulosis, seminibus triquetris punctulatis. ① in agris rariis. — Fl. fr. 3. p. 543. Fl. ochroleuci. (v. v.)
3. *G. LADANUM* (Vill. dauph. 2. p. 386.) caulibus erectis ramosis, foliis lineari-

lanceolatis lanceolatisve subobtusis dentatis aut subintegris, verticillis distantibus, calycibus pubescentibus, seminibus obsoletè triquetris punctulatis. ① in agris, cultis. — Fl. fr. 3. p. 543. (v. v.)

4. *G. PARVIFLORA* (Lam. dict. 2. p. 600.) caulibus erectis pubescentibus ramosis, foliis lanceolatis aequaliter serratis petiolatis subacutis, calycis hirsuti laciniis subulato-pungentibus corollâ brevioribus. ① in cultis, agris. — Fl. fr. 3. p. 544. *G. intermedia* Vill. dauph. 2. t. 9. (v. s.)

#### XVI. LAMIUM L. gen. 716.

Calyx 5-dentatus aristatus nudus, apice patens; corolla longior fauce inflatâ, limbo labiato suprâ fornicato integro, inferiore 3-lobo lobis lateralibus minimis reflexis, medio majore emarginato; antheræ glabræ; semina triquetra lævia.

1. *L. LEVIGATUM* (L. sp. 808.) caule sublavi, foliis cordatis rugosis grossè serratis petiolatis breviter pubescentibus, verticillis sub 6-floris, calycis glabri dentibus acuminatis inæqualibus longiore tubo corollæ subsequente. ② in sepibus et incultis montanis Galloprov., Delph., Alsatiæ ex Mapp. — Fl. fr. 3. p. 541. — Bocc. mus. 2. t. 23. Corolla ampla purpurea fauce dilatato. (v. s.)
2. *L. ALBUM* (L. sp. 371.) caulibus erectis apice pilosulis, foliis cordatis acuminatis serratis petiolatis, verticillis 12-20-floris, calycis dentibus subulatis ciliatis. ② in incultis. — Fl. fr. 3. p. 540. — Cam. epit. 865. ic. Fl. albi. (v. v.)
3. *L. MACULATUM* (L. sp. 809.) caulibus decumbentibus, foliis cordatis petiolatis acuminatis aut subobtusis grossè serratis, verticillis remotis, calycis obliqui laciniis inæqualibus setaceo-acuminatis ciliatis. ② in montanis passim. — Fl. fr. 3. p. 540. *L. hirsutum* Lam. dict. 3. p. 410. Fl. fr. 3. p. 541. *L. stoloniferum* Lapeyr. abr. 333 et supp. 82. — Garid. aix. t. 58. Folia interdum maculata. (v. v.)
4. *L. PURPUREUM* (L. sp. 809.) caulibus decumbentibus glabriusculis, foliis rotundato-cordatis obtusis aequalibus crenatis petiolatis, verticillis confertis 8-10-floris, calycis dentibus subæqualibus lineari-subulatis. ① in cultis. — Fl. fr. 3. p. 541. — Blackw. t. 152. f. 1. (v. v.)
5. *L. BIFIDUM* (Cyr. fasc. neap. 1. t. 7.) caule procumbente à basi ramoso, foliis hirsutis petiolatis oblongo-ovatis cordatis obtusis inciso-dentatis, floralibus superioribus sessilibus, corollæ labio superiore bifido laciniis divaricatis. ① in Corsicâ (cl. Soleirol). (v. s.)
6. *L. HYBRIDUM* (Vill. dauph. 2. p. 385.) caulibus decumbentibus glabriusculis, foliis cordatis acutiusculis inciso-dentatis petiolatis, corollis calyce angustioribus, calycis obliqui dentibus subinæqualibus acuminato-subulatis, seminibus sublævibus. ① in cultis. — Fl. fr. 3. p. 541. *L. incisum* Wild. — Pluck. t. 41. f. 3. (v. v.)
7. *L. AMPLEXICAULE* (L. sp. 809.) caule decumbente glabro, foliis rotundatis obtusissimis inciso-crenatis, caulinis sessilibus, calycis hirsuti dentibus subæqualibus lineari-subulatis, corollæ tubo gracili. ① in cultis, ruderalis. — Fl. fr. 3. p. 542. — Lob. ic. t. 463. f. 2. (v. v.)

#### XVII. ORVALA L. gen. 715.

Calyx 5-dentatus nudus apice patens; corollæ faux inflata utrinquè appendice 3-lobo, labio superiore fornicato dentato, inferiore 3-lobo lobis lateralibus minimis reflexis, medio majore emarginato, antheræ glabræ.

1. *O. LAMIOIDES* (DC. fl. fr. 3. p. 539.) foliis cordatis inæqualiter argutèque serratis. ② in umbrosis Pyrenæorum ex Lapeyr., circâ Nannetes ex Bon. — *Lamium Orvala* L. sp. 808. *O. garganica* L. sp. 807. — Scop. carn. t. 27. — Mich. gen. t. 17. (v. s. c.)

XVIII. GLECHOMA *L. gen.* 714.

*Calyx striatus cylindricus fructiferus nudus; corolla calyce duplò longior 2-labiata, labio superiore 2-fido, inferiore 3-fido laciniâ intermedia majore emarginatâ; antheræ per paria in crucem conniventes; semina læviuscula cylindrico-ovata.*

1. *G. HEDERACA* (*L. sp.* 807.) foliis reniformibus grossè crenatis petiolatis.  $\mathcal{U}$  in sepibus umbrosis ad muros. — *Fl. fr.* 3. p. 538. — *Moris. s. 11. t. 21. f. 10. n. 1. (v. v.)*

XIX. STACHYS *L. gen.* 719.

*Calyx angulatus 5-fidus aut 5-dentatus acuminatus; corolla tubo brevi, limbo suprâ fornicato, infrâ trilobo lateribus reflexis; stamina deflorata ad latus utrinquè reflexa; semina obsolete triquetra ovoidea aut subrotunda.*

1. *L. CORNICA* (*Pers. ench. 2. p. 124.*) hirsuta, caulibus procumbentibus debilibus, foliis ovatis lobato-crenatis obtusis, floribus axillaribus solitariis oppositis, calycis 5-fidi laciniis acuminato-pungentibus, seminibus subrotundis leproso-punctulatis. (1) in incultis montosis Corsicæ. — *Glechoma grandiflora* *Fl. fr.* 3. p. 538. (v. s.)
2. *S. ARVENsis* (*L. sp.* 814.) hirsuta, caule debili, foliis ovatis cordatis obtusis crenatis, verticillis 6-floris, calycis 5-fidi laciniis linearibus acuminato-pungentibus, seminibus subrotundis. (1) in arvis, vineis. — *Fl. fr.* 3. p. 551. (v. v.)
3. *S. ANNUA* (*L. sp.* 813.) pubescens, caule 4-gono erecto, foliis ovato-lanceolatis trinerviis lævibus obtusis crenatis serratisve petiolatis, verticillis 6-floris in apicam subdigestis, calycis 5-fidi laciniis lineari-subulatis, seminibus subrotundis nitidis. (1) in cultis, ad agrorum margines. — *Fl. fr.* 3. p. 551. *Jacq. aust. t. 360.* — *Moris. s. 11. t. 12. f. 8. n. 14. (v. v.)*
4. *S. MARUBIIFOLIA* (*Viv. cors. app. p. 2.*) hirta, foliis cordatis subrotundo-ovatis crenatis obtusis, floralibus ovatis ellipticisque integerrimis apice spinulosis, verticillis 6-floris laxis, calycis 5-lobi laciniis lanceolatis mucronato-pungentibus, seminibus ovoideis punctulato-leprosis. — In Corsicâ (cl. Viviani.) — *Fl. purpurei. (v. s.)*
5. *S. SYDENHIS* (*Vill. dauph. 2. p. 375.*) hirsuta, caulibus ascendentibus, foliis ellipticis lanceolatisve sessilibus acutis crenatis scabris, verticillis sub 6-floris subpicatis, calycis semi-5-fidi laciniis lanceolatis acutis pungentibus, seminibus subrotundis nitidis rugulosis.  $\mathcal{U}$  in arenosis, sterilibus. — *Fl. fr.* 3. p. 560. *S. recta* *L. mant. 82. Jacq. aust. t. 359. (v. v.)*
6. *S. HIRTA* (*L. sp.* 812.) hirsuta, caulibus 4-gonis prostratis, foliis ovatis cordatis crenatis petiolatis, verticillis 6-floris, calycis 5-fidi laciniis lanceolatis acuminatis spinosis, seminibus subrotundis leproso-punctulatis.  $\mathcal{U}$  in saxosis collibus Nicæe, Monspelii ex Gouan, Bayonæ ex Lois, Lotharingiæ ex Mong. — *Fl. fr.* 3. p. 550. *All. ped. t. 2. f. 3.* — *Dalech. hist. 684. ic. (v. s.)*
7. *S. MARITIMA* (*L. mant. 82.*) hirsuta, caulibus tetragonis ascendentibus, foliis oblongo-lanceolatis obtusis tomentosis crenatis, inferioribus petiolatis, superioribus sessilibus ovatis integerrimis, verticillis 2-6-floris apicatis, calycis sub 5-fidi laciniis lanceolatis acutis, seminibus subrotundis nitidis.  $\mathcal{U}$  in Calliæ mediterraneæ maritimis. — *Fl. fr.* 3. p. 549. — *Dill. elth. t. 42. f. 50. (v. v.)*
8. *S. HERACLEA* (*All. ped. t. 84. f. 1.*) hirsutissima, caule simpliciori ascendente hirta, foliis oblongo-cordatis crenatis serratisve petiolatis, superioribus sessilibus, floralibus ovato-acuminatis integris, verticillis 10-floris subpicatis, calycis 5-lobi laciniis lanceolatis mucronatis pungentibus, corollis extus hirsutissimis, seminibus subrotundis nitidis.  $\mathcal{U}$  in siccis collibus Nicæe, Gallo-

- provincia circà Grasse, agri Ruscimonensis, et circà Milhan (el. Prost). — Fl. fr. 5. p. 401. *S. barbata* Lapeyr. (v. s.)
9. *S. ALPINA* (L. sp. 812.) tomentoso-pubescent, caule erecto, foliis cordatis oblongo-ovatis crenato-dentatis petiolatis, verticillis subsexfloris, calycis 5-lobi laciniis ovatis mucronatis, corollæ labio superiore plano, seminibus ovoides nitidis.  $\mathcal{L}$  in umbrosis. — Fl. fr. 3. p. 548. Lapeyr. fl. pyr. 1. p. 14. t. 8. (v. v.)
10. *S. GERMANICA* (L. sp. 812.) lanata, caule erecto, foliis ovato-lanceolatis lanceolatisve acutis crenato-serratis, superioribus sessilibus, inferioribus petiolatis, verticillis multifloris, calycis 5-dentati dentibus rigidis acuminatis, seminibus subrotundis.  $\mathcal{L}$  ad margines viarum, agrorum. — Fl. fr. 3. 549. — Moris. s. 11. t. 10. f. 4. n. 1. (v. v.)
11. *S. SYLVATICA* (L. sp. 811.) caule erecto hirsuto, foliis cordatis ovatis petiolatis grosse dentatis, verticillis sub 6-floris subspicatis, calycis sub 5-fidi laciniis lineari-lanceolatis rigidis acuminatis, seminibus subrotundis nitidis.  $\mathcal{L}$  in nemoribus umbrosis. — Fl. fr. 3. p. 547. — Riv. t. 26. f. 2. (v. v.)
12. *S. FALUSTRIS* (L. sp. 811.) caule erecto hispidio, foliis lanceolatis sessilibus semiamplexicaulibus serratis, verticillis sub 6-floris subspicatis, calycis 5-lobi laciniis linearibus acuminatis, seminibus ovoides lævibus.  $\mathcal{L}$  in humidis aquosis. — Fl. fr. 3. p. 548. — Riv. t. 26. f. 1. (v. v.)
13. *S. GLUTINOSA* (L. sp. 813.) glabra ramosissima, ramis oppositis rigidis glutinosi apice tandem spinescentibus, foliis sublinearibus, floribus axillaribus subsessilibus subsolitariis 2-bracteatis, calycis sub 5-fidi laciniis lanceolatis acuminato-spinosis, labio superiore abbreviato, seminibus subrotundis rugulosis.  $\mathcal{L}$  in Corsicâ. — Fl. fr. 3. p. 549. (v. s.)

## XX. SIDERITIS L. gen. 712.

Calyx 5-lobus; corolla calyci æqualis aut sæpius longior 2-labiata, superius integra aut emarginata, inferius triloba lobo intermedio latiore crenulato; stamina non exserta; stigma alterum longius brevior altero involutum.

## Sect. I. BURGSDORFIA Mœnch. meth. 392.

Calyces villis clausi bilabiati, labio superiore lato ovato, inferiore 4-dentato; bractæ nullæ.

1. *S. ROMANA* (L. sp. 802.) caulibus herbaceis decumbentibus hirsutis, foliis lanceolato-ovatis, calycibus spinosis corollam subæquantibus.  $\textcircled{I}$  in Galliæ australis aridis. — Fl. fr. 3. p. 529. Cav. ic. 2. t. 187. (v. v.)

## Sect. II. HESIODIA Mœnch. meth. 392.

Calyces villis clausi bilabiati, labio superiore tridentato, inferiore 2-lobato, bractæ nullæ.

2. *S. MONTANA* (L. sp. 802.) caulibus herbaceis ascendentibus hirsutis, foliis lanceolatis, calycibus corollâ majoribus spinosis.  $\textcircled{I}$  in Galliæ australis montanis. — Fl. fr. 3. p. 530. Jacq. aust. t. 434. — Moris. s. 11. t. 4. f. 10. n. 12. (v. v.)

## Sect. III. SIDERITIS DC. fl. fr. 3. p. 531.

Calyx 5-fidus nudus; flores bracteati.

3. *S. SCORPIOIDES* (L. sp. 803.) hirsuta, caulibus ascendentibus herbaceis subvillois, foliis lanceolatis ovatisve dentatis aut dentato-incisis, bracteis ovatis dentato-spinosis, floribus verticillato-spicatis, calycis campanulati laciniis æqualibus acuminato-spinosis corollam subæquantibus.  $\mathcal{L}$   $\mathcal{Z}$  in Galliæ australis aridis. — Fl. fr. 3. p. 532. — Lob. ic. 523. f. 1 et 2. — Clus. hist. 2. p. 40. f. 2. (v. v.)
4. *S. MYRSINIFOLIA* (L. sp. 807.) caulibus herbaceis ascendentibus, foliis lanceolatis subintegris dentatisve, bracteis cordatis dentato-spinosis, floribus

verticillato-spicatis, verticillis superioribus approximatis, calycis campanulati laciniis aequalibus acuminato-pungentibus corollâ brevioribus.  $\mathcal{L}$  in Alpihus et Pyrenæis. — Fl. fr. 3. p. 532. 8. crenata Lapeyr. — Clus. hist. 2. p. 41. f. 2. — Barr. ic. 171. (v. v.)

5. *S. PARVOLIATA* (L. sp. 802.) herbacea piloso-hispida, caulibus erectis, foliis superioribus lanceolatis amplexicaulibus denticulatis acuminatis, bracteis cordatis acuminatis reticulato-nervosis margine pilosis.  $\mathcal{L}$  in arenosis maritimis circâ Monspelium ex Magn. et Gouan, sed civitas admodum dubia. — Fl. fr. 3. p. 531. (v. v. c.)

XXI. *NEPETA DC. fl. fr. 3. p. 526. — Nepetæ sp. L.*

Calyx cylindricus fauce nudâ; corollæ tubus longus, faux patens, limbus 2-labiatus, suprâ emarginatus, infrâ 3-lobus lateralibus brevissimis reflexis medio majore crenato concavo.

1. *N. CATANIA* (L. sp. 796.) caule pubescente, foliis ovatis cordatis petiolatis grossè dentatis superioribus serratis, floribus verticillatis verticillis subpedicellatis in spicas dispositis, seminibus ovoideis lævibus.  $\mathcal{L}$  in lapidosis campis passim. — Fl. fr. 3. p. 526. Bull. herb. t. 287. (v. v.)
2. *N. NEPETELLA* (L. sp. 797.) caule breviter pubescente, foliis oblongo-lanceolatis cordatis profundè serratis crenatis, cymis pedunculatis in spicas digestis, calyce hirsuto, seminibus ovoideis punctato-tuberculatis.  $\mathcal{L}$  in aridis montosis Delph., Galloprovinciæ et Pyren. centralium. — Fl. fr. 3. p. 527. (v. v. DC.)

$\beta$ . *lanceolata*, foliis ovato-lanceolatis. — *N. lanceolata* Lam. fl. fr. 2. p. 399. DC. fl. fr. 3. p. 526. — All. ped. t. 2. f. 1. (v. s.)

3. *N. PANNONICA* (L. sp. 797.) caule subglabro, foliis oblongo-lanceolatis cordatis petiolatis crenatis serratisve glabris, florum glomerulis pedicellatis cymosis remotis in racemum elongatum dispositis, calyce subpiloso dentibus ovato-lanceolatis acutis, semin. punctato-tuberculosus.  $\mathcal{L}$  in montosis Corsicæ. — Moris. s. 11. t. 6. f. 10. n. 6. (v. s.)
4. *N. VIOLACEA* (L. sp. 797.) caule pubescente, foliis subpetiolatis glabresculis inferioribus cordatis oblongis superioribus lanceolatis, florum glomerulis pedicellatis cymosis in spicam verticillis distantibus dispositis, corollæ lobis lateralibus patentibus, seminibus punctato-tuberculosus.  $\mathcal{L}$  in Pyrenæis orientibus. — Fl. fr. 5. p. 396. — Barr. ic. 601. (v. s.)
5. *N. NUDA* (L. sp. 797.) caule breviter pubescente paniculato, foliis cordato-oblongis sessilibus serratis, superioribus linearibus minimis, florum glomerulis pedicellatis in spicas elongatas interruptè verticillatas digestis, seminibus ovoideo-triquetris sublævibus.  $\mathcal{L}$  in collibus Delph. ex Vill. — Fl. fr. 3. p. 527. Jacq. aust. t. 24. (v. v. c.)
6. *N. LATIFOLIA* (DC. fl. fr. 3. p. 528.) caule villosa apice paniculato, foliis sessilibus ovatis obtusè dentatis, florum glomerulis cymosis pedicellatis remotis, calyce hirsuto striato sub 5-fido, laciniis lineari-subulatis.  $\mathcal{L}$  in Pyrenæis orientalibus. — *N. grandiflora* Lapeyr. (v. v. DC.)

XXII. *LAVANDULA L. gen. 711. de Gingins lav. hist. 124.*

Calyx ovatus intus nudus terdecim nervatus, dentibus 4 subæqualibus quinto summo, apice appendiculatim producto; corollæ galea biloba, labium inferius 3-lobum; semina gynobasi tergo adnata; stigmata carnosa complanata.

Sect. I. *STACHAS Clus. hist. 1. p. 344. Ging. l. c. p. 128.*

Flores spicati ebracteolati, bractes imbricatæ, apica comosa.

- x. *L. STACHAS* (L. sp. 800. excl. var.  $\alpha$ .) caule folioso, foliis oblongo-lanceolatis

canis, calycibus dentatis ovatis canis, spicâ coarctatâ comosâ, seminibus ovatis reticulatis.  $\zeta$  in provinciis australibus. — Fl. fr. 3. p. 599. exol. var.  $\beta$ . de Ging. tab. 1. f. A. — Black. herb. t. 241. — Lob. ic. 429. f. 2.

$\alpha$ . *brachystachya* (de Ging. l. c. t. 3. f. 1.) spicis brevioribus 4-gonis, comâ rarâ, foliis angustis. (v. v.)

$\beta$ . *macrostachya* (de Ging. l. c. t. 3. f. 2-4.) spicis crassioribus 8-gonis, comâ speciosâ, foliis latioribus. — In Corsicâ (Ph. Thomas). (v. s.)

#### Sect. II. SPICA Ging. l. c.

Flores verticillato-spicati bracteolati, bracteæ remotæ, spica non comosa; semina nitida oblonga.

2. *L. PYRENAICA* (DC. fl. fr. 5. p. 398.) caule supernè nudo, foliis angustè lanceolatis, bracteis late cordatis subfoliaceis calyces æquantibus, calyce oblongo velutino, dentibus validis acutiusculis, appendice reniformi-dilatato.  $\zeta$  in Pyrenæis orientalibus. — De Ging. t. 1. f. F. et tab. 6. f. 1. *L. Spica* Lapeyr. abr. pyr. p. 39a. Semina tergo trinervia? (v. v.)

3. *L. VERA* (DC. fl. fr. 5. p. 398.) caule supernè nudo, foliis lanceolatis, bracteis cordato-acuminatis scariosis calycibus brevioribus, calyce ovato tomentoso, dentibus obsoleteis, appendice ovato.  $\zeta$  vulgaris in collibus et declivibus montium Gallie meridionalis et propè Lugdunum in loco dicto *Montou*. — De Ging. t. 1. f. G. et t. 6. f. 2. *L. spica*  $\alpha$  Fl. fr. 3. p. 520. — Blackw. herb. t. 294. — J. Bauh. hist. 3. p. 281. f. 2. Calyces cærulei; semina enervia. Variat angustifolia et latifolia, cærulea et alba. Folia nunquam spatulata. (v. v.)

4. *L. SPICA* (DC. fl. fr. 5. p. 397.) caule supernè nudo, foliis spatulato-lanceolatis, bracteis linearibus velutinis, calyce ovato cano dentibus obsoleteis, appendice rhomboideo-ovato.  $\zeta$  in aridis Gallie australioris in regione Olearum. — De Ging. t. 2. f. H. et t. 7. *L. spica*  $\beta$  Fl. fr. 3. p. 528. — Blackw. herb. t. 295. — J. Bauh. hist. 3. p. 280. f. 1. Variat caule ramoso, foliis latè et angustè spatulatis. (v. v.)

#### XXIII. SATUREIA Lin. gen. 707.

Calyx campanulatus cylindricusve 5-fidus laciniis æqualibus; corolla limbo 5-lobo subæquali; stamina distantia.

#### Sect. I. SABATTIA Mæench. meth. 388.

Calyx cylindricus 10-striatus, fructifer villis clausus.

1. *S. GRÆCA* (L. sp. 794.) caulibus ramosis, foliis ovatis ovato-lanceolatisve pubescentibus subtus venosis, pedunculis axillaribus 3-6-floris bracteis brevioribus, calyce hispido, laciniis subulatis ciliatis.  $\mathcal{U}$  in aridis Corsicæ, Nicææ ex All. — Fl. fr. 3. p. 525. — Moris. s. 11. t. 17. f. 10. n. 2. (v. s.)

2. *S. JULIANA* (L. sp. 793.) caulibus subsimplicibus, foliis lanceolato-linearibus margine revolutis pubescentibus, pedunculis axillaribus 3-6-floris, calyce hispido, laciniis subulatis ciliatis.  $\mathcal{U}$  in aridis Corsicæ, Nicææ ex All. — Fl. fr. 3. p. 524. — Moris. s. 11. t. 17. f. 7. n. 4. (v. s.)

#### Sect. II. THYMARIA.

Calyx 3-queter 2-labiatus, fauce villosâ.

3. *S. CAPITATA* (L. sp. 795.) caule suffruticoso ramoso, foliis lineari-lanceolatis punctatis ciliatis, inferioribus plicatis, floribus capitato-spicatis, bracteis floribus punctatis brevioribus calycem 2-labiatum superantibus, calycis labio superiore dentibus 3 conniventibus, inferiore laciniis 2 setaceis ciliatis, fauce villosâ.  $\zeta$  in montosis Galloprovinciæ ex Garid. — Fl. fr. 3. p. 523. — Cam. epit. 485. ic. Dalech. hist. 900. ic. (v. s. ex Oriente.)

#### Sect. III. SATUREIA Mæench. meth. 386.

Calyx campanulatus, ore nudo.

4. *S. THYMARIA* (L. sp. 794.) caule pubescente, foliis obovato-oblongis acumi-

natis punctatis hispidis, verticillis multifloris subrotundis hispidis, calyce hispido laciniis subulatis ciliatis.  $\frac{1}{2}$  ad colles lapideos agri Nicomensis ex All. — Fl. fr. 3. p. 523. — Black. t. 318. (v. s. ex Oriente.)

5. *S. MONTANA* (L. sp. 795.) caule ramoso, foliis lineari-lanceolatis planis in petiolum subatennatis, pedunculis axillaribus 1-3-floris, calycis campanulati laciniis lanceolatis aequalibus acutis. (1) in Gallia australis aridis et agris. — Fl. fr. 3. p. 523. — Cam. epit. 487. ic. (v. v. c.)
6. *S. MONTANA* (L. sp. 794.) caule ramoso, foliis punctatis ciliatis lineari-lanceolatis mucronatis sessilibus, pedunculis subtrifloris axillaribus subcymosis, calycis campanulati striati laciniis acuminatis ciliatis.  $\frac{1}{2}$  in montosis Gallia australis et Pyrenæorum vallibus. — Fl. fr. 3. p. 523. — Cam. epit. 717. B. (v. v.)

XXIV. MENTHA L. gen. 713.

Corolla calyce paulò longior 4-loba; subæqualis, lobo superiore latiore sæpius emarginato; stamina distantia.

Sect. I. MENTHASTRUM. — *Mentha Tournef. t. 89.*

Calyx campanulatus intus nudus, lobus superior corollæ emarginatus.

1. *M. sylvestris* (L. sp. 804.) caulibus erectis tomentosis, foliis ovatis oblongisve sessilibus tomentosis argutè inæqualiter serratis acutis, spicis oblongis subcontinuis, bracteis subulatis, calycis hirsuti 5-dentati campanulati dentibus acuminatis.  $\frac{1}{2}$  in ruderatis, humidis. — Fl. fr. 3. p. 533. — *M. nemorosa*, et *M. gratissima* Wild. sp. 3. p. 75. — Lob. ic. t. 508. f. 2 et 509. f. 2. (v. v.)
2. *M. rotundifolia* (L. sp. 805.) caule erecto hirsuto apice ramoso, foliis subrotundis rugosis serrato-crenatis sessilibus obtusis, bracteis lanceolatis acuminatis, spicis elongatis basi subinterruptis, calyce campanulato 5-dentato, dentibus acutis.  $\frac{1}{2}$  in aquosis. — Fl. fr. 3. p. 534. *M. crispa* L. sp. ed. 1. p. 576. — Riv. t. 51. f. 2. Lob. ic. t. 506. f. 2. (v. v.)
3. *M. viridis* (L. sp. 804.) caulibus erectis flexuosis ramosis glabris, foliis lanceolatis glabris argutè serratis sessilibus acutis, bracteis lineari-subulatis, spicis interruptis gracilibus acutis, pedicellis glabris, calycis campanulati 5-lobi glabri laciniis acuminatis ciliatis.  $\frac{1}{2}$  in humidis provinc. borealium, Parisiorum, Occitanie ex Benth. et Pyren. ex Lapeyr. — Fl. fr. 3. p. 534. — Lob. ic. t. 507. f. 2. (v. s.)
4. *M. piperita* (Huds. ang. 251.) glaberrima, caulibus erectis flexuosis apice ramosis, foliis petiolatis subovatis glabriusculis serratis acutis, bracteis lanceolatis lineari-lanceolatisve acuminatis, spicis obtusis infernè interruptis, pedunculis axillaribus oppositis multifloris, calycis punctato-glandulosi campanulato-cylindrici sub 5-lobi laciniis acuminatis.  $\frac{1}{2}$  in hortis sæpè colitur. — Fl. fr. 3. p. 534. eng. bot. t. 687. (v. v. c.)
5. *M. hirsuta* (Sm. trans. lin. 5. p. 193.) caulibus hirsutis flexuosis ascendentibus, foliis ovatis petiolatis subsessilibusve serratis pubescentibus subacutis, floribus capitatis, pedicellis hirsutis, calycis subcampanulati hirsuti 5-dentati dentibus acutis.  $\frac{1}{2}$  in aquosis. — Fl. fr. 3. p. 535. eng. bot. t. 447. *M. hirsuta* L. mant. 81. *M. aquatica* L. sp. 805. — Lob. ic. 509. f. 1. (v. v.)
6. *M. sativa* (L. sp. 805.) caulibus flexuosis basi radicanlibus, foliis ovatis acutiusculis subserratis petiolatis, floribus verticillatis axillaribus pedicellatis, calycibus subcylindricis 5-dentatis hirsutis, dentibus acutis.  $\frac{1}{2}$  in humidis. — Fl. fr. 3. p. 535. eng. bot. t. 448. *M. procumbens* Thail. *Nimium* sequenti affinis. (v. v.)
7. *M. arvensis* (L. sp. 806.) caulibus hirsutis ramosis, foliis ovatis acutis serratis subsessilibus, floribus verticillatis axillaribus, calycis campanulati 5-dentati hirsuti dentibus acutis.  $\frac{1}{2}$  in campis humidis frequens post menses. — Fl. fr. 3. p. 535. Fl. dan. t. 512. (v. v.)

8. *M. GENTILIS* (L. sp. 805.) caulibus erectis flexuosis hirsutis ramosis, foliis ovatis acutis serratis petiolatis subacutis, floribus verticillatis axillaribus, calycibus subcampanulatis margine hirsutis basi et pedicellis glabris, calycis dentibus acutis. ♀ in aquosis. — Fl. fr. 3. p. 536. — Moris. s. 11. t. 7. f. 5. (v. v.)
9. *M. RUBRA* (Smith. brit. 2. p. 619.) caulibus erectis flexuosis, foliis ovatis subsessilibus serratis subacutis, floribus verticillatis axillaribus, calyce subcampanulato hirsuto, dentibus acutis. ♀ in aquosis circa Parisios. — Fl. fr. 3. p. 536. — Lob. ic. t. 507. Varietas præcedentis? (v. s.)

Sect. II. *PULEGIUM* *Bauh. pin.* 222.

Calyx cylindricus villis clausus, lobus superior corollæ integer.

10. *M. PULEGIUM* (L. sp. 807.) caule repente, foliis ovatis obtusis subcrenatis subtus punctatis, floribus verticillatis, calycis subcylindrici 5-dentati dentibus 3 acutis, dentibus 2 inferioribus sublongioribus acuminatis. ♀ in inundatis, ad ripas. — Fl. fr. 3. p. 537. — Lob. ic. 500. f. 2. (v. v.)
11. *M. CERVINA* (L. sp. 807.) foliis linearibus integerrimis obtusis subtus punctatis, floribus verticillatis, calycis 5-dentati cylindrici dentibus macronatis. ♀ in Galliæ australis humidis. — Fl. fr. 3. p. 537. — Blackw. t. 304. Moris. s. 11. t. 7. f. 9. n. 7. (v. v.)

XXV. *THYMUS* *Scop. carn.* 1. p. 424. — *Thymus* et *Melissæ* sp. L.

Calyx striatus fauce villis clausa, limbus labiatus supra 3-dentatus, infra bisetus aut bifidus; corolla brevis labiata supra emarginata infra 3-loba lobo medio latiore integro aut emarginato; semina lævia.

Sect. I. *PULEGIASTRUM*.

Calyx campanulatus subbilabiatus; corolla 4-fida regularis.

1. *T. PARVIFLORUS* (Req. in ann. sc. natur. t. 5. p. 386.) caulibus cespitosis decumbentibus, foliis subrotundis glabris subnerviis petiolatis subintegris, pedunculis solitariis 1-3-floris axillaribus terminalibusve, calycis campanulati dentibus subæqualibus acutis, corollis 4-fidis, seminibus subglobosis. — In montibus Corsicæ. (v. s.)

Sect. II. *SERPYPILLUM* *Pers. ench.* 2. p. 130.

Calyx campanulatus, corollæ labii inferioris lobus medius integer, pedunculi subuniflori.

2. *T. HERBA-BARONA* (Lois. gall. 2. p. 360. t. 9.) pumilus procumbens glaber suffruticosus, ramis erectis, foliis lanceolato-linearibus uninerviis integerrimis subacutis glanduloso-punctatis, floribus subverticillato-capitatis, calycis campanulati glanduloso-punctati labio superiore 3-dentato dentibus acutis, inferiore 2-fido laciniis subulatis ciliatis, seminibus subglobosis. ♀ in montibus Corsicæ. — Fl. fr. 5. p. 402. T. Marschallianus Viv. cors. 9 non Wild. (v. s.)
3. *T. VULGARIS* (L. sp. 825.) erectus suffruticosus, foliis lineari-lanceolatis margine revolutis fasciculatis obtusis punctato-glandulosis, floribus capitatis, calycis punctato-glandulosi labii superioris dentibus acutis, inferioris bifidi laciniis subciliatis, seminibus globosis. ♀ in Galliæ australis montosis saxosis. — Fl. fr. 3. p. 561. — Blackw. t. 211. (v. v.)
4. *T. ZYGIS* (L. sp. 826.) caulibus suffruticosis procumbentibus ramosis canescentibus, foliis oppositis fasciculatisve linearibus obtusissimis enerviis sessilibus utriusque punctato-glandulosis margine revolutis basi ciliatis, floribus capitatis, calycis campanulati labio superiore 3-dentato erecto, inferiore subbifido laciniis subulatis, seminibus subglobosis. ♀ in ericetis Pyrenæorum ex Lapeyr., Monspelii, Nicæ (Stire). — Fl. fr. 3. p. 560. — Clus. hist. 1. p. 358. f. 2. (v. s.)



5. *T. SCARPELUM* (L. sp. 825. excl. var.  $\delta$ .) caulibus repentibus, foliis ovatis oblongisve planis obtusis integris basi ciliatis petiolatis, floribus verticillato-capitatis, calycis campanulati labio superiore tridentato erecto, inferiore subfido, laciniis subulatis, seminibus subglobosis.  $\mathcal{U}$  in aridis apricis. — Fl. fr. 3. p. 559. — Moris. s. 11. t. 17. f. 1 et 2. (v. v.)

$\beta$ . *intermedius*, caulibus suffruticosis procumbentibus, foliis glabris subsessilibus nervosis linearibus obtusissimis planis fasciculatis basi ciliatis. — In Pyrenæis. (v. v.)

6. *T. LANGEVINOSUS* (Schk. bot. 2. t. 164.) hirsutus, caulibus repentibus, foliis subrotundis obtusis villosis integerrimis subpetiolatis, floribus capitatis, calycis campanulati labio superiore 3-dentato dentibus acutis, inferiore 2-fido laciniis subulatis, seminibus subglobosis.  $\mathcal{U}$  in sylvaticis sterilibus passim. — Fl. fr. 3. p. 560. An prioris ut vult Linnæus var? (v. s.)
7. *T. CORSICUS* (Pers. ench. 2. p. 131.) caule tortuoso suffruticoso procumbente, foliis orbiculatis pilosis glabrisve petiolatis, margine calloso integerrimo, floribus subsessilibus axillaribus, calyce cylindrico, labii superioris dentibus acutis, inferioris subulatis, corollis calyce duplò majoribus.  $\mathcal{U}$   $\delta$  in montibus editioribus Corsicæ, Pyrenæorum ex Lapeyr. — Fl. fr. 5. p. 402. Lois. gall. 2. t. 9. f. 2. (v. s.)

Sect. III. *ACYNOS* Moench. meth. 407.

Calyx basi gibbus; corollæ labii inferioris lobus medius subinteger.

8. *T. ALPINUS* (L. sp. 826.) caulibus diffusis basi glabris apice pubescentibus, foliis subrotundis obtusis subserratis, petiolis glabris, verticillis 6-8-floris, calyce cylindrico basi dilatato gibbo, dentibus superioribus acutis, inferioribus longioribus acuminate, corollis inflatis calyce subduplò longioribus, seminibus obsolete triquetris ovoideis.  $\mathcal{U}$  in Alpestribus saxosis. — Fl. fr. 3. p. 562. Jacq. aust. t. 97. (v. v.)
9. *T. ACYNOS* (L. sp. 826.) caulibus subramosis suberectis hirsutis, foliis ovatis oblongisve acutis subserratis integrisve inferioribus subpetiolatis, pedunculis unifloris, verticillis 6-floris, calyce cylindrico basi gibbo dilatato, labii superioris 3-dentati dentibus acuminate, inferioris subbifidi laciniis subulatis, seminibus ovoideis. (1) in glareosis, cretaceis, siccis. — Fl. fr. 3. p. 561. — Lob. ic. t. 506. f. 1. (v. v.)

Sect. IV. *CALAMINTHA* Lam. fl. fr. 2. p. 394.

Calyx cylindricus, corollæ labii inferioris lobus medius emarginatus; pedunculi multiflori.

10. *T. CRETICUS* (DC. fl. fr. 3. p. 564.) adpressè tomentoso-canescens, caulibus ramosis erectis ascendentibusve, foliis ovatis subintegris obtusis subpetiolatis, pedunculis subsolitariis submultifloris, bracteis minimis linearibus, calycis subcylindrici dentibus subæqualibus ovatis mucronatis, seminibus ovoideis.  $\delta$  circa Monspelium ex Lin., Massiliam ex Villen. statist. — *Melissa Cretica* L. sp. 828. — Lob. ic. t. 514. f. 1. (v. s. c.)
11. *T. GLANDULOSUS* (Req. in ann. sc. nat. t. 5. p. 386.) caulibus erectis suffruticosis pubescentibus, foliis ovatis argutè serratis aut subintegris obtusis utrinquè punctato-glandulosis, pedunculis axillaribus multifloris dichotomocorymbosis, calycis subcylindrici corollâ paulò brevioris dentibus subæqualibus acuminatis, seminibus subovoideis.  $\delta$  in Corsicâ. — *Th. clandestinus* Salzm. exsicc. (v. s.)
12. *T. NEPETA* (Sm. brit. 2. p. 642.) caulibus pubescentibus decumbentibus, foliis ovatis obtusis subserratis, pedunculis axillaribus multifloris dichotomia corymbosis, calycis dentibus subæqualibus superioribus duobus suberectis, inferioribus acuminatis longioribus, seminibus subglobosis.  $\mathcal{U}$  in aggeribus glareosis. — Fl. fr. 3. p. 563. *Melissa nepeta* L. sp. 828. An satis à sequente distincta? (v. v.)

13. *T. CALAMINTHA* (Scop. carn. ed. 2. n. 733.) caulibus pubescentibus erectis, foliis ovatis obtusis serratis integrisve petiolatis hirsutis, pedunculis axillaribus multifloris dichotomis corymbosis, bracteis linearibus minimis, calycis subcylindrici corollâ duplò brevioris dentibus ciliatis labii superioris lanceolatis acutis, inferioris acuminatis longioribus, seminibus ovoideis obsolete 3-quetris.  $\frac{1}{2}$  in collibus saxosis. — Fl. fr. 3. p. 563. *Melissa Calamintha* L. sp. 827. — Lob. ic. t. 513. f. 1. (v. v.)
14. *T. GRANDIFLORUS* (Scop. carn. ed. 2. n. 732.) caulibus erectis pilosulis, foliis ovatis acutis argutè serratis petiolatis, pedunculis axillaribus 3-4-floris, bracteis linearibus lanceolatis pedicellos subsuperantibus, calyce cylindrico corollâ subtriplo minore laciniis dentibusque ciliatis, seminibus subglobosis.  $\frac{1}{2}$  in montosis umbrosis passim. — Fl. fr. 3. p. 562. *Melissa grandiflora* L. sp. 827. — Fl. maximi. (v. v.)

XXVI. MELISSA *Manch. meth. p. 408.*

Calyx apice patens fauce nudâ 2-labiatus, suprâ planus 3-dentatus infrâ 2-lobus; corolla tubo cylindrico 2-labiata superius fornicata emarginata, inferius triloba lobo medio obcordato.

1. *M. OFFICINALIS* (L. sp. 827.) caule ramoso, foliis ovatis grossè serratis obtusis mucronatis, verticillis dimidiatis, calycis labii superioris dentibus minimis ovatis mucronatis.  $\frac{1}{2}$  in montosis, sepibus passim; in hortis sæpè colitur. — Fl. fr. 3. p. 564. — Lob. ic. 514. f. 2. (v. v.)
2. *M. PYRÆNICA* (Jacq. vind. 2. t. 183.) caule subnudo simplici, foliis oblongis obtusè dentatis, floribus verticillatis, calycis campanulati labii superioris dentibus acuminatis.  $\frac{1}{2}$  in editioribus Pyrenæis. — Fl. fr. 3. p. 565. *Horminum Pyrenæicum* L. sp. 831. — Magn. hort. t. 133. (v. v. DC.)

XXVII. MELITTIS *L. gen. 731.*

Calyx magnus 3-fidus suprâ emarginatus inæqualiter 2-labiatus tubo corollæ amplior; corolla calyce duplò longior limbo dilatato patente labiato, suprâ integro plano, infrâ 3-lobo, lobis magnis inæqualibus; semina subrotundo-triangularia extus villosa.

1. *M. MELISSOPHYLLUM* (L. sp. 832.) caulibus foliosis, foliis ovalibus dentatis crenatisve.  $\frac{1}{2}$  in sylvis umbrosis. — Fl. fr. 3. p. 565. Jacq. aust. t. 26. Fl. maximi. (v. v.)

XXVIII. DRACOCEPHALUM *L. gen. 729.*

Calyx 5-fidus 2-labiatus vel subæqualis; corolla fauce inflatâ 2-labiata superius fornicata aut emarginata, inferius 3-loba, lobis lateralibus brevibus erectis, medio integro aut emarginato.

1. *D. AUSTRIACUM* (L. sp. 829.) hirsutum, foliis sessilibus linearibus mucronatis, caulibus 3-7-partitis, ramis simplicibus, floralibus tripartitis, floribus spicatis.  $\frac{1}{2}$  in Alpibus Delph. ex Vill. et Pyrenæis orient. propè *Font de Comps*. — Fl. fr. 3. p. 566. — Clus. hist. 2. p. 185. f. 1. (v. s.)
2. *D. RUSCHIANA* (L. sp. 830.) glabrum, foliis bracteisque lanceolatis linearibusve indivisis muticis, floribus spicatis.  $\frac{1}{2}$  in Alpibus Delphinatâ propè Vapincum ex Vill. — Fl. fr. 3. p. 567. — Moris. s. 11. t. 5. f. 9. (v. s. ex Helv.)

XXIX. CLINOPODIUM *Tourn. inst. t. 92. L. gen. 725.*

Calyx labiatus fauce nudâ, limbo suprâ 3-fido, infrâ bipartito; corolla tubo brevi, fauce sensim ampliata, labio superiore erecto emarginato, inferiore 3-fido laciniâ mediâ majore emarginatâ.

1. *C. VULGARÆ* (L. sp. 821.) caule subsimplici foliisque remotè dentatis pilosis,

bracteis setosis.  $\mathcal{N}$  ad oras sylvarum. — Fl. fr. 3. p. 557. — Lob. ic. t. 504. f. 2. (v. v.)

XXX. ORIGANUM *L. gen.* 726.

Calyx varius nunc 5-dentatus, nunc 2-labiatus aut 2-partitus; corolla tubo compresso labiata, superius erecta emarginata, inferius 3-fida subæqualis; semina subrotunda.

Sect. I. ORIGANUM *Tourn. t.* 64.

Calyx cylindricus 5-dentatus, fructifer villis clausus.

1. *O. vulgare* (L. sp. 824.) foliis ovatis, bracteis calyce longioribus ovatis acutis, apicis subrotundis paniculatis conglomeratis.  $\mathcal{N}$  in rupestribus. — Fl. fr. 3. p. 558. — Fuchs. hist. 552. ic. (v. v.)
2. *O. caritum* (L. sp. 823.) foliis ovatis, bracteis membranaceis calyce duplò longioribus subobtusis, inferioribus ovatis, superioribus lanceolatis, apicis aggregatis longis prismaticis rectis.  $\mathcal{N}$  Monspelii. — Fl. fr. 3. p. 558. — Cam. epit. 468. ic. (v. s.)

Sect. II. MAJORANA *Tourn. t.* 94.

Calyx fructifer nudus bilabiatus, labio superiore majore tridentato, inferiore bilobo.

3. *O. majoranoides* (Wild. sp. 3. p. 137.) caule suffruticoso, foliis petiolatis ellipticis obtusis tomentosis, apicis subrotundis pluribus glomeratis pedunculatis.  $\mathfrak{z}$  colitur in hortis. — Fl. fr. 3. p. 558. — Lob. ic. t. 498. f. 1. (v. v. c.)

XXXI. OXYMUM *Tourn. t.* 96. *L. gen.* 732.

Calyx labiatus suprà latior orbiculatus infrà 4-fidus; corolla resupinata tubo brevi, 2-labiata labio altero longiore indiviso crenato, altero 4-lobo; filamenta declinata, 2 breviora basi appendiculata.

1. *O. basilicum* (L. sp. 833. excl. var.  $\delta$ .) foliis ovatis obtusis glabris integris, calycibus ciliatis. (1) colitur in hortis. — Fl. fr. 3. p. 570. — Cam. epit. 308. ic. — Lob. ic. t. 503. f. 2. (v. v. c.)
2. *O. bullatum* (Lam. dict. 1. p. 384.) foliis ovatis concavis bullatis integris incisisque. (1) colitur in hortis. — Fl. fr. 3. p. 570. — Barr. t. 1053. 1054 et 1072. (v. v. c.)
3. *O. murinum* (L. sp. 833.) glaberrimum, foliis subrotundo-ovatis integerrimis planis. (1) colitur in hortis. — Fl. fr. 3. p. 571. — Barr. ic. t. 1068. 1075. 1077. (v. v. c.)

XXXII. CLEONIA *L. gen.* 736.

Calyx fauce villis clausa, corolla labiata superius concava inferius 3-loba, lobo medio majore emarginato, filamenta apice bifurca seu bidentata, dente altero antherifero altero nudo, stigma 4-fidum.

1. *C. lusitanica* (L. sp. 837.) (1) in Occitania inter Carcassonam et Sorricinium ex Dom Fourmeault in Herb. Lamarck. — Fl. fr. 3. p. 569. — Barr. ic. t. 561. (v. s. ex Hisp.)

XXXIII. BRUNELLA *Tourn. inst. t.* 84. *Juss. gen.* 116. — *Frunella L.*

Calyx labiatus fauce nuda, suprà planus subtruncatus 3-fidus, infrà brevior 2-fidus; corolla labiata, superius concava integra aut 2-loba, inferius 3-loba, lobo medio majore emarginato; filamenta apice bifurca seu bidentata, dente altero antherifero, altero nudo; semina ovoidea nitida.

1. *B. vulgaris* (Moench. meth. 414.) caule ascendente, foliis petiolatis oblongo-ovatis obtusis basi dentatis integrisve, bracteis ciliatis rotundatis mucronatis, calycis labio superiore truncato tridentato, dentibus minimis mucronatis. ♀ in pascuis. — Fl. fr. 3. p. 368. excl. var. s. — Blackw. t. 24. — Tabern. ic. p. 553. (v. v.)
2. *B. laciniata* (Lam. fl. fr. 2. p. 366.) caule ascendente, foliis petiolatis ovato-oblongis obtusis, superioribus lanceolatis dentatis pinnatifidisque, bracteis rotundatis acuminatis ciliatis, calycis labio superiore truncato dentibus brevissimis emarginato-retusis mucronatis. ♀ in pascuis sylvaticis. — Fl. fr. 3. p. 568 et *B. vulgaris* s. p. 567. — Cins. hist. 2. p. 43. f. 2. — Vaill. bot. t. 5. f. 1. (v. v.)
3. *B. grandiflora* (Moench. meth. 414.) caule ascendente, foliis petiolatis oblongo-ovatis obtusis basi dentatis, bracteis rotundatis ciliatis acuminatis, calycis labio superiore tridentato, dentibus ovatis mucronatis, galeâ in medio flexâ. ♀ in siccis, rupestribus, montosis. — Fl. fr. 3. p. 568. Prun. grandiflora Jacq. aust. t. 377. *P. vulgaris* fl. L. sp. 837. — Moris. s. 11. t. 5. n. 4. f. 5. (v. v.)
4. *B. myrsinifolia* (Lam. fl. fr. 2. p. 366.) caule ascendente, foliis sessilibus lanceolatis integerrimis scabris ciliatis acutis, bracteis rotundatis ciliatis acuminatis, calycis labio superiore 3-dentato, dentibus mucronatis. ♀ in Galliæ australis pascuis humidis argillosis. — Fl. fr. 3. p. 569. — Moris. s. 11. t. 5. n. 7. f. 6. (v. v.)

## XXXIV. SCUTELLARIA L. gen. 734.

Calyx brevis labio utroque integro, squamulâ concavâ superius incumbente; corolla longior basi flexa, limbo labiato, suprâ compressa fornicata ad ortum bidentata, inferius latior emarginata; semina calyce clauso tecta.

1. *S. alpina* (L. sp. 834.) foliis ovatis cordatis crenatis, superioribus ovatis ciliatis flore duplò brevioribus, spicis imbricatis rotundatis. ♀ in saxosis Alpium, Cebennarum, Pyrenæorum. — Fl. fr. 3. p. 572. All. ped. t. 26. f. 3. (v. v.)
2. *S. hastifolia* (L. sp. 835.) foliis inferioribus hastatis, superioribus lanceolatis petiolatis obtusis, floribus axillaribus secundis. ♀ in agro Andegavensi, Aurelianensi. — Fl. fr. 5. p. 403. (v. s.)
3. *S. minor* (L. sp. 835.) foliis cordatis ovatis integris obtusis, superioribus lanceolatis flores subsuperantibus, floribus axillaribus geminis. ♀ in paludibus inundatis. — Fl. fr. 3. p. 572. (v. v.)
4. *S. galericulata* (L. sp. 835.) foliis cordato-lanceolatis crenatis subobtusis, floribus axillaribus geminis. ♀ in palustribus. — Fl. fr. 3. p. 572. — Lob. ic. t. 344. f. 2. (v. v.)

## XXXV. PRASIUM L. gen. 737.

Calyx bilabiatus labio superiore bilobo inferiore trilobo; corollæ labium superius emarginatum, inferius 3-lobum; semina 4 baccata.

1. *P. majus* (L. sp. 838.) caule suffruticoso, foliis ovato-lanceolatis subcordatis dentatis glabris. ♂ in Corsicâ. — Fl. fr. 5. p. 403. — Barr. ic. 895. (v. s.)

Ordo LXXIV. VERBENACEÆ *Juss. in ann. mus. 7. p. 63.* —  
*Vitices Juss. gen.* — *Pyrenaceæ Vent. tabl. DC. fl. fr. 3.*  
*p. 501.*

Calyx tubulosus sæpè persistens. Corolla hypogyna monopetala tubulosa decidua sæpiùs irregularis. Stam. 4 didynama, rariùs 2 aut 6. Ovarium liberum 2-4-loculare, ovulis erectis solitariis geminisve. Stylus 1. Stigma simplex aut bilobum; pericarpium drupaceum seu baccatum 1-4 pyrena monosperma continens. Semina aut pyrena interdum membranà utriculâ inclusa. Embryo rectus; radícula infera; albumen nulum. — *Herbæ aut frutices, foliis oppositis exstipulatis.*

I. VITEX *Tourn. inst. t. 373. L. gen. 790.*

Calyx brevis 5-dentatus; corolla tubo tenui, limbo plano subbilabiato 5-6-lobo inæquali; stamina 4 didynama; drupa sicca tenuis putamine osseo 4-loculari 4-spermo.

1. *V. Aëvus-caeruus* (L. sp. 890.) foliis digitatis septenatis quinatiseve, foliolis lanceolatis subintegerrimis, spiciis verticillatis paniculatis. 3 in humidis Gallis australioris passim. — *Fl. fr. 3. p. 502. Lam. ill. t. 541. f. 1. (v. v.)*

II. VERBENA *Tourn. inst. t. 94. L. gen. 32.*

Calyx 5-fidus; corolla limbo subbilabiato 5-lobo inæquali; stamina 4 didynama; textus utricularis nectens semina 4.

1. *V. officinalis* (L. sp. 29.) caule erecto solitario, foliis multifido-laciniatis, spiciis filiformibus paniculatis. ① in ruderatis. — *Fl. fr. 3. p. 503. Lam. ill. t. 17. f. 1. (v. v.)*

2. *V. suffruta* (L. sp. 29.) caulibus ramosissimis decumbentibus, foliis bipinnatifidis, spiciis filiformibus solitariis. ① in sterilibus Galloprovinciæ ex Gar., *Pyren. centralium ex Lapeyr.* — *Fl. fr. 3. p. 503. — Clus. hist. 2. p. 46. f. 1. (v. v. c.)*

Ordo LXXV. ACANTHACEÆ *Juss. gen. p. 102.*

Calyx divisus persistens, sæpè bracteatus. Corolla monopetala hypogyna decidua irregularis. Stamina 2 aut 4 didynama. Ovarium disco glanduloso basi cinctum 2-loculare, loculis polyspermis; stylus 1; stigma bilobum. Capsula 2-locularis, loculis 2 polyspermis abortu 1-spermis, elasticè bivalvis, dissepimento valvis opposito iisdem mediis innascente, scissili ab apice ad basim in placentas 2 utrinquè seminiferas et continuas valvis indè semibilocularibus. Semina processibus subulatis dissepimenti plerumque subtensa. Albumen nullum; radícula infera; cotyledones foliaceæ. — Herbæ (in nostris).

I. ACANTHUS *Tourn. inst. t. 80. L. gen. 793.*

Calyx 4-partitus subbilabiatus; corolla tubo brevi villis clauso, labio unico inferiore maximo 3-lobo; stamina 4 didynama; antheræ antice villosæ; stigma bifidum; capsula ovata, loculia 1-2-spermis.

1. *A. mollis* (L. sp. 891.) foliis sinuatis inermibus.  $\mathcal{V}$  in Galliæ mediterraneæ humidis et umbrosis. — Fl. fr. 3. p. 493. Lam. ill. t. 550. f. 2. (v. v.)
2. *A. urinosus* (L. sp. 891.) foliis pinnatis margine spinosis.  $\mathcal{V}$  in Galliæ mediterraneæ humidis et umbrosis. — Fl. fr. 3. p. 493. Lam. illust. t. 550. f. 1. — Moris. s. 15. t. 2. f. 2. (v. v.)

Ordo LXXVI. LENTIBULARIÆ *Rich. in flor. paris. 1. p. 26.*  
*Brown prod. p. 429. — Utriculineæ Hoffm. et Link. fl. portug.*

Calyx divisus persistens. Corolla monopetala hypogyna irregularis calcarata bilabiata. Stam. 2 inclusa imæ corollæ inserta; antheræ simplices nunc medio constrictæ. Ovarium 1-loculare. Stylus 1 brevissimus; stigma bilabiatum. Caps. 1-locularis polysperma, placentâ centrali magnâ. Semina parva exalbuminosa. Embryo nunc dicotyledoneus nunc indivisus. — Herbæ aquaticæ seu paludosæ, foliis radicalibus, scapis 1-∞-floris, floribus unibracteatis vel rariùs ebracteatis.

I. PINGUICULA *Tourn. inst. t. 74. L. gen. 30.*

Calyx campanulatus 5-fidus; corolla bilabiata superius triloba, inferius 2 loba brevior basi calcarata; stigma bilamellatum.

1. *P. vulgaris* (L. sp. 25.) scapo glabro, calcare conico recto longitudine corollæ, labio superiore bilobo, inferiore tripartito.  $\mathcal{V}$  in uliginosis. — Fl. fr. 3. p. 575. Lam. ill. t. 14. f. 1. (v. v.)
2. *P. lusitanica* (L. sp. 25.) foliis obtusissimis, scapo gracili subpubescente,

flore minimo nutante, calcare elongato conico obtuso.  $\mathcal{U}$  in humidis turfosis provinc. occidentaliū. — Fl. fr. 5. p. 405. eng. bot. t. 145. Lois. gall. t. 1. (v. v.)

3. *P. ALPINA* (L. sp. 25.) calcare conico ascendente corollā breviorē, labio superiore emarginato, inferiore lacinii truncatis, capsulā rostratā.  $\mathcal{U}$  in Alpibus et Pyrenæis. — Fl. fr. 3. p. 576. — Cus. hist. 1. p. 310. f. 2. (v. v.)
4. *P. GRANDIFLORA* (Lam. ill. t. 14. f. 2.) calcare subulato recto longitudine floris, corollæ labio superiore patente emarginato maximo, inferiore trilobo.  $\mathcal{U}$  in editioribus montibus. — Fl. fr. 3. p. 575. (v. v.)  
 $\beta$ . *longifolia* (DC. fl. fr. 5. p. 404.) foliis elongatis scapum subsuperantibus. — In Pyrenæis. — *P. longifolia* Fl. fr. 3. p. 728. Lapeyr. abr. p. 12. (v. v.)

II. UTRICULARIA L. gen. 31.

Calyx 2-partitus labiis æqualibus indivisis; corolla personata labio inferiore basi calcarato; stam. 2 filamentis apice intus antheriferis; stigma 2-labiatum. — Herbæ aquaticæ foliis radiciformibus demersis multifidis vesiculis numerosis instructis.

1. *U. MINOR* (L. sp. 26.) foliis 3-partitis, lacinii capillaribus dichotomis, calcare carinato, corollæ labio superiore integro.  $\mathcal{U}$  in aquis stagnantibus rariū. — Fl. fr. 3. p. 574. Lam. ill. t. 14. f. 2. eng. bot. t. 254. — Petiv. herb. t. 36. f. 12. (v. s.)
2. *U. INTERMEDIA* (Hayn. term. t. 26. f. 6.) foliis tripartito-dichotomis, ampullis radicibus (non foliis) adfixis, calcare conico, labio superiore integro palato duplō longiore.  $\mathcal{U}$  in aquis stagnantibus Alsatia? Lutetia ex Merat. — Fl. fr. 5. p. 404. (v. s.)
3. *U. VULGARIS* (L. sp. 26.) foliis pinnatifido-multifidis, lacinii capillaribus, calcare conico, labio superiore integro.  $\mathcal{U}$  in aquis stagnantibus. — Fl. fr. 3. p. 574. Lam. ill. t. 14. f. 1. (v. v.)

Ordo LXXVII. PRIMULACEÆ Vent. tabl. 2. p. 285. DC. fl. fr. 3. p. 430. Duby prim. monog. ined. — Lysimachia Juss. gen. 95.

Calyx monosepalus persistens 4-5-lobatus. Corolla 1-petala hypogyna regularis staminifera, limbo plus minus profundē diviso. Stamina corollæ inserta, ejus lobis numero æqualia et iisdem opposita. Ovarium liberum; stylus 1; stigma simplex. Capsula 1-locularis, polysperma; placenta centrali liberā. Embryo rectus intrā albumen carnosum; radícula infera. — Herbæ, foliis oppositis rariū alternis aut radicalibus simplicibus.

I. HOTTONIA L. gen. 203.

Calyx 5-partitus; corolla tubo brevi, limbo plano 5-lobato; stamina 5 subsessilia; capsula globosa stylo duplō longiore persistente coronata.

1. *H. PALUSTRIS* (L. sp. 208.) foliis pectinatis verticillatis, floribus pedunculatis verticillatis, calycis laciniis linearibus acutis tubum corollæ aequantibus.  $\mathcal{U}$  in aquis stagnantibus. — Fl. fr. 3. p. 436. Lam. ill. t. 100. eng. bot. t. 364. (v. v.)

## II. LYSIMACHIA L. gen. 205.

Calyx 5-partitus; corolla 5-partita rotata tubo minimo, calyce longior; stamina 5 rarius 6; filamenta in quibusdam basi connata; capsula globosa, 5-10-valvis, apice dehiscens polysperma.

### Sect. I. NAUMBURGIA Mœnch. meth. supp. p. 23.

Flores thyrsoidei, racemi axillares multiflori; stam. 6 rarius 5.

1. *L. THYRSIFLORA* (L. sp. 209.) foliis oppositis lanceolatis acutis, calyce 5-partito laciniis lineari-lanceolatis acutis, corollæ lobis lineari-spathulatis.  $\mathcal{U}$  in paludibus Galliae septentrionalis. — Fl. fr. 3. p. 435. Fl. dan. t. 517. (v. s. ex Helvetiâ.)

### Sect. II. EPHEMERUM.

Flores racemosi, racemi terminales, stamina exserta.

2. *L. EPHEMERUM* (L. sp. 209.) foliis elongato-lanceolatis glabris sublaevibus sessilibus, corollæ lobis obovatis patulis.  $\mathcal{U}$  in Pyrenæis orientalibus. — Fl. fr. 5. p. 381. — Till. pis. 106. t. 40. f. 2. — Moris. s. 6. t. 10. n. 16. Fl. albi. (v. s.)

### Sect. III. LYSIMASTRUM.

Flores paniculati aut solitarii axillares, stamina corollâ breviora.

3. *L. VULGARIS* (L. sp. 209.) caule erecto, foliis oppositis ternisque ovato-lanceolatis acuminatis, floribus paniculatis, calycis laciniis ovatis, corollæ lobis ovatis obtusis.  $\mathcal{U}$  ad ripas et paludes. — Fl. fr. 3. p. 434. eng. bot. t. 761. — Blackw. t. 278. (v. v.)
4. *L. NUMMULARIA* (L. sp. 211.) caule repente, foliis subrotundis subcordatis integerrimis obtusis petiolatis, pedunculis solitariis folio longioribus, laciniis calycinis ovato-lanceolatis acutis corollâ duplò brevioribus.  $\mathcal{U}$  in pascuis humidis et umbrosis. — Fl. fr. 3. p. 435. Fl. dan. t. 493. — Moris. s. 5. t. 26. f. 5. n. 1. (v. v.)
5. *L. NEMORUM* (L. sp. 211.) caule procumbente, foliis integerrimis ovatis acutis petiolatis, pedunculis solitariis filiformibus per maturationem recurvis, laciniis calycinis linearibus acuminatis corollâ brevioribus.  $\mathcal{U}$  in nemoribus montanis. — Fl. fr. 3. p. 435. Fl. dan. t. 174. Lerouxia nemorum Mer. fl. par. ed. 11. p. 149. — Moris. s. 5. t. 26. f. 6. n. 2. (v. v.)

### III. ASTEROLINUM Link et Hoff. fl. portug. 333. Duby prim. mon. ined.

Calyx 5-partitus; corolla calyce triplo quadruplo brevior, tubo ovato limbo campanulato tubum subaequante 5-loba lobis rotundatis; capsula globosa 5-valvis 2-3-sperma; placenta globosa.

1. *A. STELLATUM* (Link et Hoff. l. c.) caule erecto ramosissimo, foliis oppositis sessilibus lanceolatis acutis, calycis laciniis lineari-lanceolatis acuminatis. ① in aëcie et herbosis Galliae australis. — Lys. linum-stellatum Fl. fr. 3. p. 436. — Magn. monsp. 163. ic.

### IV. CENTUNCULUS L. gen. 145.

Calyx 4-lobus; corolla rotata 4-loba; stamina 4; capsula globosa circumscissa.

1. *C. MINIMUS* (L. sp. 169.) foliis alternis ovatis, floribus sessilibus. ① in ali-



ginosis, pascuis et ericetis inundatis passim. — Fl. fr. 3. p. 431. — Vaill. bot. t. 4. f. 2. (v. v.)

V. ANAGALLIS Tourn. inst. t. 59. L. gen. 206.

Calyx 5-partitus; corolla rotata 5-loba; capsula globosa circumscissa.

1. A. ARVENSIS (L. sp. 211.) caulibus ramosis subprocumbentibus, foliis oppositis ternisve ovatis sessilibus subacutis, pedunculis folio longioribus, calycis laciniis lanceolato-linearibus acuminatis corollæ subæqualibus. ① in campis et cultis. — A. cærulea et phænicea. Fl. fr. 3. p. 431 et 432. — Cam. epit. 394 et 395. ic. Variat floribus purpureis carneis cæruleisve. (v. v.)
2. A. ARVEN (DC. syn. gall. p. 205.) caule ramosissimo ramisque repentibus, foliis sessilibus ovatis oppositis, pedicellis folio subæqualibus, calycis laciniis lanceolato-linearibus acuminatis. ½ in montibus Galloprovinciæ Sedonensibus detexit cl. Clarion. — Fl. fr. 5. p. 381. (v. s. spec. imperf.)
3. A. CRASSIFOLIA (Thore chl. land. p. 62.) caule repente, foliis alternis subrotundis crassis subpetiolatis, pedunculis folio brevioribus, calycis laciniis lanceolato-linearibus acuminatis corollâ brevioribus. ½ in uliginosis turfosis circa Aquas Tarbellicas. — Fl. fr. 3. p. 433. (v. v.)
4. A. TENELLA (L. mant. 335.) caule prostrato filiformi, foliis subrotundis ovatisve mucronulatis oppositis petiolatis, pedunculis folio longioribus, calycis laciniis lanceolato-linearibus acuminatis corollâ quadruplò brevioribus. ½ in humidis et paludosis provinciæ australium occidentaliumque et circa Parisios. — Fl. fr. 3. p. 432. Lysimachia tenella L. sp. 211. — C. Bauh. prod. 136. ic. (v. v.)

VI. CORIS L. gen. 243.

Calyx ventricosus 5-dentatus dentibus basi spinosis spinis exsertis; corolla tubulosa limbo 5-fido inæquali; stam. 5; capsula globosa 5-valvis 5-sperma.

1. C. MONSPELIENSIS (L. sp. 252.) caulibus sublignosis, foliis linearibus, floribus racemoso-capitatis. ½ in Galliæ mediterraneæ maritimis aridis. — Fl. fr. 3. p. 437. Lam. ill. t. 102. — Cam. epit. 699. ic. (v. v.)

VII. TRIENTALIS Tourn. L. gen. 461. DC. fl. fr. 5. p. 382.

Calyx 7-partitus, corolla rotata 7-partita; stamina 7; bacca exsucca suturis dehiscens.

1. T. EUROPEA (L. sp. 488.) caule erecto simplici basi nudo, foliis confertis lanceolatis, calycis laciniis linearibus acuminatis. ½ in Arduennâ sylvâ circa Sanctum Hubertum, in Delphinatû ex Dalech. — Fl. fr. 1. c. eng. bot. t. 15. — Moris. s. 12. t. 10. f. ult. (v. s. ex Germ.)

VIII. ANDROSACE Tourn. inst. t. 46. DC. fl. fr. 3. p. 437. Aretia et Androsace L.

Calyx persistens quinquefidus; corolla hypocrateriformis limbo 5-lobo lobis plerumque integris, fauce ad limbum coarctatâ, tubo ovato calyce brevior; antheræ obtusæ; stylus brevissimus; capsula globosa 5-valvis, valvis ex apice ad basin dehiscentibus; semina plerumque pauca (3-5).

Sect. 1. ARETIA L. gen. 195.

Flores solitarii.

1. A. SAXONICA (DC. fl. fr. 3. p. 440.) densè cespitosa, foliis imbricatis confertis

- lanceolatis obtusis pilosis pilis simplicibus, floribus terminalibus sessilibus, calyce piloso 5-fido, lobis obtusis, capsula valvis post maturationem erectis.  $\frac{1}{2}$  in Alpibus Sedenensibus (cl. Clarion), Pyren. editoribus ex cl. Rentham. — DC. ic. rar. t. 6. — Hall. helv. t. 11. (v. v. in Helv. D.)
2. *A. IMBRICATA* (Lam. ill. t. 98. f. 4.) cespitosa, foliis subverticillatis lanceolato-oblongis obtusis apice tomentosis tomento brevi conferto albo pilis ramosis et radiantibus, floribus axillaribus pedicellatis, calycis tomentosi 5-fidi lobis acutis, capsula valvis post maturationem reflexis et patentibus.  $\frac{1}{2}$  in saxosis editiorum Pyrenæorum, Alptum Delphinatûs ex Vill. — Fl. fr. 3. p. 439. *A. argentea* Gært. carp. 3. t. 198. f. 4. (v. v. DC.)
3. *A. PYRENAICA* (Lam. ill. n. 1953.) cespitosa, foliis imbricatis carinatis recurvis linearibus obtusis ciliatis pubescentibus pilis simplicibus, pedunculis folio longioribus incurvatis glabris, calyce campanulato 5-fido subglabro lobis obtusis.  $\frac{1}{2}$  in editioribus Pyrenæis ad rupes frigidiores muscosas versus Secalejo et Averan. — Fl. fr. 3. p. 438. — *A. dispensioides* Lapeyr. fl. pyr. t. 3. (v. s.)
4. *A. CYLINDRICA* (DC. fl. fr. 3. p. 439.) cespitosa, caule ramoso ramis cylindricis, foliis dense imbricatis lanceolato-ovatis obtusissimis pilosis pilis simplicibus antiquioribus persistentibus reflexis, pedunculis folio longioribus pilosis, calyce 5-lobato lobis lanceolatis acutis.  $\frac{1}{2}$  in rupibus calcareis sylva dicta *St.-Bertrand*, propè *la Oule du Marbord* in editioribus Pyrenæis. — *A. frutescens* Lapeyr. abr. pyr. p. 92. (v. s.)
5. *A. ALPINA* (Lam. ill. t. 98. f. 3.) caulibus elongatis procumbentibus, foliis laxè imbricatis patentibus caducis lanceolatis obtusis ciliatis pubescentibus pilis apice ramosis, pedunculis folio longioribus pilosis ad florem incrassatis, calyce sub 5-partito, lobis subpatentibus lanceolato-linearibus obtusis.  $\frac{1}{2}$  in Alpibus Delphin. ex Vill. — Fl. fr. 3. p. 440. (v. v. in Sabaudia.)
6. *A. PUBESCENS* (DC. fl. fr. 3. p. 438.) caulibus erectis confertis, foliis dense imbricatis persistentibus spathulatis obtusis ciliatis apice pubescentibus pilis simplicibus rarè bifurcatis, pedunculis folio sub brevioribus ad florem non incrassatis, calyce 5-fido pubescente pilis simplicibus, lobis lanceolato-ovatis erectis acutis.  $\frac{1}{2}$  in rupestribus Alpium Delphin. ex Vill., Pyrenæorum ex Lapeyr. — Ic. rar. t. 5. *A. Alpina junior* Lapeyr. An var. sequentis cui primo aspectu simillima. (v. s. ex Helv.)
7. *A. CILIATA* (DC. fl. fr. 3. p. 441.) caulibus laxis, foliis imbricatis lanceolato-oblongis persistentibus glabris venosis ciliatis ciliis simplicibus rarissimè subbifurcatis, pedunculis folio sublongioribus incurvatis ad calycem incrassatis, calyce 5-fido pubescente pilis simplicibus, lobis erectis lanceolato-linearibus acutis.  $\frac{1}{2}$  in editioribus Pyrenæis, *Port d'Oo*, *Maladetta*. — *A. Alpina adultior* Lapeyr. (v. s.)

Sect. II. *ANDRASPIIS Duby primul. monog. ined.* — Androsace  
L. 196.

Flores in umbellam digesti, involuero scapoque muniti.

8. *A. VILLOSA* (L. sp. 203.) caulibus confertis, foliis rosulatis lineari-oblongis subobtusis utrinquè pilis longis albis obteatis, involucri foliolis acutis linearilanceolatis villosissimis pedicellos inæquales superantibus, calycis sub 5-partiti laciniis subobtusis.  $\frac{1}{2}$  in Pyrenæis monte Ventoso, Alpibus Delphin. et Jurassi summo cacumine *la Dole* dicto. — Fl. fr. 3. p. 441. Jacq. coll. 1. t. 12. f. 3. bot. mag. t. 743. (v. v.)
9. *A. CHAMÆJASME* (Wild. sp. 1. p. 799.) caulibus erectis laxis, foliis rosulatis ovato-lanceolatis acutis in petiolum subcoarctatis utrinquè subglabris margine ciliatis, scapis villosis pilis ramosis, involucri foliolis subobtusis lanceolato-ovatis villosis ciliatisque, calycis sub 5-partiti pilosi laciniis acutis.  $\frac{1}{2}$  in Alpibus Delphinatûs. — Fl. fr. 3. p. 443. exol. var. *A. villosa* Jacq. austr. t. 332. non L. — Clns. hist. 2. p. 62. ic. (v. v. in Helv.)

10. *A. obtusifolia* (All. ped. t. 46. f. 1.) caulibus erectis confertis, foliis rosulatis lanceolatis obtusis sessilibus glabris margine brevissimè ciliatis, scapis tomentosis, involucri foliolis lanceolato-ovatis acutis pedicellis subduplò minoribus, calycis 5-lobi brevissimè tomentosi lacinii acutis. 2/ in editioribus Alpibus Delphin. ex Vill. — *A. chamæjasme* var.  $\beta$  et  $\gamma$ . — Fl. fr. 3. p. 443. *A. brevifolia* Vill. dauph. 2. t. 15. (v. v. in Helv.)
11. *A. lactea* (L. sp. 204. excl. syn. All.) caulibus procumbentibus, foliis rosulatis lanceolato-linearibus acutis-sessilibus pilis brevibus ciliatis, scapis erectis paucifloris glaberrimis, involucri foliolis lineari-lanceolatis acutis glaberrimis pedicellis multo brevioribus, calycis 5-fidi glabri lacinii lanceolatis acutis. 2/ in Cebennis, Alpibus Delph. Galloprov. — Fl. fr. 3. p. 442. bot. mag. t. 981. *A. pauciflora* Vill. dauph. 2. t. 15. (v. s.)
12. *A. carnea* (L. sp. 204.) caulibus erectis brevissimis, foliis rosulatis linearibus acutis subulatisve subciliatis, scapis flexuosis multifloris pilosis, involucri foliolis acutis exterioribus subappendiculatis pedicellis subbrevioribus, calycis 5-fidi glabri lacinii lanceolatis acutis. 2/ in Vogesis, Alpibus, montibus Arvernæ, Pyrenæis. — Fl. fr. 3. p. 442. — Hall. helv. t. 17. (v. v.)
13. *A. septentrionalis* (L. sp. 203.) caulibus nullis, foliis rosulatis lanceolato-ovatis denticulatis subciliatis in petiolum alatum subcoarctatis, scapis numerosis multifloris, involucri polyphyllò minimo foliolis lanceolatis acutis, pedicellis filiformibus elongatis, calycis vix 5-lobi lacinii acuminatis. ① in frigidioribus montium Delph., Galloprov., Cebennarum ex Gouan. — Fl. fr. 3. p. 444. Lam. ill. 98. f. 2. — Mill. ic. 20. t. 30. f. 2. (v. s.)
14. *A. maxima* (L. sp. 203.) caulibus nullis, foliis rosulatis ovatis denticulatis sessilibus acutis, scapis multifloris, involucri maximi foliolis obovatis obtusis pedicellis longioribus, calycis sub 5-partiti urceolati corollà longioris lacinii acutis. ① in cultis Delphin., Galloprov., Occitanis, Ligeris, Pictaviæ et Andegaviæ. — Fl. fr. 3. p. 444. — Moris. s. 5. t. 24. f. 6. (v. v.)

IX. *GREGORIA* Duby prim. mon. ined.

Calyx persistens campanulatus 5-fidus; corolla hypocrateriformis limbo 5-fido lobis ovatis, fauce ovatâ non coarctatâ, tubo tereti calyce multo longiore; capsula ovalis 5-valvis, valvis ex apice ad basin dehiscentibus, disperma, placenta centralis compressa.

1. *G. Vitaliana* (Dub. l. c.) caulibus numerosis prostratis sublignosis, foliis imbricatis linearibus angustissimis acutis, floribus subsessilibus solitariis. 2/ in Alpibus Delphin., monte Ventoso, Pyrenæis. — *Primula Vitaliana* L. sp. p. 206. — Sess. epist. t. 10. f. 1. — Column. cephr. 2. t. 65. f. 1. Flores lutei desiccatione virescentes. (v. v.)

X. *PRIMULA* L. gen. 197. Duby l. c.

Calyx plus minus profundè 5-dentatus; corollâ hypocrateriformis limbus 5-lobus lobis plerumque emarginatis, fauce ad limbum dilatâ, tubo tereti calycem æquante aut superante; antheræ sæpè acuminatæ; capsula ovata 5-aut 10-valvis, valvulis apice tantum dehiscentibus, seminibus minimis numerosis.

Sect. I. *PRIMULASTRUM* Dub. l. c.

Calycis corollæ tubum æquantis aut subæquantis lacinia acuta; involucri foliola subæqualia sublinearia acuta. — Folia rugosa mollia crenata aut dentata.

1. *P. officinalis* (Jacq. misc. 1. p. 159.) foliis ovato-oblongis in petiolum subcoarctatis undulato-crenatis obtusis, scapo multifloro, involucri foliolis linearibus acutis, floribus nutantibus secundis, calycis campanulati ampliati 5-dentati dentibus lanceolato-ovatis. 2/ in sylvis pascuisque. — Fl. fr. 3. p. 446. — P. veris Cam. epit. 833. ic. — Fuchs. hist. 850. ic. (v. v.)

2. *P. ELATIOR* (Jacq. misc. 1. p. 158.) foliis ovatis in petiolum subcoarctatis infernè tomentosis sinuatis aut denticulatis obtusis, scapis pubescentibus multifloris, pedicellis divaricatis, calycis pubescentis subventricosi adpressi dentibus lanceolatis acutis.  $\frac{1}{2}$  in pascuis et sylvis submontosis. — Fl. fr. 3. p. 445. *P. veris elatior* L. — Fuchs. hist. p. 845. t. 851. ic. (v. v.)
3. *P. GRANDIFLORA* (Lam. fl. fr. 2. p. 248.) foliis ellipticis subsessilibus denticulatis obtusis superne glabris subtus pilosis, scapo plerumque abortivo, pedunculis unifloris foliis minoribus, calycis campanulati 5-costati sub 5-fidi lobis lanceolatis acutis.  $\frac{1}{2}$  in pratis nemoribus, etc. — Fl. fr. 3. p. 445. *P. vulgaris* Huds. *P. acutis* Fl. dan. t. 194. *P. brevistyla* DC. fl. fr. 5. p. 383. — Lob. ic. 568. Scapi cum adest involucrium foliolis linearibus acutissimis. (v. v.)

## Sect. II. AURICULA Tourn. 46.

Calycis campanulati corollæ tubo duplò aut triplò minoris lacinis subacutis; involucri foliola inæqualia ovata aut ovato-lanceolata obtusa. — Folia plana crassa obtusa.

4. *P. AURICULA* (L. sp. 205.) foliis ovato-spathulatis sessilibus glabris glaucescentibus farinaceis, scapo farinoso, involucri foliolis obtusis ovatis, calycis 5-lobi lacinis ovato-lanceolatis subobtusis.  $\frac{1}{2}$  in Alpibus Delphinatùs spontanea, in hortis culta. — Fl. fr. 3. p. 448. Jacq. austr. t. 415. — Lob. ic. 569 et 570. Variat spontanea flor. rubris et luteis, foliis integerrimis aut crenatis. (v. v. in Alp. Helv.)
5. *P. MARGINATA* (Curt. bot. mag. t. 191.) foliis lanceolato-ovatis obtusis serrato-dentatis margine albo farinaceo cinetis in petiolum subcoarctatis, involucri foliolis ovatis obtusis inæqualibus, calycis sub 5-fidi parvi intus farinacei lacinis lanceolatis acutis.  $\frac{1}{2}$  in Alpibus Delph. Galloprovinc., Pyrenæis ex Lapeyr. — *P. crenata* Lam. ill. t. 98. f. 3. DC. fl. fr. 3. p. 449. — Fl. purpurei. (v. v. DC.)
6. *P. VILLOSA* (Jacq. aust. 5. app. t. 27.) foliis spathulatis inæqualiter serratis in utraq. superficie glanduloso-villosis margine ciliatis in petiolum subcoarctatis, involucri foliolis linearibus acutis, calycis villosa-pubescentis lacinis subacutis.  $\frac{1}{2}$  in Alpibus et Pyrenæis. — *P. viscosa* et *P. hirsuta* Vill. dauph. 2. p. 467 et 469. Fl. fr. 3. p. 449. *P. pubescens* Jacq. misc. 1. t. 18 f. 2. *P. latifolia* Lapeyr. (v. v.)

## Sect. III. ARTHRITICA Dub. l. c.

Calycis corollæ tubo minoris lacinis obtusis, involucri foliola caduca oblonga acuta pedicellis majora. — Folia plana obtusa crassa.

7. *P. INTERFOLIA* (L. sp. 205.) foliis ellipticis integerrimis ciliatis nitidis sessilibus, involucri foliolis lineari-lanceolatis tomentosis, calycis 5-fidi lacinis obtusis.  $\frac{1}{2}$  in Alpibus, Corbariensibus, Cebennis et Pyrenæis. — Fl. fr. 3. p. 450. Jacq. austr. t. 327. — Clus. hist. p. 304. ic. (v. v.)

## Sect. IV. ALEURITIA Dub. l. c.

Calycis tubum corollæ subæquantis glabri lacinis acutis, involucri foliola lanceolato-linearis acuminata. — Folia plana subobtusata. farinâ plerumque aspersa.

8. *P. FARINOSA* (L. sp. 205.) foliis in petiolum subcoarctatis obtusissimis in paginâ inferiore pulvere luteo farinaceo obteetis, involucri farinacei foliolis lanceolato-linearibus basi subappendiculatis, calycis 5-fidi lacinis acutis.  $\frac{1}{2}$  in pratis humidis Jurassi, Alpium, Pyrenæorum. — Fl. fr. 3. p. 446. — Lob. ic. 570 et 571. (v. v.)

## XI. CORTUSA L. gen. 198.

Calyx 5-partitus; corolla tubo minimo, limbo campanulato sub 5-partito; stamina 5 basi limbi inserta; antheræ conniventes acuminatæ; caps. 5-valvis apice dehiscens polysperma.

1. *C. MATTHIOLI* (L. sp. 206.) foliis petiolatis villosis 7-9-lobis obtusis, calycibus corollâ brevioribus.  $\frac{1}{2}$  in Pyrenæis ex Lapeyr. P — Fl. fr. 3. p. 451. bot. mag. t. 987. Valdè dubia civis. (v. s. ex Pedem.)

XII. SOLDANELLA L. gen. 199.

Corolla campanulata lacero-multifida; stamina 5; capsula multivalvis spiraliter striata oblonga cylindrica polysperma.

1. *S. ALPINA* (L. sp. 206.) foliis petiolatis cordato-orbiculatis.  $\frac{1}{2}$  in Jurasso, Alpibus, Pyrenæis propè nives deliquescentes. — Fl. fr. 3. p. 451. — Moris. s. 3. t. 15. f. 8. (v. v.)

XIII. DODECATHEON L. gen. 200.

Calyx 5-fidus; corolla tubo minimo, limbo rotato 5-partito lobis reflexis; stam. 5 filamentis minimis, antheris magnis acutis apice conniventibus; capsula oblongo-ovata 5-valvis, valvis exapice ad basin dehiscentibus, polysperma.

1. *D. MEADIA* (L. sp. 207.) foliis ovato-oblongis irregulariter dentatis, scapo multifloro, floribus natis, involneri foliolis exterioribus ovato-spathulatis, interioribus lanceolatis.  $\frac{1}{2}$  ex Virginia orta, in hortis colitur. — Fl. fr. 3. p. 452. Curt. mag. 12. — Catech. var. 3. p. 1. f. 1. (v. v. c.)

XIV. CYCLAMEN L. gen. 201.

Calyx campanulatus 5-fidus; corolla tubo ovato ad limbum dilatato, limbo 5-partito reflexo; stam. 5 imo tubo inserta; antheræ sessiles; capsula coriacea aut subcarnosa globosa polysperma.

1. *C. EUROPEUM* (L. sp. 207.) foliis reniformibus crenatis basi cordatis sinibus rotundis, calycis lobis ovatis acutis.  $\frac{1}{2}$  in sylvis umbrosis et lapideosis montium. — Fl. fr. 3. p. 452. — Moris. s. 13. t. 7. f. 1. v. 2. (v. v.)
2. *C. HEDERAEVOLVUM* (Ait. kew. 1. p. 196.) foliis oblongis crenatis subhaustatis angulatis basi cordatis sinibus angulatis divaricatis, calycis 5-fidi lobis lanceolatis acuminatis.  $\frac{1}{2}$  in montosis Corsicæ, Galloprovinciæ et Occitaniæ? — Folia sæpè zonata. Flores sæpè antè folia nascuntur. — *C. linearifolium* DC. fl. fr. 3. p. 453. ic. rar. t. 8. nihil aliud mihi videtur quam hujus speciei specimen, cujus pedunculi aliquot floribus aliquo casso orbatu foliorum linearium dessicatione speciem induerunt; aliter cepset summus De Candolle qui (organogr. 1. p. 181.) hæc organa habet ut phyllodia aut pedunculos floribus orbatos. Dub. in bibl. univ. genev. mai 1827. sc. et art. p. 76. (v. s.)

XV. GLAUX L. gen. 201. Dut. de Salv. et St.-Hil. mem. mus. 2. p. 393.

Calyx campanulatus 5-fidus coloratus; cor. nulla; stam. 5 hypogyna; stigma capitatum; capsula 1-locularis 5-valvis; semina placentæ centrali globoso affixa, embryo rectus umbilico parallelus.

1. *G. MARITIMA* (L. sp. 301.)  $\frac{1}{2}$  in Oceani maritimis et Nicæ ex All. — Fl. fr. 4. p. 411. Lam. ill. t. 141. Folia opposita coriacea ovato-lanceolata, flores sessiles axillares. (v. v.)

XVI. SAMOLUS L. gen. 222.

Calyx semi-adhærens persistens; corolla hypocrateriformis 5-loba, squamis 5 filiformibus introflexis è corollæ sinibus ortis; stamina 5 immè corollæ inserta; ovarium semi-inferum; capsula semi-5-valvis.

1. *S. VALERANDI* (L. sp. 243.) caule erecto, foliis ovato-spathulatis obtusis.  $\frac{1}{2}$  in paludosis. — Fl. fr. 3. p. 454. — Moris. s. 3. t. 24. n. 28. f. 10. (v. v.)

Ordo LXXVIII. GLOBULARIÆ DC. fl. fr. 3. p. 427. *Cambess. glob. mon. in ann. sc. nat. 9. p. 15.*

Flores capitati, involucro polyphylo cincti, receptaculo paleaceo impositi. — Calyx 1-sepalus tubulosus 5-lobus. Corolla hypogyna, receptaculo inserta, tubulosa inæqualiter 5-loba. Stamina 4-5 summo tubo inserta et corollæ lobis alterna. Antheræ medio dorso insertæ, uniloculares. Ovarium liberum ovoideum 1-loculare ovulo unico pendulo. Stylus bifidus. Fructus ovoideus calyce persistente tectus 1-spermus. Embryo rectus; radícula supera; albumen carnosum. — Herbari aut suffrutices, foliis alternis. Ordinem vicinum Dipsaceis affirmat cl. Cambessedes l. c.

I. GLOBULARIA *Tourn. t. 265. L. gen. n. 112.*

Vide caracterem ordinis.

\* *Fruticosæ.*

1. *G. ALPUM* (L. sp. 139.) caule erecto fruticoso, foliis lanceolato-ovatis mucronatis integris apice interdum tridentatis, capitulis terminalibus, involucri squamis imbricatis lanceolatis margine ciliatis, calycibus hirsutis. ♂ in lapidosis Galliæ mediterraneæ. — Fl. fr. 3. p. 427. — Lob. ic. t. 371. f. 2. Garid. six. t. 42. (v. v.)
2. *G. CORDIFOLIA* (L. sp. 139.) caulibus prostratis suffruticosis, foliis canei-formibus petiolatis integerrimis apice emarginatis, dente intermedio minimo, scapis unifloris subnudis, involucri squamis imbricatis lanceolatis acuminatis, calyce hirsuto. ♂ in saxosis et apricis Alpium, Vogesorum, Jurassi. — Fl. fr. 3. p. 429. Lam. ill. t. 56. f. 2. (v. v.)
3. *G. NANA* (Lam. dict. 2. p. 723.) caule fruticoso prostrato ramoso, foliis ovato-spathulatis integerrimis petiolatis mucronatis, scapis nudis brevissimis, squamis imbricatis lanceolatis acuminatis, calycibus hirsutis. ♂ in collibus et montibus apricis à Nicæa usque ad Pyrenæos passim. — Fl. fr. 3. p. 429. *G. cordifolia* ≠ *nana* Cambess. l. c. p. 27. *G. punctata* Lspeyr. (v. v.)

\*\* *Herbaceæ.*

4. *G. NUDICAULIS* (L. sp. 140.) foliis radicalibus integerrimis ovatis oblongis obtusis petiolatis, scapo nudo, involucri squamis lineari-lanceolatis acuminatis, calycibus glabris. ♀ in Alpibus meridionalibus et in Pyrenæis. — Fl. fr. 3. p. 428. Jacq. aust. t. 230. (v. v.)
5. *G. VULGARIS* (L. sp. 139.) caule herbaceo, foliis radicalibus petiolatis ovato-spathulatis apice tridentatis, caulinis alternis ovato-lanceolatis lanceolatisve acutis, squamis lanceolatis acuminatis, calycibus hirsutis. ♀ in sterilibus et pratis siccis. — Fl. fr. 3. p. 428. — Lob. ic. t. 478. f. 2. (v. v.)

## Subclassis IV. MONOCHLAMYDEÆ.

Perigonium simplex, petalis nempè aut nullis aut cum calyce coalitis.

Ordo LXXIX. PLUMBAGINEÆ *Juss. gen. 92.*

Perigonium duplex. persistens; externum (involucrum? calyx ex R. Brown?) 1-sepalum tubulosum integrum aut dentatum; interius corollaceum hypogynum mono-aut polypetalum. Stamina 5, in monopetalis receptaculo inserta, in polypetalis imis petalis adnata. Ovarium simplex liberum monospermum, ovulo inverso pendulo ab apice funiculi è fundo ovarii orti. Styli plures, aut unicus stigmata plura gerens. Capsula monosperma. Semen inversum. Embryo compressus albumine farinaceo cinctus. — Herbæ aut suffrutices; folia simplicia integra alterna aut radicalia; flores hermaphroditi capitati aut spicati.

I. PLUMBAGO *Tourn. inst. t. 5. L. gen. 213.*

Perigonium externum hirtum tubulosum 5-dentatum; internum petaloideum infundibuliforme monopetalum 5-lobum; stamina 5 hypogyna basi dilatata; stylus 1; stigmata 5; capsula apice dehiscens, 5-valvis.

1. *P. EUROPEA* (L. sp. 215.) foliis amplexicaulibus lanceolatis scabris, caule stricto. 2 $\frac{1}{2}$  in Gallia australi. — Fl. fr. 3. p. 424. — Moris. s. 15. t. 1. f. 2. (v. v.)

II. STATICE *L. gen. 388.*

Perigonium externum scariosum plicatum integrum; internum monopetalum aut 5-petalum coloratum persistens; stamina 5, petalis inserta; styli 5; capsula non dehiscens duplici perigonio oblecta.

Sect. I. LIMONIASTRUM *Mœnch. meth. 428.*

Flores monopetali, sessiles, secus ramos longâ serie dispositi; folia caulina sparsa.

1. *S. MONOPETALA* (L. sp. 396.) caule fruticoso folioso, foliis lanceolatis vaginantibus tuberculosi, floribus solitariis alternis, ramis paniculato-fasciculatis, squamis cucullatis vaginantibus mucronulatis dimidio brevioribus perigonio externo 5-dentato dentibus linearibus acuminatis, perigonio interno 1-petalo 5-lobato. 3 in arenis maritimis propè Narbonam. — Fl. fr. 3. p. 423. — Bocc. sic. t. 17. (v. v.)

Sect. II. LIMONIUM *Tourn. inst. t. 177.*

Flores 5-petali, sessiles, secus ramos longâ serie dispositi, squamis scariosis cincti; folia radicalia rosulata.

\* *Perigonio externo bractea internâ longiore.*

2. *S. LIMONIUM* (L. sp. 394.) foliis obovato-oblongis margine undulatis glabris obtusis subtus mucronatis petiolatis, scapo erecto paniculato, paniculâ ramosâ, floribus solitariis geminisve fasciculato-spicatis unilateralibus, squamis obovatis imbricatis, perigonio externo denticulato dentibus acutis.  $\mathcal{L}$  in maritimis Mediterranei et Oceani. — Fl. fr. 3. p. 421. Fl. dan. t. 315. Folia scapo vix triplò breviora. (v. v.)
3. *S. AURICULAEFOLIA* (Vahl. symb. 1. p. 25.) foliis spatulatis glaucis obtusis aut mucronatis petiolatis margine membranaceis, scapo tereti paniculato, paniculâ simplici, floribus solitariis geminisve dense spicato-fasciculatis secundis 2-bracteatis, squamis foliaceis obtusissimis rotundato-ovatis, perigonio externo subdentato dentibus obtusis.  $\mathcal{L}$  in arenosis maritimis raritis. — Fl. fr. 3. p. 421. Squama interna maxima flores involvens. Folia scapo octuplo decuplove breviora. (v. v.)
4. *S. RETICULATA* (L. sp. 394.) caulibus diffusis, foliis spatulato-lanceolatis petiolatis obtusis aut subtus mucronulatis, scapis teretibus paniculatis, floribus solitariis geminisve laxè spicatis unilateralibus 2-bracteatis, squamis subscariosis obtusissimis rotundato-ovatis, perigonio externo 5-dentato dentibus obtusis.  $\mathcal{L}$  in arenosis maritimis Corsicæ, Mompellii ex Gouan et Benth. — Fl. fr. 3. p. 422. — Pluckn. t. 42. f. 4. (v. s.)
5. *S. OLAEFOLIA* (Pourr. ex DC. fl. fr. 3. p. 422. non Scop.) caulibus erectis, foliis spatulato-lanceolatis mucronato-cuspidatis margine cartilagineis, scapis paniculatis ramis angulatis, floribus geminis ternis quaternisve laxè spicatis unilateralibus, squamis rotundato-ovatis externâ minimâ sæpè mucronulatâ, internâ obtusâ, perigonio externo 5-dentato dentibus obtusis.  $\mathcal{L}$  in maritimis Gallie mediterraneæ. — Nimum præcedenti affinis. an satis diversa? (v. v.)
6. *S. GLOBULARIFOLIA* (Desf. atl. 1. p. 274.) caulibus erectis, foliis obovato-spatulatis mucronatis in petiolum attenuatis, scapis erectis dichotomis, floribus unilateralibus ternatim aggregatis et simul floridis laxè spicatis, bracteis subscariosis obtusis, perigonii externi 5-dentati dentibus ovatis obtusis.  $\mathcal{L}$  in maritimis Arelatis et Seti. — Fl. fr. 5. p. 379. — Barr. ic. t. 793. Bractea interna flores amplectens sed non involvens. (v. s.)
7. *S. DICHOTOMA* (Cav. ic. 1. p. 37. t. 50.) foliis lanceolato-spatulatis mucronulatis petiolatis, scapis teretibus dichotomis ramosis ramis tuberculatis, floribus geminis secundis laxè spicatis spicis paniculatis, bracteis obtusis scariosis, perigonii externi dentibus subacutis.  $\mathcal{L}$  in arenosis maritimis agri Syrtici propè la Testa reperiit cl. Des Moulins. (v. s. D.)
8. *S. BELLIDIFOLIA* (Gou. monsp. 231.) caulibus diffusis, foliis oblongis spatulatis obtusis petiolatis, scapis teretibus dichotomis ramosissimis, ramis tuberculato-leprosis, floribus geminis secundis spicato-capitatis, spicis longè pedicellatis subcorymbosis, bracteis obtusis scariosis externâ membranaceo-pellucidâ, perigonii externi dentibus obtusis.  $\mathcal{L}$  in maritimis Gallie mediterraneæ. — Fl. fr. 3. p. 421. — Bocc. mus. t. 103. (v. v.)
9. *S. MINUTA* (L. mant. 59.) caule fruticoso ramo ramoso, foliis cuneatis margine revolutis obtusis, scapis pubescentibus ramosis, floribus geminis solitariisve laxè spicatis, spicis paniculato-corymbosis, bracteis coriaceis margine scariosis mucronulatis, perigonii externi dentibus subobtusis.  $\mathcal{L}$  in rupibus maritimis Galloprovinciæ, Corsicæ. — Fl. fr. 3. p. 423. — Pluck. t. 200. f. 3. (v. v.)
10. *S. PUBESCENS* (DC. fl. fr. 5. p. 380.) caule fruticoso, foliis scapis bracteis perigonioque externo pubescentibus, foliis ovato-spatulatis obtusis petiolatis, scapis ramosissimis dichotomis, floribus solitariis distichis spicatis, spicis confertis subcorymbosis, bracteis coriaceis non scariosis confertis obtusis, perigonii externi dentibus subobtusis.  $\mathcal{L}$  circâ Nicæam et in Galloprov. orientali. — Bocc. sic. 25. t. 13. (v. v.)



11. *S. ARTICULATA* (Lois. gall. 2. p. 723. t. 6.) caule à basi ramoso, foliis lanceolato-spathulatis mucronulatis petiolatis, scapis dichotomis ramosissimis, ramis flexuoso-articulatis tuberculosis, floribus solitariis geminisve fastigiato-corymbosis à basi articulorum nascentibus 3-bracteatis, squamis 2 externis minimis, internâ ovatâ margine scariosâ mucronulatâ, perigonii externi interno paulò minore dentibus obtusis.  $\frac{1}{2}$  ad rupes maritimas Corsicæ. — Fl. fr. 5. p. 380. (v. s.)

12. *S. xeriodora* (L. sp. 394.) foliis bracteis calycibusque tuberculato-leprosis, foliis obovato - spathulatis obtusissimis, scapis teretibus articulatis, floribus laxissimè spicatis 3-bracteatis, squamis 2 exterioribus minimis, internâ coriaceâ, perigonio externo dentibus 10, externis 5 subulatis apice hamatis. ① ② in arenis maritimis Galloprovinciæ, Occitanis. — Fl. fr. 3. p. 422. Gou. ill. t. 2. f. 4. — Magn. monsp. 157. io. Squama interna flores involvens. (v. v.)

\* \* *Perigonio externo bracteâ internâ breviorè.*

13. *S. VERULACRA* (L. sp. 396.) scapis ramosissimis paniculatis basi aphyllis? squamosis, ramulis confertis densè fasciculatis multi-squamosis squamis scariosis basi latis apice longè setaceo-cuspidatis, florum bracteâ internâ apice scariosâ obtusâ, perigonii externi 5-fidi perigonio interno duplò brevioris lacinii setaceo - subulatis.  $\frac{1}{2}$  in pratis salsis circâ Narbonam. — Fl. fr. 5. p. 380. — Pluckn. t. 28. f. 3 et 4. — Moris. s. 15. t. 1. f. 5. n. 23. Fl. rosei. (v. v.)

14. *S. DIFFUSA* (Pourr. sect. tol. 3. p. 330.) scapis ramosissimis diffusis basi aphyllis squamosis, ramulis laxis in paniculam secundam dispositis, bracteis argenteo-membraneis imbricatis breviter cuspidatis, floribus spicatis, perigonii externi interno duplò brevioris dentibus acuminatis.  $\frac{1}{2}$  in maritimis circâ Narbonam. — Fl. fr. 3. p. 423. — Pluck. t. 42. f. 5. (v. v.)

#### Sect. III. *ARMERIA* DC. fl. fr. 3. p. 419.

Flores terminales capitato-involucrati, involuero communi imbricato scarioso scapum nudum vaginante; folia subradicalia conferta.

15. *S. FASCICULATA* (Vent. cels. t. 38.) caule fruticoso supernè folioso, foliis fasciculatis canaliculatis linearibus acutis, scapo glaberrimo simpliciter foliis triplò quadruplòve longiore, squamis obtusis scariosis obovatis, perigonii externi dentibus obsoletis mucronulatis perigonio interno brevioribus.  $\frac{1}{2}$  in Corsicâ. — Fl. fr. 3. p. 420. (v. s.)

16. *S. PLANTAGINEA* (All. ped. n. 1606.) foliis oblongo-lanceolatis 3-5-nerviis acuminatis, scapo glaberrimo elongato foliis 6-10-plò longiore, involucri squamis exterioribus linearibus lanceolatisve acuminatis flores sæpè superantibus, interioribus obovatis scariosis mucronulatis, perigonii externi dentibus aristato - acuminatis interno monopetalo brevioribus.  $\frac{1}{2}$  in arenosis præsertim montanis Gallie australis, mediæ et occidentalis passim. — Fl. fr. 3. p. 420. — Jacq. vind. t. 42. (v. v.)

17. *S. ARMERIA* (Pers. ench. 1. p. 332.) foliis sublinearibus trinerviis rigidulis subacutis, scapis elongatis foliis 4-6-plò longioribus, involucri squamis exterioribus lanceolatis acuminatis flores sæpè superantibus, interioribus obovatis scariosis mucronulatis, perigonio externo basi piloso dentibus aristato-acuminatis interno monopetalo subæqualibus.  $\frac{1}{2}$  in collibus aridis Gallie mediæ australisque. — *S. armeria*  $\beta$  Fl. fr. 3. p. 419. — Fl. dan. t. 1092. (v. v.)

18. *S. ARMERIA* (L. sp. 394.) foliis linearibus planis obtusis, scapo foliis duplò quadruplòve longiore, involucri squamis scariosis obovatis obtusissimis floribus brevioribus, perigonio externo basi piloso 5-dentato dentibus acutis interno monopetalo brevioribus.  $\frac{1}{2}$  in arenosis. — Fl. fr. 3. p. 419. excl. var.  $\beta$ .

*a pubescens* (DC.) foliis angustissimis, scapis pubescentibus. — In mari-

timis. — *S. linearifolia* Laterr. fl. bord. ed. 11. p. 189. *Armeria maritima* Willd. enum. h. ber. — Eng. bot. t. 226. (v. v.)  
*β. alpina* (DC.) scapis glaberrimis. — In *Alpibus* et *Pyrenæis*. — *Armeria Alpina* W. l. c. (v. v.)

### Ordo LXXX. PLANTAGINEÆ Juss. gen. 89.

Flores hermaphroditi rariùs monoïci. Perigonium duplex; exterius (calyx ?) 4-partitum persistens; interius (corolla ?) monopetalum tubulosum hypogynum scariosum persistens basi staminifer, limbo 4-partito. Stamina 4 tubo inserta, laciniis alterna; filamentis exsertis. Antheræ biloculares, loculis longitudinaliter dehiscentibus. Ovarium liberum simplex. Stylus unicus capillaris; stigma 1 hispidulum indivisum raro semibifidum. Capsula circumscissa, in 2-4 loculamenta divisa ope dissepimenti longitudinalis mobilis nunc plani nunc tetragoni. Semina parietibus dissepimenti affixa sessilia peltata, sæpiùs definita nunc indefinita. Embryo rectus intrā albumen subcornueum. Radicula infera. — In monoïcis stamina receptaculo inserta. *Fem.* perigonium externum 3-partitum, internum urceolatum indivisum; ovarium 1-spermum. — Herbæ nunc acaules nunc caulescentes, flores sessiles spicati vel spicato-câpitati 1-bracteati.

#### I. LITTORELLA L. mant. 295.

Monoïca. *Masc.* pedicellati 4-partiti; stamina receptaculo inserta. *Fem.* Fl. sessiles; perig. externum 3-partitum; internum urceolatum ore coarctatum obsolete dentatum; ovarium monospermum.

1. *L. LACUSTRIS* (L. mant. 160 et 295.)  $\frac{1}{2}$  in herbosis humidis inundatis, circa lacus stagna et aquilegia passim. — Fl. fr. 3. p. 417. — Juss. acad. 1742. t. 7. — Moris. s. 8. t. 9. f. ult. n. 30. (v. v.)

#### II. PLANTAGO L. gen. 142.

Flores hermaphroditi, capsula 2-4-locularis, 2-aut  $\infty$ -sperma.

##### Sect. I. CORONOPUS Tourn. t. 49.

Dissepimentum 3-4-laterale, semen unicum in quaque facie gerens.

1. *P. CORONOPUS* (L. sp. 166.) foliis subhirsutis pinnatifidis, scapo tereti pubescente, spicâ cylindricâ, bracteis lanceolatis ovatisve perigonio externo brevioribus. (1) in graminosis et sterilibus. — Fl. fr. 3. p. 417. Fl. dan. t. 272. — Moris. s. 8. t. 17. f. 3. n. 31. (v. v.)

##### Sect. II. PSYLLIUM Tourn. inst. t. 49.

Dissepimentum planum, semen unicum utrinque gerens.

\* *Caulibus elongatis foliosis, pedunculis axillaribus.*

2. *P. ARENARIA* (Walds. et Kit. hung. t. 51.) caule herbaceo ramoso erecto pubescente, foliis linearibus aut lineari-lanceolatis pubescentibus viscidis integris aut denticulatis basi pilosis, capitulis ovatis, bracteis pubescentibus

- lineari-lanceolatis perigonii externi lacinias lanceolatas acutas superantibus, exterioribus foliaceis obovatis longè acuminatis capitulum subæquantibus.  $\frac{1}{2}$  in sterilibus et arenosis passim. — Fl. fr. 3. p. 416. P. Psyllium Bull. herb. t. 363. non L. (v. v.)
3. P. PSYLLIUM (L. sp. 167.) caule herbaceo ramoso pubescente, foliis lineari-bus subdentatis basi pilosis, capitulis ovato-globosis, bracteis linearibus aut lineari-lanceolatis perigonii externi lacinias lanceolatas margine scariousas æquantibus. ① in arenis et ægetibus provinciarum australiorum. — Fl. fr. 5. p. 378. — Moris. s. 8. t. 17. f. 8. n. 2. (v. v.)
4. P. CYNOPS (L. sp. 167.) caule ramoso basi suffruticoso, foliis linearibus strictis basi ciliatis, capitulis globosis, bracteis rotundis acutis exterioribus acuminatis perigonii externi lacinias scariousas superantibus.  $\frac{3}{5}$  in incultis prov. australiorum, mediarum usque ad Lugdunum, austro-occidentaliun propè Burdigalam ex Laterr. — Fl. fr. 3. p. 415. P. Genevensis Fl. fr. 3. p. 416. — Lob. ic. t. 437. f. 1. (v. v.)
- \*\* *Caule subnullo, pedunculis et foliis axillaribus.*
5. P. SUBULATA (L. sp. 166.) dense cespitosa, foliis linearibus subulatis daris, scapo erecto aut subflexuoso, spicâ cylindricâ strictâ, bracteis lanceolatis carinatis acutis perigonii externi lacinias ovatis subcarinatis dorso ciliato-asperis submucronatis perig. interno subbrevioribus.  $\frac{1}{2}$  in lapidosis et arenosis Gallie australis et occidentalis. — Fl. fr. 3. p. 415. P. pungens Lapeyr. — Lob. ic. t. 439. f. 2 et 438. f. 2. Folia sæpè basi lanata. (v. v.)
6. P. SERPENTINA (Lam. ill. n. 1686.) cespitosa, foliis linearibus glabris acutis, scapo ascendente tereti, spicâ cylindricâ sæpius curvatâ, bracteis lineari-lanceolatis flore longioribus, perigonii externi lacinias ovatis submucronulatis.  $\frac{1}{2}$  in Gallia australi et in Andegaviâ (Bast.). — Fl. fr. 3. p. 414. (v. v.)
7. P. GRAMINEA (Lam. ill. n. 1685.) foliis sublinearibus planis subdentatis sub-acutis, scapo tereti ascendente, spicâ elongatâ cylindricâ densâ, bracteis rotundo-ovatis ovatisve glabris acutis perigonii externi lacinias lanceolatas obtusas margine scariousas subæquantibus.  $\frac{1}{2}$  in Gallie australis montosis, Cebennis, Arvernâ et paludosis ad ostium Sequanæ. — Fl. fr. 3. p. 413. P. coronopus J. integralis. — Fl. fr. 5. p. 378. — Dod. pempt. 108. ic. (v. v.)
8. P. MARITIMA (L. sp. 165.) foliis semicylindraceis integerrimis carnosis sub-acutis, scapo tereti pubescente erecto vel basi ascendente foliis longiore, spicâ cylindricâ elongatâ densâ, bracteis rotundatis glabris obtusis perigonii externi lacinias ovatis obtusis margine scariosis duplò brevioribus.  $\frac{1}{2}$  in maritimis Mediterranei et Oceani. — Fl. fr. 3. p. 412. Fl. dan. t. 243. (v. v.)
9. P. ALPINA (L. sp. 165.) foliis sublinearibus elongatis integerrimis acutis utrinque attenuatis glabris mollibus, scapo tereti foliis duplò longiore, spicâ cylindricâ, bracteis ovatis subacutis perigonii externi lacinias ovato-lanceolatas glabras superantibus.  $\frac{1}{2}$  in pascuis Jurassi, Alpium, Montis-Aurei, Pyrenæorum. — Fl. fr. 3. p. 413. Jacq. vind. t. 125. (v. v.)
10. P. INCANA (Ram. in DC. fl. fr. 3. p. 414.) foliis sublinearibus elongatis sub-denticulatis integrisve acutis pubescentibus planis basi lanatis, scapis incurvis pubescentibus folio duplò longioribus, spicâ oblongo-cylindricâ, bracteis rotundatis subacutis ciliatis perigonii externi lacinias ovatis obtusis glabriusculis subbrevioribus.  $\frac{1}{2}$  in humidis Pyrenæorum et Cebennarum. (v. v. DC.)
- $\beta$ . *capitellata*, foliis linearibus scapo triplò quadruplòve brevioribus subglabris, spicâ cylindrico-globosâ pauciflorâ. — In Pyrenæis et editoribus montibus Corsicæ. — P. capitellata Fl. fr. 3. p. 414. (v. v. DC.)
11. P. ALBICANS (L. sp. 165.) foliis lineari-lanceolatis sericeo-pubescentibus utrinque attenuatis acuminatis, scapis ascendentibus teretibus sublanatis, spicâ elongatâ laxiusculâ subhirsutâ, bracteis rotundato-ovatis obtusis perigonii externi lacinias ovatas obtusas margine scariousas superantibus.  $\frac{1}{2}$  in sterilibus agri Ruscinonensis, Occitanie, Galloprov., Delph. australioris. — Fl. fr. 3. p. 411. — Clus. hist. 2. p. 110. ic. (v. v.)

12. *P. FILOSA* (Pourr. act. toul. 3. p. 324.) foliis linearibus lineari-lanceolatisve hirtis acuminatis basi lanuginosis, scapo tereti hirtio foliis sublongiore, spicâ elongato ovatâ, bracteis lanceolatis subulatis perigonii externi lacinias ovatas acutas superantibus, inferioribus flore longioribus. (1) In sterilibus maritimis provinciarum australium. — Fl. fr. 3. p. 412. *P. Bellardi* All. ped. t. 85. f. 3. (v. v.)
13. *P. LAGORUS* (L. sp. 165.) foliis lanceolatis subdenticulatis pubescentibus acutis, scapo tereti pubescente, spicâ ovato-globosâ hirsutâ, bracteis ovatis lanceolatisve acutis perigonii externi lacinias lanceolatas pilis coronatas subsuperantibus. 2/ in siccis et petrosis Gallie australis. — Fl. fr. 3. p. 410. *P. intermedia* Lapeyr. non Gil. — Moris. s. 8. t. 16. f. 3. u. 13. (v. v.)
14. *P. MONTANA* (Lam. ill. n. 1670.) foliis ovato-lanceolatis lanceolatisve 5-nerviis subpilosis planis integris utrinque attenuatis, scapo hirsuto foliis subduplò longiore, spicâ ovato-globosâ, bracteis rotundatis scariosis obtusis ciliatis perigonii externi lacinias ovatas obtusas scariosas apice ciliatas superantibus. 2/ in Alpibus, Pyrenæis. — Fl. fr. 3. p. 410. (v. v.)
15. *P. VICTORIALIS* (Poir. dict. enc. 5. p. 377.) foliis lineari-lanceolatis hirsutis acuminatis, scapo tereti pubescente pilis rufescentibus basi circumdato, spicâ ovato-globosâ 12-20-florâ basi hirsutâ, bracteis obovatis hirsutis perigonium externum superantibus internoque brevioribus. 2/ in Alpibus et monte Victorie Galloprovincie, circa Vapincum ex Vill. — Fl. fr. 3. p. 410. — Ger. gallop. t. 12. (v. v. DC.)
16. *P. ARGENTEA* (Lam. ill. n. 1660 exel. syn. Ger.) foliis linearibus oblongis sericeo-argenteis sessilibus, scapo erecto tereti pubescente, spicâ ovato-sphærica densâ, bracteis obovatis obtusis dorso sericeis margine scariosis perigonium externum scariosum superantibus interno brevioribus. 2/ in montosis siccis Galloprovincie, Pyrenæorum. — Fl. fr. 3. p. 411. (v. v.)
17. *P. LANCEOLATA* (L. sp. 164.) foliis oblongo-lanceolatis nervosis utrinque attenuatis, scapis pubescentibus erectis ascendentibusve folio 2-6° longioribus, spicâ subovatâ densâ, bracteis glabris scariosis subacutis perigonii externi lacinias ovatas subacutis sequantibus. 2/ in pratis siccis frequens. — Fl. fr. 3. p. 409. — Moris. s. 8. t. 15. f. 9. (v. v.)
18. *P. MEDIA* (L. sp. 163.) foliis ovatis pubescentibus subsessilibus, scapo tereti foliis 4-8° longiore, spicâ cylindricâ abbreviatâ, bracteis obtusis perigonii externi lacinias ovatis obtusis abbrevioribus. 2/ in aridis frequens. — Fl. fr. 3. p. 409. Gærtn. fruct. 1. p. 237. t. 51. (v. s.)

### Sect. III. *PLANTAGO* DC. fl. fr. 3. p. 408.

Dissepimentum planum utriusque polyspermum.

19. *P. CORNUTI* (Gou. ill. p. 6. non Jacq.) foliis ovatis glabris carnosius utrinque attenuatis longe petiolatis scapo nudo duplò triplòve brevioribus, collo tomentoso, spicâ multiflorâ elongatâ, bracteis ovatis subobtusis perigonii externi lacinias ovatis obtusis subduplò brevioribus, capsulâ 4-spermâ. 2/ in pratis salis circa Monspelium. — Fl. fr. 5. p. 376. — Corn. can. p. 163. ic. (v. v. DC.)
20. *P. INTERMEDIA* (Gil. elem. 1. p. 125. t. 1.) foliis ovatis subsessilibus inæqualiter grossè serratis, scapo ascendente, spicâ elongatâ, bracteis ovatis acutis perigonii externi lacinias ovatas obtusas subsequantibus. 2/ in humidis arenosis Lugduni, Monspelii. — Fl. fr. 5. p. 376. Varietas probabiliter subsequens. (v. s.)
21. *P. MAJOR* (L. sp. 163.) foliis ovatis septemnerviis integris vel subdentatis in petiolo subattenuatis, spicâ elongatâ multiflorâ, bracteis ovatis acutis perigonii externi lacinias ovatis obtusis subbrevioribus. 2/ in pratis, aridis, ad vias. — Fl. fr. 3. p. 408. — Moris. s. 8. t. 15. f. 2. (v. v.)
- β. minima* (DC. fl. fr. 5. p. 376.) foliis 3-5-nerviis, spicâ 3-10-florâ. — In humidis pascuis. — *P. minima* Fl. fr. 3. p. 408. Planta vix pollicaris. (v. s.)

Ordo LXXXI. NYCTAGINEÆ *Juss. gen. 90.*

Involucrum calyciforme monophyllum uni-aut multiflorum. Perigonium monopetalum coloratum basi dilatatum suprâ ovarium coarctatum et in tubum dein ampliatus, persistens, cum ovario non adhærens. Stamina definita, inserta disco glanduloso ovarium ambiens; filamenta cum perigonii parte contractatâ adhærentia. Ovarium 1 perigonio tectum. Stylus 1. Stigma capitatum. Capsula monosperma indehiscens disco staminifero et perigonio tecta. Albumen farinaceum embryone involutum. — Herbæ aut frutices, folia sæpius opposita.

I. NYCTAGO *Royen. Juss. gen. p. 90. — Mirabilis L.*

Involucrum campanulatum 5-lobum uniflorum; perigonium campanulatum, limbo patulo 5-gono 5-lobo; stam. 5; capsula oblecta basi perigonii coriacea et densâ, ejusdem parte superiore marcescente et post inflorescentiam deciduâ.

1. *N. JALAPA* (DC. fl. fr. 3. p. 426.) floribus congestis pedunculatis foliisque glabris. ① ② è Peruvia orta, in hortis culta. — *Mirabilis Jalapa L. sp. 252. (v. v. c.)*
2. *N. LONGIFLORA* (DC. l. c.) floribus congestis sessilibus foliisque pubescentibus. ② ① ex editioribus Mexici montibus orta, in hortis culta. — *M. longiflora L. l. c. (v. v. c.)*

Ordo LXXXII. AMARANTHACEÆ *Juss. gen. 87.*

Perigonium (calyx?) liberum 1-sepalum persistens 4-5-lobum sæpè coloratum. Stamina 3 aut 5 hypogyna libera aut monadelphæ. Ovarium unicum 1-loculare rariùs 2-loculare uni-aut rariùs pluri-ovulatum. Stylus et stigma simplex aut multiplex. Capsula 1-locularis circumscissa aut indehiscens. Semina solitaria aut rariùs plurima receptaculo centrali adfixa. Albumen farinaceum embryone curvato involutum. — Herbæ; folia alterna, integra; flores parvi interdum sexu distincti sæpè squamis coloratis cincti, spicati paniculati aut capitati.

I. AMARANTHUS *L. gen. 1064.*

Monoicus. Perigonium 3-5-lobum. *Masc.* Stamina 3 aut 5. *Fem.* Styli 3; stigmata 3; capsula 1-sperma 3-rostris circumscissa.

1. *A. ALBUS* (L. sp. 1404.) caule ramosissimo erecto, foliis ovato-oblongis setulâ mucronatis, bracteis subulatis, floribus triandris glomeratis axillaribus, glomerulis 2-partitis. ① Monspelii in vineis, in agro Ruscinonensi ex cl. Benth. — Fl. fr. 3. p. 726. (v. v. DC.)
2. *A. STYLIS* (Desf. cat. 44.) caule ramoso erecto in sculo striato, foliis ovatis integris petiolatis obtusiusculis, floribus triandris glomeratis axillaribus, bracteis lanceolato-linearibus acuminatis, perigonii laciniis linearibus

mucronulatis. ① æchus vias Galliæ mediæ, occidentalis australisque. — Fl. fr. 5. p. 374. (v. s.)

3. *A. BIRTUM* (L. sp. 1405.) caule diffuso ramoso, foliis petiolatis obovatis retusis aut emarginatis, floribus triandris glomeratis lateralibus terminalibusque, bracteis perigonique lobis lanceolatis acutis. ① ad muros in vicis pagorum. — Fl. fr. 3. p. 401. — Cam. epit. 236: ic. (v. v.)
4. *A. PROSTRATUS* (Balb. misc. p. 44. t. 10.) caule prostrato ramoso striato, foliis petiolatis ovato-subrhomboideis obtusiusculis retusis, floribus triandris dense spicatis lateralibus terminalibusve, bracteis sublinearibus acutis. ② in Corsicâ, agro Nicæensi, Occitanico, Narbonensi, Nannetensi et Andegavensi. — Fl. fr. 3. p. 727. (v. v. DC.)
5. *A. RETROFLEXUS* (L. sp. 1407.) caule erecto subramoso sulcato, foliis ovato-oblongis petiolatis obtusis subemarginatis, floribus pentandris dense spicatis terminalibus, bracteis lanceolatis acuminatis, perigonii laciniis lanceolatis mucronatis. ① in agris sterilibus, lapideis, ruderatis passim. — Fl. fr. 5. p. 374. *A. spicatus* Lam. diet. 1. p. 117. DC. fl. fr. 3. p. 401. (v. v.)

Ordo LXXXIII. CHENOPODEÆ *Vent. tabl. 2. p. 253. DC.*  
*fl. fr. 3. p. 380. — Atriplicæ Juss. gen. 83.*

Perigonium liberum monosepalum 5-partitum, æstivatione imbricatâ. Stamina imo perigonio inserta, ejusdem laciniis numero æqualia. Ovarium 1. Stylus simplex aut multiplex. Fructus indehiscens, nunc bacca multilocularis polysperma, nunc cariopsis baccata seu perigonio demum pulposo oblecta, nunc cariopsis nuda aut perigonio membranaceo oblecta. Albumen sæpius farinaceum centrale. Embryo circularis aut spiraliter tortus. Radicula infera. — Herbæ aut rariùs suffrutices; folia alterna simplicia stipulis vaginisque destituta; flores parvuli virescentes sæpius hermaphroditi.

Trib. I. *ATRIPLICEÆ*. — Chenopodæ *Rob. Brown congo* 33.

Stamina definita perigonii laciniis opposita, rarissimè indefinita.

† FRUCTUS CAPSULARIS, ALBUMEN CARNOSUM.

I. THELIGONUM *L. gen. 1068.*

Monoicum, *Masc.* Perigonium turbinatum 2-fidum, laciniis revolutis; stamina 12-19. *Fem.* Perigonium 2-fidum persistens; stylus unicus, capsula 1-sperma globosa coriacea; albumen globulosum bifidum carnosum; embryo filiformis incurvus; radicula infera.

1. *T. CYNOCRAME* (L. sp. 1411.) ① in rimis saxorum umbrorum Corsicæ, Nicæ, Stæchadum, Mompeli. — Fl. fr. 3. p. 399. Lam. ill. t. 777. — C. Bauh. prod. p. 59. ic. Caules prostrati, folia ovalia subobtusâ petiolata. (v. v.)

II. POLYCNUM *L. gen. 53.*

Perigonium 5-partitum; stamina 3; stylus 2-fidus; capsula 1-sperma non dehiscens.

## CHENOPODEÆ.

395

1. *P. ARVENSE* (L. sp. 50.) caule diffuso, foliis subulatis triquetris. ① in arvis. — Fl. fr. 3. p. 398. Lam. ill. t. 23. (v. v.)

### III. CAMPHOROSMA L. gen. 164.

Perigonium urceolatum 4-partitum laciniis alternis majoribus; stam. 4 exserta; stylus 2-fidus; capsula monosperma.

1. *C. MONSPELIACA* (L. sp. 178.) foliis hirsutis linearibus. ♂ in sabulosis ad vias in Galloprovincia, Occitania. — Fl. fr. 3. p. 398. Lam. ill. t. 86. (v. v.)

++ FRUCTUS PERIGONIO TECTUS; ALBUMEN FARINACEUM.

### IV. CORISPERMUM L. gen. 12.

Perigonium 2-partitum; stamina 1-5; semen ovatum compressum hinc planum inde convexum nudum margine membranaceum.

1. *C. HYSSOPIFOLIUM* (L. sp. 6.) foliis linearibus muticis. ① in Occitania, Agatha, Monspelii. — Fl. fr. 3. p. 397. Gært. fruct. 1. t. 76. f. 7. (v. s.)

### V. SALICORNIA Tourn. inst. t. 485. L. gen. 10.

Perigonium tubuloso-ovoideum compressum vix 5-dentatum; stam. 1-2 exserta; stylus 1 brevis; stigmata 2 exserta papillosa; utriculus perigonio tectus; semen cochleatum.

1. *S. HERBACEA* (L. sp. 5.) caule herbaceo carnoso ramoso articulo articulis apice compressis emarginato-bifidis, ramis patentibus, sterilibus acutis, spicis axillaribus oppositis pedunculatis, squamis obtusis. ① in paludosis maritimis et in salsis Lotharingæ. — Fl. fr. 3. p. 396. Lam. ill. t. 4. f. 1. (v. v.)
2. *S. FRUTICOSA* (L. sp. 5.) caule erecto fruticoso, ramis ascendentibus obtusis, articulis ramulorum ancipitibus, superioribus sterilibus crassitie triplo longioribus, spicis cylindricis breviter pedunculatis, squamis floralibus truncatis membranaceis. ♂ in maritimis Mediterranei et rarius Oceani. — Fl. fr. 3. p. 397. (v. v.)
3. *S. MACROSTACHYA* (Moric. fl. venet. 1. p. 2.) caule fruticoso ramis ascendentibus, articulis superioribus vix crassitie longioribus, spicis crassis subclavato-cylindricis sessilibus. ♂ in arenosis maritimis Corsicæ, Occitanicæ. (v. v.)

### VI. SALSOLA Bieb. DC. fl. fr. 3. p. 394. — Salsolæ sp. L.

Perigonium 5-partitum persistens, laciniis post inflorescentiam appendice scarioso (peraphyllo) dorso instructis; stam. 5; stigmata 2-3; semen solitarium; embryo periphericus.

Sect. I. *KALI* Tourn. inst. t. 128. — *Salsola* Roth. in Schrad. journ. 1800. 1. p. 307.

Perigonium 5-partitum, embryo spiralis.

1. *S. KALI* (L. sp. 322.) caule herbaceo prostrato, foliis subulatis spinosis scabris, inferioribus linearibus linearibus triquetris, superioribus ovalibus, floribus solitariis axillaribus, perigonii fructiferi appendicibus latis scariosis. ① in maritimis Mediterranei et Oceani. — Fl. fr. 3. p. 396. Gært. fruct. 1. t. 75. f. 4. S. decumbens Lam. ill. t. 181. f. 2. (v. v.)
2. *S. TRAGUS* (L. sp. 322.) caule herbaceo ascendente ramoso, foliis carnosis lineari-subulatis spinosis lævibus subtriquetris, superioribus lanceolatis, floribus axillaribus solitariis, perigonii fructiferi appendicibus brevissimis. ① in maritimis Gallicæ australis, circa Nannetes, ad littora Calvadosii (de Bréb.), ad ostium Sequanæ (Le Drév.). — Fl. fr. 3. p. 396. (v. v.)

3. *S. SONA* (L. sp. 323.) caule erecto ascendente herbaceo glaberrimo ramoso, foliis linearibus carnosius mucronulatis, floribus hermaphroditis axillaribus solitariis, perigonii sepalis lanceolatis acutis fructiferis auctis rotundatis, appendicibus nullis. ① in maritimis Gallia australis. — Fl. fr. 3. p. 395. Lob. ic. t. 394. f. 1. (v. v.)

SECT. II. *KOCHIA* Roth. in Schrad. journ. l. c. t. 2.

Perigonium 5-fidum, embryo curvatus non spiralis.

4. *S. PROSTRATA* (L. sp. 318.) caule frutescente prostrato, foliis fasciculatis linearibus inerimibus, floribus polygamis, antheris purpurascensibus. ③ in locis subsalsis agri Ruscinonensis, Narbonensis, Avenionensis, Tarasconensis. — Fl. fr. 3. p. 394. Kochia prostrata Schrad. — All. ped. t. 38. t. 4. (v. v. DC.)

VII. *CHENOPODIUM* Bieb. DC. fl. fr. 3. p. 388. — *Chenopodium* et *Salsola* sp. L.

Perigonium 5-partitum nunc 3-4-partitum, persistens nec tuberosum nec post inflorescentiam accretum; stam. 5 vel pauciora; stylus 2-fidus; stigmata 2-4; semen orbiculare integumento duplici externo crustaceo.

\* *Folii linearibus integris.*

1. *C. FRUTICOSUM* (L. sp. ed. 1. p. 221.) caule fruticoso glabro ramis erectis, foliis glabris carnosius teretibus obtusis laxè imbricatis, floribus subsolitariis axillaribus. ③ in maritimis provinc. Mediterran. et anastro-occidentium. — Fl. fr. 3. p. 393. *S. fruticosa* L. sp. 324. — Lob. ic. t. 381. f. 2. (v. v.)
2. *C. HIRSUM* (L. sp. ed. 1. p. 221.) caule herbaceo diffuso, foliis teretibus obtusis tomentosis, floribus axillaribus solitariis. ① in maritimis Occidentis. — Fl. fr. 3. p. 394. *S. hirsuta* L. sp. 323. Fl. dan. t. 187. (v. v. DC.)
3. *C. MARITIMUM* (L. sp. 321.) foliis glabris obtusis carnosius semiteretibus, floribus glomeratis axillaribus. ① in paludosis maritimis Oceani et Mediterranei. — Fl. fr. 3. p. 393. — Lob. ic. t. 394. f. 2. — Dod. pempt. 81. f. 2. (v. v.)
4. *C. SETIGERUM* (DC. cat. monsp. 94.) caule ramosissimo, foliis teretibus crassis glabris glaucis setâ rectâ longiusculâ terminatis, floribus glomeratis axillaribus. ① in salis subcultis Monspelii et Narbon. — Fl. fr. 5. p. 372. (v. v.)
5. *C. SCOPARIA* (L. sp. 321.) caulibus erectis ramosissimis, ramis strictis, foliis planis lineari-lanceolatis margine ciliatis, racemis axillaribus foliosis hirsutis. ① circa Nicmam? — Fl. fr. 3. p. 393. — Dod. pempt. 151. (v. v. c.)

\*\* *Folii ovatis aut rhomboideis, sæpius dentatis aut lobatis.*

6. *C. POLYPERNUM* (L. sp. 321.) caule decumbente, foliis ovatis integerrimis petiolatis acutis utrinque viridibus, floribus glomeratis in racemos breves divaricatos aphyllas axillares terminales dispositis. ① in cultis. — Fl. fr. 3. p. 392. *C. arrectum* Desmaz. cat. pl. bot. belg. p. 60. — Lob. ic. t. 256. f. 1. (v. v.)
7. *C. VULVARIA* (L. sp. 321.) caule diffuso, foliis glaucis rhombo-ovatis subintegerrimis, floribus glomeratis in racemos glaucos axillares breves dispositis. ① ad vias, muros et in hortis. — Fl. fr. 3. p. 392. — Moris. s. 5. t. 31. f. 10. n. 6. (v. v.)
8. *C. GLAUCUM* (L. sp. 320.) foliis ovato-oblongis obtusis repandis glabris sub-  
tibus glaucis, floribus glomeratis in racemos glabros laxos axillares simplices compositosve digestis. ① in arvis et cultis. — Fl. fr. 3. p. 392. — Tab. ic. 247. (v. v.)
9. *C. AMAROIODES* (L. sp. 320.) foliis lanceolatis remote dentatis sessilibus, floribus glomeratis in racemos foliosos laxos simplices dispositis. ① in Gallia austro-occidentali ex agro Ruscinonensi ad Nannetes. — Fl. fr. 3. p. 391. — Moris. s. 5. t. 31. f. 2. n. 8. (v. v.)



10. *C. BORAVE* (L. sp. 320.) foliis oblongis pinnatifido-sinuatis viscidis subhirsutis, floribus in racemos numerosissimos axillares breves hirsutos dispositis. ① in arenosis Gallie australis et austro-occidentalis. — Fl. fr. 3. p. 391. — Black t. 314. (v. v.)
11. *C. HYALIDUM* (L. sp. 319.) foliis cordatis petiolatis angulato-dentatis acuminatis, floribus glomeratis in racemos ramosissimos subcymosos divaricatos aphyllis dispositis, seminibus punctulatis. ① in arvis et cultis. — Fl. fr. 3. p. 391. — Vaill. par. t. 7. f. 2. (v. v.)
12. *C. BLITOIDES* (Lejeune fl. spa 126.) foliis lanceolatis acuminatis basi cuneatis inciso-dentatis lævibus subcarnosis, floribus glomeratis in racemos foliosos axillares confertos erectos dispositis. ① secus muros Parisiis. — Fl. fr. 5. p. 372. An satis à sequente distinctum? (v. s.)
13. *C. RIVARUM* (L. sp. 318.) foliis cordato-triangularibus subacutis sinuato-dentatis subtus glaucis, superioribus lanceolatis acuminatis, florum glomerulis minimis in racemos compositos graciles longissimos foliosos dispositis, staminibus stigmatibusque subexsertis. ① in ruderalis et fœmis. — Fl. fr. 3. p. 389. — Tab. ic. 427. (v. v.)
14. *C. VICIFOLIUM* (Sm. brit. 1. p. 276.) foliis hastato-sinuatis erosis postice integris, superioribus oblongis integerrimis, floribus glomeratis in racemos simplices confertos axillares terminalesve dispositis, staminibus stigmatibusque inclusis, seminibus subpunctatis. ① in cultis. — Fl. fr. 3. p. 390. An satis à sequente distinctum? (v. v.)
15. *C. LÆIOSPERMUM* (DC. fl. fr. 3. p. 390.) foliis rhomboideo-ovatis eroso-dentatis postice integris, superioribus oblongis integerrimis, florum glomerulis laxis in racemos subsimplices axillares terminalesve dispositis, stigmatibus inclusis, seminibus lævibus. ① frequens ad vias et in agris. — *C. album* et *C. viride* L. sp. 319. (v. v.)
16. *C. OPULEPOLIUM* (Schrader ex DC. fl. fr. 5. p. 372.) foliis rhomboideo-ovatis inæqualiter dentatis mucronatis subtus glaucis, florum glomerulis in racemos glaucos confertos subsimplices axillares terminalesve dispositis, staminibus exsertis, stigmatibus inclusis, seminibus lævibus. ① in locis submontanis agri Parisiensis, Andegaviensis, Monspeliensis. — Fl. fr. 5. p. 372. — Vaill. bot. paris. t. 7. f. 1. *C. erosum* Bast. (v. s.)
17. *C. MURALE* (L. sp. 318.) caule ramoso subpatulo, foliis ovato-deltoides subinciso-dentatis acutis nitidis, floribus solitariis binis aut ternis in racemos corymbosos undos compositos terminales axillaresve digestis, staminibus stigmatibusque subexsertis, seminibus lævibus. ① ad muros et vias. — Fl. fr. 3. p. 389. — Tab. ic. 428. (v. v.)
18. *C. URARUM* (L. sp. 318.) foliis triangularibus inæqualiter dentato-sinuatis, floribus glomeratis confertis in racemos compositos strictissimos axillares terminalesve cauli approximatos dispositis, stigmatibus inclusis. ① circa urbes et pagos. — Fl. fr. 3. p. 389. eng. bot. t. 717. (v. v.)
19. *C. BONUS-HABITUS* (L. sp. 318.) foliis triangulari-sagittatis integris, floribus glomeratis in spicas compositas terminales erectas aphyllas digestis, stigmatibus exsertis. ② in ruderalis, secus vias. — Fl. fr. 3. p. 388. Fl. dan. t. 579. (v. v.)

VIII. ATRIPLEX *Tourn. inst. t. 286. L. gen. 1153.*

Polygama aut sæpius monoica. *Hermaphrod.* Perigonium 5-partitum; stam. 5; pistillum mancum rarè perfectum. *Fem.* Perigonium 2-partitum, laciniis adpressis, post inflorescentiam crescentibus et semen tegentibus; stylus 2-fidus; semen erectum integumento duplici, exteriori crustaceo.

1. *A. HALIMUS* (L. sp. 1492.) caule fruticoso, foliis alternis deltoides utrinque argenteis integris aut basi obsolete dentatis in petiolum decurrentibus superioribus acutis, summis lanceolatis. ♂ ad sepes et in fuscis circa Nannetes ex

- Bon., in arenosis maritimis Nicæ, Antipolis (Stire.) — Fl. fr. 3. p. 384. — J. Bauh. hist. 1. p. 2. p. 227. f. 1. (v. s.)
2. *A. PORTULACOIDES* (L. sp. 1493.) caule fruticoso, foliis oppositis utrinque argenteis integerrimis ovato-lanceolatis lanceolatisve obtusis. ♂ in paludosis maritimis à Porta-Gratiæ ad Nicæam. — Fl. fr. 3. p. 385. — Cus. hist. 1. p. 54. ic. — Dalech. hist. 552. ic. (v. v.)
3. *A. PEDUNCULATA* (L. sp. 1675.) caule herbaceo flexuoso subramoso, foliis lanceolato-ovatis obovatisve obtusis in petiolum attenuatis integerrimis utrinque argenteis, floribus femineis cuneatis bicornibus pedunculatis. ① in maritimis circa Abbavillam. — Fl. fr. 3. p. 385. eng. bot. t. 232. — Pluck. t. 36. f. 1. (v. s.)
4. *A. ROSA* (L. sp. 1493.) caule diffuso herbaceo, ramis patulis, foliis incanis subtus argenteis rotundato-ovatis inæqualiter sinuato-dentatis, floribus spicato-glomeratis axillaribus terminalibusque, valvis seminalibus rhomboides angulato-dentatis acutis. ① in maritimis Nicæ ex All., Corsicæ, Narbonæ et Galliæ occidentalis passim. — Fl. fr. 3. p. 386. (v. v.)  
*β. alba* (DC. herb.) caule erecto, foliis glaucis, valvis seminalibus serratis. — In Arvernâ propè Claromontem. (v. s.)
5. *A. LACINIATA* (L. sp. 1494.) caule herbaceo erecto summitatibus pubescentibus, foliis subtus argenteis elongato-deltoides laciniatis obtusis mucronulatis petiolatis, inferioribus oppositis, floribus glomeratis in spicas simplices densas axillares terminalesve dispositis, valvis seminalibus rhomboides angulatis obtusis. ① in maritimis Corsicæ, Galloprov., Occitanis. — Fl. fr. 3. p. 386. *A. tatarica* Lois. gall. 2. p. 694. non L. — Dod. pempt. 615. f. 4. (v. v.)
6. *A. HASTATA* (L. sp. 1494.) caule herbaceo erecto, ramis diffusis, foliis latius triangularibus hastatis glaberrimis utrinque viridibus acuminatis petiolatis, floribus glomeratis in racemos laxos elongatos axillares terminalesve dispositis, valvulis seminalibus hastato-deltoides acuminatis basi argutè inciso-serratis. ① ad muros et sepes. — Fl. fr. 3. p. 386. — Moris. s. 5. t. 32. f. 5. n. 14. (v. v.)
7. *A. MONTENSIS* (L. sp. 1493.) caule herbaceo erecto, foliis hastato-deltoides, superioribus ovato-lanceolatis obtusis integerrimis mucronulatis, valvulis seminalibus rotundo-ovatis integerrimis obtusis mucronulatis. ① ex Asia orta in oleraceis culta. — Fl. fr. 3. p. 388. — Black. t. 99 et 552. (v. v. c.)
8. *A. PROSTRATA* (Bouch. fl. abb. 76.) caule herbaceo prostrato ramosissimo, foliis triangularibus auriculatis glaberrimis utrinque viridibus obtusis petiolatis, superioribus lanceolatis acutis, valvulis seminalibus triangularibus acutis integerrimis. ① in maritimis Galliæ boreali-occidentalis usque ad Ruppellam passim. — Fl. fr. 3. p. 387. (v. v. DC.)
9. *A. PATULA* (L. sp. 1494.) caule herbaceo ramoso patulo, foliis deltoides subglaucis subhastatis denticulatis subacutis petiolatis, floribus glomeratis in racemos laxos elongatos ramos terminantes dispositis, valvulis seminalibus triangularibus acutis dorso exasperatis basi serratis. ① in maritimis Oceani et Mediterraneæ passim. — Fl. fr. 5. p. 371. non 3. p. 387. *A. oppositifolia* DC. fl. fr. 5. p. 371. (v. s.)
10. *A. ANGUSTIFOLIA* (Sm. fl. brit. 3. p. 102.) caule herbaceo patulo ramosissimo, foliis viridibus lanceolatis integerrimis acutis in petiolum attenuatis, infimis subhastatis denticulatis, floribus glomeratis in racemos graciles laxos breves axillares terminalesque digestis, valvulis seminalibus hastatis lævisculis. ① ad vias et margines agrorum. — DC. syn. 196. et Fl. fr. 5. p. 371. *A. patula* Fl. fr. 3. p. 387. non L. (v. v.)
11. *A. LITTORALIS* (L. sp. 1494.) caule herbaceo erecto ramoso, foliis lanceolato-linearibus linearibusve dentatis integerrimis acutis subpetiolatis utrinque viridibus, floribus glomeratis in racemos ramos terminantibus digestis, val-

vulvis seminalibus ovatis acutis sinuatis disco muricatis. ① in maritimis septentrionalibus, etiam circà Parisios. — Fl. fr. 3. p. 387. eng. bot. t. 708. — Booc. sic. t. 15. f. 1. (v. v.)

β. *erecta*, foliis lanceolatis, inferioribus sinuato-dentatis. — Propè Nannes. — A. *erecta* Sm. brit. 3. p. 1093. Fl. fr. 5. p. 371. (v. s.)

#### IX. SPINACIA *Tourn. inst. t. 308. L. gen. 1112.*

Dioica. *Masc.* Perigonium 5-partitum, stamina 5. *Fem.* Perigonium 2-4-partitum; styli 4; fructus tectus perigonio persistente et post inflorescentiam increcente.

1. *S. INERMIS* (Mœnch. meth. 318.) foliis oblongo-ovatis, fructibus inermibus. ① ② in hortis culta. — Fl. fr. 3. p. 384. — Moris. s. 5. t. 30. f. 5. n. 2. (v. v. c.)

2. *S. SPIROSA* (Mœnch. l. c.) foliis sagittatis, fructibus cornutis. ① ② in hortis oleraceis culta. — Fl. fr. 3. p. 385. — Moris. s. 5. t. 30. f. 6. n. 1. (v. v. c.)

#### X. BETA *Tourn. inst. t. 286. L. gen. 310.*

Perigonium 5-partitum, basi ovario semi-adhærens; stamina 5; styli 2; fructus reniformis basi perigonii capsulari involutus.

1. *B. VULGARIS* (L. sp. 322.) caule erecto, floribus ternis aut quaternis. ② colitur in hortis oleraceis. — Fl. fr. 3. p. 383. — Moris. s. 5. t. 30. f. 2 et 3. n. 3 et 6. (v. v. c.)

2. *B. MARITIMA* (L. sp. 322.) caule basi prostrato, floribus solitariis aut binis. ② in maritimis passim. — Fl. fr. 3. p. 382. (v. v.)

#### XI. BLITUM *L. gen. 14.*

Perigonium 3-partitum; stamen unicum; styli 2; fructus perigonio baccato tectus.

1. *B. VIRGATUM* (L. sp. 7.) floribus glomeratis, capitulis sparsis lateralibus. ① in humidis et cultis. — Fl. fr. 3. p. 381. — Moris. s. 5. t. 32. f. 2 et 3. n. 10 et 11. (v. v.)

2. *B. CAPITATUM* (L. sp. 7.) floribus glomeratis, capitulis spicatis terminalibus. ① in humidis et cultis. — Fl. fr. 3. p. 382. — Moris. s. 5. t. 32. f. 1. n. 9. (v. v.)

#### Trib. II. PHYTOLACEÆ *Rob. Brown congo 35.*

Stamina indefinita aut cum perigonii laciniis alterna; fructus baccatus; albumen farinaceum.

#### XII. PHYTOLACCA *Tourn. inst. t. 154. L. gen. 588.*

Perigonium 5-partitum; stamina 8-20; ovarium 8-10-striatum; stigmata 8-10; bacca 8-10-locularis, loculis monospermis.

1. *P. DECANDRA* (L. sp. 631.) floribus decandris decagynis. ② in Pyrenæis et Galliâ austro-occidentali. — Fl. fr. 3. p. 381. Lam. ill. t. 393. f. 1. (v. v.)

Ordo LXXXIV. POLYGENÆ *Juss. gen.* 82.

Perigonium liberum simplex persistens monosepalum profundè divisum, laciniis duplici serie dispositis, internis ovarii lateribus externis ovarii angulis oppositis. Stamina definita imo perigonio inserta. Antheræ 2-loculares quadrisulcatæ lateraliter rimâ duplici dehiscentes. Ovarium 1. Styli plures aut stigmata plurima sessilia. Cariopsis perigonio plus minus tecta 1-sperma. Embryo lateralis aut centralis sæpè curvatus; albumen farinaceum. — Herbæ (in nostris); caules nodosi, folia alterna basi vaginantia aut vaginæ intrafoliaceæ adnata, juniora subtus revoluta.

I. RUMEX *Campd. rum. mon. p.* 60. — Rumicis *sp. L.* D6.

Involucellum infundibuliforme vel campanulatum 3-bracteolatum; perigonium 6-partitum laciniis 3 interioribus post anthesin sæpè auctis, 3 exterioribus involucelli bracteolis adnatis; stamina 6; styli 3 reflexi; stigmata totidem dissecta; cariopsis acute trigona; embryo unilateralis; radícula sup̄era.

Sect. I. LAPATHUM *Tourn. inst. — Campd. l. c. p.* 63.

Involucellum è pedicelli articulatione ortum bracteolis nunquam spontè reflexis; styli liberi. — Sapor minimè acidus.

\* *Flores hermaphroditi; laciniis internis demùm dentatis basi nunquam cordatis, apice acuminatis vel subacutis.*

1. R. MARITIMUS (L. sp. 478.) foliis infimis ovato-lanceolatis acutis planis, superioribus oblongo-lanceolatis, ramis floriferis alternis simplicibus, verticillis confertis multifloris omnibus axillaribus, laciniis internis demùm valdè granu-  
latis ovatis acuminatis setaceo-dentatis dentibus laciniis longioribus. ① in  
paludosis præsertim maritimis. — Fl. fr. 3. p. 375. excl. var. *Campd. mon.*  
t. 1. f. 3. *J. Bauh. hist.* 2. p. 387. ic. (v. s.)
2. R. PALUSTRIS (Smith. brit. 1. p. 394.) foliis infimis lineari-lanceolatis acutis  
suberispis, ramis floriferis alternis vel geminatis simplicibus vel divisis, verti-  
cillis confertis omnibus axillaribus, laciniis demùm granulatis ovato-lanceo-  
latis acutiusculis 3-dentatis dentibus laciniis brevioribus. 2/ in cænosis et  
palustribus Parisiorum, Andegaviæ. — Fl. fr. 5. p. 368. R. maritimus 2. Fl.  
fr. 3. p. 375. — *Lob. ic.* 286. (v. s.)
3. R. PULCHRA (L. sp. 477.) foliis infimis cordato-ovatis obtusiusculis nunc  
penduriformibus, ramis floriferis alternis, verticillis remotis 1-lateralibus  
paucifloris omnibus axillaribus, laciniis internis demùm deltoideo-ovatis  
acutis obtusiusculis serrato-dentatis dentibus laciniis longè brevioribus,  
omnibus altero præcipue granulatis. 2/ ad sepes et vias. — Fl. fr. 3. p. 374.  
excl. var. *eng. bot. t.* 1576. — *Moris. s.* 5. t. 27. f. 10. n. 13. (v. v.)
4. R. PUSPUREUS (Poir. dict. 5. p. 63.) foliis infimis cordato-ovatis acutiusculis  
undulosis, ramis floriferis alternis rarè geminatis simplicibus, verticillis con-  
fertis, superioribus apylyis, laciniis internis demùm granulatis triangulari-  
ovatis acutiusculis 2-3-dentatis dentibus laciniis brevioribus. 2/ circà Late-  
tiam ex Merat. — *Campd. mon. t.* 1. f. 4. (v. s. c.)
5. R. OBTUSIPOLUS (L. sp. 478.) foliis infimis cordato-ovatis obtusis rarè  
rotundatis suberispis, superioribus ovato-lanceolatis utrinque attenuatis  
obtusis longè petiolatis, ramis floriferis alternis rariùs geminatis simplicibus,

verticillis subconfertis superioribus aphyllis, laciniis internis demum granulatis ovatis obtusiusculis 2-raro 4-dentatis dentibus laciniis brevioribus subulatis.  $\mathcal{U}$  in pratibus, ad sepes. — Fl. fr. 3. p. 375. eng. bot. t. 1999. — Lob. ic. 285. — Cam. epit. 228. ic. (v. v.)

6. *R. acutus* (L. sp. 478.) foliis infimis cordato-lanceolatis acutis margine sæpe leviter crispis, superioribus lanceolatis acuminatis, ramis floriferis alternis vel rariùs geminatis, verticillis distantibus multifloris axillaribus, laciniis internis demum ovatis acutiusculis integris vel leviter 2-3-dentatis, omnibus granulo maximo onustis.  $\mathcal{U}$  in humidis et paludosis, ad rivulos, fossas et stagna. — Fl. fr. 3. p. 375. eng. bot. t. 924. — Dod. pempt. 637. ic. — Fuchs. hist. 461. ic. (v. v.)

\* \* *Flores hermaphroditi vel polygami, laciniis internis demum integris raro serratis sæpius basi cordatis apice obtusis.*

7. *R. nemolapathum* (L. f. supp. 212.) foliis infimis cordato-lanceolatis, superioribus lanceolatis undulatis serratis acutis, ramis floriferis rectis, verticillis distantibus paucifloris superioribus aphyllis, laciniis internis demum granulatis ovatis obtusis integris.  $\mathcal{U}$  in sylvaticis, humidis et paludosis. — Fl. fr. 3. p. 373. *R. divaricatus* Thail. fl. per. excl. syn. (v. v.)

8. *R. nemorosus* (Schröd. cat. göett. ex Wild. enum. 1. p. 397.) foliis superioribus subsessilibus planis infimisque? lanceolatis, ramis floriferis rectis, verticillis distantibus fere omnibus aphyllis, laciniis internis demum integris, uno granulato.  $\mathcal{U}$  in Gallia septentrionali ex cl. Desmazières. — Fl. fr. 5. p. 367. *Nimium* præcedenti affinis. (v. s. ex Germaniâ.)

9. *R. sanguineus* (L. sp. 476.) foliis acutis leviter crispis purpurascensibus petiolatis infimis cordato-lanceolatis, ramis floriferis alternis simplicibus, verticillis distantibus paucifloris superioribus aphyllis, laciniis internis demum ovato-lanceolatis obtusiusculis integris, uno sæpius valde granulato.  $\mathcal{U}$  in paludosis et secus rivos. — Fl. fr. 3. p. 374. eng. bot. t. 1533. — J. Bauh. hist. 2. p. 988. ic. (v. v.)

10. *R. crispus* (L. sp. 476.) foliis undosis crispis acutis infimis oblongo-lanceolatis petiolatis, ramis floriferis alternis binis vel ternis simplicibus vel divisis, verticillis subconfertis multifloris superioribus aphyllis, laciniis internis demum cordato-orbicularibus integris acutiusculis profuse granulatis.  $\mathcal{U}$  in humidis et ruderalibus. — Fl. fr. 3. p. 373. eng. bot. t. 1998. — Tabern. ic. 436. — Munt. brit. t. 190. (v. v.)

11. *R. aquaticus* (L. sp. 479.) foliis infimis cordatis undosis, superioribus oblongis utrinque acutis, ramis floriferis alternis binisve, verticillis subconfertis superioribus aphyllis, laciniis internis demum deltoideo-ovatis acutiusculis integris granulo magno oblongo notatis.  $\mathcal{U}$  in stagnorum, fossarum aquaticarum oris. — Fl. fr. 3. p. 373. eng. bot. t. 2014. — Lob. ic. 285. f. 2. (v. v.)

12. *R. patientia* (L. sp. 476.) foliis inferioribus cordato-ovatis acutiusculis undatis, superioribus oblongo-lanceolatis peracutis, ramis floriferis sæpe ternis, laciniis internis demum cordatis integris acutiusculis planis, altero granulo globoso notato.  $\mathcal{U}$  Pedemontii Alpibus indigena, in hortis culta et ex cl. Merat circa Parisios subsponsanea. — Fl. fr. 3. p. 372. Campd. mon. t. 1. f. 7. — Fuchs. hist. 462. ic. (v. v. c.)

13. *R. alpinus* (L. sp. 480.) foliis infimis caulinisque cordato-ovatis obtusis petiolatis, omnibus minoribus ovato-lanceolatis, verticillis multifloris confertis sæpius aphyllis, floribus longè pedicellatis, laciniis internis demum cordatis acutiusculis apice subcontortis sinuatis uno sæpius granulato.  $\mathcal{U}$  in Alpibus, Pyrenæis et Arvernæ montibus circa stabula. — Fl. fr. 3. p. 372. 5. p. 367. Campd. mon. t. 2. f. 1. *Rheum Rhaponticum* Delarb. non L. — Moris. s. 5. t. 27. f. 2. n. 2. (v. v. DC.)

## Sect. II. ACETOSA Tourn. Campd. mon. p. 67.

Involucellum longè vel è pedicelli articulatione ortum, bracteolis demùm spontè reflexis; styli ovarii angulis supernè accreti. — Sapor acidus.

\* *Involucello longè è pedicelli articulatione orto; ramis et verticillis ferè omnibus aphyllis.*

14. R. *ARIFOLIUS* (All. ped. 2. p. 204. non Ait. nec L. f.) foliis deltoideis hastatis auriculis brevibus obtusis rectè divaricatis, inferioribus acutiusculis longè petiolatis, superioribus subacuminatis sessilibus, floribus dioicis, verticillis subconfertis paucifloris, florum fem. laciniis demùm cordato-ovatis obtusis.  $\mathcal{L}$  in summis jugis Alpium, Arvernæ, Pyrenæorum. — Fl. fr. 3. p. 377. — Boec. mus. t. 125. (v. v.)
  15. R. *AMPLEXICAULIS* (Lapeyr. fl. pyr. p. 200.) foliis infimis cordato-ovatis obtusis petiolatis auriculis obtusis nervo medio parallelis, caulinis acutiusculis sessilibus amplexicaulibus, ramis floriferis terminalibus aphyllis, floribus dioicis, verticillis subconfertis paucifloris, laciniis demùm cordato-orbicularibus obtusis subgranulatis.  $\mathcal{L}$  in Pyrenæis et Arvernæ summo jugo Cantal. — Fl. fr. 5. p. 369. — Boec. mus. 2. t. 126. (v. v. DC.)
  16. R. *TUBEROSUS* (L. sp. 481.) radice tuberosâ, foliis petiolatis infimis lanceolato-sagittatis, auriculis linearibus acuminatis patentibus, floribus dioicis.  $\mathcal{L}$  in pratis circâ Nicæam. — Fl. fr. 3. p. 376. 5. p. 368. — Tabern. ic. 449. f. 1. (v. s.)
  17. R. *ACETOSA* (L. sp. 481.) foliis subtus glaucescentibus, infimis longè petiolatis oblongo-sagittatis, auriculis retrorsum porrectis, superioribus oblongo-lanceolatis basi cordatis amplexicaulibus, verticillis nudis paucifloris, floribus dioicis, laciniis fem. internis demùm cordato-orbicularibus obtusis granulo sphæroideo parvo notatis.  $\mathcal{L}$  in pratis et pascuis spontanea, in hortis culta. — Fl. fr. 3. p. 377 excl. var. eng. bot. t. 127. — Fuchs hist. 461. ic. — Tabern. ic. 440. (v. v.)
  18. R. *INTERMEDIUS* (DC. fl. fr. 5. p. 369.) foliis infimis elongatis longè petiolatis hastato-lanceolatis acutis undosis, auriculis acutis bifidis incurvis, summis amplexicaulibus revolutis, floribus dioicis, verticillis paucifloris subconfertis, laciniis internis orbicularibus profundè cordatis obtusis demùm granulo parvo sphæroideo instructis.  $\mathcal{L}$  in siccis Galliæ meridionalis. — Campd. mon. t. 2. f. 3. (v. v.)
  19. R. *ACETOSELLA* (L. sp. 481. excl. var. f.) foliis infimis hastato-lanceolatis subovatis petiolatis, auriculis divaricatis acutis integris, summis oblongo-linearibus in petiolum subdecurentibus, ramis aphyllis filiformibus elongatis, floribus dioicis, laciniis fem. internis demùm ovatis acutiusculis cariosidi accretis.  $\mathcal{L}$  in cultis, sabulosis. — Fl. fr. 3. p. 378. eng. bot. t. 1674. Campd. mon. t. 2. f. 4. — Tabern. ic. 440 et 441. f. 1 et 2. (v. v.)
  20. R. *TINGITANUS* (L. sp. 479.) caule herbaceo, foliis hastato-ovatis acutis erosis petiolatis, floribus hermaphroditis, verticillis aphyllis paucifloris distantibus, laciniis internis demùm cordato-orbicularibus integris.  $\mathcal{L}$  in maritimis ab Arelate ad Narbonam. — Fl. fr. 5. p. 370. — Bauh. prod. 56. ic. — Moris. s. 5. t. 28. f. 10. n. 8. (v. v.)
  21. R. *SCUTATUS* (L. sp. 480.) caule herbaceo tereti, foliis cordato-hastatis obtusis longè petiolatis, verticillis paucifloris, floribus hermaphroditis, laciniis internis demùm cordato-orbicularibus obtusissimis integris.  $\mathcal{L}$  in paludosis Galliæ occidentalis, lapidosis montosis Pyren., montium Lapi, Cantal, Aurei. — Fl. fr. 3. p. 378. Campd. mon. t. 2. f. 6. — J. Bauh. hist. 2. p. 991. ic. — Mant. brit. t. 74. (v. v.)
- \*\* *Involucello è pedicelli articulatione orto, ramis floriferis et verticillis ferè omnibus axillaribus.*
22. R. *BUCEPHALOPHORUS* (L. sp. 479.) foliis infimis obovatis, floribus hermaphroditis, laciniis internis demùm deltoideo-ovatis acutiusculis 3-dentatis. (X)

in sabulosis maritimis Gallie mediterranæ. — Fl. fr. 3. p. 376. Cav. ic. t. 41. Campd. mon. t. 3. f. 2. — Col. ecphr. 1. t. 150. (v. v.)

II. OXYRIA Hill. *veg. syst.* 10. p. 24. *Campd. mon.* 153. — Rumicis sp. L. DC.

Flores hermaphroditi, involucellum nullum; perigonium 4-partitum laciniis post anthesin auctis; styli 2 reflexi; cariopsis 2-gona; embryo centralis.

1. O. DIGYNA (Campd. mon. 155. t. 3. f. 3.) 2/ in Alpibus, montibus Arvernæ, Pyrenæis, in editioribus. — Rumex digynus L. sp. 480. Fl. fr. 3. p. 379. eng. bot. t. 910. — Pluck. alm. 8. t. 252. f. 2. Folia radicalia subreniformia longè petiolata. (v. v.)

III. POLYGONUM L. gen. 495. Meisner *polyg. mon.* p. 3.

Perigonium 4-6-partitum persistens, stamina 5-9 sæpius 8; ovarium 2-3 stylos totidemque stigmata gerens; cariopsis ovata aut triangularis; embryo lateralis vel centralis, radícula supera.

Sect. I. FAGOPYRUM Tourn. t. 290. Meisn. *mon. polyg.* p. 61.

Cariopsis triquetra acuminata, embryo incurvus centralis, cotyledones induplicatæ. — Folia cordata, flores racemoso-corymbosi.

1. P. FAGOPYRUM (L. sp. 522.) caulibus erectis, stipulis amplexicaulibus abbreviatis integris basi vaginantibus, foliis cordato-sagittatis petiolatis, floribus pedicellatis racemoso-corymbosis, cariopside lævi perigonio longiore. ① ex Asiâ ortum in arvis cultum. — Fl. fr. 3. p. 370. — Moris. s. 5. t. 29. f. 1. (v. v. c.)

Sect. II. TINIARIA Meisn. l. c. p. 62.

Cariopsis triquetra acuminata, embryo lateralis incurvus, cotyledones angustæ lineares albumini accumbentes (1). — Caulis volubiles, flores laxè racemosi axillares.

2. P. DOMESTICUM (L. sp. 522.) caule volubili tereti, stipulis brevibus amplexicaulibus subacutis, foliis ovatis acuminatis basi cordatis petiolatis, racemis axillaribus foliosis, floribus verticillatis remotis pedicellatis, cariopsidibus triquetris lævibus perigonio persistente triptero arcuè obvallatis pendulis. ① ad sepes et in umbrosis. — Fl. fr. 3. p. 371. — Lob. ic. t. 624. f. 1. (v. v.)

3. P. CONVOLVULUS (L. sp. 522.) caule volubili angulato, stipulis amplexicaulibus subtruncatis brevibus, foliis hastato-cordatis acuminatis petiolatis, racemis axillaribus, floribus subverticillatis remotis, cariopside triquetra granulato-striatâ perigonio persistente involutâ. ① in agris frequens. — Fl. fr. 3. p. 371. Fl. dan. t. 744. (v. v.)

Sect. III. ACONOGONUM Meisner. l. c. p. 55.

Cariopsis acute triquetra acuminata, embryo lateralis incurvus, cotyledones planæ ovatæ albumini accumbentes. — Stipulæ simplices, flores racemoso-paniculati.

4. P. ALPINUM (All. ped. t. 68. f. 1.) caule erecto ramoso, ramis divaricatis, stipulis brevibus vaginatis truncatis hirsutis demum laceris et deciduis, foliis oblongo-lanceolatis acuminatis undulatis margine ciliatis lævibus basi in petiolum brevem decurrentibus, floribus racemoso-paniculatis. 2/ in pascuis Alpium, Pyren. orient., Corsicæ montium ex Wild. — Fl. fr. 3. p. 369. (v. v.)

(1) Cotyledones albumini *incumbentes* appellamus, cum commissura cotyledonum axi albuminis parallela est (|| o), *accumbentes* cum commissura axin albuminis spectat (=o).

Sect. IV. BISTORTA *Tourn. t. 291. Meisn. polyg. p. 50.*

*Cariopsis triquetra acuminata*, embryo lateralis, cotyledonibus accumbentibus angustis. — Spicæ densæ cylindricæ solitariae terminales, radix tuberosa.

5. *P. viviparum* (L. sp. 516.) caule simplicissimo monostachyo erecto, stipulis cylindricis dilatatis in parte superiori lateraliter dehiscentibus amplexicaulibus oblique truncatis, foliis lanceolatis acutis margine revolutis, inferioribus ellipticis in petiolum attenuatis, cariopside obtusè triquetra.  $\mathcal{U}$  in pascuis Vogesorum, Jurassii, Alpium, Pyrenæorum. — Fl. fr. 3. p. 364. — Pluck. t. 151. f. 2. (v. v.)
6. *P. bistorta* (L. sp. 516.) caule erecto simplicissimo monostachyo, stipulis dilatatis glabris truncatis demum lateraliter dehiscentibus basi vaginantibus, foliis ovato-lanceolatis acuminatis, superioribus sessilibus, radicalibus basi subcoarctatis in petiolum decurrentibus, cariopside lævi acute triquetra.  $\mathcal{U}$  in pratis et pascuis montosis. — Fl. fr. 3. p. 364. Fl. dan. t. 421. — Moris. s. 5. t. 28. f. 11. n. 2. (v. v.)

Sect. V. AMBLYGONUM *Meisn. p. 53.*

*Cariopsis globosa compressa vix mucronulata*, embryo lateralis incurvus, cotyledonibus linearibus albumini incumbentibus. — Stipulæ cylindricæ liberæ, flores densè spicati.

7. *P. orientale* (L. sp. 519.) caulibus erectis altissimis villosis supernè ramosis, stipulis vaginantibus hypocrateriformibus truncatis ciliatis, foliis ovatis acuminatis pubescentibus petiolatis, spicis elongatis densis nutantibus, cariopside lævi nitidâ. (1) in hortis cultum. — Fl. fr. 3. p. 367. Mill. ic. t. 201. Meisn. t. 3. f. k. (v. v. c.)

Sect. VI. PERSICARIA *Tourn. t. 290. Meisn. polyg. p. 66.*

*Cariopsis compressa lenticularis acuta aut acuminata*, embryo lateralis incurvus, cotyledonibus linearibus planis albumini accumbentibus. — Stipulæ teretes vaginantes simplices, flores spicati.

8. *P. amphibium* (L. sp. 517.) caulibus radicanibus procumbentibus ascendentibus, stipulis tubulosis vaginantibus caulem vestientibus illicque adnatis, foliis lanceolato-ellipticis petiolatis acutis margine scabris basi plus minus cordatis, floribus densè spicatis spicâ terminali ovato-oblongâ.  $\mathcal{U}$  in paludibus, fossis, aquaticis. — Fl. fr. 3. p. 365. eng. bot. t. 436. — Moris. s. 5. t. 29. f. 4. n. 2. (v. v.)
9. *P. hydropteris* (L. sp. 517.) caule erecto ramoso, stipulis dilatatis infernè vaginantibus supernè amplexicaulibus truncatis ciliatis, foliis lanceolatis apice attenuatis obtusiusculis glabris margine ciliato-scabriusculis, spicis elongatis filiformibus laxissimis, floribus inferioribus remotis axillaribus, cariopside obsolete 3-quetra acuminatâ nitidâ tenuissimè punctato-striatâ. (1) in aquosis et ad fossas humidas. — Fl. fr. 3. p. 365. eng. bot. t. 989. — Moris. s. 5. t. 29. f. 11. n. 6. (v. v.)
10. *P. persicaria* (L. sp. 518 var.  $\alpha$ .) caulibus procumbentibus aut ascendentibus ramosis, stipulis vaginantibus truncatis ciliatis, foliis oblongo-lanceolatis acuminatis margine brevissimè ciliatis asperis, floribus in paniculas spiciformes laxiusculas congestis, cariopside lenticulari aminatâ lævi. (1) in humidis, ad margines fossarum et viarum. — Fl. fr. 3. p. 366. eng. bot. t. 756. Folia sæpius maculata. (v. v.)

$\beta$ . *incana*, caulibus suberectis, foliis immaculatis antiquioribus subtus pubescenti-incanis. — Cum priore passim. — *P. incanum* Willd. sp. 2. p. 446. Fl. fr. 3. p. 366.

$\gamma$ . *lapathifolia* (Meisn.) caule erecto, stipulis integris, foliis ovato-lanceolatis, floribus in spicas multas axillares terminalesve densas oblongas dis-



positis. — In paludosis uberioribus. — *P. lapathifolium* L. Fl. fr. 3. p. 367. — Lob. ic. t. 315. f. 1. (v. v.)

11. *P. fusillum* (Lam. fl. fr. 3. p. 235.) caulibus ascendentibus ramosis, stipulis vaginantibus truncatis ciliato-fimbriatis, foliis lineari-lanceolatis acumine glabris margine scabriusculis, spicis filiformibus elongatis laxiusculis, cariopside lenticulari acutâ lævi nitidâ. (1) in humidis et sabulosis passim. — Fl. fr. 3. p. 366. eng. bot. t. 1043. *P. minus* Ait. Kew. 2. p. 31. *P. strictum* All. ped. t. 68. f. 2. — Lob. ic. t. 316. f. 1. (v. v.)

Sect. VII. AVICULARIA Meisn. l. c. p. 85.

*Cariopsis triquetra*, embryo lateralis, cotyledonibus angustis linearibus planis albumini incumbentibus. — Stipulæ plus minus bilobæ, flores subsolitarii axillares.

12. *P. aviculare* (L. sp. 519.) caule procumbente herbaceo ramoso folioso, stipulis 2-partitis ovato-lanceolatis acutis demum fimbriato-laceris, foliis lanceolatis subobtusis margine sæpè scabriusculis, floribus axillaribus 2-3 pedicellatis, cariopside granulato-striatâ. (1) ad vias, in agris et incultis frequens. Fl. fr. 3. p. 368. — Moris. s. 5. t. 29. f. 12. n. 1. (v. v.)

ℓ. *Bellardi*, caulibus erectis, foliis angustioribus acutis. — In agris passim. — *P. Bellardi* All. ped. t. 90. f. 2. Fl. fr. 3. p. 369. (v. v.)

13. *P. maritimum* (L. sp. 519.) caule procumbente ramoso folioso basi sub lignoso, stipulis magnis 2-partitis ovato-lanceolatis acutis demum fimbriato-laceris, foliis subcarnosis lanceolatis acutis, floribus axillaribus 2-4 pedicellatis, cariopside lævissimâ. 2 ♂ in arenosis maritimis. — Fl. fr. 3. p. 368. — Cam. epit. 691. ic. (v. v.)

15. *P. equisetiforme* (Sibth. et Smith. fl. græc. p. 266.) caule erecto ramoso aphylo basi sublignoso, stipulis apice sub 2-lobis demum dentato-fimbriatis internodiis multo brevioribus, floribus axillaribus solitariis geminativè sessilibus aut breviter pedicellatis in spicam laxam flagelliformem dispositis. 3 acutis torrentes Corsicæ. — Bocc. mus. t. 41. (v. s.)

Ordo LXXXIV. THYMELEÆ Juss. gen. 76.

Perigonium liberum coloratum monopetalum tubulosum, limbo 4-fido rarè 5-fido, æstivatione imbricatâ. Stamina definita fauci tubove inserta, perigonii laciniis numero dupla. Antheræ 2-loculares, loculis medio longitudinaliter dehiscen- tibus. Ovarium unicum, ovulo 1 pendulo. Stylus 1 sæpè lateralis; stigma 1. Fructus unicus monospermus perigonio tectus, siccus aut baccatus. Albumen nullum vel tenue carnosum. Embryo rectus; radicula supera brevis; cotyledones plano-convexæ. — Frutices, folia simplicia integra alterna aut opposita exstipulata; flores sæpius hermaphroditi interdum abortu dioici.

I. STELLERA L. gen. 488.

Perigonium tubulosum 4-lobum; stamina 8; stylus brevis; stigma capitatum; capsula dura nitida rostrata. }

1. *S. PASSERINA* (L. sp. 512.) foliis linearibus, floribus axillaribus sessilibus 4-fidis. (1) in agris. — Fl. fr. 3. p. 361. Gou. monsp. p. 44. t. 3. (v. v.)

## II. PASSERINA L. gen. 489.

Perigonium 4-lobum; stamina 8; stylus filiformis lateralis; capsula 1-locularis 1-sperma.

1. *P. dioica* (Ram. bull. phil. n. 41.) foliis subtus punctulatis lineari-lanceolatis acutis glabris confertis, floribus dioicis aut hermaphroditis axillaribus geminis ebracteatis, perigonio tubuloso, lacinii lanceolatis. ♂ in Pyrenæis. — Fl. fr. 3. p. 359. *Daphne dioica* Gou. ill. 27. t. 17. f. 1. *P. empetrifolia* Lapeyr. (v. v.)
2. *P. nivalis* (Ram. bull. phil. n. 41. t. 9. f. 4.) caulibus tortuosis prostratis ramosis, foliis nitidis lineari-lanceolatis obtusis glabris, floribus dioicis aut hermaphroditis axillaribus solitariis bracteatis, perigonio tubuloso, lacinii ovatis. ♂ in summis Pyrenæis. — Fl. fr. 3. p. 359. *P. calycina* Fl. fr. 3. p. 360. *Daphne calycina* Lam. ill. t. 290. f. 3. *P. janiperifolia* Lapeyr. abr. pyr. (v. v. DC.)
3. *P. thomasi* (Duby mss.) caulibus erectis ramosis, foliis coriaceis lineari-lanceolatis subspathulatis subacutis glaberrimis, floribus axillaribus sessilibus geminis ternisve bracteatis, perigonio tubuloso hirtio, lacinii ovatis. ♂ in montosis Corsicæ reperiit Ph. Thomas. (v. s. D.)
4. *P. tarton-raira* (DC. fl. fr. 5. p. 367.) canle hirsuto, foliis obovatis aut elliptico-lanceolatis utrinque sericeis, floribus sessilibus lateralibus aggregatis basi squamis imbricatis, perigonio hirsuto, lacinii ovatis. ♂ in rupestribus Niceæ, Galloprovinciæ, Corsicæ. — *Daphne tarton-raira* L. sp. 510. Fl. fr. 3. p. 357. — Lob. ic. t. 371. f. 2. (v. v.)
5. *P. nirsuta* (L. sp. 513. excl. Breyn. syn.) canlibus erectis ramosis tomentosis, foliis ovatis carnosis intus tomentosis extus glabris, floribus aggregatis axillaribus, perigonio extus hirsuto subcampanulato, lacinii ovatis. ♂ in lapidosis sterilibus provinc. mediterr. — Fl. fr. 3. p. 360. — Lob. ic. 2. p. 217. f. 1. (v. v.)  
*P. polygalæfolia*, foliis confertis utrinque tomentosis. — Circà Massiliam.  
*P. polygalæfolia* Lapeyr. (v. v. D.)
6. *P. thymelæa* (DC. fl. fr. 5. p. 366.) ramis simplicibus, foliis ovato-lanceolatis glaucis acutis, floribus axillaribus sessilibus infernè solitariis superne aggregatis, perigonio longè tubuloso piloso, lacinii linearibus. ♂ in provinciis mediterraneis. — *Daphne Thymelæa* L. sp. 509. Fl. fr. 3. p. 356. Gér. gallop. 2. 17. f. 2. (v. v.)

## III. DAPHNE L. gen. 485.

Perigonium 4-lobum; stamina 8; stylus brevis terminalis; bacca 1-locularis 1-sperma.

\* *Floribus terminalibus.*

1. *D. genkium* (L. sp. 511.) canle ramosissimo, foliis coriaceis lineari-lanceolatis glabris acuminato-cuspidatis, racemis terminalibus paniculatis, floribus sessilibus, tubo tomentoso, lacinii ovato-lanceolatis. ♂ in aridis provinc. mediterr. et occidentalium. — Fl. fr. 3. p. 358. — Lob. ic. t. 369. f. 1. (v. v.)
2. *D. cneorum* (L. sp. 511.) foliis elliptico-lanceolatis obtusis mucronulatis nitidis coriaceis glabris, floribus fasciculatis terminalibus sessilibus, tubo pubescente, lacinii ovato-lanceolatis. ♂ in montibus Alsatix, Lotharingiæ propè Commercy (Monnier), Jurasso, Alpibus, Cebennis, Pyrenæis, in syrtibus circà Aginum ex St.-Am., Aquas Tarbellicas ex Thore, Burdigalam ex Laterr. — Fl. fr. 3. p. 358. Bull. herb. t. 121. (v. v.)
3. *D. oleoides* (L. mant. 66.) caulibus ramosis, foliis coriaceis elliptico-lanceolatis acutis supra nitidis glabris subtus adpressè subpubescentibus, floribus ternis geminisve terminalibus sessilibus, perigonii tubo hirsuto basi dilatato, lacinii linearibus. ♂ in montosis Corsicæ. — Schreb. dec. t. 7. — Alp. exot. 44. t. 43. (v. s.)

\*\* *Floribus axillaribus lateralibus.*

4. *D. ALPINA* (L. sp. 510.) foliis elliptico-lanceolatis obtusis subtus pubescentibus, floribus sessilibus aggregatis in extremitate ramorum lateralibus, perig. tubo hirsuto, laciniiis ovatis lanceolatisve. 5 in lapideis Alpium, Cebennarum, Pyren. ex cl. Benth. — Fl. fr. 3. p. 357. — Lob. ic. t. 370. f. 1. (v. v.)
5. *D. MISERUM* (L. sp. 509.) foliis lanceolatis acutis glabris post flores evolutis deciduis, floribus ternis lateralibus sessilibus, tubo pubescente, laciniiis ovatis. 5 in sylvis montosis. — Fl. fr. 3. p. 356. Lam. ill. t. 290. f. 1. (v. v.)
6. *D. LAUREOLA* (L. sp. 510.) foliis coriaceis lanceolatis glabris acutis perennantibus, floribus axillaribus pedicellatis quinis, tubo glabro, laciniiis ovato-lanceolatis. 5 in sylvis montosis. — Fl. fr. 3. p. 357. Bull. herb. t. 37. (v. v.)

Ordo LXXXV. LAURINEÆ DC. fl. fr. 3. p. 361. — Lauri  
*Juss. gen.*

Perigonium liberum 1-sepalum persistens 6-fidum aut 6-partitum, æstivatione imbricatâ. Stamina imis laciniiis inserta, nunc 6 ordine simplici nunc 12 ordine duplici digesta. Antheræ filamento adnatæ à basi ad apicem dehiscentes. Ovarium 1; stylus 1; stigma simplex aut divisum. Drupa aut bacca 1-locularis 1-sperma. Albumen nullum; embryo rectus, radícula supra. — Arbores aut frutices; folia alterna; flores hermaphroditi aut abortu dioici.

1. *LAURUS* Fourn. inst. t. 367. L. gen. 508.

Perigonium 4-6-fidum æquale; stamina 6-9-12 duplici serie disposita; exteriora omnia fertilia, interiora alterna sterilia alterna fertilia et basi 2-appendiculata aut 2-glandulosa; drupa carnosa.

1. *L. NOBILIS* (L. sp. 529.) foliis lanceolatis vasis perennantibus, floribus 4-fidis dioicis. 5 in provinciis mediterraneis et in occidentalibus usque ad Armoraciam sub dio crescit. — Fl. fr. 3. p. 362. Duham. arb. 2. t. 134 et 135. (v. v. c.)

Ordo LXXXVI. SANTALACEÆ R. Brown. *prod. nov. holl.*  
p. 350.

Perigonium cum ovario adhærens, 4-5-fidum semicoloratum, æstivatione valvatâ. Stamina 4-5 perigonii laciniarum basi inserta iisque opposita. Ovarium uniloculare 2-4-spermum, ovulis juxtâ apicem placentæ centralis affixis pendulis. Stylus 1. Stigma sæpè lobatum. Fructus monospermus, nucamentaceus vel drupaceus. Albumen carnosum. Embryo inversus teres. — Herbæ aut suffrutices (in nostris), foliis alternis aut suboppositis exstipulatis indivisis, floribus parvis subspicatis rarò umbellatis vel solitariis.

I. THESIUM L. *gen.* 292.

Perigonium 4-5-fidum; stamina 4-5 perigonii lobis opposita; capsula monosperma non dehiscens perigonio persistente coronata.

1. T. LYOPHYLLUM (L. sp. 301.) floribus pedunculatis pentandris, caulibus erectiusculis.  $\mathcal{U}$  in collibus, pratis siccis et montosis. — Fl. fr. 3. p. 352. — Moris. a. 15. t. 1. f. 9. n. 3. (v. v.)

$\beta$ . *humifusum*, caulibus numerosis humifusis. — in arenosis maritimis Gallie occidentalis. — T. *humifusum* Fl. fr. 5. p. 366. (v. v.)

2. T. ALPIMUM (L. sp. 301.) floribus subsessilibus tetrandris, ramis subsimpli-  
cibus.  $\mathcal{U}$  in editioribus montibus. — Fl. fr. 3. p. 366. Gér. galloprov. t. 17.  
f. 1. (v. v.)

II. OSYRIS L. *gen.* 1101.

Flores abortu polygami; perigonium 3-fidum. *Masc.* Stamina 3 brevía; ovarii rudimentum. *Herm.* Stam. 3 aliquando abortiva; ovarium 1; stigma triplex; hacca globosa sicca umbilicata; nucleus 1-spermus.

1. O. ALBA (L. sp. 1450.)  $\mathfrak{f}$  in rupestribus et sterilibus Delph. et prov. austr. et austro-occidentalium passim. — Fl. fr. 3. p. 353. — Cam. epit. 26. ic. Flores parvi pedicellati in parte superiore ramorum aggregati. (v. v.)

Ordo LXXXVII. ELÆAGNEÆ *R. Brown in prod. nov. holl.*  
*p. 350. A. Rich. in mém. soc. hist. nat. paris. t. 1. p. 374. —*  
*Eleagnorum gen. Juss. gen.*

Flores dioici rariùs hermaphroditi. *Masc.* subamentiformes 3-4-8-andri, staminibus introrsis subsessilibus bilocularibus. *Fem.* Perigonium tubulosum monosepalum persistens, limbo integro aut 2-4-fido ad faucem disco glandulari prominenti variè lobato coronatus (in Hippophaë discus nullus). Ovarium è fundo perigonii assurgens nec cum illo coalitum uniloculare uniovulatum, ovulo ascendenti subpedicellato. Stylus brevissimus. Stigma linguiforme. Nux calyce baccato et akenio crustaceo constans, semen erectum. Albumen carnosum tenue; embryo rectus, radícula infera; cotyledones planæ carnosæ. — Arbores aut arbusculæ foliis alternis aut oppositis integris exstipulatis.

I. ELÆAGNUS *L. gen. 159.*

Flores hermaphroditi; perigonii tubus gracilis, limbus campanulatus 4-5-fidus, discus annularis aut 2-fidus; stamina 4-5; nux calyce baccato intus osseo et akenio constans.

1. *E. ANGUSTIFOLIA* (*L. sp. 176.*) foliis lanceolatis acutis integerrimis utrinque squamulato-argenteis nitidis, floribus in axillis foliorum 1-3 pedicellatis erectis. ♂ in humidis Galloprov. ex Gér. — *Fl. fr. 3. p. 354. Lam. ill. t. 73. f. 1. (v. v. c.)*

II. HIPPOPHAË *Nutt. gen. amer. 1. p. 97. A. Rich. l. c. p. 402.*

Flores dioici. *Masc.* amentiformes, tetrandri. *Fem.* axillares solitarii; perigonium tubulosum apice bifidum clausum, disco nullo; fructus calyce baccato et akenio constans.

1. *H. RHAMNOIDES* (*L. sp. 1452.*) ♂ in sabulosis humidis secus Mediterraneum, secus flumina in prov. orient. et littora Neustris ex cl. de Bréb. et Leprév. — *Fl. fr. 3. p. 353. Lam. ill. t. 808. Folia alterna lanceolata subtus squamato-argentea. (v. v.)*

Ordo LXXXVIII. CYTINÆ *R. Brown in linn. trans.* 13.*Ad. Brongn. in ann. sc. nat.* 1. p. 39.

Fl. monoici vel dioici. Perigonium adhærens aut ( in exoticis ) liberum, 4-5-partitum, æstivatione imbricatâ. Stam. 8, 16, vel numerosiora columnæ centrali inserta extrorsa. Antheræ 2-loculares longitudinaliter dehiscentes. Ovarium 1-loculare ( in nostris ), placentæ 4 vel 8 parietales. Stylus cylindricus vel nullus. Stigma lobatum lobis numero placentarum æqualibus. Semina numerosa. Albumen carnosum; embryo rectus 2-cotyledoneus.

I. CYTINUS *L. gen.* 123a.

Fl. monoici. Perigonium superum tubuloso-campanulatum limbo 4-fido basi squamosum. *Masc.* Columna centralis antherifera apice tuberculis conicis circiter 8 coronata; antheræ 8 circum columnæ apicem sessiles. *Fem.* Ovarium inferum 1-loculare; placentæ 8 parietales; stylus cylindricus simplex; stigma capitatum 8-sulcatum; sem. . . . .

1. *C. HYCISTIS* ( *L. gen.* p. 566. ) ? Parasitica in radicibus Cistorum, in Galloprov., Occitaniâ, Corsicâ, propè Burdigalam ex Laterr.—*Fl. fr.* 3. p. 350. *Lam. ill.* t. 737. Caulis simplex squamis oblongis ovatisve imbricatis undique tectus, flores spicati subcapitati, fem. inferiores masculi superiores, utriusque 3-bracteati. ( *v. v. DC.* )

Ordo LXXXIX. ARISTOLOCHIÆ *Juss. gen.* 74. excl.*Cytinc.*

Flores hermaphroditi. Perigonium ovario adhærens, monosepalum limbo mox 3-loba, mox tubuloso in parte superiore irregulariter dilatato; æstivatione valvatâ. Stamina definita numero ternario, mox libera et distincta, mox cum stylo et stigmate coherencia epigyna. Ovarium 3-6-loculare; stylus brevis; stigma divisum. Capsula aut bacca coriacea 6-locularis polysperma, placentæ laterales. Embryo minimus ad basin albuminis cartilaginei. — Herbæ aut suffrutices sæpius sarmentosi, foliis alternis simplicibus petiolatis.

I. ARISTOLOCHIA *Tourn. inst.* 71. *L. gen.* 1022.

Perigonium tubulosum basi ventricosum apice dilatatum et in ligulam extensum; antheræ 6 subsessiles stylo insertæ; stigma 6-fidum; capsula 6-gona 6-locularis.

\* *Floribus solitariis.*

1. *A. ROTUNDA* ( *L. sp.* 1364. ) foliis ovatis cordatis subsessilibus, radice rotundâ. ? ad vineas et in arvis Galliæ australis et austro-occidentalis usque ad Burdigalam ex Laterr. — *Fl. fr.* 3. p. 348. — *Lob. ic.* t. 606. f. 2. ( *v. v.* )

2. *A. LONGA* (L. sp. 1364.) foliis petiolatis cordatis integerrimis ovatis, radice longa simplici.  $\mathcal{L}$  in arvis, ad sepes et vineas Gallie australis. — Fl. fr. 3. p. 348. Mill. ic. t. 51. f. 2. — Cam. epit. 420. ic. (v. v.)
3. *A. PISTOLOCHIA* (L. sp. 1364.) foliis breviter petiolatis cordatis ovatis crenatis scabris, radice fasciculata.  $\mathcal{L}$  in sterilibus Galloprov., Oecit., agri Ligericini (Prost), Ruscinonensis. — Fl. fr. 3. p. 348. — Moris. s. 12. t. 17. f. 7. n. 12. (v. v.)

\* \* *Floribus confertis.*

4. *A. CLEMATITEA* (L. sp. 1364.) foliis petiolatis subrotundo-cordatis obtusiusculis, floribus axillaribus confertis.  $\mathcal{L}$  in lapideis sterilibus Gallie austro-orientalis, australis, medii occidentalisque. — Fl. fr. 3. p. 349. — Moris. s. 12. t. 17. f. 5. n. 5. (v. v.)

## II. ASARUM Tourn. inst. 286. L. gen. 589.

Perigonium campanulatum 3-lobum; stamina 12 ovario imposita; antheræ mediis filamentis adnatæ; stylus brevis; stigma stellatum 6-lobatum; capsula 6-locularis.

1. *A. EUROPEUM* (L. sp. 633.) foliis reniformibus obtusis subhirsutis.  $\mathcal{L}$  in umbrosis saxosis. — Fl. fr. 3. p. 349. Lam. ill. t. 394. f. 1. (v. v.)

## Ordo XC. EUPHORBIACEÆ Juss. gen. 384. Adr. Juss. gen. euph. p. 10.

Flores monoici aut dioici. Perigonium monosepalum laciniis definitis, interdum nullum, sæpissimè intus appendicibus variis squamiformibus aut glandulosis auctum. *Masc.* Stamina indefinita aut sæpius definita, centro floris nonnunquam sub pistilli rudimenta inserta. Antheræ 2-loculares, loculis extus in longum dehiscentibus interdum distinctis. *Fem.* Ovarium superum sessile aut stipitatum 2-3-∞-loculare, loculis circa placentam centalem in orbem dispositis; ovula solitaria geminatave, angulo interiori sub apice suspensa. Styli tot quot loculi nunc distincti, nunc coaliti, nunc nulli. Stigma multiplex aut pluri-lobatum. Capsula 2-3-coeca coecis distinctis elasticè bivalvibus. Semina solitaria aut geminata placentæ centrali superius annexa arillata. Embryo albumine carnoso involutus; cotyledones planæ; radícula supera. — Herbæ aut frutices, sæpissimè lactescentes, foliis plerumque stipulaceis alternis aut rarissimè oppositis, floribus axillaribus aut terminalibus plerumque bracteatis, bracteis quibusdam interdum majoribus involucriformibus.

### I. CROTON L. gen. 1083.

Monoicum. Perigonium 5-10-partitum, laciniis alternis 5 minoribus petaloideis. *Masc.* Stam. 8-15 filamentis coalita; glandulæ 5 receptaculo affixæ. *Fem.* Styli 3; stigmata 6 vel plura; capsula 3-coeca 3-sperma.

- \* 1. *C. tinctorum* (L. sp. 1425.) foliis ovato-rhomboides repandis basi integerrimis utrinque canis, pedunculis terminalibus sub 5-floris, capsulis squamato-pubescentibus pendulis. ① Nicæ, in Occitania, agro Ruscitonensi et Corsicâ. — Fl. fr. 3. p. 345. Lam. ill. t. 790. f. 4. (v. v.)

#### II. RICINUS Tourn. inst. t. 307. L. gen. 1085.

Monoicus. *Masc.* Perigonium 5-fidum; stamina plurima, filamentis coalita quasi ramosa. *Fem.* Perigonium 3-partitum; styli 3 bifurci; capsula tuberculis spinosis hirta, 3-locularis loculis 1-spermis.

1. *R. communis* (L. sp. 1430.) foliis peltatis palmatis, lobis lanceolatis serratis, caule herbaceo pruinoso, stigmatibus 3 apice 2-fidis, capsulis echinatis. ① ex Barbariâ et Oriente ortus, in hortis cultus. — Fl. fr. 3. p. 346. Lam. ill. t. 792. (v. v. c.)

#### III. BUXUS Tourn. inst. t. 345. L. gen. 1053.

Fl. monoici. Perigonium 3-4-partitum. *Masc.* Squama 2-loba; stam. 4 germinis rudimento inserta. *Fem.* Squamulae 3 minimae; styli 3; stigmata 3 obtusa; capsula 3-cornis 3-locularis 6-sperma.

1. *B. sempervirens* (L. sp. 1394.) foliis ovatis, petioliâ margine pilosiusculis, antheris ovato-sagittatis. 3 ad sepes et in sterilibus. — Fl. fr. 3. p. 345. (v. v.)

#### IV. EUPHORBIA L. gen. 243. Ræper enum. euph. germ. et pann. p. 11. (1)

Flores monoici masculi et feminei in eadem inflorescentiâ cinctâ involucri monophyllo 5-fido laciniis integris aut ciliato-fissis, exterius appendiculato glandulis 5-1 magnis cum laciniis involucri alternis truncato-peltatis. *Masc.* plures (10-36) verticillato-umbellati, articulati cum pedicellis persistentibus, bracteis ciliato-laceris stipati, abortu monandri; perigonium nullum. *Fem.* solitarii centrales, perigonio destituti; ovarium pedicellatum; stigmata 3 bifurca; capsula exserta 3-cocca, 3-sperma.

#### Sect. I. ANISOPHYLLUM Ræper ined.

Stipulatæ, involucri glandulae processibus membranaceis externe suffultæ, semina carunculâ arillari destituta.

1. *E. CHAMÆSYCE* (L. sp. 652.) ramis filiformibus foliisque anorbiculatis basi hinc leviter cordatis membranaceis, fructibus subrotundo-trigonis lævibus vel ad ovariorum dorsum punctulato-scabris, seminibus subobovato-tetragonis rugulosis albidis. ① in arenosis provinciarum australiorum.

a. *glabra* (Ræp.) foliis caulibus capsulisque glabris. — *E. Chamæsyce* Fl. fr. 3. p. 330. — Lob. ic. t. 363. f. 2. (v. v.)

β. *canescens*, foliis caulibus capsulisque pilosis. — In littoribus Galloprov. — *E. canescens* L. sp. 652. *E. Massiliensis* Fl. fr. 5. p. 357. (v. v.)

2. *E. PERLIS* (L. sp. 652.) ramis firmis, foliis oblongis basi hinc profundè cordatis integerrimis vel in baseos latere subdentatis membranaceo-carnosis glaberrimis, fructibus ovoideo-trigonis lævissimis glaberrimisque, seminibus obovato-subtetragonis lævibus albidis. ① in arenosis maritimis. — Fl. fr. 3. p. 330. — Lob. ic. t. 363. f. 1. (v. v.)

#### Sect. II. TITHYMALUS Tourn. inst. 18.

Exstipulatæ, processus membranacei nulli sub involucri glandulis, semina carunculâ arillari destituta (exceptâ *E. ptericoccâ*).

(1) Hocce genus mecum elaboravit cl. et amiciss. Ræper, auctor excellentissimæ euph. germanicæ et pannonicæ enumerationis.



§. 1. *Helioscopia* (Rœp. ined.) — *Involucris glandulæ integræ subrotundæ, cotyledones (subrotundæ) radicula latiores. — Rami floriferi sæpius trifidi.*

\* *Helioscopia veræ. — Capsulæ læves, semina reticulato-exsculpta.*

3. *E. HELIOSCOPIA* (L. sp. 658.) foliis obovato-cuneatis obtusis emarginatisve apicem versùs serratis glabris vel rariùs sparse pilosis, verticillo 5-fido rariùs 4-3-fido, ovariis dorso convexis lævibus glabris, seminibus obovatis reticulato-exsculptis opacis fuscis. ① in cultis ubique. — Fl. fr. 3. p. 331. Fl. dan. t. 725. — Matth. comm. 864. ic. (v. v.)

4. *E. PERISCOCCA* (Brot. fl. lus. 2. p. 312.) foliis spathulatis lanceolatis serrulatis obtusiusculis laxis, verticillo 5-fido, glandulis integris, ovariis glabris lævibus in dorso convexo hialatis aliis subdistantibus, seminibus ovatis subnitidis reticulatis ecarunculatis. ① in umbrosis circà Ajaccio reperiit Salzmänn et subnomine *E. stellulata* divulgavit. (v. s.)

\*\* *Helioscopia spuria. — Capsulæ verrucosæ rariùs læves, semina lævia rarissimè scabriuscula.*

5. *E. PLATYPHYLLOS* (L. sp. 660.) foliis membranaceis lanceolatis plerumque acutis serrulatis glabris vel piloso-pubescentibus, verticillo 5-fido rariùs 4-3-fido, ovariis dorso convexis læviusculis vel plus minùs verrucosis glabris, seminibus lævibus obovatis fuscis nitidis. ① 2 in siccioribus campis et secus vias. — Fl. fr. 3. p. 343.

a. *micrantha*, involucro extùs glabro, verticillo plerumque 3-fido, ovariis verrucosis glabris. — *E. micrantha* Willd. ex Rœp. (v. v.)

β. *stricta*, involucro extùs plerumque piloso, verticillis plerumque 5-rariùs 3-fidis, ovariis verrucosis glabris aut pilosis. — *E. stricta* L. E. Coderiana Fl. fr. 5. p. 365. (v. v.)

6. *E. PUBESCENS* (Vahl. symb. 2. p. 55.) foliis lanceolatis plerumque acutis serrulatis pilosis, involucro extùs piloso, verticillo plerumque 5-fido; ovariis dorso convexis verrucosis pilosis, seminibus obovatis fuscis nitidis punctis elevatis plus minùs scabris. 2 in maritimis Gallie mediterraneæ. — Fl. fr. 3. p. 344. *E. platyphyllos* γ Rœp. l. c. p. 60. An ut vult. cl. amicus præcedentis var? (v. v.)

β. *paniculata*, foliis glabriusculis. — *E. paniculata* Lois. gall. 728. DC. fl. fr. 5. p. 365. non Desf. (v. v.)

7. *E. HYPERBA* (L. sp. 662.) foliis lato-lanceolatis sessilibus obtusis vel acutis integerrimis pubescentibus vel anbiglabris, verticillo 5-fido rariùs 6-fido, ovariis dorso convexis muricato-verrucosis glabris, seminibus obovatis lævibus subnitidis nigrofuscis. 2 in montibus Corsicæ, Arvernæ, Pyrenæis. — Fl. fr. 3. p. 344. — Dill. elth. 287. t. 290. f. 374. (v. s.)

8. *E. DULCIS* (L. sp. 656. non DC. fl. fr. 3. p. 341.) foliis lato-lanceolatis subpetiolatis petiolatisque obtusis vel acutis integerrimis vel serrulatis, glabris vel piloso-pubescentibus, verticillo 5-rariùs 4-fido, ovariis in dorso convexo verrucis obsitis glabris aut pilosis, seminibus obovatis lævibus subopacis cinereo-fuscis. 2 in sylvaticis. — *E. purpurata* Thuil. fl. par. ii. 1. p. 235. Fl. fr. 3. p. 342. *E. carniolica* Jacq. Host. Fl. fr. 3. p. 342. — Scop. carn. 1. t. 21. Magnoperè variat. (v. v.)

9. *E. VERRUCOSA* (L. sp. 658.) foliis lato-lanceolatis sessilibus obtusiusculis mucronulatis vel acutis serrulatis, verticillo 5-fido rariùs 4-fido, ovariis dorso convexis undiquè verrucosis glabris, seminibus obovatis lævibus rufogriseis. 2 in humidiusculis sylvarum et varum. — Fl. fr. 3. p. 343. *E. flavicoma* Fl. fr. 5. p. 364. — Moris. s. 10. t. 3. f. 2. n. 3. (v. v.)

10. *E. SPINOSA* (L. sp. 655.) caule fruticoso, ramis senescentibus subspinosis, foliis ovato-vel lineari-lanceolatis sessilibus obtusis vel acutis integerrimis vel rariùs apicem versùs subserrulatis glabris, verticillo 5-4-3-fido, ovariis

dorso convexo verruculoso - mucronatis glabris, seminibus obovatis lævibus.  $\mathcal{U}$  in maritimis Corsicæ, Nicæ, Galloprovinciæ, agri Ruscinsonensis ex Lapeyr. — Fl. fr. 3. p. 342. — Moris. s. 10. t. 1. f. 9. n. 8. (v. v.)

11. *E. PALUSTRIS* (L. sp. 662.) foliis lanceolatis ovato - lanceolatisve sessilibus acutis vel obtusiusculis subintegerrimis vel serrulatis glabris, ramis florigeris sub verticillo in pseudo-umbellam multifidam aggregatis vel rarius verticillo 5-fido, ovariis dorso convexis undique tuberculato-verrucosis glabris, seminibus obovatis lævibus nitidis nigro-fuscis.  $\mathcal{U}$  in paludibus secus rivos. — Fl. fr. 3. p. 344. — Moris. s. 10. t. 2. f. 12. n. 1. Rami caulini inferiores steriles. (v. v.)
12. *E. PILOSA* (L. sp. 659.) foliis lato - lanceolatis sessilibus acutiusculis vel obtusis serrulatis pubescenti-pilosis, verticillo 5-fido, ovariis dorso convexis lævibus vel verrucosis plus minusve pilosis, seminibus obovatis nitidis nigro-fuscis.  $\mathcal{U}$  in Galloprovinciâ, Oecitaniâ, agro Syrtico et Burdigalensi. — Fl. fr. 3. p. 341. — Gm. sib. 2. t. 93. (v. v.)

§. 2. *Esula* (Rœp. ined.) — *Glandulæ involucri triangulares lunatæ aut lunato-bicornes, cotyledones (sublineares) latitudine radiculæ; rami florigeri, ferè semper bifidi.*

\* *Esula veræ. — Folia ramorum floriferorum libera.*

13. *E. DEMORORDIS* (L. sp. 662.) foliis lineari - lanceolatis utrinque attenuatis integerrimis obtusis sessilibus glabris, verticillo 5-10-fido, glandulis reniformibus, ovariis dorso carinato - convexis punctis elevatis scabriusculis glabris, seminibus lævibus cylindraceo-ovatis? obtusè tetragonis.  $\mathfrak{f}$  in Corsicâ, insulis Stæchadum, Nicæ. — Fl. fr. 3. p. 340. — Moris. s. 10. t. 1. f. 5 et 6. n. 11 et 12. (v. v.)
14. *E. ESULA* (L. sp. 660.) foliis lanceolatis sessilibus obtusiusculis mucronatis integerrimis vel supernè denticulato - scabris glabris, ramis florigeris sub verticillo in pseudo-umbellam multifidam congestis vel rarius verticillo 5-fido, glandulis lunatis subbicornibus, ovariis in dorso convexo punctato - scabris glabris, seminibus obovatis lævibus opacis griseo-fuscis.  $\mathcal{U}$  in Galliâ occidentali et boreali. — *E. Esula* æ et  $\gamma$ . Fl. fr. 5. p. 361. *Esula minor* Lob. ic. 357. (v. v.)
15. *E. LUCIDA* Waldest et Kit. hung. 1. t. 54.) foliis lanceolatis sessilibus obtusis mucronulatis vel acutis integerrimis glabris, ramis florigeris sub verticillo in pseudo-umbellam multifidam congestis vel rarius verticillo 5-fido, glandulis lunato-bicornibus, ovariis in dorso convexo punctato - scabris glabris, seminibus obovatis lævibus cinereo-fuscis.  $\mathcal{U}$  in demissioribus Corbatiensibus propè Villesèque ex cl. Benth. (v. s. ex Pann.)
16. *E. SALICIFOLIA* (Host. syn. p. 267.) foliis lanceolatis sessilibus acutiusculis mucronulatis integerrimis pubescentibus, ramis florigeris sub verticillo in pseudo-umbellam multifidam congestis vel rarius verticillo 5-fido, glandulis lunatis vel lunato-bicornibus, ovariis in dorso convexo punctato - scabris glabris, seminibus obovatis lævibus subopacis griseo-fuscis vel albidis.  $\mathcal{U}$  ad ripas Rhodani ex Lois, Guardinis (le Gardon), circâ Monspelium — *E. salicifolia* æ. Fl. fr. 5. p. 363. Præcedenti nimis affinis. (v. s.)
17. *E. TENUIFOLIA* (Lam. dict. 2. p. 428.) foliis linearibus acutis glaberrimis lucidis sessilibus integerrimis, verticillo terminali 5-fido, bracteis reniformi-subcordatis, glandulis lunulatis breviter bicornibus, ovariis glabris lævibus dorso convexis.  $\mathcal{U}$  in herbis Galloprovinciæ. — Fl. fr. 3. p. 332. *E. gracilis* Lois. Semina non vidi. Affinem *E. virgatæ* Wald. et Kit. ait cl. Rœp. (v. s.)
18. *E. CYPRASSIAS* (L. sp. 661.) foliis linearibus sessilibus obtusis vel acutiusculis integerrimis glabris, ramis florigeris sub verticillo in pseudo-umbellam multifidam aggregatis vel rarius verticillo 5-fido, glandulis lunatis subbicornibus, ovariis in dorso convexo punctato - scabris glabris, seminibus obovatis lævibus opacis griseo-fuscis vel albidis.  $\mathcal{U}$  in sterilibus frequens. — Fl. fr. 3. p. 337. Jacq. anat. 5. t. 435. *E. Esula* Fl. fr. 3. p. 337. non L. (v. v.)

19. *E. GERARDIANA* (Jacq. aust. 5. t. 436.) foliis lanceolatis sessilibus acutis vel obtusiusculis mucronatis integerrimis glabris, ramis florigeris sub verticillo in pseudo-umbellam multifidam aggregatis vel rarius verticillo 5-fido, glandulis obtusè triangularibus, ovariis dorso convexis punctis elevatis minutissimis obsitis glabris, seminibus obovato-cylindraceis lævibus opacis albidis.  $\frac{1}{2}$  in sterilibus et glareosis. — Fl. fr. 3. p. 337. *E. cajogala* Ehrh. (v. v.)  
*ℓ. minor*, 2-3-pollicaris, foliis oblongis obovatisve. — in monte Ventoso. — *E. saxatilis* Lois. not. p. 77. DC. fl. fr. 5. p. 363. non Bieb. nec Jacq. (v. s.)
20. *E. NICKENSI* (All. ped. 1. p. 285. t. 69. f. 1.) foliis lanceolatis vel ovato-lanceolatis sessilibus acutis vel obtusis mucronatis margine scabris glabris, ramis florigeris sub verticillo in pseudo-umbellam multifidam aggregatis vel rarius verticillo 5-fido, glandulis lunatis vel lunato-bicornibus, ovariis dorso convexis punctis elevatis minutissimis obsitis vel lævibus glabris vel lanatis pilosisve, seminibus obovato-cylindraceis lævibus vel rarissime subexsculptis opacis griseis.  $\frac{1}{2}$  in sterilibus Gallie australis, propè Parisios ex Thail. — Fl. fr. 3. p. 338. *E. oleæfolia* Gou. — Barr. ic. 823. (v. v.)
21. *E. PITREUSA* (L. sp. 656.) suffruticosa, foliis rigidis lineari-lanceolatis confertis acuminatis glaucis margine revolutis scabris inferioribus reflexis, superioribus latioribus, verticillo 4-8-fido, glandulis reniformibus, ovariis dorso convexo lævibus glabris, seminibus ovatis lævibus.  $\frac{1}{2}$   $\delta$  in arenosis maritimis Galloprovinciæ, insulæ S. Lucie propè Narbonam ex cl. Benth. — Fl. fr. 3. p. 334. — Matth. comm. 867. ic. (v. v.)
22. *E. PARALIAS* (L. sp. 657.) foliis crasso-coriaceis lanceolatis sessilibus acutis vel obtusiusculis integerrimis glabris, verticillo 5-4-3-fido, glandulis lunatis auberosis, ovariis in dorso convexo sulco longitudinali exaratis punctis elevatis minutissimis obsitis rugulosis glabris, seminibus obovato-globosis lævibus vel exsculptis cinereis.  $\frac{1}{2}$  in arenosis maritimis. — Fl. fr. 3. p. 335. Jacq. vind. t. 188. — Moris. s. 10. t. 1. f. 8. n. 24. (v. v.)
23. *E. MYRSINITES* (L. sp. 661.) foliis ovato-lanceolatis spathulatisve glaucis carnosius glaberrimis mucronatis margine scabris confertis, verticillo 5-fido, glandulis 2-cornibus, cornubus compressis dilatatis, ovariis lævibus glabris carinato-convexis, seminibus cylindraceo-ovatis obsolete tetragonis tuberculato-rugosis opacis griseis.  $\frac{1}{2}$  in Corsicâ, circâ Nicæam ex All. — Fl. fr. 3. p. 338. — Lob. ic. t. 355. f. 1. (v. s.)
24. *E. SERPATA* (L. sp. 658.) foliis lineari-lanceolatis lanceolatisve sessilibus acutis mucronulatisve glabris superioribus latioribus bracteis dentato-serratis, verticillo 3-5-fido, glandulis suborbiculatis vix lunulatis, ovariis dorso convexis læviusculis glabris, seminibus obovato-rotundis lævibus opacis carunculâ magnâ antheræformi instructis.  $\frac{1}{2}$  secus arva et vias in prov. australioribus. — Fl. fr. 3. p. 336. Jacq. ic. rar. 3. p. 384. — Moris. s. 10. t. 1. f. 3. n. 6. (v. v.)
25. *E. RUMBELLATA* (Poir. voy. barb. 2. p. 174. ic.) foliis lineari-lanceolatis mucronulatis glabris apicem versus scabris patulis, superioribus latioribus, bracteis cordato-reniformibus mucronatis, verticillo 5-9-fido, glandulis lunulatis bicornibus cornubus clavatis, ovariis in dorso convexo punctis elevatis scabriusculis glabris, seminibus ovoideis tuberculato-rugosis.  $\frac{1}{2}$  in agro Roscinonensi ex Benth. Occitaniâ, circâ Avenionem, Telonem. — Fl. fr. 5. p. 360. — Plerumque sub verticillo terminali secundus tertiusque invenitur. (v. s.)
26. *E. SEGETALIS* (L. sp. 657.) foliis rigidulis lineari-lanceolatis sessilibus sen in petiolum attenuatis mucronatis integerrimis glabris, verticillo 5-fido, glandulis lunatis longissimè bicornibus, ovariis in dorso convexo punctis elevatis scabris glabris, seminibus obovatis reticulato-exsculptis albidis. (1) inter segetes prov. australiorum occidentaliumque et propè Parisios ex Mer. — Fl. fr. 3. p. 335. exel. var. Jacq. aust. t. 450. *E. pinifolia* Lam. ex cl. Reper. Fl. fr. 3. p. 336. — Moris. s. 10. t. 2. f. 10. n. 3. *E. Artandiana* Fl. fr. 5. p. 361. est *E. segetalis* caulocarpica. (v. v.)  
*ℓ. longibracteata*, foliis ramorum florigerorum lineari-lanceolatis. — Propè Montpellierum. — Fl. fr. 5. p. 359. (v. s.)

- γ. maritima*, bracteis latis subcordatis mucronulatis. — In maritimis passim. — E. Portlandica Fl. fr. 5. p. 359 an L. sp. 656? (v. s.)
27. E. PROVINCIALIS (Wild. sp. 2. p. 914.) foliis lanceolatis spathulati-ve obtusis mucronulatis minutissimè serrulatis sessilibus glabris, bracteis cordato-reniformibus, verticillis 5-4-3-fidis, glandulis lunatis longissimè bicornibus, ovariis dorso carinato-convexis glaberrimis lævibus, seminibus subovatis lævibus opacis griseo-albis. ①  $\frac{1}{2}$  in maritimis Corsicæ, Galloprovinciæ, et agri Ruscignoniensis ex cl. Benth. — E. Terracina Fl. fr. 3. p. 333 an L? E. affinis Fl. fr. 5. p. 363. — Barr. ic. 831. (v. v.)
- β. seticornis*, foliis linearilanceolatis acutis, bracteis cordatis ovatis longè acuminatis. — In collibus propè Digne. — E. seticornis Poir. voy., 2. p. 178. Fl. fr. 5. p. 359. (v. v. DC.)
28. E. EXIGUA (L. sp. 654.) foliis rigidulis linearibus vel linearilanceolatis sessilibus acutiusculis submucronatis integerrimis glabris, verticillo 3-4-5-fido, glandulis longissimè bicornibus, ovariis in dorso convexo punctis elevatis scabris glabris, seminibus obovato-cylindraceis subtetragonis tuberculato-reticulatis opacis albidis vel cinereo-fuscis. ① in arvis. — Fl. fr. 3. p. 332. — Lob. ic. t. 357. f. 2. (v. v.)
- β. retusa*, foliis retusis. — E. retusa Cav. ic. 1. t. 34. f. 3. Fl. fr. 5. p. 358. E. rubra Fl. fr. 5. p. 359. non Cav. — Magn. bot. p. 259. ic. 258. (v. v.)
29. E. FALCATA (L. sp. 654.) foliis rigidis lanceolatis sessilibus acutis mucronati-ve margine denticulato-scabris glabris; verticillo 3-4-5-fido, glandulis lunatis vel lunato-bicornibus, ovariis dorso convexis lævibus glabris, seminibus obovato-cylindraceis subtetragonis, lateribus transversè rugosis, opacis albidis cinereis vel fuscis. ① in cultis passim. — Fl. fr. 3. p. 331. Jacq. aust. t. 121. (v. v.)
- β. obicura*, foliis bracteisque acuminatis. — In arvis Galloprovinciæ. — E. obscura Lois. not. 76. t. 5. f. 2. Fl. fr. 5. p. 358. (v. v.)
30. E. PEPLUS (L. sp. 653.) foliis suborbiculatis in petiolum attenuatis obtusissimis integerrimis glabris, verticillo 3-fido rarissimè 5-fido, glandulis lunatis longissimè bicornibus, ovariis in dorso bicarinato alato ruguloso-scabris glabris, seminibus obovato-cylindraceis obtusè hexagonis, lateribus quatuor punctorum serie exsculptis, duobus sulco longitudinali exaratis, opacis griseo-albidis. ① in cultis, vineis, sepibus. — Fl. fr. 3. p. 331. E. peploides Gou. fl. 174. Fl. fr. 5. p. 358. E. rotundifolia Lois. not. t. 5. f. 1. (v. v.)
31. E. LATYRIS (L. sp. 655.) foliis linearibus sessilibus acutiusculis vel obtusis mucronatis integerrimis glabris, verticillo 4-fido rariùs 2-fido rarissimè 5-fido, glandulis lunatis bicornibus cornibus dilatatis obtusis, ovariis dorso convexis sulco longitudinali profundo exaratis lævibus glabris, seminibus obovatis basi truncatis scabris opacis fuscis. ② in cultis et secùs vias, vix indigena. — Fl. fr. 3. p. 333. (v. v.)
- \* \* *Esulæ spuria*. — Folia ramorum floriferorum connata.
32. E. SYLVATICA (L. sp. 663.) foliis lato-lanceolatis in petiolum attenuatis acutis vel obtusis mucronulatis integerrimis pubescentibus, verticillo 5-fido vel ramis florigeris subverticillo in pseudo-umbellam sex-octofidam aggregatis, glandulis lunatis bicornibus, ovariis dorso convexis punctis minutissimis elevatis obtusis glabris, seminibus subrotundo-ovatis lævibus opacis fuscis.  $\frac{1}{2}$   $\frac{3}{4}$  in sylvaticis, sepibus. — Fl. fr. 3. p. 339. Jacq. aust. t. 275. E. amygdaloides L. — Moris. s. 10. t. 1. f. 2. n. 3. (v. v.)
33. E. CHARACIAS (L. sp. 662.) foliis linearilanceolatis in petiolum attenuatis acutis vel obtusiusculis mucronulatis integerrimis pubescentibus vel glabris, ramis florigeris sub verticillo in pseudo-umbellam multifidam aggregatis, glandulis lunatis, ovariis dorso convexis punctis minutissimis elevatis obtusis, seminibus oblongo-ovatis lævibus opacis nigricantibus.  $\frac{3}{4}$  in montosis lapideosis provinc. Mediterranearum. — Fl. fr. 3. p. 340. — Jacq. ic. rar. 1. t. 89. (v. v.)

V. MERCURIALIS *Tourn. inst. t. 3o8. L. gen. 1125.*

Dioica aut rariùs monoica. Perigonium 3-partitum. *Masc.* Stam. 9-12. *Fem.* Ovarium 2-gibbosum 2-sulcatum, filamentis 2 sterilibus brevibus ex quoque sulco prodeuntibus cinctum; styli 2 bifurcati, capsula 2-coeca 2-sperma.

1. *M. TOMENTOSA* (L. sp. 1465.) caule suffruticoso, foliis ovatis tomentosis, capsulis hirsutis.  $\S$  in Occitaniâ, agro Ruscinonensi, Corbariensibus. — Fl. fr. 8. p. 329. — Clus. hist. 2. p. 48. f. 1. 2. (v. v.)
  2. *M. PERENNIS* (L. sp. 1465.) caule simplicissimo, foliis ovatis lanceolatis lanceolatisve scabris crenato-serratis pubescentibus, floribus longè pedunculatis, capsulis pubescentibus.  $\mathcal{L}$  in sylvis. — Fl. fr. 3. p. 328. Fl. dan. 400. Moris. s. 5. t. 84. f. 3 et 4. (v. v.)
  3. *M. ELLIPTICA* (Lam. dict. vol. 4. p. 119.) dioica, caule suffruticoso brachiato, foliis lanceolatis serratis glaberrimis, floribus masculis verticillato-glomeratis longè pedunculatis, femineis verticillatis axillaribus breviter pedunculatis, capsulis glabris.  $\S$  in montosis Corsicæ. — Vent. cels. t. 22. (v. s.)
  4. *M. ANNUA* (L. sp. 1465.) caule brachiato, foliis lanceolatis serratis glabris subpetiolatis, floribus masculis spicato-glomeratis pedunculatis, femineis verticillatis axillaribus, capsulis echinato-pubescentibus. ① in cultis frequens. — Fl. fr. 3. p. 328. Lam. ill. t. 820. — Moris. s. 5. t. 34. f. 1 et 2. (v. v.)
- $\rho$ . *ambigua*, floribus verticillatis femineis masculisque intermixtis. — In Corsicâ, Galloprovinciâ. — *M. ambigua* Lin. f. dec. 1. t. 8. Fl. fr. 5. p. 357. (v. s.)

---

Ordo XCI. URTICEÆ *Juss. gen. 400. excl. Piperis. DC. fl. fr. 3. p. 517.*

Flores parvuli virescentes monoïci aut dioici, solitarii amentacei aut involucri monophyllo cincti. Perigonium monosepalum 3-5-lobum persistens. *Masc.* Stamina definita, basi perigonii inserta. *Fem.* Ovarium simplex liberum; styli 2 aut 1 bifurcatus. Fructus Achenium aut Drupa perigonio persistente tecta, solitaria aut inserta receptaculo carnosio dilatato. Semina pendula albuminosa aut exalbuminosa. Embryo rectus curvatus aut spiralis. Radicula sæpiùs supera. — Herbæ aut Arbores foliis sæpiùs hispidis et spathulatis. Fl. capitati aut racemosi.

Trib. I. URTICEÆ *DC. fl. fr. l. c.*

Flores solitarii, amentacei vel spicati, fructus nunquam carnosii. Albumen nullum. Embryo sæpiùs rectus.

I. CANNABIS *Tourn. inst. t. 3o8. L. gen. 1115.*

Dioica. *Masc.* Perigonium 5-partitum; stamina 5. *Fem.* Perigonium oblongum latere fissum; ovarium 1; styli 2; capsula crustacea 2-valvis, valvis subglobosis perigonio obtectis, embryo incurvus.

1. *C. SATIVA* (L. sp. 1457.) (1) ex Indiâ orta, circa pagos culta. — Fl. fr. 3. p. 325. — Lob. ic. t. 526. f. 1-2. Folia petiolata digitata. (v. v. c.)

II. *HELXINE* Req. in ann. sc. nat. 5. p. 384.

*Monoica. Masc.* Flores solitarii perigonio campanulato 4-partito. *Fem.* Perigonium persistens ventricosum suprâ 3-partitum; semen unicum. (Req.)

1. *H. SOLZINOLII* (Req. l. c.) — in muris umbrosâ Corsicæ. — Cespitosa, caulibus radicantibus, folia obliqua ambrotunda glabresculâ subsessilia. (v. s. s. fl.)

III. *PARIETARIA* Tourn. inqt. t. 289. L. gen. 1152.

Fl. polygami, involucri multipartito cincti. *Herm.* Perigonium 4-partitum; stam. 4, filamentis primo incurvis elasticè dehiscentibus; ovarium 1; stylus 1; stigma 1; semen perigonio elongatâ et suprâ connivente tectum. *Fem.* Stamina 0; cætera hermaphroditis similia.

1. *P. LUSTRANICA* (L. sp. 1492.) caule filiformi procumbente ramoso pubescente, foliis ovatis ambrotundis obtusis pilosis petiolo longioribus, floribus axillaribus subsessilibus subternatis aggregatis. (1) in humidis montosis Corsicæ, ad muros propè Telonem et in agro Ruscinonensi circa Banyuls ex Lapeyr. — Fl. fr. 5. p. 366. Viv. cors. app. f. 1. — Bocc. rar. t. 24. f. B. (v. s.)
2. *P. OFFICINALIS* (L. sp. 1492.) caule ascendente ramoso, foliis longè petiolatis lanceolato-ovatis suprâ sublucidis subtus hirsutis nervosis. 2/ ad muros et sepes. — Fl. fr. 3. p. 324. Lam. ill. t. 853. f. 1. (v. v.)
3. *P. JUDEICA* (L. sp. 1492.) caule prostrato ramoso, foliis longè petiolatis ovatis pubescentibus, floribus maribus elongato-tubulosis. 2/ circa Vapincum ex Vill., in provinciis australibus, in muris Chantenai propè Nannetes, circa Aureliam, Aroneil propè Parisios ex Lois. — Fl. fr. 3. p. 824. Lam. ill. t. 853. f. 2. — An prioria varietas aut potius deformatio? (v. v. DC.)

IV. *URTICA* Tourn. inst. t. 308. L. gen. 1054.

*Monoica* rarius dioica. *Masc.* racemosi; perigonium 4-partitum; stamina 4; filamenta antè florescentiam curvata. *Fem.* capitato-racemosi; perigonium 2-valve; ovarium 1; stigma 1; semen 1 perigonio cinctum.

1. *U. MELLIFERA* (L. sp. 1395.) foliis oppositis ovatis dentatis dentibus obtusis, apiculis fructiferis globosis pedunculatis. (1) in arvis Gallicæ australis, circa Aginnum ex St.-Am., ad ostium Ligeris, propè Parisios ex Mérat, Falaise (cl. de Bréb.). — Fl. fr. 3. p. 322. Lam. ill. t. 761. f. 2. eng. bot. t. 148. — Moria. t. 26. f. 8. n. 5. (v. v.)
2. *U. ANTHRASTIS* (Poir. enc. 4. p. 638.) monoica, foliis oppositis ovatis acutis grossè dentatis longè petiolatis, spicis pedunculatis simplicibus, masculularum geminarum petiolo longiorum rachi utrinque alata. 2/ in cultis Telonis, Arelatis, Perpiniani, Armoracis infer. — Fl. fr. 5. p. 355. (v. s.)
3. *U. MURINA* (DC. fl. fr. 5. p. 355.) dioica, caulibus foliisque hispidis, foliis petiolatis grossè dentatis acuminatis ovato-orbicularibus, spicis axillaribus geminatis, masculis ramosis. 2/ in Corsicâ, agro Ruscinonensi propè Prades (Corder). (v. s.)
4. *U. DIOICA* (L. sp. 1396.) dioica, foliis oppositis cordatis ovato-lanceolatis grossè serratis, spicis paniculatis glomeratis geminatis petiolo longioribus. 2/ ad sepes et in hortis. — Fl. fr. 3. p. 322. Lam. ill. t. 761. f. 1. fl. dan. t. 746. (v. v.)
5. *U. VAGANS* (L. sp. 1396.) monoica, foliis oppositis ellipticis sub 5-nerviis argutè serratis subincisis, spicis glomeratis geminatis. (1) in cultis, ad pagos

areas frequens. — Fl. fr. 3. p. 323. Fl. dan. t. 739. — Moris. s. 11. t. 25. f. 6. n. 4. (v. v.)

V. HUMULUS *L. gen.* 1116.

Dioica. *Masc.* Perigonium 5-partitum; stamina 5. *Fem.* Strobili constantes squamis magnis persistentibus concava axillâ unifloris; ovarium 1; stylus 2; semen 1 arillatum; embryo spiraliter tortus.

1. *H. LUPULUS* (*L. sp.* 1457.) 2½ ad sepes, sæpè colitur. — Fl. fr. 3. p. 322. Lam. ill. t. 815. Caules scandentes, folia petiolata cordata serrata simplicia aut 3-loba. (v. v.)

Trib. II. *ARTOCARPEÆ DC. fl. fr.* 3. p. 318.

Flores receptaculo communi insiti. Fructus carnosus. Semina albuminosa. Embryo curvatus.

VI. MORUS *Lam. DC. fl. fr.* 3. p. 320. — *Mori sp. L.*

Monoica. Amenta unisexualia; perigonium 4-lobum, lobis concavis. *Masc.* Stamina 4 perigonii laciniis alterna. *Fem.* Ovarium liberum; stigma 2; semina 1-2, perigonio pulposo oblecta.

1. *M. ALBA* (*L. sp.* 1398.) foliis profunde cordatis basi inæqualibus ovatis lobatisve serratis lævibus tenuioribus. 5 ex Oriente orta, nunc subsponsanea, secus rivos Gallie australis. — Fl. fr. 3. p. 321. Gærtn. fruct. 2. t. 126. f. 6. Fructus albi aut rubelli. (v. v. c.)
2. *M. NIGRA* (*L. sp.* 1398.) foliis cordatis ovatis lobatisve inæqualiter dentatis scabris crassiusculis. 5 ex Persia aut China orta, in hortis culta. — Fl. fr. 3. p. 320. — Duham. arb. fruit. 2. p. 161. t. 1. Fructus atropurpurei. (v. v. c.)

VII. FICUS *Tourn. inst. t.* 410. *L. gen.* 1168.

Monoica. Flores numerosi pedicellati, receptaculo carnoso apice umbilicato intus cavo inclusi; perigonium 3-5-lobum, lobis acuminatis. *Masc.* umbilico vicini; stamina 3-5. *Fem.* Ovarium liberum (Deaf.), semi-adnatum (Gærtn.); stylus 1; stigma 2; drupa sive utriculus 1-spermus, in receptaculi pulpâ nidulans; nuclei cortex fragilis, crustaceus.

1. *F. CARICA* (*L. sp.* 1513.) foliis cordatis palmatis supra scabris subtus pubescentibus. 5 3 in siccis et lapideis Gallie australis. — Fl. fr. 3. p. 318. (v. v.)

Ordo XCII. JUGLANDEÆ DC. *theor. ed.* 1. p. 215. *Kunth. tereb. gen. p.* 11.

Fl. monoici. *Masc.* amentacei. Perigonium squamosum lateraliter plus minus profundè 2-vel 6-lobum. Stamina hypogyna numero indefinita; filamenta brevissima libera; antheræ 2-loculares basi affixæ. *Fem.* Perigonium duplex aut simplex cum ovario adhærens, externum 4-divisum, internum quandò adest 4-sepalum. Ovarium 1-loculare, ovulo erecto. Styli nunc 1-2 brevissimi et stigmata 2 crassa, aut nulli et stigma magnum discoideum 4-lobum. Drupa carnosa nucem 2-4-valvim fovens. Semen cerebriformè plus minus 4-lobum, integumento membranaceo instructum. Embryo magnus exalbuminosus. Cotyledones carnosæ 2-lobæ. Radicula supera. — Arbores, folia alterna imparipinnata; stipulæ nullæ. Fl. masculi et feminei in eadem vel in distinctis gemmis; feminei terminales 1-3 vel plures laxè spicati; masculi remoti arcuè spicati.

I. JUGLANS Nutt. *gen. amer.* 2. p. 220. — Juglandis sp. L.

~ *Masc.* Amentum imbricatum; perigonium simplex singulæ bractæ internè adnatum 5-6-partitum, stamina numerosa (14-36); antheræ crassæ. *Fem.* Perigonium duplex; styli 2 brevissimi; stigmata 2 magna; drupa fæta nuce osseâ bivalvi extus rugosâ et irregulariter profundè sulcatâ.

1. J. REGIA (L. sp. 1415.) foliolis subnovenis ovalibus glabris subserratis subæqualibus, fructibus glohosis. Ex Persia? orta, nunc in totâ Galliâ frequens. — Fl. fr. 4. p. 618. — Blackw. t. 247. (v. v. c.)

Ordo XCIII. AMENTACEÆ Juss. *gen.* 407.

Flores dioici monoici aut rariùs hermaphroditi. *Masc.* capitati aut in amentum dispositi, squamâ aut perigonio squamifero instructi. Stamina squamæ inserta ferè nunquàm monadelphæ; antheræ biloculares. *Fem.* solitarii, fasciculati aut julacei, squamâ aut perigonio instructi. Ovarium liberum simplex aut rariùs multiplex; stigmata plurima. Pericarpia tot quot ovaria ossea aut membranacea. Albumen nullum aut tenue. Embryo rectus aut curvatus planus. Radicula sæpiùs supera. — Arbores aut frutices, foliis alternis caducis, junioribus basi stipulis binis stipatis.



Trib. I. *CELTIDEÆ* Duby mss.

Fl. hermaphroditi polygami vel monoici. Perigonium liberum campanulatum 5-6-partitum. Stam. 5-6 imo perigonio inserta ejusque lobis opposita libera. Ovarium unicum simplex. Stigmata 2 ex ovarii apice distincta. Drupa globosa. Putamen osseum 1-loculare. Semen unicum pendulum. Albumen membranaceo-carnosum tenue; embryo curvatus; cotyledones foliaceæ contortuplicatæ tenuissimæ; radícula ad hylum versa. — Arbores, folia alterna petiolata simplicia. Flores axillares solitarii racemosi vel rariùs paniculati.

I. *CELTIS* Tourn. inst. t. 383. L. gen. n. 1143.

Character isdem ac ordinis.

1. *C. AUSTRALIS* (L. sp. 1478.) foliis ovato-lanceolatis subhirsutis. 5 in Occitania, Galloprovincia, circa Aginnum et Burdigalam. — Fl. fr. 3. p. 315. Lam. ill. t. 844. f. 1. — Duham. arb. ed. secunda. 2. p. 34. t. 8. (v. v.)

Trib. II. *BETULINEÆ* Rich. in Kunth nov. gen. am. 2. p. 21. DC. mss.

Flores hermaphroditi polygami aut monoici, dispositione variâ. Perigonium liberum campanulatum 4-5-lobum. Stam. 4-12, sæpiùs tot quot perigonii lobi et tunc iis opposita aut eorum numero dupla triplave, perigonii basi inserta libera. Ovarium 1 simplex. Stigmata 2 ex ovarii apice distincta. Pericarpium indehiscens 2-loculare membranaceum aut subcoriaceum compressum, interdum lateraliter in alam expansum. Semina in quoque loculo solitaria pendula exalbuminosa. Radícula ad hylum versa. Cotyledones planæ foliaceæ. — Arbores et frutices. Folia alterna petiolata simplicia penninervia.

Subtr. I. *ULMEÆ* Mirb. elem. 2. p. 905.

Flores in capitula parva aggregati laxiusculi pedicellati hermaphroditi aut abortu polygami.

II. *ULMUS* Tourn. inst. t. 372. L. gen. 316.

Hermaphrodita. Perigonium campanulatum 4-5-dentatum coloratum persistens; stamina 3-6; ovarium compressum; stigmata 2; samara suborbiculata latè marginata membranacea 1-sperma.

1. *U. CAMPESTRIS* (L. sp. 327.) foliis duplicato-serratis basi inæqualibus, floribus subsessilibus conglomeratis. 5 5 ad vias, in sylvis montosis. — Fl. fr. 3. p. 315. Lam. ill. t. 185. (v. v.)

a. *microphylla*, ramis erectis, foliis scabris, samaris obovatis glabris. — *U. campestris* eng. bot. t. 1886. Host. fl. aust. 1. p. 330. (v. v.)

β. *suberosa*, cortice suberoso, ramis patulis, foliis scabris, inferioribus ovatis, superioribus oblongis, samaris suborbiculatis glabris. — *U. suberosa* Willd. eng. bot. t. 2161. Host. 1. c. 328. (v. v.)

γ. *major*, ramis patulis, foliis oblongis scaberrimis basi cordatis, samaris

subrotundis glabris. — U. major eng. bot. t. 2542. Host. l. c. An hæc varietates reverà ut sp. distinctæ habendæ? (v. v.)

2. U. *xyvusa* (Wild. sp. 1. p. 1325.) foliis duplicato-serratis basi inæqualibus, floribus longè pedunculatis effatis. § Lutetiæ in horto Armamentarii, in agro Bellovacensi ex Graves, ad mœnia Suessionum cultus, in agro Andegavensi ex Bast, in sylvis propè Argentoratam ex Nestler. — Fl. fr. 3. p. 316. U. octandra Schk. bot. handb. 178. t. 57. (v. s.)

Subtr. II. *BETULÆ DC. mos.*

Fl. amentacei, squamis ad axillam flores 1-3 sessiles gerentibus, monoici in amentis distinctis.

III. *BETULA Tourn. inst. 350. Hall. herb. p. 298. — Betula sp. L.*

Monoica. Amenta elongato-cylindrica. *Masc.* Squamæ ternatæ, media staminifera. *Fem.* Squamæ 3-lobatæ membranacæ deciduæ; styli 2; ovarium compressum 2-loculare, loculo uno abortivo; nuculæ compressæ margine membranacæ uniloculares.

1. B. *WARA* (L. lapp. t. 6. f. 4.) foliis orbiculatis crenatis glabris subtus reticulato-venosis, strobilis ovatis subsessilibus, squamis cuneatis, nuculis ovatis subcompressis. § in humidis Jurassi. — Fl. fr. 3. p. 302. — Fl. dan. t. 91. (v. v.)
2. B. *PUBESCENS* (Ehr. arb. n. 67.) foliis deltoidibus acutis subcordatis duplicato-serratis subtus ramulisque pubescentibus, strobilorum squamis lobis lateralibus rotundatis. § in turfosis Jurassi, circa Valletriam, Marly propè Parisios ex cl. Ram. et Mérat, Beauvais ex Graves. — Fl. fr. 3. p. 302. (v. s.)
3. B. *ALBA* (L. sp. 1303.) foliis deltoidibus acutis duplicato-serratis glabris; strobilorum cylindricorum pedunculorum squamis lobis lateralibus rotundatis, petiolis glabris pedunculo longioribus. § in sterilibus arenosis frigidioribus. — Fl. fr. 3. p. 301. — Duh. arb. 1. p. 100. t. 39. (v. v.)

IV. *ALNUS Tourn. inst. t. 359. Hall. herb. p. 300. — Betula sp. L.*

Monoica. *Masc.* Amenta elongato-cylindrica; squamæ pedicellatæ cordiformes, subtus squamulis ternis instructæ basi floriferæ. *Fem.* Amenta ovato-globosa pedicellis ramosis; squamæ 2-floræ coriaceæ persistentes; ovarium compressum; stigmata 2; nuculæ compressæ ovatæ nudæ biloculares dispermæ.

\* *Stamina* 12.

1. A. *SUAVOLENS* (Req. in ann. sc. nat. 5. p. 381.) foliis subrotundis argutè duplicato-serratis obtusis valdè resinosis glabris margine crispis, strobilis ovatis minoribus, squamis cuneatis. § in montibus Corsicæ. (v. s.)
2. A. *VIRIDIS* (DC. fl. fr. 3. p. 304.) foliis ovatis argutè duplicato-serratis glabris subtus in nervis pubescentibus acutis, strobilis ovatis minoribus, squamis cuneatis, seminibus lanceolatis compressis. § in montibus editioribus Delphinatæ. — Betula ovata Schrank. DC. syn. gall. p. 180. (v. s.)

\*\* *Stamina* 4 ex urceolo 4-lobæ.

3. A. *INCANA* (DC. fl. fr. 3. p. 304.) foliis ovatis acutis subtus tomentoso-pubescentibus duplicato-serratis, stipulis lanceolatis, amentis masculis cylindricis elongatis, strobilis ovato-globosis. § 5 in montosis. — Betula incana Reich. syst. 4. p. 127. (v. v.)
4. A. *GLUTINOSA* (Gært. fruct. 2. t. 90. f. 2.) foliis subrotundo-cuneatis obtusis subretusis glutinosis duplicato-serratis, axillis venarum subtus villosis, amentis masculis cylindricis, strobilis subglobosis. § 5 secus rivos et in humidis. — Fl. fr. 3. p. 303. Betula Alnus L. (v. v.)

5. *A. cordata*, foliis subrotundis obtusissimis cordatis æqualiter serratis, axillis venarum subtus villosis, strobilis grossis subglobosis valde glutinosis, squamis dense imbricatis, nucibus lanceolato-ellipticis. 5 in Corsicâ. — *Betula cordata* Lois. not. 139. (v. s.)
6. *A. elliptica* (Req. in ann. sc. nat. 5. p. 381.) foliis ellipticis obtusis lucidis glutinosis regulariter serratis, axillis venarum subtus villosis, fructibus paucis majoribus. 5 in Corsicâ ad ripas Salenarzæ (Req.) An satis à priori distincta? (v. s. sine fr.)

Trib. III. *SALICINÆ* Rich. l. c. DC. *ms.* — *Salicin. sect.* 1. *Mirb. elem.* 2. p. 905.

Flores dioici ad ejusque squamæ axillam solitarii, amentacei. *Masc.* in amentum cylindricum dispositi. Perigonium parvum glanduliforme. Stam. 2-30 glandulæ subadnata sæpius libera rarissimè monadelpha. *Fem.* in strobilum ovatum seu cylindricum dense dispositi. Perigonium liberum simplex sæpè persistens aut minimum. Ovarium 1-loculare. Stylus simplex. Stigm. 2 sæpè bifida. Caps. 1-locularis 2-valvis, loculis ∞-spermis. Semina minima pendula exalbuminosa crinita undique aut apice comosa. Embryo rectus. Radicula ad umbilicum tendens. Cotyledones planiusculæ foliaceæ. — Arbores aut frutices, foliis alternis simplicibus, stipulis nunc minimis subnullis nunc foliaceis. Gemmæ squamis foliaceis constantes.

V. *SALIX* Tourn. *inst.* t. 364. *L. gen.* 1098.

Flores dioici aut rarissimè monoici julecei, squamæ imbricatæ; glandula genitalia cingens. *Masc.* Stamina 2-5 sæpius 2 interdum in unum coalita et tunc anthera 4-locularis. *Fem.* Semina comosa; radícula infera.

Sect. I. *CINERELLA* Ser. *sal. rev. ined.* (1)

Amenta præcocia primo ovato-cylindrica, fem. demum elongata; ovaria tomentosa; stylus brevis. *Masc.* diandria; stam. libera vel monadelpha. — Folia ovata obovata vel lanceolata plerumque tomentosa.

1. *S. CAPREA* (L. sp. 1448.) arborea aut subarborea, foliis ovatis rugosis subtus tomentosis undatis supernè denticulatis acutis aut mucronatis petiolatis, amentis ovato-cylindricis, femineis sessilibus pilosis, ovarii tomentosis basi ventricosis longè conicis pedicellatis. 3 in sylvaticis et sepibus. — Fl. fr. 3. p. 290. L. fl. lapp. t. 8. f. 3. Hoffm. sal. t. 3. f. 1 et 2. t. 5. f. 4. t. 21. f. 1. a. b. c. d. *S. tomentosa* Ser. *ess.* 14. (v. v.)
2. *S. CINEREA* (L. sp. 1449.) subarborea, foliis ovato-oblongis acuminatis acutisve subrenatis aut subintegris reticulato-nervosis subtus tomentosis petiolatis, stipulis rotundatis subserratis, amentis femineis sessilibus elongatis pilosis, ovarii tomentosis longè conicis pedicellatis. 3 in pratis et sepibus. — *S. acuminata* Mill. Fl. fr. 3. p. 290 et 5. p. 342. Hoffm. sal. t. 6. f. 1. 2. t. 22. f. 2. *S. rufinervis* Fl. fr. 5. p. 341. *S. aurigerana* et incerta Lapeyr. (v. v.)
3. *S. CINERASCENS* (Willd. sp. 4. p. 706.) arborea aut subarborea, ramulis subglabris, foliis oblongo-lanceolatis dentatis integrisve acutis petiolatis anprè viri-

---

(1) In sal. helv. collect. exsicc. 1824. cum notis è quibus plures hic laudatos et imprimis Smithii et Hoffmanni synonymos desumpsi.

ribus glabris subtus subœneis, stipulis magnis dimidiato-cordatis serratis, stylo brevissimo, amentis femineis subcœtaneis sessilibus demum elongatis pilosis, ovariis oblongo-conicis tomentosis pedicellatis.  $\zeta$   $\zeta$  secus rivos et in tur-  
fosis Jurassi, Alpium, Pyrenæorum. — *S. grandifolia* Ser. ess. p. 20. Fl.  
fr. 3. p. 343. (v. v.)

4. *S. AURITA* (L. sp. 1446.) frutex, foliis obovatis rugosis subtus villosis reti-  
culato-nervosis subobtusis aut breviter mucronatis petiolatis, stipulis dimi-  
diato-cordatis, staminum filamentis basi coalitis, amentis femineis sessilibus  
pilosis, ovariis conico-oblongis tomentosis pedicellatis.  $\zeta$  ad sepes, locis  
umbrosis. — Fl. fr. 3. p. 291 et 5. p. 342. eng. bot. 1487. — L. lapp. t. 8. f. 9.  
*S. rugosa* Ser. ess. p. 18. (v. v.)

5. *S. REPENS* (L. sp. 1447.) arbuscula, foliis ovatis aut ovato-oblongis acutis  
integerrimis petiolatis subtus sericeo-pubescentibus, amentis femineis ses-  
silibus ovato-cylindricis, ovariis confertis sessilibus sericeis.  $\zeta$   $\zeta$  in humidis.  
— *S. depressa* Fl. fr. 5. p. 346. *S. arenaria* Fl. fr. 3. p. 293. non L.  
nec Willd.

a. *argentea* (Ser.) foliis ovatis supernè pubescentibus mucronatis, amen-  
tis femineis sessilibus. — *S. argentea* Sm. brit. 1059. eng. bot. t. 1354.  
*S. incubacea* L. sp. 1447. Fl. fr. 3. p. 293. (v. v.)

ß. *elliptica* (Ser.) foliis ellipticis supernè glabris, amentis femineis sub-  
pedunculatis. — *S. repens* eng. bot. t. 183. Vill. dauph. 4. t. 50. f. 10.  
*S. depressa* Hoffm. sal. t. 15 et 16. Fl. fr. 3. p. 294. (v. v.)

6. *S. VERAEFOLIA* (Wahl. lapp. t. 18. f. 2.) suffrutex, foliis ovatis lanceolatisve  
integris aut denticulato-serratis subrevolutis supernè subglabris aut adpressè  
pubescentibus subtus velutinis reticulatis discoloribus petiolatis acutis,  
stipulis lanceolatis ovatisve subacutis, amentis cœtaneis sessilibus, femineis  
primum ovatis densis subvillosis demum ovato-cylindricis, ovariis tomen-  
tosis ovato-oblongis demum pedicellatis.  $\zeta$  in turfosis Jurassi, Alpium. —  
Fl. fr. 5. p. 345. *S. spathulata* Willd. sp. 4. p. 700. (v. s.)

#### Sect. II. DAPHNELLA Ser. sal. rev. ined.

Amenta præcocia primum ovato-cylindrica, ovaria glabra raro tomen-  
tosa; stylus elongatus; Masc. diandri liberi. — Folia lanceolata vel ob-  
ovata.

7. *S. DAPHNOIDES* (Vill. dauph. 4. t. 50. f. 7.) frutex, foliis elliptico-acuminatis  
serratis glabris subtus glaucis supernè viridibus nitidis junioribus subvillosis,  
amentis sessilibus crassis dense tomentosis, ovariis glabris.  $\zeta$  in Alpibus  
Delphinatûs. — Fl. fr. 3. p. 286. excl. syn. L. *S. præcox* Ser. ess. 53. (v. v.  
in Helv.)

8. *S. PHYLLICIFOLIA* (L. sp. 1442.) frutex, foliis ovatis lanceolatisve dentato-  
serratis acuminatis subtus pubescentibus, junioribus tomentosis, stipulis  
magnis dentatis, amentis subcœtaneis ovatis demum cylindricis, femineis  
subpedunculatis, stylo elongato, capsulis longè conicis pedicellatis glabris  
aut tomentosis.  $\zeta$  secus rivos et torrentes Alpium. — Ser. rev. ined. exsicc.  
n. 10. non Fl. fr. 3. p. 286.

a. *stylosa*, capsulis laxis glabris aut vix tomentosis. — *S. stylosa* Fl. fr. 5.  
p. 339. *S. stylaris* Ser. ess. 65. *S. amaniana* Willd. sp. 4. p. 663. *S. phylli-  
cifolia* Wahl. lapp. t. 17. f. 2. — L. lapp. t. 8. f. d. (v. v.)

ß. *nigricans*, capsulis subconfertis tomentosis. — *S. nigricans* Wahl. lapp.  
t. 17. f. 3. Fl. fr. 5. p. 345. Smith. eng. bot. t. 1223.

#### Sect. III. VIMINELLA Ser. sal. rev. ined.

Amenta cylindracea subcœtanea; ovaria tomentosa raro glabra; stylus  
elongatus; Masc. diandri; stamina libera vel monadelphæ. — Folia lanceolata  
vel linearia.

9. *S. LANCEOLATA* (Ser. ess. 37. t. 1.) arbuscula, ramis junioribus subpubes-  
centibus, foliis lato-lanceolatis denticulatis acutis breviter petiolatis supernè

- subglabris subtus cano-velutinis reticulatis, stipulis ovatis acutis obtusive, amentis subpræcoccibus cylindricis subcurvis breviter pilosis, ovariis tomentosis elongato-conicis pedicellatis.  $\bar{\sigma}$  in udibus circa Parisios. — Fl. fr. 5. p. 348. (v. s.)
10. *S. MOLLISSIMA* (Ebr. dec. 8. n. 79.) arbuscula, foliis lineari-lanceolatis subintegerrimis aut tenuissimè glanduloso-serratis supernè glabris subtus adpressè tomentosis discoloribus acuminatis breviter petiolatis, amentis subpræcoccibus oblongis, femineis subpedunculatis lanatis, pilis squamarum stigmatum longitudine, ovariis lanatis sessilibus confertis.  $\bar{\sigma}$  secus rivos in valle Rhenanâ. — Fl. fr. 5. p. 349. (v. s.)
11. *S. VIMINALIS* (L. sp. 1448.) frutex, foliis lanceolato-linearibus longissimis integerrimis acuminatis subtus argenteis, junioribus supernè glabris subtus adpressè tomentosis discoloribus acuminatis breviter petiolatis, amentis femineis sessilibus elongato-cylindricis, ovariis sessilibus sericeo-pubescentibus.  $\bar{\sigma}$  in humidis. — Fl. fr. 3. p. 297. eng. bot. t. 1898. (v. v.)
12. *S. VITÆA* (Hoffm. sal. 6r. t. 13. f. 2. t. 14. f. 3. 4.) arbuscula, foliis linearilanceolatis denticulatis aut undulato-crenatis acuminatis breviter petiolatis supernè glabris subtus pubescentibus, stipulis lineari-lanceolatis acuminatis petiolo subbrevioribus, amentis præcoccibus cylindricis sessilibus, staminibus a basi subcoalitis, ovariis sessilibus sericeis confertis.  $\bar{\sigma}$  in humidis. — Fl. fr. 5. p. 349. *S. rubra* eng. bot. t. 1145. *S. virescens* Vill. dauph. 3. t. 5r. f. 30. (v. v.)
13. *S. MONANDRA* (Ard. mem. 1. t. 11.) subarborea, foliis lanceolato-linearibus apice glanduloso-serratis sessilibus acutis glabris, junioribus adpressè pubescentibus, staminibus in unum coalitis, amentis femineis cylindricis sessilibus, ovariis ovato-globosis confertis tomentosis.  $\bar{\sigma}$  ad ripas et in humidis. — Fl. fr. 3. p. 297. Hoffm. sal. t. 1. f. 1. 2. et t. 5. f. 1. *S. purpurea* et *S. Helix* L. (v. v.)
14. *S. INCANA* (Schrank. fl. bev. 1. p. 230 non Mich.) frutex, foliis linearibus aut lineari-lanceolatis margine revolutis glanduloso-serratis sessilibus subtus tomentosis, staminum filamentis bifidis, amentis femineis cylindricis sessilibus, ovariis confertis sessilibus glabris.  $\bar{\sigma}$  in montosis Gallie orientalis, australis et austro-occidentalis. — Fl. fr. 3. p. 284 et 5. p. 337. *S. angustifolia* Poir. in Duham. arb. 2. ed. 2. t. 29. non Willd. *S. lavandulæ-folia* Lapeyr. *S. riparia* Willd. (v. v.)

Sect. IV. *ALBELLA* Ser. l. c.

Amenta cylindrica elongata coctanea vel æspius serotina, ovaria glabra vel villosa, stylus brevis. *Masc.* 2-3-andri liberi. — Folia lanceolata vel linearia.

15. *S. TRIANDRA* (L. sp. 1442.) arborea, foliis oblongo-lanceolatis glabris subintegris serrulatisve acuminatis acutisve petiolatis, stipulis parvulis obtusis ovato-cordatis serrulatis, staminibus ternis, amentis femineis elongatis sessilibus pubescentibus, ovariis glabris.  $\bar{\sigma}$  in arenosis secus flumina. — Fl. fr. 3. p. 285 et 5. p. 337. eng. bot. t. 1435. *S. amygdalina* L. sp. 1443. Fl. fr. 3. p. 285. eng. bot. t. 1936. *S. Villarsiana* Willd. (v. v.)
16. *S. FRAGILIS* (L. sp. 1443.) subarborea, foliis oblongo-lanceolatis serratis glabris acuminatis petiolatis, inferioribus minoribus obovatis, amentis femineis elongatis pubescentibus pedunculatis, ovariis laxis glabriusculis pedicellatis.  $\bar{\sigma}$  secus rivulos et flumina. — Fl. fr. 3. p. 288 et 5. p. 340. eng. bot. t. 1807. *S. decipiens* Hoffm. sal. 2. t. 3r. eng. bot. t. 1397. *S. pendula* Ser. ess. 79. (v. v.)
17. *S. ALBA* (L. sp. 1449.) arborea, foliis lanceolato-acuminatis serratis subpubescentibus subsessilibus serraturis infimis glandulosis, amentis femineis elongatis pedunculatis pubescentibus, ovariis laxis glabris sessilibus.  $\bar{\sigma}$  Ad vias, ripas et prata humida. Fl. fr. 3. p. 283. Hoffm. sal. t. 7. f. 1. t. 8. f. 2. t. 24. f. 3. *S. vitellina* L. Fl. fr. 3. p. 283. Hoffm. sal. t. 11. f. 1; t. 12. f. 2 et 3. Variat cortice virescente aut vitellino. (v. v.)

18. *S. BANYLONICA* (L. sp. 1443.) arborea, ramis elongatis flexilibus pendulis, foliis linearilanceolatis oblongo-lanceolatisve integris aut subserratis longe acuminatis glabris, amentis femineis subpedunculatis elongatis, ovarii confertis.  $\zeta$  ex Oriente orta, in humidis culta. — Fl. fr. 3. p. 286. *S. propendens* Ser. ess. (v. v. c.)

#### Sect. v. HERBELLA Ser. l. c.

Amenta ovata coëtanæ; ovaria nuda vel villosa; stylus brevissimus; *Masc.* 2-andri, filamenta libera. — Folia elliptica vel subrotunda, caulis prostratus.

19. *S. HERBACEA* (L. sp. 1445.) suffruticosa, caulibus procumbentibus subterraeis, foliis orbiculatis serratis glabris petiolatis, amentis paucifloris.  $\zeta$  in pascuis editiorum Alpium et Pyrenæorum. — Fl. fr. 3. p. 288. Lin. lapp. t. 7. f. 3 et 4 et t. 8. f. 12. Hoffm. sal. t. 20. f. 1-4. (v. v.)
20. *S. ARTICULATA* (L. sp. 1446.) suffruticosa procumbens, foliis ovato-rotundis obtusissimis integerrimis petiolatis subtus albidis reticulato-nervosis glabris aut lanâ caducâ villosis, amentis femineis longè pedunculatis, ovarii ovato-globosis tomentosissimis sessilibus.  $\zeta$  in lapideis editiorum Alpium et Pyrenæorum. — Fl. fr. 3. p. 289. Hoffm. sal. t. 25, 26 et 27. — Lin. lapp. t. 7. f. 1. 2. et t. 8. f. L.

#### Sect. vi. ARBUSCULA Ser. l. c.

Amenta coëtanæ ovata obovata vel elliptica raro præcocia; ovaria tomentosa vel glabra, stylus elongatus; *Masc.* 2-rarè 5-8-andri, filamenta libera. — Folia ovata vel elliptica.

21. *S. MOOROSA* (Ehr. dec. 11. n. 118.) suffrutex, foliis ellipticis integris supernè glabris atroviridibus, subtus pubescentibus glaucis, amentis præcocibus, masculis ellipticis sericeis, femineis gracilibus, ovarii tomentosissimis. —  $\zeta$  in montibus Arvenis. — Fl. fr. 5. p. 344. (v. v. c.)
22. *S. ARBUSCULA* (L. sp. 1445.) arbuscula, foliis ovatis glanduloso-serrulatis glabris lucidis acutis aut subobtusis breviter petiolatis, junioribus subtus pubescentibus, amentis ovato-cylindricis sessilibus, squamis lanato-villosis, ovarii tomentosissimis subsessilibus ovato-oblongis.  $\zeta$  in vallibus editioribus Alpium. — Fl. fr. 3. p. 295. et 5. p. 347. L. lapp. t. 8. f. E. *S. fetida* Schleich. Fl. fr. 3. p. 296. et 5. p. 347. *S. vacciniifolia* Eng. bot. t. 2341. *S. venulosa* Eng. bot. t. 1362. *S. prunifolia* Ser. ess. 49. (v. v.)
23. *S. CASSA* (Vill. dauph. 3. t. 50. f. 11.) arbuscula, foliis ovali-lanceolatis lanceolatisve acutis integerrimis glabris subsessilibus, amentis cylindricis brevibus sessilibus, squamis glabriusculis, ovarii confertis sessilibus lanato-sericeis.  $\zeta$  secus rivulos in editioribus Alpibus Galloprovinciæ. — Fl. fr. 3. p. 294. *S. prostrata* Ehr. pl. select. 159. Ser. ess. 24. (v. v.)
24. *S. MYRSINITES* (L. sp. 1445.) arbuscula, foliis ovatis aut lanceolato-ovatis acutis subobtusisve glabris serratis reticulato-venosis sessilibus, amentis ovatis, femineis demum cylindricis subpedunculatis, ovarii hirsutis oblongo-ovatis sessilibus.  $\zeta$  in Alpibus editioribus Delphinatus. — Fl. fr. 3. p. 296. Lin. lapp. t. 7. f. 6. Eng. bot. t. 1360. *S. arbutifolia* Willd. Ser. ess. 44. (v. v. in Helv.)
25. *S. PYRÆNICA* (Gou. ill. 77.) suffruticosa repens, foliis obovatis subobtusis aut breviter mucronulatis integerrimis subsessilibus ciliatis supernè adpressè pubescentibus subtus pilosis, junioribus sericeis, amentis femineis subpedunculatis elongatis, ovarii sessilibus tomentosissimis ovatis.  $\zeta$  in Pyrenæis ad nives perpetuas. — Fl. fr. 3. p. 292. *S. ciliata* Fl. fr. 3. p. 292 excl. syn. — Clas. hist. 1. p. 85. ic. (v. v.)
26. *S. ARENARIA* (L. sp. 1447 non 2. fr. 293.) arbuscula, foliis oblongo-lanceolatis petiolatis obtusis aut subacutis integerrimis supernè primo pubescentibus demum glabris atroviridibus subtus tomento albo lanatis, amentis femineis primùm ovatis demum cylindricis subpedunculatis, ovarii tomen-

- tosis sessilibus ovato-oblongis. ♂ in summo jago Cantal repetit cl. DC. in Alpibus Delphinatūs? — Eng. bot. t. 10. t. 1809. 8. Helvetica Vill. Fl. fr. 3. p. 291 et 5. p. 343. S. limosa Wahl. lapp. t. 16. f. 4. — Hall. Helv. t. 14. f. 2. (v. v. DC.)
27. S. GLAUCA (L. sp. 1446.) arbuscula basi procumbens, foliis oblongo-lanceolatis acutis integerrimis petiolatis utrinque sericeo-villosis, amentis femineis demum cylindrico-elongatis lanatis pedunculatis, ovariis sessilibus sericeo-lanatis. ♂ in montosis subhumidis Montis Aurei, Alpium Delphinatūs. — Linn. lapp. t. 7. f. 6. S. sericea Vill. dauph. t. 41. f. 27. Fl. fr. 3. p. 292 et 5. p. 344. S. Lapponum Linn. lapp. t. 8. f. t. (v. s.)
28. S. RETUSA (L. sp. 1445.) suffruticosa procumbens, foliis obovatis obtusissimis glabris integris aut subserratis sessilibus, amentis paucifloris, femineis sessilibus glabris, ovariis laxis glabris. ♂ in Jurasso, Alpibus, montibus Arvernæ, Pyrenæis. — Fl. fr. 3. p. 289. Schk. handb. t. 317. a. (v. v.)
- ♂. *serpillifolia* (Wahlb.) foliis ellipticis integerrimis interdum subacutis. — S. *serpillifolia* Scop. carn. t. 61. (v. v.)
29. S. HASTATA (L. sp. 1443.) frutex, foliis ovatis lanceolatisve subpubescentibus serratis acutis aut subacuminatis, amentis sericeo-villosissimis ovatis demum elongato-cylindricis et glabrescentibus, femineis subpedunculatis, ovariis glabris longe conicis breviter pedicellatis. ♂ secus torrentes Alpium, Pyren. (cl. L. Dufour). — Fl. fr. 5. p. 338. Wahl. lapp. t. 16. f. 5. S. *malifolia* Eng. bot. t. 1617. (v. v.)
- 30? S. PENTANDRA (L. sp. 1442) arborea, foliis elliptico-lanceolatis tenuiter serrato-crenatis glabris subviscosis supernè lucidis acutis acuminatisve, staminibus 5-7, amentis femineis cylindricis subpedunculatis, ovariis laxis glabris subsessilibus. ♂ in Jurasso, Alpibus, montibus Arvernæ, Ligèris (Lozère), Pyrenæis. — L. lapp. t. 8. f. z. Fl. fr. 3. p. 287. — Gm. sib. 1. t. 34. f. 1. (v. v.)

VI. POPULUS Tourn. inst. t. 365. L. gen. 1123.

Dioica. Amenta cylindrica, squamis apice laceris. Masc. Stamina 8-30, squamularum basi ex urceolo oblique truncato prodeuntia. Fem. Ovarium 1; stigmata 4; capsula 2-valvis marginibus introflexis sub-2-locularis polysperma; semina comosa; radícula supera.

Sect. I. LEUCE.

Turiones tomentosi vel hirti non glutinosi, stam. 8.

4.

1. P. ALBA (L. sp. 1463.) foliis cordato-subrotundis lobatis dentatis subtus tomentoso-niveis supra glabris, amentis ovato-oblongis. ♂ ubique, præcipuè in humidis. — Fl. fr. 3. p. 298. — Lob. ic. 2. p. 193. f. 1. (v. v.)
2. P. CANADENSIS (Smith. brit. 3. p. 1060.) foliis subrotundis angulato-dentatis subtus subtomentosis supra glabris, amentis cylindraceis longis. ♂ in nemorosis humidis. — Fl. fr. 3. p. 299. — Lob. ic. 2. p. 192. f. 2. (v. v.)
3. P. TREMULA (L. sp. 1464.) foliis suborbiculatis dentatis utrinque glabris, petiolis compressis, amentis oblongis. ♂ in frigidioribus subhumidis. — Fl. fr. 3. p. 299. (v. v.)

Sect. II. ALGIDIOS.

Turiones leves et glabri glutinosi, stamina 12-30.

4. P. NIGRA (L. sp. 1464.) foliis deltoidibus acuminatis serratis utrinque glabris, ramis patulis. ♂ in aquosis et cultis. — Fl. fr. 3. p. 299. — Blackw. t. 248. (v. v.)
5. P. VIRGINIANA (Desf. cat. 242.) foliis deltoidibus latis acuminatis grossè crenatis longè petiolatis utrinque glabris, ramis patulis. ♂ ex America boreali orta, nunc in provinciis plurimis et præsertim in Neustriâ et circa Parisios subsponsanea. (v. v. c.)

6. *P. FASTIGIATA* (Poir. dict. 5. p. 235.) foliis subdeltoidibus inæqualiter serratis utrinque glabris, ramis erectis strictis. § ex Oriente? orta, in ambulacris subhumidis culta. — Fl. fr. 3. p. 300. (v. v.)

Trib. IV. *QUERCINEÆ* Juss. dict. sc. nat. 2. sup. DC. mss. — *Corylaceæ* Mirb. elem. p. 906.

Fl. monoici. *Masc.* in amentum cylindricum dispositi. Perigonium parvum aut squamiforme. Stamina 5-20 perigonii basi adnata, filamentis liberis rariùs basi coadunatis. *Fem.* Involucrum varium 1-∞-florum. Perigonium ovario adhærens pluridentatum. Ovarium unicum ∞-loculare ∞-ovulatum. Stylus 2-3-∞-fidus, stigmatibus distinctis. Involucrum (Calybio Mirb.) post anthesin excrescens pericarpia 1-∞ obtegens aut fovens. Glandes aut nuces abortu 1-loculares 1-spermæ. Semen pendulum exalbuminosum. Embryo rectus. Radicula ad umbilicum tendens. Cotyledones crassæ aut foliaceæ. — Arbores aut rariùs frutices. Folia alterna simplicia. Stipulæ deciduæ gemmam constituentes.

Sect. I. INVOLUCRUM HISPIDUM CLAUSUM, DEMUM DEHISCENS.

VII. *FAGUS* Tourn. inst. 351. Lam. dict. 3. p. 125. — *Fagi* sp. L.

Monoica. *Masc.* Amenta pendula globosa densa; perigonium 6-lobatum; stamina 8. *Fem.* bini in involucri 4-lobo extus spinis mollioribus muricato; perigonium ovario adnatum tomentosum 6-lobum; stigmata 3; ovarium 3-gonum 3-loculare, nux triangularis abortu 1-locularis 1-2-sperma.

1. *F. SYLVATICA* (L. sp. 1416) foliis ovatis glabris obsolete dentatis margine ciliatis. § sylvas constituit præcipuè in calciorum montium declivibus. — Fl. fr. 3. p. 305. — Gært. fruct. 1. t. 37. f. 2. (v. v.)

VIII. *CASTANEA* Tourn. inst. t. 352. Lam. dict. 1. p. 708. — *Fagi* sp. L.

Polygama. *Masc.* Amenta longissima cylindrica floribus hinc inde glomeratis; perigonium 6-partitum; stamina 5-20. *Herm.* Involucrum subtriflorum 4-lobum, extus spinis durioribus et ramosis muricatum; perigonium ovario adnatum 5-6-lobum tomento rigido intus obtectum; stamina 12 rubra abortiva, ovarium 6-loculare loculis dispermis, 5 abortivis; styli 6; nux 1-locularis subrotunda; semina 1-3 rugosa.

1. *C. VULGARIS* (Lam. dict. 1. p. 708.) foliis oblongo-lanceolatis acuminatis mucronato-serratis utrinque glabris. § in montosis et glareosis. — Fl. fr. 3. p. 306. — Duham. ed. sec. 3. p. 65. t. 19. (v. v.)

Sect. II. INVOLUCRUM PATENS.

IX. *QUERCUS* Tourn. inst. 394. L. gen. 1070.

Monoica. *Masc.* Amentum laxum et pendulum, perigonium lacerum; stamina 5-10. *Fem.* Involucrum squamosum squamis plurimis imbricatis, in cupulam hemisphæricam et coriaceam coadunatis constans; perigonium ovario adnatum 6-lobatum; ovarium 3-loculare, loculis 2 abortivis; stigmata 3; nux (glands) 1-locularis 1-sperma cupulâ basi cincta.



\* *Foliis deciduis.*

1. *Q. CERRIS* (L. sp. 1415.) foliis oblongis pinnatifidis glabris aut subtus hirtis basi angustatis, lobis oblongo-lanceolatis dentatis, cupulâ hemisphæricâ squamis longis acutis patulis echinâtâ.  $\frac{5}{5}$  circâ Bellovacum ex cl. Graves, Cœnomanum, Andegavum, Nannetes. — Fl. fr. 5. p. 354 et 3. p. 311. excl. var.  $\beta$ . et  $\gamma$ . (v. v. DC.)
2. *Q. TOZA* (Bosc. journ. hist. nat. 2. p. 155. t. 32. f. 3.) foliis petiolatis subtus tomentosis oblongis sinuato-lobatis pinnatifidisve, lobis obtusis, fructibus pedunculatis, cupulâ hemisphæricâ squamis oblongis subacutis basi adpressis apice subdivergentibus.  $\frac{5}{5}$  et  $\frac{5}{5}$  in provinciis occidentalibus vulgaris. — Fl. fr. 5. p. 352. *Q. Cerris*  $\gamma$ . Fl. fr. 3. p. 311. *Q. humilis* Fl. fr. 3. p. 312. non L. Foliorum formâ, pedunculi longitudine et habitu mire variat. (v. v.)
3. *Q. PUBESCENS* (Willd. sp. 4. p. 450.) foliis petiolatis oblongo-obovatis sinuatis subtus pubescentibus lobis obtusis angustatis basi subcordatis inæqualibus, fructibus subsessilibus, cupulâ hemisphæricâ glande vix duplò breviorè, squamis ovatis adpressis obtusis.  $\frac{5}{5}$  in collibus siccis. — Fl. fr. 5. p. 352. (v. v.)
4. *Q. ASPENINA* (Lam. dict. 1. p. 725.) foliis subpetiolatis obovatis sinuato-lobatis lobis obtusis subtus petiolisque pubescentibus, fructibus pedunculatis spicatis, cupulâ hemisphæricâ glande vix duplò breviori, squamis lanceolatis subobtusis.  $\frac{5}{5}$  in collibus Alsatie ex cl. Nestler, Lugduni et provinc. australium ex cl. Bosc. — Fl. fr. 5. p. 352. (v. s.)
5. *Q. RACEMOSA* (Lam. dict. 1. p. 725.) foliis subsessilibus glabris oblongis apice dilatatis sinuato-lobatis, lobis rotundatis, fructibus oblongis longè pedicellatis, cupulâ hypocrateriformi glande triplò breviori, squamis ovatis subobtusis.  $\frac{5}{5}$  in sylvis vulgaris. — Fl. fr. 3. p. 309. — Duham. arb. 2. t. 47. *Q. microcarpa* Lapeyr. (v. v.)
6. *Q. FASTIGIATA* (Lam. dict. 1. p. 725.) foliis subsessilibus glabris oblongis sinuato-lobatis, lobis obtusis, ramis erectis, fructibus 3-5 in extremitate longi pedunculi, cupulâ hemisphæricâ glande quadruplò quintuplò breviorè, squamis ovatis obtusis.  $\frac{5}{5}$  in Pyrenæis occidentalibus et agro Syrtico culta? — Fl. fr. 5. p. 351. Vix à priore distincta? (v. v.)
7. *Q. SESSILIFLORA* (Sm. brit. 3. p. 1026.) foliis petiolatis glabris oblongis sinuato-lobatis, lobis rotundatis, fructibus oblongis subsessilibus, cupulâ hypocrateriformi fructu triplò quadruplò breviori, squamis lanceolatis obtusis.  $\frac{5}{5}$  in sylvis. — Fl. fr. 3. p. 310. excl. var.  $\alpha$ . et  $\zeta$ . (v. v.)

\*\* *Foliis persistentibus.*

8. *Q. ILEX* (L. sp. 1412.) cortice integro, foliis oblongis indivisis serratisque acuminatis subtus incanis, cupulis pedunculatis glande ovato-oblongo triplò quadruplò brevioribus.  $\frac{5}{5}$  in Galliâ australiori frequens, in Galliâ occid. usque ad Nannetes rarior. — Fl. fr. 3. p. 313. — Duham. arb. 1. t. 123. — Pluck. t. 197. f. 1. (v. v.)
9. *Q. SUBER* (L. sp. 1413.) cortice rimoso fungoso, foliis ovato-oblongis indivisis serratisve subtus tomentosis obtusis aut mucronatis.  $\frac{5}{5}$  in Galliâ australiori et austro-occidentali. — Fl. fr. 3. p. 313. — Duh. arb. 2. p. 291. t. 80 et 81. (v. v.)
10. *Q. COCCIFERA* (L. sp. 1413.) foliis oblongis indivisis spinoso-dentatis basi cordatis utrinque glabris, cupulâ subsessili, squamis lineari-lanceolatis acutis patulis.  $\frac{5}{5}$  in Galliâ australiori et Nigri-Monasterii insulâ. — Fl. fr. 3. p. 313. — Garid. aix. t. 53. (v. v.)

X. *CORYLUS* Tourn. inst. 347. L. gen. 1074.

*Monoica. Masc. Amenta cylindrica, squamæ 3-lobatæ deltoideæ laciniâ mediâ latiore laterales tegente, stamina 8, antheræ 1-loculares:*

*Fem.* Flores plurimi in gemmâ squamosâ inclusi; stigmata 2; ovarium 1 primo denudatum, dein involucri calyciformi monophyllo margine lacero cinctum; nux ovata lævis basi involucriata 1-sperma.

1. *C. AVELLANA* (L. sp. 1417.) stipulis oblongis obtusis, involucri campanulati apice patulis lacero-dentatis, foliis ovato-cordatis acuminatis. 3 in æpibus. — Fl. fr. 3. p. 308. Lam. ill. t. 780. — Lob. ic. 2. t. 192. (v. v.)

XI. *CARPINUS* Mich. 104. DC. fl. fr. 3. p. 304. — *Carpini* sp. L.

Monoica? *Masc.* Amenta elongato-cylindrica; squamæ basi ciliatæ; stamina 8-14 apice subbarbata. *Fem.* Strobili laxi; involucri squamiforme 3-lobum biflorum; ovarium apice denticulatum 2-loculare, loculo uxo antè maturitatem abortivo, stigmata 2; nux ossea.

1. *C. BETULYA* (L. sp. 1416.) strobilorum squamis 3-partitis, laciniis integris lanceolatis, intermediâ elongatâ. 3 5 in sylvis. — Fl. fr. l. c. Gærtn. fruct. 2. t. 89. f. 2. (v. v.)

Trib. V. *PLATANÆ* Juss. dict. sc. nat. DC. mss.

Fl. monoici in amenta globosa aut oblonga sexu distincta dense aggregati. Involucri commune 6 aut 4-phyllum ad basin amenti. *Masc.* Squamulæ lineares parvæ innumeræ cum staminibus intermixtæ et squamarum aut perigoniorum vice gerentæ. *Fem.* Squamæ floribus intermixtæ aut nullæ? Perigonium ovario adhærens? nunc cupuliforme nunc in setas piliformes minimas desinens, nunc vicinis adglutinatum nunc ab iis liberum. Carpella 1-2 unilocularia cum perigonio coalita oblonga aut subturbinata apice cornuta indehiscencia aut dehiscencia subcoriacea. Semen in quoque carpello solitarium pendulum exalbuminosum. Embryo rectus. Radicula supera. Cotyledones planæ foliaceæ. — Arbores, folia alterna petiolata palminervia palmatiloba; stipulæ nullæ aut foliaceæ. Gemmæ intrâ petioli basin latentes.

XII. *PLATANUS* Tourn. inst. t. 363. L. gen. 1075.

Amenta basi nuda. *Fem.* Squamæ spathulatæ; ovaria filiformia apice incrassata stigmate aduncò terminata; carpella solitaria clavata basi papposa.

1. *P. ORIENTALIS* (L. sp. 1417.) foliis 5-lobis palmatis laciniis lanceolatis sinuatis, stipulis subintegerrimis. 3 ex Oriente orta, in ambulacris culta. — Fl. fr. 3. p. 314. Lam. ill. t. 783. (v. v. c.)  
2. *P. OCCIDENTALIS* (L. sp. 1418.) foliis 5-angularibus obsolete lobatis dentatis basi cuneatis subtus pubescentibus. 3 ex America septentrionali orta, in ambulacris culta. — Catesb. carol. 1. t. 56. (v. v. c.)

Trib. VI. *MYRICEÆ* DC. mss.

Flores rariùs hermaphroditi sæpiùs diclini monoici dioice amentacei, amentis unisexualibus. Squamæ ovatæ acutæ ad axillam 1-floræ. *Masc.* Squamulæ aut perigonii partes 2 oppositæ. Stam. 4, uno interdum incompletò, libera. Anthere 2-loculares longitudinaliter dehiscences. *Fem.* Squama post anthesin exeres-

cens. Squamulæ seu perigonii partes 3-6 minutissimæ. Ovarium simplex liberum. Stigmata 2 filiformia. Drupa sessilis globosa extus granulis ceraceis plus minus oblecta, matura subexsiccata stylo persistente superata aut apice submutica. Putamen osseum evalve. Albumen nullum aut amygdalino-carnosum. Embryo rectus inversus. Radicula supera. Cotyledones carnosæ plano-convexæ. — Frutices in partibus plurimis punctis resinosis instructi aromatici. Folia alterna, simplicia.

XIII. MYRICA L. *gen.* 1197.

Dioica. Amenta ovata squamis lunulatis, *Masc.* Stam. 4-6; antheræ 4-valves. *Fem.* Ovarium 1; stigmata 2; drupa 1-locularis 1-sperma, semen exalbuminosum.

1. M. GALE (L. sp. 1453.) caule fruticoso, foliis lanceolatis subserratis. 3 in erictis uliginosis Gallie occidentalis et boreali-occidentalis. — Fl. fr. 3. p. 301. Fl. dan. t. 327. (v. v.)

Ordo XCIV. CONIFERÆ Juss. *gen.* p. 411. *Rich. conif.* 89.

Fl. diclini monoici aut dioici in amentum formâ varium sæpius dispositi. *Masc.* squamæ plurimæ varie dispositæ antheras nunc gerentes nunc foveantes. Perigonium nullum. Antheræ numero variæ 1-∞-loculares, nunc squamis bracteiformibus insertæ, nunc pedicello proprio donatæ. *Fem.* Squamæ bracteiformes varie dispositæ interdum post anthesin accretæ succulentæ aut fructu adnatæ. Cupula interdum duplex, rarius simplex 1-flora ovarium cingens (an Perigonium?). Ovarium unicum. Stylus in Ephedra longus exsertus stigmate obliquo terminatus; in cæteris sessilis simplex parvus. Pericarpium proprium (bracteis cupulisque quibuscum sæpè adhæret aut quibus occultatur exelulis) indehiscens, 1-loculare coriaceum osseumve. Semen 1 pendulum albuminosum. Embryo axillis rectus. Radicula ad umbilicum tendens sæpè apice attenuata et cum albumine adnata. Cotyledones 2 oppositæ aut plurimæ verticillatæ. — Arbores aut frutices resinose, Folia alterna aut verticillata aut rarius opposita sæpius acerosa sempervirentia, ramulis abortivis sæpè fasciculata.

Trib. I. TAXINEÆ *Rich. ann. mus.* 16. p. 296. *conif.* 124.

Gemmæ florales uni-aut rarius bifloræ, squamis plurimis cruciatim oppositis et imbricatis constantes. Fl. femineæ squamis minimè innati.

I. TAXUS *Tourn. inst.* 362. *L. gen.* 1135.

Flores dioici vel monoici, squamis pluribus cincti. *Masc.* Stamina 8-10, filamentis monadelphis, antheris peltatis 6-8-locularibus, localis subtus dehiscentibus. *Fem.* Stylus nullus; stigma concavum; drupa carnosa apice hians; nucleus 1-spermus.

1. *T. BACCATA* (L. sp. 1472.) foliis approximatis linearibus acutis, floribus axillaribus sessilibus. § in subalpinis et Jurasso frequens, Neustriæ indigena ex cl. Le Prévost. — Fl. fr. 3. p. 280. Rich. conf. t. 2. f. 1. — Cam. epit. 840j ic. (v. v. c.)

II. EPHEDRA *Tourn. inst.* t. 477. *L. gen.* 1136.

Flores dioici in amentum breve dispositi. *Masc.* Perigonium 2-lobum; stamina 6-8, filamentis monadelphis, antheris 1-locularibus, extus dehiscentibus. *Fem.* Squamæ 4-5 persistentes concavæ truncatæ imbricatæ post inflorescentiam carnosæ; ovaria 2, monostyla; bacca ovata 2-sperma, seminibus hinc planis inde convexis.

1. *E. DISTACHYA* (L. sp. 1472.) pedunculis oppositis, amentis geminis. § in sabulosis maritimis Galliae australis à Niczâ ad Nannetes usquæ. — Fl. fr. 3. p. 281. *E. vulgaris* Rich. conf. t. 4. f. 1. — Barr. ic. 731. (v. v.)

Trib. II. CUPRESSINÆ *Rich. l. c. conf.* 137.

Amenta feminea squamis paucis constantia, squamis demum inter se agglutinatiss in galbulos subglobosos mutata. Flores erecti stigmatem nempè sursum directo.

## Sect. I. SQUAMÆ DEMUM CARNOSE BACCAM SPURIAM CONSTITUENTES.

III. JUNIPERUS *L. gen.* 1134.

Dioica rariùs monoica. *Masc.* Amenta ovata, squamis verticillatis peltato-pedicellatis; antheræ 4-8, uniloculares. *Fem.* Amenta globosa squamis 3 concavis coadunatis; stigma hians; bacca conflata ex tribus cariopsidibus osseis 1-spermis cinctis squamis coalitis et carnosiss.

1. *J. PRAENICKA* (L. sp. 1471.) foliis ternis densissimè imbricatis obtusis ovatis convexis. § in Gallia australi. — Fl. fr. 3. p. 279. — Duham. arb. t. 52. — Lob. ic. 2. p. 221. (v. v.)
2. *J. SABINA* (L. sp. 1472.) foliis ovatis convexis densissimè imbricatis erectis decurrentibus oppositis, oppositionibus pyxidatis. § in Alpium vallibus calidis. — Fl. fr. 3. p. 279. Bull. herb. t. 139. (v. v. c.)
3. *J. OXYCEDRUS* (L. sp. 1470.) foliis ternis patentibus mucronatis bacca brevioribus. § in Gallia australi. — Fl. fr. 3. p. 278. Rich. conf. t. 6. f. 1. — Lob. ic. 2. p. 223. f. 2. (v. v.)
4. *J. COMMUNIS* (L. sp. 1470.) foliis ternis patentibus mucronatis bacca longioribus. § in collibus aridis et siccis. — Fl. fr. 3. p. 278. Rich. conf. t. 6. — Duh. arb. 1. t. 127. — J. Bauh. hist. 1. p. 301. ic. p. 302. (v. v.)

## Sect. II. SQUAMÆ SICCÆ GALBULUM DEHISCENTEM CONSTITUENTES.

IV. CUPRESSUS *Tourn. inst.* t. 358. *L. gen.* 1079. *DC. fl.* 5. p. 335.

Monoica. *Masc.* Amentum imbricatum; antheræ 4 sessiles. *Fem.* Amentum strobilaceum, squamis lignosis pedicellatis; galbulus angulatus.

1. *C. FASTIGIATA* (DC. cat. monsp. 22.) ramis strictis apice quadrangulis, foliis quadrifariam imbricatis obtusis, strobili squamis muticis. § in prov. australibus culta. — Fl. fr. 5. p. 336. *C. sempervirens* a L. sp. 1422. Duham. arb. ed. 2 vol. 3. t. 1. (v. v. c.)
2. *C. HORIZONTALIS* (Mill. dict. n. 2.) ramis patulis apice quadrangulis, foliis quadrifariam imbricatis obtusis, strobili squamis muticis. § in prov. australibus culta. — Fl. fr. 5. p. 336. — Black. herb. t. 127. (v. v. c.)

Trib. III. *ABIETINÆ* Rich. l. c. conif. 145.

Amenta feminea squamis plurimis constantia, squamis imbricatis demum distinctis in strobilos conosve conicos dispositis. Fl. inversi bracteis nempe per totam ferè longitudinem adnatis, stigmate deorsum directo.

V. *PINUS* Tourn. inst. 355. Juss. gen. 414. — Pini sp. L.

**Monoica. Masc.** Amenta racemosa compacta et terminalia, squamosa squamis apice staminiferis; stamina 2; antheræ 1-loculares. **Fem.** Amenta simplicia imbricata squamis acuminatis; ovaria 2; stigmata glandulosa; coni squamæ oblongæ clavatæ ligneæ, apice umbilicato anguloso; cariopsides geminæ 1-spermæ membranâ apiculatâ obtectæ; cotyledones digitato-partitæ. — Folia bina aut plurima ex eadem vaginâ.

1. *P. SYLVESTRIS* (Mill. dict. n. 1.) foliis geminis rigidis, strobilis ovato-conicis acutis longitudine foliorum subgeminis basi rotundatis, embryone 5-lobo. § in montosis sylvas formans. — Fl. fr. 3. p. 271. Lamb. pin. t. 1. Rich. conif. t. 11. *P. rubra* Mill. Fl. fr. 3. p. 272. — Duham. 2. t. 30. — Blackw. t. 180. (v. v.)
2. *P. UNGUINATA* (Ram. in DC. fl. fr. 3. p. 726.) foliis geminis longioribus strictis subglaucis, strobilis ovato-oblongis obtusis, squamis umbilicato-uncinatis, embryone 7-lobo. § in Pyrenæis editioribus, Alpibus, Jurasso. — *P. sanguinea* Lapeyr. abr. 587. (v. v. DC.)
3. *P. MUÑO* (Mill. dict. n. 5.) foliis geminis trinervi tenuioribus rigidis, strobilis pyramidalis acutis basi rotundatis, squamis acutis. § in editioribus montibus Delphinatûs ex Vill. — Fl. fr. 3. p. 273. Duham. arb. 2. p. 134. t. 31. n. 6. Dubia civis.
4. *P. MARITIMA* (Lam. fl. fr. 2. p. 201.) foliis geminis prælongis rigidis pungentibus squamâ reflexâ basi donatis, strobilis oblongo-conicis obtusis glaberrimis lucidis foliis brevioribus, squamis echinatis. § in arenosis maritimis Galliæ austro-occidentalis, Narbonæ ex Benth. Galloprov., Niceæ. — Fl. fr. 3. p. 273. et 5. p. 335. *P. Pinaster* Lamb. pin. t. 4 et 5. — Duham. arb. 2. t. 29. n. 4. (v. v.)
5. *P. PUMILIO* (Waldst. et Kit. hung. t. 149.) foliis geminis adpressis, strobilis ovatis subrotundis erectis, truncis ascendentibus. § in turfosis Jurassi. — Lamb. pin. t. 2. Fl. fr. 5. p. 334. (v. s.)
6. *P. PINEA* (L. sp. 1419.) foliis geminis primordialibus ciliatis, strobilis ovatis obtusis subinermibus foliis longioribus, nucibus duris. § in montosis Galliæ australis frequens. — Fl. fr. 3. p. 273. Lamb. pin. t. 6, 7 et 8. Rich. conif. t. 12. f. 1. — Duham. arb. 2. t. 27. n. 1. (v. v.)
7. *P. ALPENSIS* (Mill. dict. n. 8.) foliis geminis filiformibus, strobilis ovato-oblongis deorsum inflexis, squamis lævibus obtusis, ramis patulis. § in regione Olearum. — Fl. fr. 3. p. 274. et 5. p. 335. Lamb. pin. t. 11. *P. maritima* Lamb. pin. t. 9 et 10. non L. (v. v.)
8. *P. LARICIO* (Poir. dict. 5. p. 339.) foliis geminis longissimis difformibus, strobilis conicis acutis brevibus nutantibus, squamis basi angustioribus apice crassissimis non angulatis. § in montibus Corsicæ. — Fl. fr. 3. p. 274. (v. s.)

9. *P. Cembra* (L. sp. 1419.) foliis quinis, strobilis ovatis obtusis, squamis adpressis ovalibus concavis. § in montibus Galloprov., Delphinatûs. — Fl. fr. 3. p. 275. Lamb. pin. t. 23 et 24. — Duham. arb. 1. t. 32. (v. s.)

VI. *ABIES* Tourn. inst. t. 353. DC. fl. fr. 3. p. 275. — *Pini* sp. L.

Monoica. Masc. Amenta solitaria non racemosa, squamis apice staminiferis; stam. 2; antheræ 1-loculares. Fem. Amenta simplicia; ovaria 2; stigmata glandulosa; strobili squamæ imbricatæ tenues apice rotundatæ nec incrassatæ nec angulosæ nec dorso umbilicatæ; cotyledones digitato-partitæ. — Folia solitaria ex vaginâ communi prodeuntia.

Sect. I. *EPICEA* DC. fl. fr. l. c.

Strobili penduli; folia sparsa.

1. *A. excelsa* (DC. l. c.) foliis 4-gonis, strobilis cylindricis, squamis rhombeis complanatis margine repandis erosis. § in Vogesis, Jurasso, Alpibus, Pyrenæis. — Rich. conf. t. 14. f. 2. t. 15. *Pinus Abies* L. sp. 1421. — Cam. epit. 47. ic. (v. v.)

Sect. II. *ABIES* DC. l. c.

Strobili erecti; folia disticha.

2. *A. pectinata* (DC. l. c.) foliis planis emarginatis, strobili squamis obtusissimis adpressis. § in montibus excelsis. — Rich. conf. t. 16. f. 2. *Pinus Picea* L. sp. 1420. — Cam. epit. 48 et 49. ic. (v. v.)

VII. *LARIX* Tourn. inst. 357. DC. fl. fr. 3. p. 277. — *Pini* sp. L.

Monoica. Omnia Abietis sed cotyledones simplices nunquam lobatæ; strobili laterales. — Folia caduca in primordio cespitoso - fasciculata demum solitaria.

1. *L. europæa* (DC. l. c.) strobilis ovato-oblongis obtusis. § in Alpibus editioribus. — *P. Larix* L. sp. 1428. *Abies Larix* Rich. conf. t. 13. — Lam. ill. t. 785. f. 2. (v. v.)

Classis Secunda. MONOCOTYLEDONEÆ  
seu ENDOGENÆ PHANEROGAMÆ DC.  
*organ.* 1. p. 138.

Truncus medullâ centrali, radiis medullaribus corticeque verâ plane destitutus, fibris sparsis nec concentricè dispositis constans, apice tantum crescens, in margine magis quam in centro induratus, interdum abortivus subterraneus radiciformis. Folia sæpe vaginata integra nervis simplicibus, aut lobata nervis ramosis, nunquam verè composita. Flores distincti et sæpius numero ternario dispositi, tegumento simplici biseriato, aut si mavis duplici utriusque seriei partibus similibus. Embryo cotyledone unico aut potius cotyledonibus alternis donatus.

---

Ordo XCV. HYDROCHARIDEÆ DC. *fl. fr.* 3. p. 265. L. C.  
*Rich. in mem. inst. sc. phys.* 1811. p. 1. et seq. — Hydrocharidum *gen. Juss.*

Flores in spathâ inclusi, dioici rariùs hermaphroditi, Spatha in masculis mox uniflora flore sessili aut pedunculato, mox multiflora floribus pedunculatis; in femineis et hermaphroditis spatha uniflora floribus sessilibus. Perigonium sexpartitum (in femineis ovario adhaerens), lobis externis foliaceis, internis petaloideis longioribus, intus (in generibus acaulibus) appendicibus variis instructum. Stamina 1-13, in hermaphroditis ovario, in masculis ovarii loco inserta; antheræ 2-loculares. Ovarium inferum; stylus sæpe 0; stigmata 3-6 internè glandulosa sæpè bifida. Fructus limbo calycis persistente interdum coronatus oblongus indehiscens. Pericarpium carnosum intus pulposum mox uniloculare, mox sepimentis plus minus prolongatis submultiloculare. Semina plurima mox parietibus mox dissepimentis adfixa, integumento membranaceo duro. Embryo cylindraceus rectus. Albumen nulum. — Herbæ aquaticæ. Folia sessilia aut petiolata integra aut tenuissimè denticulata.

## 436 MONOCOTYL. SEU ENDOGENÆ.

### + FRUCTUS UNILOCULARIS.

#### I. VALLISNERIA *L. gen.* 1097.

**Dioica. Masc.** Spadix terminalis conicus parvus, spatha 2-3-4-partita, floribus minimis sessilibus tectus; singulis perigonium 3-partitum; stamina 2 (ovario abortivo imposita?) **Fem.** Scapus spiralis longissimus; spatha tubulosa, 2-fida, 1-flora; perigonium elongatum 6-partitum, lacinii alternis linearibus; stylus 0; stigmata 3 ovata 2-fida medio appendiculata; capsula elongata cylindrica 3-dentata 1-locularis polysperma ad parietes seminifera.

1. *V. spiralis* (*L. sp.* 1441.) in canali Occitanico, in Rhodano propè Arelatem, Arausionem, in Axonâ propè Suessiones, circâ Domfront. — *Fl. fr.* 3. p. 267. *Mish. gen.* 12. t. 10. f. 1. 2. (v. s.)

### + + CAPSULA SEXLOCULARIS.

#### II. STRATIOTES *L. gen.* 687.

**Spatha compressa persistens profundè bipartito-carinata 1-flora; perigonium tubulosum limbo 6-partito, lacinii 3 exterioribus minimis virescentibus, 3 interioribus magnis coloratis; stamina circiter 20, apici tubi vel ovarii margini inserta; styli 6 bifidi; capsula carnea apicibus attenuata 6-gona 6-locularis; semina subangulosa valvarum sepimentis alligata.**

1. *S. aloides* (*L. sp.* 1754.) foliis ensiformi-triangularibus ciliato-aculeatis. 2/ circâ Burdigalam ex Laterr. — *Fl. fr.* 3. p. 266. *eng. bot.* t. 379. (v. s. ex Belgio.)

#### III. HYDROCHARIS *L. gen.* 1126.

**Dioica. Masc.** Spatha 2-partita 3-flora; perigonium 6-partitum petaloideum, lacinii tribus interioribus majoribus; stam. 12 abortu 9, trino ordine ovario abortivo imposita. **Fem.** Spatha sessilis uniflora, perigonium masculi ovario adnatum, intus appendicibus 6 filiformibus instructum; stigmata 6 cuneiformia 2-fida; capsula coriacea ovoidea 6-locularis polysperma.

1. *H. morsus-ranæ* (*L. sp.* 1466.) in aquis stagnantibus Parisiorum, Abbavillæ, Aureliæ, Andegaviæ, Arelatis, Lugduni, Bressiæ. — *Fl. fr.* 3. p. 266. *eng. bot.* t. 808. — *Tourm. act. acad.* 1765. t. 4. Folia petiolatâ orbicularia. (v. v.)



Ordo XCVI. ALISMACEÆ *Juss. dict. sc. nat.* 1. p. 474. —  
*Alismoideæ DC. fl. fr.* 3. p. 188.

Perigonium liberum 6-partitum coloratum; stamina 6-9 rarò plura. Ovaria 3-6- $\infty$ , styli et stigmata totidem. Capsulæ indehiscentes 1-spermæ aut  $\infty$ -spermæ 2-valves. Embryo rectus aut incurvus. Albumen o. — Herbæ aquaticæ; foliis radicalibus alternis vaginantibus. Flores spicati vel umbellati, hermaphroditi, rariùs monoici.

Trib. I. *BUTOMEÆ Rich. mém. du mus.* 2. p. 365.

Perigonii segmenta 3 interna petaloidea. Semina plurima venis per totam capsularum parietem reticulatim discurrentibus adnexa ascendentiâ; embryo in directione seminis.

I. *BUTOMUS Tourn. inst.* 143. *L. gen.* 507. *Rich. l. c.*

Stamina 9, 3 interiora; pistilla 6 longissimè rostrata; capsulæ introso margine dehiscentes salvo laterali connexû; semina lineari-oblonga recta longitudinaliter striata.

1. *B. UMBELLATUS* (L. sp. 532.)  $\mathcal{U}$  secùs aquas lentè fluentes passim. — *Fl. fr.* 3. p. 191. *Lam. ill. t.* 324. — *B. floridus* Gærtn. *fruct.* 1. t. 19. f. 3. (v. v.)

Trib. II. *ALISMOIDEÆ*. — *Alismaçæ Rich. mém. du mus.* 2. p. 365.

Perigonii segmenta 3 interna petaloidea, capsulæ indehiscentes. Semina 1-2 suturæ annexa, erecta s. ascendentiâ, embryo flexus; radícula ad hylum versa.

II. *ALISMA L. gen.* 460.

Stamina 6; ovaria 6-25; capsulæ distinctæ sæpiùs 1-spermæ caducæ, non dehiscentes.

Sect. I. *ALISMOIS*. — *Alisma Juss. gen.* p. 46.

Capsulæ numerosæ 1-2-spermæ.

1. *A. MANUNCULOIDES* (L. sp. 487.) foliis lineari-lanceolatis, capsulis angulatis subpentagonis incurvis acutis globoso-aggregatis.  $\mathcal{U}$  in aquosis. — *Fl. fr.* 3. p. 189. — *Lob. ic. t.* 300. f. 2. (v. v.)

*A. repens*, caule decumbente radicante, scapis paucifloris. — In arenosis humidis provinciarum occidentalium. — *A. repens* Cav. *ic.* 1. t. 55. *Fl. fr.* 5. p. 312. (v. v.)

2. *A. NATANS* (L. sp. 487.) foliis ellipticis obtusis, pedunculis solitariis, capsulis striatis.  $\mathcal{U}$  in aquis Gallis mediæ et occidentalis. — *Fl. fr.* 3. p. 189. *eng. bot. t.* 775. — *Vaill. act. t.* 4. f. 9. (v. v.)

3. *A. PLANTAGO* (L. sp. 486.) foliis acutis oblongo-ovatis lanceolatisve 5-nerviis, capsulis compressis muticis obtusè 3-gonis.  $\mathcal{U}$  in fossis aquaticis, stagnis. — *Fl. fr.* 3. p. 189. *Lam. ill. t.* 272. — *Barr. ic. t.* 1157. (v. v.)

4. *A. PARHASSIFOLIA* (L. mant. 371.) foliis cordatis nervis longitudinalibus 9 notatis, capsulis aristatis.  $\mathcal{U}$  in Delphin., Bressiâ ex Latourr. — *Fl. fr.* 3. p. 189. — *Till. pis. t.* 46. f. 1. (v. s. c.)

## 438 MONOCOTYL. SEU ENDOGENÆ.

### Sect. II. DAMASONIUM *Juss. gen. p. 46.*

Capsulæ 6, 2-spermæ.

5. A. DAMASONIUM (L. sp. 486.) foliis cordato-oblongis, capsulis 6 stellatim patentibus subulatis.  $\mathcal{U}$  in stagnorum oris passim. — Fl. fr. 3. p. 188. Damasonium stellatam Rich. — Lob. ic. t. 301. f. 1. (v. v.)

### III. SAGITTARIA *L. gen. 1067.*

Monoica. Perigonium 6-partitum laciniis 3 exterioribus persistentibus calycinis, 3 interioribus coloratis petaloideis. *Masc.* Stamina circiter 24. *Fem.* Ovaria numerosa receptaculo globoso imposita; capsulæ compressæ marginatæ 1-spermæ.

1. S. SAGITTIFOLIA (L. sp. 1410.) foliis lanceolatis acuminatis sagittatis, lobis lanceolatis rectis, scapo simplici.  $\mathcal{U}$  in stagnis, fossis et ad oras fluminum Galliæ australis, mediæ et occidentalis. — Fl. fr. 3. p. 190. Lam. ill. t. 776. Specimina 5-8-pedalia foliis ovatis plerumque obtusis in Garumnâ propè Burdigalam observavit al. Des Moulins. (v. v.)

### Trib. III. JUNCAGINEÆ *Rich. l. c.*

Perigonium subuniforme rarò nullum, semen 1 aut 2 basi approximata erecta. Embryo rectus, radiculâ ad hylum seminis versâ. — Folia linearia rigida; flores spicati non colorati.

### IV. SCHEUCHZERIA *L. gen. 452.*

Stamina 6, antheris longis; ovaria 3-6; capsulæ compressæ inflatæ 2-valves 1-2-spermæ.

1. S. PALUSTRIS (L. sp. 482.) in paludosis turfosis Vogesorum, Jurassi et Alpium. — Fl. fr. 3. p. 191. Lam. ill. t. 268. (v. v.)

### V. TRIGLOCHIN *L. gen. 453.*

Perigonium caducum; stamina 6 brevissima; antheræ extrorsæ subsessiles; styli nulli; ovaria 3-6 conniventia; capsulæ 3-6 erectæ 1-spermæ conniventes.

1. T. MARITIMUM (L. sp. 482.) capsulis strictis 6-locularibus sulcatis ovatis.  $\mathcal{U}$  in maritimis, et Lotharingiæ pratis salis. — Fl. fr. 3. p. 192. Lam. ill. t. 270. f. 2. — Lob. ic. t. 16. f. 2. (v. v.)
2. T. PALUSTRE (L. sp. 482.) radice fibrosâ, scapo foliis subduplò longiore, capsulis strictis 3-locularibus lævibus ovato-linearibus basi attenuatis.  $\mathcal{U}$  in paludibus et pratis humidis. — Fl. fr. 3. p. 192. Lam. ill. t. 270. f. 1. — Lob. ic. t. 17. f. 1. (v. v.)
3. T. BARBELLINI (Lois. gall. 2. p. 725.) radice bulbosâ, scapo foliis longiore, capsulis 3-locularibus lævibus linearibus subpatentibus basi non attenuatis.  $\mathcal{U}$  in salis Mediterranei et Oceani propè la Teste de Buch (Des Moulins). — Fl. fr. 5. p. 313. T. palustre  $\beta$ . Fl. fr. 3. p. 192. — Barr. ic. t. 271. (v. v.)

Ordo XCVII. POTAMEÆ *Juss. dict. sc. nat. t. 43. p. 93.* —  
*Fluviales Vent. tabl. 2. p. 80. Rich. mém. du mus. 2. p. 365.*

Flores hermaphroditi aut dielines. Spatha perigonii vicem gerens, aut perigonium plus minus profundè divisum. Ovaria numero definita receptaculo communi aut spadici centrali inserta. Stylus 1 aut nullus; stigma simplex. Stamina numero definita receptaculo aut spadici inserta. Capsulæ indehiscentes 1-loculares 1-spermæ. Semen inversum pendulum. Albumen o. Embryo rectus aut incurvus. Radicula ad punctum hylo seminis contrarium versa. — *Herbæ aquaticæ; foliis simplicibus alternis aut rariùs oppositis. Flores axillares aut terminales, solitarii aut spicati.*

I. POTAMOGETON *Tourn. inst. 103. L. gen. 174.*

Flores hermaphroditi inserti spadici spathis 2 basi instructo; perigonium 4-partitum; antheræ 4 sessiles perigonii laciniis alternæ; ovaria 4; nuces 1-spermæ sessiles.

1. *P. NATANS* (L. sp. 182.) foliis natantibus petiolatis ellipticis lanceolatisve acutis aut obtusis basi rotundatis subcordatis, stipulis membranaceis lanceolato-linearibus acuminatis, spicis densis, nucibus lævibus ovato-compressis acutis.  $\mathcal{U}$  in aquis stagnantibus. — Fl. fr. 3. p. 183. Lam. ill. t. 89. (v. v.)  
*P. fluitans*, foliis elliptico-oblongis utrinque attenuatis. — *P. fluitans* Roth germ. 1. p. 72. Fl. fr. 5. p. 310. non fl. fr. 3. p. 184. (v. s.)
2. *P. OBLONGUM* (Viv. fragm. t. 2.) foliis omnibus petiolatis ovato-oblongis utrinque acutis, stipulis lineari-lanceolatis membranaceis acuminatis, spicis densis.  $\mathcal{U}$  in pratis humidis nec inundatis Andegaviæ, Soloniæ, Vogesorum. — Fl. fr. 5. p. 310. *P. plantago* East. An satis à priore distinctum? (v. s.)
3. *P. LUCENS* (L. sp. 183.) foliis immersis ovato-lanceolatis planis subsessilibus nitidis venosis acutis acuminatisve, stipulis lineari-lanceolatis subobtusis intermedio æqualibus, spicis densis.  $\mathcal{U}$  in lacubus et fluminibus. — Fl. fr. 3. p. 185. fl. dan. t. 195. — J. Bauh. hist. 3. p. 769. ic. (v. v.)
4. *P. OSCURUM* (DC. fl. fr. 5. p. 311.) foliis alternis subsessilibus obtusis ovali-oblongis multinerviis, summis subcoriaceis subimmersis, omnibus fusco-rufescentibus, spicis densis cylindraceis, nucibus ovatis subcompressis sublævibus breviter obtusè rostratis.  $\mathcal{U}$  in rivis Palatinatûs et probabiliter Alatiæ. — *P. rufescens* Koch. ined. (v. s.)
5. *P. TERRESTRIUM* (L. sp. 182.) foliis immersis ovatis oblongisve obtusis cordatis amplexicaulis nitidis nervosis, spicis densis brevibus.  $\mathcal{U}$  in stagnis, lacubus et fluviiis. — Fl. fr. 3. p. 185. — J. Bauh. hist. 3. p. 778. ic. (v. v.)
6. *P. CAMPESTRIS* (L. sp. 183.) foliis immersis lanceolatis alternis oppositisve undulatis, stipulis apice lacero-ciliatis.  $\mathcal{U}$  in fossis, rivis. — Fl. fr. 3. p. 186. eng. bot. t. 1012. — Lob. ic. t. 266. f. 2. (v. v.)
7. *P. DENSEM* (L. sp. 182.) foliis immersis ovatis lanceolatisve sessilibus acuminatis oppositis confertis, caulibus dichotomis, spicâ pauciflorâ, nucibus ovatis subcompressis rostratis.  $\mathcal{U}$  in rivis et fluminibus. — Fl. fr. 3. p. 185. eng. bot. t. 397. — J. Bauh. hist. 3. p. 777. ic. (v. v.)
8. *P. OPPOSITIFOLIUM* (DC. fl. fr. 3. p. 186.) foliis ovatis lanceolatisve oppositis sessilibus obtusis subundulatis, stipulis minimis integris, spicâ pauciflorâ

## 440 MONOCOTYL. SEU ENDOGENÆ.

- densâ, caulibus longè ramosis.  $\frac{1}{2}$  in rivis passim. — *P. serratum* L. sp. 183. An satis à priore distinctum? (v. s. sine fructu.)
9. *P. КЕТАОРНЫЛЛУМ* (Schreb. spic. p. 21.) foliis superioribus natantibus petiolatis ellipticis oblongo-acutis membranaceis utrinque attenuatis, inferioribus immersis confertis sessilibus sublinearibus aut oblongo-lanceolatis acuminatis, stipulis lineari-lanceolatis acuminatis, spicis densis, nucibus striatis ovatis subcompressis acutis.  $\frac{1}{2}$  in aquis stagnantibus circâ Parisios, Andegaviâ. — Fl. fr. 3. p. 184. *P. palustre* Teesd. trans. linn. 5. p. 43. (v. s.)
10. *P. VARIIFOLIUM* (Thore chl. 47.) foliis superioribus natantibus longè petiolatis lanceolatis basi attenuatis acuminatis, inferioribus axillaribusque linearibus, spicâ densâ.  $\frac{1}{2}$  in rivulis ericetorum Vasconiz. — Fl. fr. 5. p. 310. *P. fluitans* Fl. fr. 3. p. 184. excl. syn. Roth. et Willd. (v. s.)
11. *P. COMPRESSUM* (L. sp. 183.) caule compresso ramoso, foliis linearibus sessilibus apice rotundatis obtusis, spicis paucifloris breviter pedunculatis foliis brevioribus, ovariis confertis ovatis subcompressis obtusè breviter rostratis.  $\frac{1}{2}$  in aquis stagnantibus. — eng. bot. t. 418. *P. gramineum* L. sp. 184. Fl. fr. 3. p. 184. eng. bot. t. 2253. — Ray. syn. ed. 3. t. 4. f. 3. (v. v.)  
*β. cuspidatum*, foliis cuspidatis, spicis 3-4-floris. — *P. cuspidatum* Schrad. germ. v. 2. ex Sm. eng. flora t. 1. p. 234. *P. compressum* Fl. fr. 3. p. 186. — Less. pruss. t. 66. An distincta species? (v. s.)
12. *P. FUSILLUM* (L. sp. 184.) caule tereti tenuissimo ramoso, foliis linearibus oppositis alternisque distinctis basi patentibus, spicis interruptis, nucibus ovatis lævibus breviter obtusè rostratis. (1) in paludibus. — Fl. fr. 3. p. 187. eng. bot. t. 215. — Vaill. bot. t. 32. f. 4. (v. v.)
13. *P. FRACTINATUM* (L. sp. 183.) foliis setaceis approximatis parallelis basi vaginantibus alternis, 2 superioribus oppositis, spicâ florum pedunculatâ gracili valdè interruptâ, nucibus sublævibus ovatis compressis brevissimè rostratis.  $\frac{1}{2}$  in fossis, paludibus, fluminibus passim. — Fl. fr. 3. p. 187. eng. bot. t. 323. *P. marinum* L. sp. 184. Fl. fr. 3. p. 187. — Vaill. bot. t. 32. f. 5. (v. v.)

### II. RUPPIA L. gen. 175.

Flores hermaphroditi distichi in spadice solitario; perigonium caducum 2-valve; antheræ 4 sessiles reniformes 1-loculares; ovaria 4 per anthesin pedicello non visibili, post anthesin pedicello longissimo imposita; capsulæ 1-spermæ ovatæ stigmate persistente coronatæ.

1. *R. MARITIMA* (L. sp. 184.) (1) in stagnis et fossis maritimis. — Fl. fr. 3. p. 183. Gärtn. carp. 2. t. 84. f. 6. — Mich. gen. t. 35. Folia linearia basi vaginantia, flores axillares et terminales. (v. v.)

### III. ZANNICHELLIA Mich. gen. t. 34. f. 1. 2. L. gen. 1034.

Flores solitarii monoici. *Masc.* Stamen 1 nudum ad basin externam perigonii fl. feminei situm. *Fem.* Perigonium campanulatum; ovaria 2-6; capsulæ 1-spermæ sessiles compressæ gibbosæ extus crenulatæ.

1. *Z. PALUSTRIS* (L. sp. 1375.) antherâ 4-loculari, stigmatibus integerrimis, seminibus dorso denticulatis. (1) in rivis et fossis aquaticis. — Fl. fr. 3. p. 182. Gärtn. carp. 1. t. 19. f. 6. (v. v.)

### IV. ZOSTERA DC. fl. fr. 3. p. 154. — Zosteræ sp. L.

Flores monoici aut dioici, nidulantes in foliis spathæ vicem gerente, unilateraliter dispositi; spadix inclusus; stylus 2-fidus, nucs monospermæ.

1. *Z. MARINA* (L. sp. 1374.) caule tereti usculo, foliis integerrimis, floribus

- monoicis, staminibus et pistillis in duas series dispositis, antheris sessilibus.  $\Psi$  in Oceano et mari Mediterraneo. — Fl. fr. 3. p. 154. Lam. ill. t. 737. Z. maritima Gaertn. carp. 1. t. 19. (v. v.)
2. Z. MEDITERRANEA (DC. fl. fr. 3. p. 154.) caule teretiusculo, foliis integerrimis, floribus dioicis, antheris filamento exserto insertis.  $\Psi$  in mari mediterraneo propè Monspelium. — Phucagrostis major. Caul. in ann. Ust. 10. p. 42. t. 3. (v. v. DC.)

V. CAULINIA DC. fl. fr. 3. p. 156. — Zosteræ sp. L.

Flores hermaphroditi; perigonium nullum; stam. 3; filaments dilatata persistentia basi extus antherifera; ovarium 1-spermum; stigma subsessile; fructus baccatus; semen unicum altero parieti adnatum exalbuminosum; radícula maxima infera.

1. C. OCEANICA (DC. l. c.)  $\Psi$  in Oceano et mari Mediterraneo sub aquâ florens. — Zostera Oceanica L. mant. 123. Caul. in ann. Ust. 6. p. 66. t. 4. Posidonia Caulini Konig in ann. bot. 2. p. 96. Kerneræ oceanica Willd. — Lob. ic. t. 248. f. 2. (v. v. sine flore.)

VI. NAYAS L. gen. 1096.

Flores monoici. *Masc.* Perigonium bilobum aut nullum, stamen unicum apice quadrivalve. *Fem.* Perigonium nullum; ovarium ovoideum; stylus 1; stigmata 2-3; capsula ovoidea 1-sperma.

1. N. MINOR (Roth. germ. II. 2. p. 500.) foliis linearibus ad apicem caulis conferis recurvis, perigonio florum masculorum nullo. (I) in lacubus et fluminibus. — Fl. fr. 2. p. 587. Lam. ill. t. 799. f. 2. Caulinia fragilis Willd. sp. 4. p. 182. — Mich. gen. t. 8. f. 3. (v. v.)
2. N. MAJOR (Roth. germ. II. 2. p. 499.) foliis lanceolatis planis, perigonio florum masculorum bilobo. (I) in fluminibus et paludibus submersa. — Fl. fr. 2. p. 587. N. marina L. sp. 1441. N. monosperma Willd. — Lam. ill. t. 799. f. 1. — Mich. gen. t. 8. f. 1. 2. (v. v.)

Ordo XCVIII. ORCHIDÆ *Juss. gen. p. 64. R. Brown prod. nov. holl. p. 310. Rich. in ann. mus. iv.*

Perigonium 1-sepalum petaloideum ovario adhærens, sex-partitum laciniis irregularibus, 3 exterioribus superiore galeâ nominatâ, 3 interioribus, 5 superioribus florem quasi verè constituentibus, inferiore (*labello*) formâ semper directione sæpiùs diversâ. Ovarium simplex; stylus columnaris simplex, stigma viscosum plus minùs orbiculare ad basin latus aut apicem styli. Stam. filamenta 3 ovario inter stylum et perigonii lobos superiores inserta; lateralialia (excepto Cypripedio) sterilia, mox plus minùs elongata, mox brevissima aut nulla; filamentum medium styli dorso plus minùs adnatum antheriferum, rarò (et tunc filamenta lateralialia antherifera) sterile. Anthera 2-locularis loculis vel discretis et lateribus styli sæpè ultrâ producti adnatis, vel approximatis et tunc anthera mox stigmati parallela immobilis persistens, mox columnam terminans sæpiùsque mobilis operculiformis decidua. Pollen granulosum granulis nunc materiâ glutinosâ elasticâ cohærentibus, nunc faciliè solubilibus, nunc homogeneis et sic coëuntibus in massas, post dehisceniam loculorum, alterâ extremitate sæpiùs attenuatâ corpusculo glandulari-viscido (*retinaculo*) subpeltatim insertas et per hoc stigmati affixas, vel immediatè, vel ope processûs filiformis à stigmate oriundi. Capsula unilocularis trivalvis tricarinata inter nervos lateraliter dehiscens. Semina numerosa scobiformia placentis tribus mediis valvis adnatis imposita. Embryo ad basin albuminis carnosus. — Herbæ, radicibus nunc fasciculatis nunc tuberosis, tuberibus ovatis aut palmatis, caulibus rariùs divisus, mox squamis foliisque instructis, mox nudis, foliis integris amplexicaulibus, floribus bracteatis spicatis aut solitariis.

I. ORCHIS *Swartz act. holm. 1800. DC. fl. fr. 3. p. 245. — Orchidis et Satyrii sp. L.*

Perigonium personatum, 6-partitum, suprà fornicatum seu galeatum, rarissimè patens, labello basi calcarato; stigma convexum anticum; anthera 2-locularis terminalis; pollinis massulæ 2 profundè in particulas numero indefinitas angulatas divisæ, basis extremitate attenuatâ retinaculo adfixâ stigmati adnexæ.

§. 1. *Radice tuberculis palmatis vel fibris cylindricis constante.*

( \* *Nigritella* Rich. de orch. annot. p. 19. — Perigonium patens, labellum scrotoformi-calcaratum indivisum; massæ polliniceæ retinaculis 2 instructæ. )

1. O. NIGRA (All. ped. n. 1845.) foliis linearibus tenuissimè glanduloso-ciliatis,

- spicâ densâ ovatâ, bracteis inferioribus flores longè superantibus, superioribus subæquantibus, perigonii segmentis linearibus acutis patentibus, labello ovato acuminato indiviso, calcare obtuso scrotiformi.  $\mathcal{U}$  in pratis montosis. — Fl. fr. 3. p. 253. *Satyrium nigrum* L. sp. 1338. *Nigritella angustifolia* Rich. p. 34. — Hall. helv. t. 27. Fl. atropurpurei aut rosei. (v. v.)
2. *O. SUAVICOLANIS* (Vill. dauph. 2. t. 1.) foliis linearibus, spicâ densâ ovatâ, floribus semiresupinatis, perigonii segmentis patentibus, labello semi-ovato bidentato, calcare ovarii longitudine.  $\mathcal{U}$  in montibus Delphinatûs propè Gratianopolim ex Vill. — Fl. fr. 5. p. 331. Fl. rubelli. (v. s. DC. in herb. Vill.) †  
(\*\* *Gymnadenia* Rich. p. 20. — Perigonium fornicatum seu galeatum, labellum calcaratum 3-fidum; retinacula 2; stigma sursum non excavatum sed in processum fissum angustatum.)
3. *O. VIRIDIS* (All. ped. n. 1846.) foliis obtusis inferioribus ovatis, superioribus lanceolatis, spicâ cylindricâ elongatâ laxâ, bracteis inferioribus florem superantibus, perigonii galeati segmentis ovato-oblongis obtusis, labello lineari apice 3-dentato dentibus exterioribus longioribus, calcare obtuso scrotiformi.  $\mathcal{U}$  in pratis humidis. — Fl. fr. 3. p. 253. Sat. viride L. sp. 1337. eng. bot. t. 94. *Gymnadenia viridis* Rich. — Hall. helv. t. 26. Fl. viridescentes. (v. v.)
4. *O. ALBIDA* (All. ped. n. 1838.) foliis ovatis mucronatis, spicâ cylindricâ elongatâ densâ, bracteis ovatis acuminatis ovario longioribus, perigonii galeati segmentis ovatis obtusis, labello 3-partito, laciniis linearibus acutis, intermediâ obtusâ, calcare obtuso ovario duplò triplòve brevior.  $\mathcal{U}$  in pratis humidis Vogesorum, Jurassi, Alpium, montium Arvernæ, Pyren., Neustriæ ex cl. Le Prév. et de Bréb. — Fl. fr. 3. p. 253. Sat. albidum L. sp. 1338. *Gymnadenia albida* Rich. — Hall. helv. n. 1270. t. 26. (v. v.)
5. *O. ODORATISSIMA* (L. sp. 1335.) foliis linearibus acutis, spicâ cylindricâ elongatâ laxâ à foliis distante, bracteis ovarium superantibus, perigonii fornicati segmentis lateralibus patentibus, labello obtusè 3-lobo, calcare recurvo ovario brevior.  $\mathcal{U}$  in pratis montosis prov. austral. propè Gratianopolim ex Vill. Monspelim ex Gouan, in Pyr. centr. ex cl. Benth., circâ Parisios ex Mérat, Château-Gaillard juxtâ flumen Ais (cl. Auger). — Fl. fr. 3. p. 252. *Gymnadenia odoratissima* Rich. — Hall. helv. n. 1274. t. 29. — Seg. ver. 3. t. 8. f. 6. Fl. purpurei. (v. v. in Ital. D.)
6. *O. CONOPSEA* (L. sp. 1335.) foliis lineari-lanceolatis aut lato-linearibus, superioribus acuminatis, spicâ cylindricâ elongatâ laxâ à foliis distante, bracteis ovarium superantibus, perigonii fornicati segmentis lateralibus patentissimis, labello obtusè 3-lobo, calcare cornuto subulato ovario duplò longior.  $\mathcal{U}$  in pratis montosis. — Fl. fr. 3. p. 252. Fl. dan. t. 224. *Gymnadenia conopsea* R. Br. kew. 5. p. 191. — Hall. helv. n. 1287. t. 29. — Vaill. bot. t. 30. f. 8. Fl. purpurei. (v. v.)  
(\*\*\* *Morio*. — Orchis Rich. p. 19. — Perigonium fornicatum seu galeatum; labellum calcaratum; retinacula 2 in apice stigmatis sursum exavati nidulantia.)
7. *O. MACULATA* (L. sp. 1335.) foliis inferioribus lineari-lanceolatis subobtusis, superioribus linearibus acuminatis, bracteis ovarium superantibus, spicâ brevi densâ conico-cylindricâ, perigonii segmentis superioribus conniventibus, lateralibus patentibus, labello subplano 3-lobo, lobis lateralibus dentatis, medio integro acuminato, calcare cylindraceo ovario brevior.  $\mathcal{U}$  in pratis montosis et sylvis. — Fl. fr. 3. p. 252. Rich. l. c. — Hall. helv. t. 32. — Vaill. bot. t. 31. f. 9 et 10. Folia sæpè maculis nigris adspersa. Fl. rosei aut purpurei. (v. v.)
8. *O. LATIFOLIA* (L. sp. 1334.) caule folioso, foliis oblongo-lanceolatis acuminatis, spicâ cylindricâ densâ, bracteis inferioribus florem superantibus, superioribus subæquantibus, perigonii segmentis superioribus ovato-lanceolatis subobtusis conniventibus, labello vix 3-lobo lateribus reflexo dentato, calcare conico ovario brevior.  $\mathcal{U}$  in pratis humidis frequens. — Fl. fr. 3. p. 251. Fl. dan. t. 266. — Hall. helv. n. 1279. t. 32. — Vaill. bot. t. 31. f. 1-5. Folia inferiora sæpè maculata. Fl. purpurei. (v. v.)

9. *O. sambucina* (L. sp. 1334.) foliis lanceolatis lineari-lanceolatisve acutis, spicâ coarctatâ ovato-cylindricâ, bracteis flores subsuperantibus, perigonii segmentis obtusiusculis patulis, labello leviter 3-lobo crenato, calcare conico ovario brevior.  $\mathcal{U}$  in montosis provinciarum austral., Pyren., Arverniz, Alsatiz, propè Abbavillam ex Bouch. — Fl. fr. 3. p. 151. *O. incarnata* L. sp. 1335. Fl. ochroleuci aut rubescentes. (v. v.)

§. 2. *Radice tuberculis duobus integris instructâ.*

10. *O. fallens* (L. mant. 292.) foliis ovatis subobtusis, spicâ coarctatâ ovato-cylindricâ, bracteis ovarium superantibus, perigonii segmentis ovatis obtusis lateralibus patentibus, labello lato obovato subtrilobo lobis obtusis, medio submarginato, calcare conico ovarium subæquante.  $\mathcal{U}$  in sylvaticis Pyren., Alpium Galloprov., Delph., circâ Parisios? ex Thuil., Aureliam? ex Dub. — Fl. fr. 3. p. 250. — Hall. helv. n. 1281. t. 30. — Seg. ver. 3. t. 8. f. 3. Fl. ochroleuci. (v. v.)
11. *O. provincialis* (Balb. misc. alt. p. 20. t. 2.) foliis lanceolatis, spicâ laxâ, bracteis ovarium subæquantibus, perigonii segmentis erectis subpatentibus lanceolatis apice callosis obtusis, labello ovato-cuneato lato 3-lobo pubescente, laciniis 2 lateralibus deflexis, intermediâ minore emarginatâ, calcare ascendente ovarium superante.  $\mathcal{U}$  in Galloprovincia ad sylvam de l'Estèrel. — Fl. fr. 5. p. 329. Fl. ochroleuci. (v. s.)
12. *O. papilionacea* (L. sp. 1331.) foliis lanceolato-linearibus acutis, superioribus vaginantibus, spicâ brevi laxâ pauciflorâ, bracteis membranaceis coloratis ovario longioribus, perigonii segmentis nervosis conniventibus lanceolato-ovatis subobtusis, labello obovato lato indiviso dentato submarginato, calcare subulato ovario brevior.  $\mathcal{U}$  in Corsicæ sylvis et pratis sterilibus. — Fl. fr. 3. p. 249. Fl. purpurei. (v. s.)
- β. rubra*, labello trapeziformi acuto. — in Corsicâ et propè Lugdunum. *O. rubra* Jacq. coll. 1. p. 60. (v. s.)
13. *O. laxiflora* (Lam. fl. fr. 3. p. 504.) foliis lanceolatis lanceolato-linearibusve acuminate, bracteis ovarium subæquantibus, spicâ elongatâ laxissimâ, perigonii segmentis ovatis obtusis apice patentibus, labello 3-partito cuneato lato, laciniâ mediâ submarginatâ, 2 lateralibus crenulatis longioribus, calcare apice submarginato ovario duplò brevior.  $\mathcal{U}$  in prov. australibus, mediis occidentalisque. — Fl. fr. 3. p. 246. — Vaill. bot. t. 31. f. 33-34. — Tab. ic. 667. Fl. purpurei. (v. v.)
14. *O. mascula* (L. sp. 1333.) foliis lanceolato-linearibus lanceolatisve acuminate, spicâ elongatâ laxâ, bracteis ovarium superantibus, perigonii segmentis obtusis, 2 superioribus erectis patentibus, labello cuneato lato trifido crenulato, laciniis lateralibus ovatis obtusis, intermediâ longiore subbilobâ, calcare obtuso suberecto ovario subbrevior.  $\mathcal{U}$  in pratis. — Fl. fr. 3. p. 247. Fl. dan. t. 457. *O. parvifolia* Chaub. in St.-Am. agen. 369. t. 7. — Hall. helv. n. 1283. t. 33. — Vaill. bot. t. 31. f. 12. Fl. purpurei rariùs albi. Folia sæpè maculata. (v. v.)
15. *O. palustris* (Jacq. ic. rar. t. 181.) foliis oblongo-linearibus acuminate, spicâ brevi laxâ, bracteis ovarium superantibus, perigonii segmentis ovatis subacutis patentibus, labello obovato latissimo leviter trilobo, lobis subæqualibus obtusis, intermedio vix emarginato, calcare ovario subbrevior.  $\mathcal{U}$  in pratis paludosis Jurassi, Vogesorum et propè Falaise ex cl. de Brébisson. — Fl. fr. 5. p. 330. Fl. purpurei. An satis à priore distincta? (v. s.)
16. *O. morio* (L. sp. 1333.) foliis lineari-lanceolatis obtusis, superioribus vaginantibus acutis, spicâ laxâ, bracteis ovarium subæquantibus, perigonii segmentis ovatis obtusis ascendentibus, labello ovato-cuneato latissimo sub 4-lobo, laciniis 2 lateralibus crenulatis reflexis, calcare conico ascendente ovario subbrevior.  $\mathcal{U}$  in graminosis et collibus aridis. — Fl. fr. 3. p. 246. Fl. dan. t. 253. — Hall. helv. n. 1282. t. 33. — Vaill. bot. 31. f. 13 et 14. Fl. purpurei rariùs rosei aut albi. (v. v.)



17. *O. VARIEGATA* (Lam. dict. 4. p. 592.) foliis lanceolatis obtusiusculis, spicâ brevi subovatâ densâ, bracteis scariosis ovarium subæquantibus, perigonii segmentis lanceolatis acuminatis conniventibus, labello tripartito lævi, laciniis lateralibus acutis, intermediâ oblongâ bifidâ serratâ cum mucrone interjecto, calcare recto subulato ovario duplò brevior.  $\mathcal{L}$  in pratis. — Fl. fr. 3. p. 248. — Hall. helv. n. 1275. t. 30. Fl. rosei, purpureo punctati. (v. v.)
18. *O. MILITARIS* (L. sp. 1333. excl. var.) foliis lato-lanceolatis oblongis obtusis, superioribus vaginantibus acutis, spicâ laxiusculâ cylindricâ, bracteis membranaceis ovario multò brevioribus, perigonii segmentis ovatis conniventibus acutis, labello 3-partito punctato-scabro, laciniis lateralibus integris linearibus, mediâ cuneatâ latâ bifidâ lobis obsoletè 4-angularibus aut rotundatis integris vel crenulatis cum mucrone interjecto, calcare assecto ovario duplò brevior.  $\mathcal{L}$  in nemoribus et pratis umbrosis. — Fl. fr. 3. p. 248. Jacq. ic. rar. 3. t. 598. *O. fusca* Jacq. aust. 4. t. 307. — Hall. helv. n. 1276. t. 31. Fl. rubri aut fusc. (v. v.)
19. *O. GALEATA* (Lam. dict. 4. p. 593.) foliis lato-ovatis obtusis, superioribus vaginantibus, spicâ conicâ densâ, bracteis membranaceis obtusis ovario multò brevioribus, perigonii segmentis conniventibus acutis, labello 3-partito lævi, laciniis lateralibus integris sublinearibus obtusis, mediâ cuneatâ latâ 2-fidâ, lobis valdè divergentibus obsoletè 4-angularibus subintegris cum mucrone interjecto, calcare ovario duplò brevior.  $\mathcal{L}$  in pratis submontosis Lutetiz, Monspelii. — Fl. fr. 3. p. 249. — Hall. helv. n. 1277. t. 28. Fl. rubelli. (v. s.)
20. *O. LONGIBRACTEATA* (Biv. sic. n. 66. t. 4.) foliis infimis lato-ovatis obtusis, superioribus membranaceis vaginantibus acuminatis, bracteis membranaceis acuminatis florem superantibus, perigonii segmentis conniventibus ovatis obtusis, labello 4-fido, laciniis oppositis oblongis integris obtusis, calcare ovario duplò brevior.  $\mathcal{L}$  in collibus siccis Arelati, Teloni, Nicæ. — Fl. fr. 5. p. 330. *O. Robertiana* Lois. gall. t. 21. Perig. rubellum punctatum, labello atropurpureum. (v. s.)
21. *O. SIMIA* (Lam. fl. fr. 3. p. 507.) foliis lato-lanceolatis obtusis, superioribus vaginantibus acutis, spicâ ovatâ densâ demùm cylindricâ laxâ, bracteis ovario brevioribus obtusis, perigonii segmentis ovatis conniventibus acutis, labello 4-partito laciniis omnibus gracilibus linearibus obtusis, mucrone lineari interdùm apice projecto, calcare ovario duplò brevior.  $\mathcal{L}$  in pratis et sylvis siccis. — Fl. fr. 3. p. 249. *O. tephrosanthos* Vill. — Hall. helv. n. 1275. t. 30. — Vaill. bot. t. 31. f. 25 et 26. (v. s.)
22. *O. CORIOPHORA* (L. sp. 1332.) foliis lanceolato-linearibus linearibusve acutis, spicâ cylindricâ subdensâ, bracteis ovarium subæquantibus superantibusve, perigonii segmentis ovatis lanceolatisve acuminatis conniventibus, labello 3-partito, laciniis 2 lateralibus crenatis obtusis aut mucronatis, calcare conico ascendente ovario duplò triplòve brevior.  $\mathcal{L}$  in pratis humidis circâ Falaise ex cl. de Breb., Parisios, Burdigalam ex cl. Later., in Pyren., prov. anstr. et circâ Mimatem (Mende). — Fl. fr. 3. p. 246. Jacq. aust. t. 122. *O. divaricata* Thuill? — Hall. helv. t. 34. f. 2. — Vaill. bot. t. 31. f. 30-31-32. Fl. rubello et viridi conspersi. (v. v.)
23. *O. USTULATA* (L. sp. 1333.) foliis lanceolatis, spicâ ovato-cylindricâ, bracteis ovarium subæquantibus, perigonii segmentis erectis acutis apice subconniventibus, labello 3-partito, laciniis obtusis, lateralibus linearibus punctato-scabris, mediâ latiore subbilobâ, calcare scrotiformi ovario triplò quadruplòve brevior.  $\mathcal{L}$  in pratis. — Fl. fr. 3. p. 247. — Fl. dan. t. 103. — Hall. helv. n. 1273. t. 28. — Vaill. bot. t. 31. f. 35-36. Spica primum atropurpurea densâ, demùm basi laxa. Fl. parvi albo et purpureo variegati. (v. v.)
24. *O. GLOBOSA* (L. sp. 1332.) foliis lanceolatis, spicâ densâ ovato-globosâ, perigonii segmentis apice aristato-clavatis, labello 3-partito, laciniâ mediâ emarginatâ, calcare ovario duplò brevior.  $\mathcal{L}$  in pascuis Vogesorum, Jurassi,

## 446 MONOCOTYL. SEU ENDOGENÆ.

*Alpium*, Pyren. ad *Port de Paillères* ex Lapeyr. — Fl. fr. 3. p. 245. Jacq. aust. t. 265. — Hall. helv. t. 27. f. 1. Fl. rosei. (v. v.)

( \* \* \* \* *Anacamptis* Rich. p. 19. — Perigonium fornicatum; labellum calcaratum; retinaculum 1 binis pollinis massulis commune in apice stigmati sursum excavati nidulans. )

25. *O. PYRAMIDALIS* (L. sp. 1332.) foliis lanceolato-linearibus acutis, superioribus vaginantibus, apicā densā ovatā aut ovato-oblongā, bracteis linearibus acuminatis ovarium subæquantibus, perigonii segmentis subovatis acutis æqualibus, labello 3-partito lacinii integris obtusis, lateralibus latioribus, calcare ovarium subæquante. ♀ in pascuis siccis passim. — Fl. fr. 3. p. 246. eng. bot. t. 110. *Anacamptis pyramidalis* Rich. — Hall. helv. t. 35. f. 1. Fl. rubelli rariis albi. (v. v.)

( \* \* \* \* *Platanthera* Rich. p. 20. — Perigonium galeatum seu fornicatum, labellum calcaratum indivisum lineari-ligulatum; antheræ loculi ab invicem valde distantes; retinacula 2 nuda. )

26. *O. BIFOLIA* (L. sp. 1331.) foliis caulinis vaginantibus brevissimis; radicalibus binis ovato-oblongis acutis basi attenuatis, apicā elongatā laxissimā, bracteis ovarium subæquantibus, labello lineari integerrimo obtuso, calcare incurvo cornuto ovario duplō longiore. ♀ in nemorosis humidis et pratis umbrosis. — Fl. fr. 3. p. 245. *O. alba* Lam. *Platanthera bifolia* Rich. — Hall. helv. t. 35. f. 2. Fl. albi odori. (v. v.)

• ( \* \* \* \* *Loroglossum* Rich. p. 16. — Perigonium galeatum, labellum lineari-tripartitum, basi breviter saccato-calcaratum, retinaculum unicum binis massulis commune, in apice stigmati sursum excavati nidulans. )

27. *O. HIRCINA* (Crantz fasc. 6. p. 484.) foliis lanceolato-ovatis, apicā elongatā laxiusculā, bracteis linearibus florem superantibus, perigonii segmentis conniventibus ovatis obtusis, labello lineari-tripartito, laciniiā intermediā prælongā bifidā, calcare conico brevissimo scrotiformi. ♀ in pratis montanis, ad marginem sylvarum passim. — L. sp. 1337. eng. bot. t. 34. *Loroglossum hircinum* Rich. — Vaill. bot. t. 30. f. 6. — Moris. s. 12. t. 12. f. 11. n. 9. Fl. viridescens et purpureo striati. (v. v.)

- 28? *O. SECUNDIFLORA* (Bert. rar. it. dec. 2. p. 42.) foliis inferioribus ovato-lanceolatis mucronatis, superioribus linearibus vaginantibus, apicā densā cylindricā elongatā, floribus secundis, bracteis ovario duplō brevioribus, perigonii galeati segmentis lanceolatis acuminatis, labello lineari 3-partito, laciniiā mediā subtridentatā aut integrā, calcare conico brevissimo. ♀ in pinetis Corsicæ et Massiliæ (cl. Solier). — *Ophrys densiflora* Desf. ann. mus. 10. t. 16. Fl. viridescens. (v. s.)

### II. OPHRYS Sw. l. c. DC. fl. fr. 3. p. 254. — *Ophrydis* sp. L.

Perigonium 6-partitum, segmentis patentibus aut galeatis; labellum ealcaratum; stigma convexum anticum; anthera 2-localaris terminalis; pollinis massulæ 2 profundè in particulas numero indefinitas angulatas divisæ, basis extremitate attenuatā retinaculo adfixā stigmati adnexæ.

( \* *Aceras* R. Brown in hort. kew. ed. 3. t. 5. p. 191. — *Loroglossi* sp. Rich. — Perigonium galeatum; labellum lineari-tripartitum, laciniiā intermediā bifidā, basi gibbulosum; retinacula 2 in apice stigmati sursum excavati nidulans. )

1. *O. ANTHROPOPHORA* (L. sp. 1343.) caule folioso, foliis ovato-lanceolatis, apicā cylindricā laxiusculā, bracteis ovario paulō brevioribus, perigonii segmentis conniventibus, 3 externis ovato-lanceolatis, 2 internis linearibus, labello ovario longiore lineari-tripartito, laciniiā mediā elongatā 2-fidā. ♀ in pratis. — Fl. fr. 3. p. 255. Fl. dan. t. 103. *Loroglossum anthropophorum* Rich. p. 32. — Hall. helv. n. 1264. t. 23. — Vaill. bot. t. 31. f. 19. 20. Perigonium albo-lutescens, labellum sulfureo et ferrugineo variegatum. (v. v.)

(\*\* *Entomea* — Ophrys Brown. prod. 313. Rich. p. 19. — Perigonium patens; labellum prorsus ecalcaratum, convexum; retinacula 2 in apice stigmatis sursum excavati nidulantia.)

2. O. LUTEA (Cav. ic. 2. t. 160.) caule folioso, foliis ovali-oblongis, spicâ laxâ pauciflorâ (3-5), bracteis ovarium æquantibus superantibusve, perigonii segmentis obtusis, 3 exterioribus ovato-ellipticis, binis interioribus linearilanceolatis duplò brevioribus, labello pubescente obovato apice sub 3-lobo, lobis subæqualibus medio majore emarginato.  $\mathcal{L}$  in pratis Corsicæ, Monspeli, Arclatis, Massiliæ, Niceæ. — Fl. fr. 5. p. 331. — Moris. s. 12. t. 13. f. 9. n. 15. Labellum in ambitu et subtus ameniacum, in medio fuscum sub 2-lineatum. (v. v.)
  3. O. PSEUDO-SPECULUM (DC. fl. fr. 5. p. 332.) caule folioso, foliis ovali-oblongis, spicâ 2-4-florâ, bracteis ovarium superantibus, perigonii segmentis obtusis patentibus, 3 exterioribus ovato-ellipticis, binis interioribus linearilanceolatis brevioribus, labello tridentato medio glaberrimo margine villosa reflexo.  $\mathcal{L}$  in pratis Monspeli rariis. — Labellum margine flavum medio fuscum. An à priore distincta species? (v. v. DC.)
  4. O. MYONIS (Jacq. ic. rar. t. 71.) caule folioso, perigonii segmentis patentibus, 3 exterioribus lanceolatis obtusis, 2 interioribus linearibus brevissimis, labello pubescente 3-partito, laciniis lateralibus linearilanceolatis, mediâ oblongâ bilobâ longiore.  $\mathcal{L}$  in pascuis montosis. — Fl. fr. 3. p. 255. O. muscifera eng. bot. t. 64. — Hall. helv. n. 1265. t. 24. Labellum cæruleo maculatum. (v. v.)
  5. O. ARANIFERA (Sm. brit. 939.) caule folioso, foliis ovali-oblongis, superioribus acutis, spicâ laxâ pauciflorâ, perigonii segmentis patentibus, 3 exterioribus oblongis obtusis, 2 interioribus lanceolatis acutis brevioribus, labello lato obovato villosa cum lineis 2 glabris vix 3-lobo, lobo medio obovato emarginato.  $\mathcal{L}$  in pascuis cretaceis glareosis. — Fl. fr. 5. p. 332. eng. bot. t. 65. O. arachnites  $\beta$  Fl. fr. 3. p. 155. — Vaill. bot. t. 31. f. 15 et 16. Labellum fuscum. (v. v.)
  6. O. ARACHNITES (Hoffm. germ. 318.) caule folioso, foliis ovali-oblongis, superioribus acutis, spicâ laxissimâ pauciflorâ, perigonii segmentis patentibus, 3 exterioribus oblongis obtusis, 2 interioribus linearilanceolatis brevissimis, labello villosa 3-lobo, lobo medio obovato apice brevissime 3-lobo.  $\mathcal{L}$  in pratis et pascuis montanis. — Fl. fr. 5. p. 332. non 3. p. 255. — Hall. helv. n. 1266. t. 24. f. 1. — Vaill. bot. t. 30. f. 10. 11. 12. 13. Perigonii segmenta albescentia, labellum fuscum basi striatum. (v. v.)
  7. O. APIFERA (Sm. brit. 938.) caule folioso, foliis ovali-oblongis acutis, perigonii segmentis patentibus, 3 exterioribus ellipticis obtusis, 2 internis linearibus brevissimis, labello villosa 3-lobo, lobis lateralibus oblongis medio obovato elongato 3-lobo, laciniâ terminali recurvatâ subulatâ.  $\mathcal{L}$  in pratis et pascuis cretaceis. — Fl. fr. 5. p. 333. eng. bot. t. 383. — Vaill. bot. t. 30. f. 9. Perigonium roseum, labellum purpureo-fuscum, flavo-striatum. (v. v.)
- (\*\*\* *Chamorchis* Rich. p. 20. — Perigonium galeatum; labellum indivisum; retinacula 2; stigma sursum non excavatum, sed in fissum processum angustatum.)
8. O. ALPINA (L. sp. 1342.) caule nudo, foliis subradicalibus linearisetaceis, spicâ pauciflorâ abbreviatâ, bracteis ovario longioribus, labello lanceolato indiviso obtuso utrinque unidentato.  $\mathcal{L}$  in Alpibus Delphinatûs. — Fl. fr. 3. p. 254. Jacq. vind. t. 9. *Chamorchis alpina* Rich. — Hall. helv. t. 22. f. 1. Fl. viridescens. (v. s.)
- (\*\*\*\* *Harminium* Brown. hort. kew. 5. p. 191. — Perigonium subcampanulatum laciniis interioribus manifeste longioribus dissimilibus, labellum basi sæcctum plurifidum; retinacula 2, nuda.)
9. O. MONORCHIS (L. sp. 1342.) caule nudo, foliis 2-3 subradicalibus lanceolatis, spicâ elongatâ tenuissimâ laxâ, bracteis ovarii longitudine, perigonii segmentis lanceolato-ovatis obtusis, labello tripartito, laciniis linearibus latera-

## 448 MONOCOTYL. SEU ENDOGENÆ.

libus divaricatis. ♀ in pratis montanis. — Fl. fr. 3. p. 254. Fl. dan. t. 102. *Herminium monorchis* Br. Rich. — Hall. helv. n. 1262. t. 22. Fl. viridescen-tes. (v. v.)

### III. SERAPIAS Sw. l. c. DC. fl. fr. 3. p. 257. Rich. de Orch. ann. p. 19. — *Serapiadis* sp. L.

Perigonium 6-partitum, segmentis 5 superioribus cucullato-coalitis, 3 exterioribus majoribus, 2 mediis tenuissimis, labellum concavum acutum ecalcaratum, columna longè cuspidata; stigma concavum anticum; anthera 2-locularis terminalis; pollinis massulæ 2 profundè in particulas numero indefinitas angulatas divisæ, basis extremitate attenuatâ retinaculo unico binis massulis communi adfixâ stigmati ad-nexæ.

1. *S. LINGUA* (L. sp. 1344.) radice tuberculosâ, tuberculo altero pedicellato, foliis linearibus oblongis, floribus 2-4, bracteis flore brevioribus, labello tripartito, laciniis lateralibus ovatis obtusissimis erectis conniventibus, mediâ oblongâ lanceolatâ acuminatâ glaberrimâ dependente, callo plano. ♀ in pratis provinc. austral. et austro-occident. usque ad Nannetes. — Fl. fr. 3. p. 256 et 5. p. 333. — Seg. veron. 3. t. 8. f. 4. (v. v.)
2. *S. CORDIGERA* (L. sp. 1345.) tuberculis 2 sessilibus, foliis oblongo-lanceolatis, bracteis flore longioribus, floribus 4-8, labello tripartito, laciniis lateralibus obtusis erectis conniventibus, mediâ ovatâ acuminatâ disco pilosâ depen-dente, callo 2-gibboso. ♀ in pratis prov. austr. et austro-occident. usque ad Nannetes. — Fl. fr. 3. p. 256 et 5. p. 233. *S. lancifera* et *cordigera* St. Am. fl. agen. p. 278 et 279. t. 9. — Rudb. elys. 2. p. 240. f. 20. (v. v.)

### IV. NEOTTIA Sw. l. c. p. 226. DC. fl. fr. 3. p. 257. — *Satyrii* et *Orchidis* sp. L.

Perigonium 6-partitum, laciniis superioribus basi conniventibus apice liberis; labellum ventricosum canaliculato-amplexans; columna acute appendiculata; stigma anticum obliquum; anthera 2-locularis postica; pollinis massæ sessiles; retinacula nulla.

(\* *Goodyera* Brown in hort. kew. ed. 3. t. 5. p. 197. — Labellum inclusum indivi-sum infernè insigniter concavo-gibbum; pollinis massulæ in particulas angulatas divisæ.)

1. *N. REPENS* (Sw. l. c. p. 226.) foliis radicalibus ovatis petiolatis reticulatis obtusis, superioribus vaginantibus acutis caule terminali floribusque pubes-centibus, floribus secundis, bracteis lineari-lanceolatis acuminatis, labello segmentisque lanceolatis. ♀ in sylvaticis montanis. — Fl. fr. 3. p. 258. *Satyrium repens* L. *Goodyera repens* Br. Rich. — Hall. helv. t. 22. f. 3. (v. v.)
- (\*\* *Spiranthes* Rich. p. 20. — Ovarium apice obliquatum; perigonium in oblongum connivens; labellum inclusum obovale indivisum; stigma in processum angusta-tum; pollen granulosum.)
2. *N. SPIRALIS* (Sw. l. c.) foliis radicalibus ovato-oblongis acutis subpetiolatis, scapo radicali axillari, spicâ spirali, floribus secundis, bracteis ovatis acu-minatis ovarium superantibus, labello ovato margine crispatulo. ♀ in gra-minosis et collibus siccis. — Fl. fr. 3. p. 257. *Ophrys spiralis* L. *Spiranthes autumnalis* Rich. *Orchidastrum autumnale* Michel. gen. 30. t. 26. — Lob. ic. t. 186. f. 1. Autumno floret. (v. v.)
3. *N. ÆSTIVALIS* (DC. fl. fr. 3. p. 258.) foliis sublinearibus, caule florifero fo-lioso, spicâ spirali, floribus secundis, bracteis ovatis acuminatis ovarium superantibus, labello ovato margine crispatulo. ♀ in pratis humidis passim. — *Spiranthes æstivalis* Rich. — Hall. helv. t. 38. — Mich. gen. t. 26. f. 2. Æstate floret. (v. v.)

V. EPIPACTIS Sw. l. a. DC. fl. fr. 3. p. 258. — Serapiadis et Ophrydis sp. L.

Perigonium erecto-patens 6-partitum, labello integro vel lobato ecalcarato; stigma obliquum terminale antheræ anticum; anthera opercularis 2-locularis ovata margini postico styli adnata, post pollinis emissionem persistens; pollinis massulæ granulosa sessiles.

§. 1. *Labello lobato.*

(\* *Listera* Br. kew. ed. 3. t. 5. p. 201. — *Neottia* Rich. p. 20 non Sw. — Perigonium in globum laxè connivens; labellum patens; stigma in processum angustatum; anthera intra cavitatem columnæ stipitata.)

1. *E. CORDATA* (All. ped. n. 1851.) caule bifolio, foliis cordatis oppositis, labello 3-fido, laciniis lateralibus brevissimis, intermedia longissima 2-partita linearis. 2/ inter muscos et pinetos Vogesorum, Juraasi, Arvernæ, Pyren. — Fl. fr. 3. p. 261. eng. bot. t. 358. *Neottia cordata* Rich. *Listera cordata* Br. — Hall. helv. t. 22. f. 4. (v. v.)
2. *E. OVATA* (All. ped. n. 1850.) caule bifolio, foliis ovatis oppositis, labello 2-fido linearis segmentis obtusis triplò longiore. 2/ in sylvis et pratis umbrosis. — Fl. fr. 3. p. 261. *Neottia ovata* Rich. *Listera ovata* Br. — Moris. s. 12. t. 11. f. 1. (v. v.)
3. *E. NIDUS-AVIS* (All. ped. n. 1849.) caule aphylo squamis membranaceis acutis vaginato, labello obcordato segmentis obtusis duplò longiore. 2/ in umbrosis et sylvis passim. — Fl. fr. 3. p. 260. *Neottia nidus-avis* Rich. — Hall. helv. n. 1290. t. 37. — Lob. ic. t. 195. f. 1. Fl. fusc. (v. v.)

§. 2. *Labello apice integro.*

(\*\* *Cephalanthera* Rich. p. 21. — Ovarium sessile; perigonium erecto-connivens; labellum genitalia amplexans interruptum; stigma in processum non angustatum; anthera subrotunda imà tantùm parte dorso stigmati incumbens, cæterâ aperte supereminens; pollen granulis simplicibus.)

4. *E. PALLENS* (Sw. l. c. p. 232.) foliis oblongo-lanceolatis sessilibus, bracteis ovario longioribus, floribus erectis, labello obtuso segmentis perigonii brevioribus, ovarii glabris. 2/ in sylvis passim. — *E. lancifolia* Fl. fr. 3. p. 261. *Serapias grandiflora* eng. bot. t. 271. *Cephalanthera pallens* Rich. Fl. ochroleuci. (v. v.)
5. *E. ENSIFOLIA* (Sw. l. c. p. 232.) foliis lanceolatis acuminatis subdistichis, bracteis minutissimis subulatis, floribus erectis, labello obtuso segmentis perigonii duplò brevioribus, ovarii glabris. 2/ in sylvis et pascuis montosis. — Fl. fr. 3. p. 259. eng. bot. t. 494. *Cephalanthera ensifolia* Rich. Fl. albescentes; labellum purpureo-striatum. (v. v.)
6. *E. RUBRA* (All. ped. n. 1857.) foliis lanceolatis sessilibus, bracteis ovarium subsuperantibus, floribus erectis, labello acuto lineis elevatis undulatis perigonii segmentis brevioribus, ovarii glabris. 2/ in umbrosis montanis. — Fl. fr. 3. p. 260. eng. bot. t. 437. *Serapias rubra* L. mant. 490. — Hall. helv. n. 1299. t. 42. Fl. rubri. (v. v.)

(\*\*\* *Helleborine*. — *Epipactis* Rich. l. c. — Ovarium pedicellatum; perigonium patulum interruptum; stigma in processum apiculare angustatum; anthera subcordata postica; pollen granulis veluti 4 globulosis.)

7. *E. LATIFOLIA* (All. n. 1855.) foliis ovatis amplexicaulibus, bracteis inferioribus flore longioribus, floribus pendulis, labello acuminato perigonii segmentis brevioribus, ovarii pubescentibus. 2/ in umbrosis et sylvis. — Fl. fr. 3. p. 259. eng. bot. t. 269. *Serapias latifolia* L. mant. 490. — Hall. helv. n. 1297. t. 40. Fl. purpurei aut viridescens. (v. v.)

ℓ. *microphylla*, foliis angustioribus internodiis brevioribus. — In collibus siccis Monspelii, Andegaviæ, Parisiorum, etc. — Fl. fr. 5. p. 334. *Serapias microphylla* Ehr. S. parvifolia Pers. — Clus. hist. 1. p. 273. f. 1. n. III. (v. s.)

## 450 MONOCOTYL. SEU ENDOGENÆ.

8. *E. PALUSTRIS* (Crantz austr. 6. p. 462. t. 1. f. 5.) foliis lanceolatis amplexicaulibus, bracteis flore brevioribus, floribus pendulis, labello ovato crenato obtuso perigonii segmentis æquali, ovariis pubescentibus.  $\mathcal{L}$  in pratis paludosis frequens. — Fl. fr. 3. p. 259. eng. bot. t. 270. *Serapias longifolia* L. — Hall. helv. n. 1296. t. 39. Fl. viridescens purpureo-variegati. (v. v.)

VI. *MALAXIS* Sw. in act. holm. 1800. p. 223. DC. fl. fr. 3. p. 261. — *Liparis* Rich. p. 21.

Perigonium 6-partitum inversum, segmenta 5 laxè patula; labellum posticum concavum columnam basi amplexens; columna gibbosa antice excavata; anthera superposita marginalis caduca hemisphærica 2-locularis loculis simplicibus; pollen solidum massis bimassulatis.

1. *M. LÆSLEII* (Sw. l. c. p. 235.) foliis binis ovato-lanceolatis, scapo trigono, labello apice ovato recurvato.  $\mathcal{L}$  in paludibus circà Parisios, Falaise, Béthune, Dunkerque, Lille et Haguenau ex Nestl. — Fl. fr. 3. p. 262. Ophrys Læselii L. *Liparis Læselii* Rich. p. 38. — Læs. pruss. 180. t. 58. (v. v. in Helv.)

VII. *CORALLORHIZA* Hall. hist. 2. p. 159. R. Br. in hort. kew. ed. 3. t. 5. p. 209. Rich. l. c. p. 21. — *Cymbidii* sp. Sw. DC.

Ovarium breviter pedicellatum; perigonium in oblongum fornicatum; labellum ecalcaratum obovale infernè canaliculatum marginibus infernè crenulâ interruptis, introrsum prominentiis longitudinalibus percursum, supernè recurvo-patens; columna libera; anthera marginalis superposita subglobosa didyma 2-locularis, loculis bilocellatis subtransversim dehiscentibus; pollen solidum, massulis 4 distinctis.

1. *C. HALLERI* (Rich. l. c. p. 39.) caule vaginato aphylo, perigonii segmentis lanceolatis, binis inferioribus linearibus deflexis, labello oblongo aento.  $\mathcal{L}$  in sylvis Jurassi, Galloprov., Occitanæ. — *C. innata* Brown l. c. *Cymbidium Corallorhiza* Sw. l. c. Fl. fr. 3. p. 263. Ophrys Corallorhiza L. — Hall. helv. n. 1301. t. 44. (v. v.)

VIII. *LIMODORUM* Tourn. inst. 250. Rich. l. c. p. 20. — *Limodori* sp. Sw. l. c.

Ovarium pedicellatum non contortum; perigonium laxiuscule erecto-connivens; labellum ascendens longè calcaratum, partis inferioris marginibus imæ columnæ adnatum; columna longissima; anthera marginalis subcordata cernua; pollen granulosum sessile.

1. *L. ABORTIVUM* (Sw. l. c. p. 243.) aphyllum, caule vaginato, labello ovato undulato, calcare subulato ovarii longitudine.  $\mathcal{L}$  in umbrosis montanis. — Fl. fr. 3. p. 263. *Orchis abortiva* L. — Hall. helv. n. 1288. t. 36. (v. v.)

IX. *EPIPOGIUM* Gmel. sib. 1. t. 2. f. 2. Brown prod. 330. Rich. l. c. p. 20. — *Limodori* sp. Sw. DC.

Ovarium pedicellatum non contortum; perigonium deorsum patens; labellum posticum vesicario-calcaratum interruptum, parte inferiore veluti in binas aurículas obtusas porrectas antrorsum desinente; anthera stipitata marginalis ferè superposita subrotunda; pollinis massæ 2 oblongo-ovoides in subulam solidam sensim desinentes.

1. *E. GMELINI* (Rich. l. c. p. 36.) aphyllum, caule vaginato paucifloro, floribus pendulis, labello concavo, calcare ascendente ovato.  $\mathcal{L}$  in faginetis Jurassi, Delphin., Monspelli. — *Limodorum epipogium* Sw. l. c. p. 243. Fl. fr. 3. p. 264. *Satyrium epipogium* L. sp. 1338. (v. v. in Helv.)

X. CYPRIPEDIUM *L. gen.* 1015.

Ovarium non contortum; perigonium laxè et subcruciatim patens; lobellum maximum obtusum inflatum calceolatum; columna appendiculata stigma obtegens, antheræ 2 distinctæ laterales basi appendice lanceolato instructæ; pollen granulosum.

1. *C. CALCEOLUS* (*L. sp.* 1346.) caule folioso, columnæ lobo elliptico obtuso, labello perigonii segmentis brevioribus compresso.  $\mathcal{U}$  in pratis umbrosis aut montanis passim. — *Fl. fr.* 3. p. 264. *liliac.* 1. t. 19. — *Hall. helv.* t. 43. (v. v.)

Ordo XCIX. IRIDEÆ *Juss. gen.* 57.

Perigonium basi tubulosum, ovario adhærens, petaloideum, sexfidum aut sexpartitum, sæpè irregulare. Stamina 3 segmentorum perigonii externorum basi inserta. Antheræ lineares extrorsum dehiscentes. Ovarium unicum triloculare multiovulatum, ovulis 2-seriatis; stylus unicus aut nullus; stigmata 3 simplicia aut laciniata, membranacea aut petaloidea. Capsula 3-locularis, trivalvis, valvulis medio septiferis, polysperma. Semina angulo loculorum interno adfixa. Embryo intrâ albumen carnosum aut corneum. — Herbæ radicibus tuberosis, foliis alternis ensiformibus aut linearibus, spathis membranaceis bivalvibus unifloris.

I. IRIS *L. gen.* 59.

Perigonium sexpartitum, laciniis tribus externis majoribus patulis, tribus interioribus minoribus erectis; stamina distincta; stylus brevis lacinias tres maximas petaloideas sæpius emarginatas gerens; stigmata 3 duplicaturæ laciniarum petaloidearum facie infera sitæ.

§. 1. *Laciniis floris exterioribus intus basi barbatis.*

1. *I. GERMANICA* (*L. sp.* 55.) barbata, foliis ensiformibus acuminatis glabris falcatis caule multifloro brevioribus, spathis scariosis, tubo ovario longiore, segmentis externis rotundo-orbicularibus.  $\mathcal{U}$  ad muros et tecta passim. — *Fl. fr.* 3. p. 236. *Bull. herb.* t. 141. — *Moris. s.* 4. t. 5. f. 10. n. 4. *Fl. violacei.* (v. v.)
2. *I. FLORENTINA* (*L. sp.* 55.) barbata, foliis ensiformibus glabris, scapo 1-3-floro, floribus subsessilibus, segmentis externis obovatis integris obtusis.  $\mathcal{U}$  in muris Galloprovinciæ orientalis. — *Fl. fr.* 5. p. 328. *Red. ill.* t. 23. — *J. Bauh. hist.* 2. p. 719. *ic.* — *Lob. ic.* t. 59. *Fl. albi.* (v. s.)
3. *I. PUMILA* (*L. sp.* 56.) barbata, scapo unifloro foliis ensiformibus glabris acutis mucronatisve brevioribus, tubo extrâ spatham exserto.  $\mathcal{U}$  in sterilibus montosis provinc. australium, ad muros Fontisbleandi. — *Fl. fr.* 3. p. 237. — *Lob. ic.* t. 63. f. 1. et t. 65. f. 1. *Clus. hist.* 1. p. 226. *Fl. violacei aut pallidi.* Floret ineunte vere. (v. v. DC.)
4. *I. LUTRACENS* (*Lam. dict.* 3. p. 297.) barbata, scapo subunifloro foliis ensiformibus acutis glabris longiore, tubo spathis scariosis incluso, segmentis

## 452 MONOCOTYL. SEU ENDOGENÆ.

exterioribus oblanceolato-ovatis.  $\frac{1}{2}$  in maritimis propè Olbias, in Occitaniâ ex cl. Benth., in agro Ruscinonensi ex Lapeyr., propè Fontembleaudi ex Mèrat. — Fl. fr. 3. p. 237. Fl. lutescentes. Floret ad finem veris. (v. v. ex Valesiâ D.)

### §. 2. *Laciniis floris externis imberbibus.*

5. *I. PSEUDACORUS* (L. sp. 56.) imberbis, foliis ensiformibus caule multiflorum sequantibus superantibusve, spathis viridibus, perigonii segmentis alternis stigmatate minoribus, externis obovatis obtusissimis.  $\frac{1}{2}$  in stagnorum oris et fossis aquaticis. — Fl. fr. 3. p. 237. Bull. herb. t. 137. — Moris. s. 4. t. 6. f. 6. n. 11. Fl. lutei. (v. v.)
6. *I. VETIDISSIMA* (L. sp. 57.) imberbis, foliis ensiformibus acuminatis caulem uniaugulatum subsuperantibus, ovario trigono, angulis sulco bifidis, seminibus rubris carnis bacciformibus.  $\frac{1}{2}$  in dumetosis ad vias passim. — Fl. fr. 3. p. 238. Black. t. 158. — Lob. ic. t. 70. f. 1. Fl. atropurpurei. (v. v. DC.)
7. *I. KYTHIODES* (Ehr. beit. 7. p. 140.) imberbis, foliis ensiformibus canaliculato-subulatis, caule bifloro, spathis viridibus, ovario trigono acutè angulato, perigonii segmentis exterioribus obovato-rotundis.  $\frac{1}{2}$  in Pyrenæis editoribus. — Fl. fr. 3. p. 238. Red. lil. t. 212. Fl. cærulei. (v. v.)
8. *I. GRAMINEA* (L. sp. 58.) imberbis, foliis linearibus acuminatis caule compresso subbifloro longioribus, spathis viridibus, segmentis externis spathulatis, ovariis hexagonis.  $\frac{1}{2}$  in collibus circà Narbonam, Carcassonam, inter Rupellam et Rochefort ex Bon, in Vogesis propè Colmars ex Nestl. — Fl. fr. 3. p. 239. Jacq. aust. t. 2. — Lob. ic. t. 69. f. 1. Fl. violaceo-cærulei, segmentis exterioribus flavo striatis. (v. v. c.)
9. *I. PRATENSIS* (Lam. dict. 3. p. 300.) imberbis, caule subtrifloro tereti fistuloso subnudo foliis radicalibus linearibus angustissimis acuminatis longiore, spathis scariosis, perigonii segmentis externis oblanceolatis, ovariis trigonis.  $\frac{1}{2}$  in pratis humidis Delphinatûs, Alsatiæ. — Fl. fr. 3. p. 239. I. sibirica L. sp. 57. Jacq. aust. t. 3. — Moris. s. 4. t. 6. f. 8. n. 13. Fl. cærulei. (v. s.)
10. *I. SISYRINCHIUM* (L. sp. 59.) imberbis, foliis lineari-canaliculatis contortis caulem 3-5-florum superantibus, spathâ scariosa, segmentis externis ovatis.  $\frac{1}{2}$  in Corsicâ propè Bonifacio reperiit cl. Soleirol. — Cav. ic. 2. t. 193. Moræa Sisyrinchium bot. mag. t. 1407. I. fugax Ten. neap. p. 15. t. 14. non Pers. Petala dilutè cærulea basi flavo notata. (v. s.)
11. *I. SPURIA* (L. sp. 59.) imberbis, caule subtrifloro tereti folioso, foliis linearibus acuminatis, spathis viridibus, perigonii segmentis externis spathulatis, ovariis hexagonis.  $\frac{1}{2}$  in pratis Occitaniz, circà Narbonam et Elne in agro Ruscinonensi ex Benth. — Fl. fr. 3. p. 239. Jacq. aust. t. 4. — Lob. ic. t. 68. f. 2. Flores cæruleo, violaceo et albo variegati. (v. v. DC.)
12. *I. TUBEROSA* (L. sp. 58.) imberbis, caule unifloro foliis tetragonis breviorè, perigonii segmentis externis lanceolatis obtusissimis medio dilatatis scariosis.  $\frac{1}{2}$  circà Telonem, Aginnum, in Pictaviâ superiore ex cl. Desvaux. — Fl. fr. 5. p. 328. Red. lil. t. 48. Curt. bot. mag. t. 531. — Lob. ic. 98. Fl. viridissimalidi, laciniis exterioribus apice atropurpureis velutinis. (v. s.)

### II. GLADIOLUS Tourn. inst. 190. L. gen. 57.

Perigonium infundibuliforme, limbo 6-fido inæquali bilabiato; stigma trilobatum patens; semina arillata.

1. *G. COMMUNIS* (L. sp. 52.) flore subringente, spathis tubo longioribus, floribus secundis, foliis ensiformibus nervosis.  $\frac{1}{2}$  in arvis Galliz australis, ad radices Pyrenæorum, in Aquitaniâ ex Laterr., Andegaviâ ex Bast. — Fl. fr. 3. p. 240. — Lob. ic. t. 99. f. 1. (v. v.)



III. IXIA *L. gen. 56.*

Perigonium tubo plus minusve elongato, limbo patente 6-partito aequali; stigma 3-partitum, laciniis filiformibus patulis sæpè bifidis.

1. *I. Bulboconium* (*L. sp. 5r.*) foliis linearibus canaliculatis angulatis, scapo unifloro, floribus campanulatis, tubo brevissimo.  $\mathcal{U}$  in maritimis Corsicæ, Occitanicæ, Vasconicæ, Armoricæ, Neustriæ (cl. Le Prévost). — *Fl. fr. 3. p. 241. Lam. ill. t. 31. f. 1.*

*a. grandiflorum*, scapis subgeminatis, stylo staminibus longiore, floribus majoribus. — *I. Bulbocodium B. Red. lil. t. 88. Romulea Bulbocodium Seb. et Mauri. fl. rom. p. 17. — Col. ecph. 2. t. 7. f. 1. Lob. ic. 141. Fl. ex albo ad roseum et rubro-violaceum variant. (v. v.)*

*β. parviflorum*, scapo subsolitario, stylo staminibus brevior, floribus minoribus. — *I. Bulboc. A. Red. lil. 88. Rom. Columnæ Seb. et Mauri rom. 18. — Col. ecphr. 1. p. 327. Lob. ic. 142. Flores ex albo-lutescentes. (v. v.)*

IV. CROCUS *Tourn. inst. 183. L. gen. 55. Gay in Féruss. bull. sc. nat. jul. 1827. p. 346.*

Perigonii tubus gracilis limbo duplò longior, limbus 6-partitus æqualis erectus; stigmata 3 crassa convoluta sæpius cristata.

1. *C. sativus* (*All. ped. n. 310.*) spathâ duplici, perigonii fauce densè barbata, stigmate tripartito longitudine perigonii pendulo, foliis cum flore nascentibus enerviis margine revolutis ciliolatis.  $\mathcal{U}$  ex Oriente ortus, in agris cultus. — *Fl. fr. 3. p. 241. C. sativus a L. Lam. ill. t. 30. f. 1. C. autumnalis. eng. bot. t. 343. (v. v. c.)*

2. *C. multividus* (*Ram. bull. phil. n. 41. t. 8. f. 1.*) stigmate trifido multifido erecto perigonio subæquali, foliis ante flores nascentibus revolutis.  $\mathcal{U}$  in Corbariensis, Pyrenæis, in agro Syrtico propè Dax. — *Fl. fr. 3. p. 242. C. nudiflorus Smith. eng. bot. t. 491. Fl. violacei. (v. s.)*

3. *C. vernus* (*All. ped. n. 309.*) spathâ simplici, perigonii fauce pilis longis barbata, stigmate trifido erecto perigonio brevior, foliis cum flore nascentibus linearibus planis.  $\mathcal{U}$  in pratis Alpium, Cebennarum, Pyrenæorum. — *Fl. fr. 3. p. 242. excl. var. γ. eng. bot. t. 344. C. sativus β. L. sp. 50. Fl. albi aut violacei. (v. v.)*

4. *C. minimus* (*Red. lil. 2. t. 81.*) spathâ plerumque simplici, perigonii fauce glabrâ, stigmate trifido erecto perigonio brevior, foliis ante flores nascentibus angustissimis filiformibus.  $\mathcal{U}$  in Corsicâ. — *Fl. fr. 3. p. 243. C. nanus DC. syn. gall. p. 108. Fl. violacei albo variegati. (v. s.)*

5. *C. varicolor* (*Gawl. in bot. mag. t. 1110.*) spathâ duplici, perigonii fauce glabrâ, stigmate erecto, foliis cum flore nascentibus canaliculatis nervosis, capsulâ sex-striatâ.  $\mathcal{U}$  circâ Nicæam et in Galloprovinciâ orientali ex cl. Gay. l. c.

## 454 MONOCOTYL. seu ENDOGENÆ.

Ordo C. AMARYLLIDEÆ *R. Brown. prod.* 196. — *Narcissi Juss. gen.* 54. *excl. gen.* — *Liliacearum tribus* 3<sup>a</sup> *DC. fl. fr.* 3. *p.* 229. *excl. gen.*

Perigonium monophyllum tubulosum, ovario adhærens, 6-partitum, æstivatione imbricatâ, laciniis tribus exterioribus interiores equitantibus. Stamina 6 filamentis liberis aut coalitis; antheræ introrsæ. Ovarium inferum 3-loculare, loculis polyspermis; stylus simplex; stigma trilobum. Capsula trivalvis, valvis medio septiferis, polysperma; aut bacca 1-3-sperma. Albumen carnosum. Embryo rectiusculus. Radicula umbilicum spectans. — Radices bulbosæ aut fibrosæ. Flores umbellati aut solitarii.

### I. AMARYLLIS *L. gen.* 400.

Perigonium infundibuliforme 6-partitum, fauce squamulis 6 instructum; stigma trifidum; stamina 6 sæpius declinata aut inæqualia.

1. *A. LUTEA* (*L. sp.* 420.) spathâ indivisâ obtusâ, flore sessili, perigonio campanulato erecto basi breviter tubuloso, staminibus erectis, alternis brevioribus.  $\mathcal{V}$  in pratis Monspelii, insulæ Nigri-Monasterii. — *Fl. fr.* 3. *p.* 229. *lil.* 3. *t.* 148. — *Lob. ic. t.* 147. *f.* 2. (*v. s.*)

### II. PANCRACTIUM *L. gen.* 404.

Perigonium infundibuliforme, limbo patente 6-partito; stamina 6 apici tubi inserta, filamenta coalita ope coronæ seu membranæ subcylindricæ; stigma simplex.

1. *P. MARITIMUM* (*L. sp.* 418.) foliis lineari-lanceolatis, spathâ multiflorâ, coronæ sinibus bidentatis.  $\mathcal{V}$  in arenosis et rupibus maritimis prov. Mediteraneanarum et austro-occidentalium. — *Fl. fr.* 3. *p.* 230. *lil.* 1. *t.* 8. *Salicab. trans. linn.* 2. *t.* 9. (*v. v.*)
2. *P. ILLYRICUM* (*L. sp.* 418.) foliis lanceolatis, spathâ multiflorâ, coronæ sinibus dentatis, dentibus acutis bifidis patentibus.  $\mathcal{V}$  in Corsicâ, et arenosis maritimis oræ meridionalis insulæ *Ré* ex *Moris*. — *Fl. fr.* 5. *p.* 319. *bot. mag.* 718. *Red. lil.* 1. 153. — *Clus. hist.* 1. *p.* 168. *f.* 1. (*v. s.*)

### III. NARCISSUS *L. gen.* 403.

Perigonium infundibuliforme, limbo patente 6-partito; corona (nec-tarium *L.*) ad faucem petaloidea cylindrica vel campanulata, integra aut divisa; stamina 6 tubo inserta, intrâ coronam latitantia.

#### Sect. 1. *Pseudo-Narcissi DC. fl. fr.* 5. *p.* 319.

Folia plana glaucescentia, scapus uniflorus; tubus brevis obconicus, corona campanulata dentata.

1. *N. PSEUDO-NARCISSUS* (*L. sp.* 414.) foliis planiusculis glaucescentibus lato-linearibus obtusis, scapo striato subcompresso ancipiti unifloro, flore in spathâ subsessili, coronæ campanulatæ limbo patulo undulato incisio lacinias perigonii concoloris planas ovatas patentes æquante.  $\mathcal{V}$  in sylvis pratisque. — *Fl. fr.* 3. *p.* 231. *N. major Curt. bot. mag.* 51 et 1301. *Fl. fr.* 5. *p.* 320. (*v. v.*)

2. *N. minor* (L. sp. 415.) foliis planiusculis glaucescentibus lato-linearibus obtusis, scapo ancipiti exstriato unifloro, flore in spathâ longè pedicellato, coronâ obconicâ subsexfidâ dentatâ laciniis oblongo-lanceolatis paulò longiore.  $\mathcal{N}$  in Pyrenæis et ericetis circâ Dax. — Fl. fr. 5. p. 320. Curt. bot. mag. t. 6. — Barr. ic. 975. — Rudb. elys. 2. p. 82. f. 11. (v. s.)
3. *N. moschatvus* (L. sp. 415.) foliis planiusculis glaucescentibus lato-linearibus obtusis, scapo ancipiti exstriato unifloro, flore in spathâ subsessili, coronæ cylindricæ limbo erecto repando-crenato perigonii cernui laciniis ovatas lanceolatasve æquante.  $\mathcal{N}$  Hispaniæ et Lusitanie indigenus, in hortis cultus. — Fl. fr. 5. p. 320. bot. mag. 1300. *N. candidissimus* Red. lil. 4. t. 188. *N. bicolor* L. sp. 415. *N. Hispanicus* Gou. ill. 23. — Barr. ic. t. 945. 946. 953. 954. — Moris. s. 4. t. 8. f. 9. — Lob. ic. t. 117. f. 1. 2. Variat flor. candidis, bicoloribus aut flavis. (v. v. c.)
4. *N. incomparabilis* (Mill. dict. n. 3.) foliis planiusculis glaucescentibus lato-linearibus obtusis, scapo subcylindrico ancipiti unifloro, flore in spathâ subsessili, tubo subcylindrico, coronâ campanulatâ erectâ margine crenulatâ subsexlobatâ laciniis ovatis dimidio brevior.  $\mathcal{N}$  in pratis Occitanie, Galloprov. — Fl. fr. 5. p. 321. Curt. bot. mag. t. 121. *N. Gouani* Red. lil. 4. t. 220. — Barr. ic. t. 927. 928. 931. 932. — Moris. s. 4. t. 8. f. 8. (v. v.)

Sect. II. *Poetici* DC. fl. fr. 5. p. 321.

Folia planiuscula glaucescentia, scapi 1-aut-multiflori; florum tubus cylindricus elongatus; corona brevis rotata margine scarioso crenato.

5. *N. poeticus* (L. sp. 414.) foliis planiusculis glaucescentibus lato-linearibus obtusis obtusè carinatis, scapo subcylindrico striato ancipiti 1-floro, coronâ brevissimâ rotatâ margine scarioso crenulato discolore, perigonii laciniis ovatis.  $\mathcal{N}$  in pratis. — Fl. fr. 3. p. 230. Bull. herb. t. 306. *N. angustifolius* bot. mag. t. 193. (v. v.)
6. *N. biflorus* (Curt. bot. mag. t. 197.) foliis planiusculis glaucescentibus lato-linearibus acutè carinatis, scapo ancipiti 1-3-floro, coronâ rotatâ brevissimâ crenulatâ margine concolore crispo, laciniis lato-ovatis.  $\mathcal{N}$  in pratis Armoraciæ, Andegaviæ, Occitanie, Galloprov. — Fl. fr. 5. p. 321. eng. bot. t. 276. *N. orientalis* L. mant. 62. non Ait. — Dod. pemp. 223. f. 2. (v. v.)

Sect. III. *Tazette* DC. l. c.

Folia planiuscula glaucescentia, scapus multiflorus; florum tubus tereti-elongatus, corona cupularis integra aut subdentata.

7. *N. tazetta* (L. sp. 416.) foliis planiusculis glaucescentibus lato-linearibus obtusis, scapo subcylindrico lævi subancipiti 3-10-floro, coronâ cyathiformi integrâ ore constrictâ laciniis subtriplo brevior, perigonii discoloris laciniis lanceolato-ovatis acutis tubo brevioribus.  $\mathcal{N}$  in pratis prov. austral. — Fl. fr. 5. p. 322. 3. p. 231. excl. var.  $\alpha$ . et  $\gamma$ . bot. mag. t. 925. — Dod. pempt. 224. f. 1. — Bar. ic. t. 943, 944, 918, 919. (v. v. DC.)
8. *N. polyanthus* (Lois. narc. 36.) foliis planiusculis subvirescentibus lato-linearibus obtusis, scapo subcylindrico ancipiti 8-20-floro, coronâ cyathiformi subintegerrimâ laciniis ovatis alternè latioribus subtriplo brevior.  $\mathcal{N}$  in pratis circâ Telonem, Nicæam ex cl. Spreng. — Fl. fr. 5. p. 323. — Lob. hist. 60. ic. — Barr. t. 915. — Moris. s. 4. t. 8. f. 2. (v. s.)
9. *N. stellatus* (DC. fl. fr. 3. p. 323.) foliis planiusculis glaucescentibus lato-linearibus obtusis, scapo compresso ancipiti 3-10-floro, pedicellis angulatis, coronâ cyathiformi denticulatâ ore vix constrictâ laciniis oblongis triplo brevior.  $\mathcal{N}$  circâ Telonem ex Lois. — *N. niveus* et *N. subalbidus* Lois. narc. 37. — Barr. ic. t. 916 et 962. — Rudb. elys. p. 50. f. 2. p. 52. f. 1x. (v. s. ex Neap.)
10. *N. chrysanthus* (DC. fl. fr. 5. p. 323.) foliis planiusculis rigidis subglaucescentibus lato-linearibus obtusis, scapo 3-10-floro, coronâ cyathiformi inte-

## 456 MONOCOTYL. seu ENDOGENÆ. NARCISSUS.

gerrimā ore subconstictā laciniis lanceolatis ovatis apiculatis discoloribus triplō breviorē.  $\mathcal{L}$  circa Grasse. — Barr. ic. t. 961. (v. s.)

11. *N. nusus* (Gou. ill. 22.) foliis planiusculis glaucescentibus linearibus obtusis, scapo lævi obtuse compressissimo 2-6-floro, coronā cyathiformi denticulatā ore vix constrictā laciniis ovalibus obtusis duplō breviorē.  $\mathcal{L}$  in rupestribus Monspelii, Gallopr. infer. — Fl. fr. 5. p. 324. Red. lil. t. 429. *N. compressus* Haw. — C. Bauh. prod. 27. ic. — Rudb. elys. 2. p. 51. f. 2. (v. s.)
12. *N. patulus* (Lois. narc. 34.) foliis subcanaliculatis angustis glaucescentibus patulis, scapo subcylindrico 2-6-floro, coronā cyathiformi integrā laciniis ovatis alterne latoribus dimidio breviorē.  $\mathcal{L}$  in Stæchadum insulis, circa Massiliam (Solier.). — Fl. fr. 5. p. 324. — Rudb. elys. 2. p. 52. f. 2. (v. s.)
13. *N. calathinus* (L. sp. 415.) foliis linearibus obtusis planiusculis glaucescentibus, scapo subcylindrico 2-4-floro, coronā campanulatā subintegrā laciniis lanceolatis subreflexas tubo longiores æquante.  $\mathcal{L}$  in Armoracis insulis Glenans. — Fl. fr. 5. p. 324. Red. lil. t. 177. — Rudb. elys. 2. p. 60. f. 5. (v. s.)

### Sect. iv. *Bulbocodii* DC. fl. fr. 5. p. 325.

Folia semi-cylindrica virentia, scapi nui-aut-multiflori; corona ampla obconica laciniis floris angustas æquans aut subsuperans.

14. *N. Bulbocodium* (L. sp. 417.) foliis semi-cylindricis angustissimis virentibus, scapo cylindrico lævi unifloro, coronā turbinatā integrā laciniis linearis lanceolatis acutis subsuperante.  $\mathcal{L}$  in Pyrenæis, ericetis Bayonæ, Aquarium - Tarbellicarum. — Fl. fr. 3. p. 231. lil. 1. t. 24. — Lob. ic. t. 118. f. 1, 2. (v. s.)

### Sect. v. *Junquillæ* DC. l. c.

Folia subulata aut semi-cylindræa plus minùve canaliculata, scapi sæpiusimè multiflori; corona campanulata segmentis duplō triplōve brevior.

15. *N. saxosinus* (Clus. hist. 1. p. 162. L. sp. 417.) foliis subulatis, scapo gracili cylindrico 1-oo-floro, tubo elongato laciniis subduplō longiore, coronā subcrenulatā brevissimā, laciniis oblongo-lanceolatis mucronatis.  $\mathcal{L}$  in Corsicā propè Calvi (cl. Soleir.). — Desf. atl. 1. t. 82. — Lob. ic. 122. (v. s.)
16. *N. Junquilla* (L. sp. 417.) foliis subulato-semicylindræis viridibus, scapo cylindrico 2-6-floro, coronā cyathiformi ore dilatatissima integrā laciniis ovatis triplō breviorē, tubo laciniis duplō longiore.  $\mathcal{L}$  in asperis Galloprov. et Occitanis. — Fl. fr. 3. p. 232 et 5. p. 325. Bull. herb. t. 334. — Moris. s. 4. t. 8. f. 6. (v. v. c.)
17. *N. intermedius* (Lois. gall. 191. t. 7.) foliis canaliculatis basi subsemi-cylindræis viridibus, scapo subcylindrico 1-3-floro, coronā cyathiformi ore dilatatā crispā subintegrā laciniis lato-ovatis quadruplō breviorē.  $\mathcal{L}$  in Pyrenæis, Bayonæ. — Fl. fr. 5. p. 325. Red. lil. t. 427. — Moris. s. 4. t. 8. f. 5. (v. s.)
18. *N. ocheroleucus* (Lois. narc. 38.) foliis canaliculatis dorso semi-cylindræis, scapo subcylindrico lævi 4-8-floro, coronā cyathiformi integerrimā laciniis ovato-subrotundis alterne latoribus dimidio breviorē.  $\mathcal{L}$  in agris circa Telonem ex Lois. — Fl. fr. 5. p. 326. †
19. *N. odoratus* (L. sp. 416.) foliis linearibus semi-cylindræis canaliculatis viridibus, scapo cylindrico 1-5-floro, coronā campanulatā 6-lobatā lævi laciniis ovatis dimidio subbreviorē.  $\mathcal{L}$  in pratis provinc. occid. et australium. — Fl. fr. 5. p. 326. Red. lil. t. 157. *N. calathinus* Gawl. bot. mag. t. 934. non L. — Clus. hist. 1. p. 158. f. 1. (v. v. DC.)
20. *N. lævis* (Salisb. prod. 224.) foliis linearibus canaliculatis apice planiusculis viridibus, scapo 1-3-floro, coronā campanulatā subsexfidā repandā crispulā

lacinii lanceolato-ovatis dimidio ferè breviorè.  $\mathcal{Y}$  circà Grasse. — Fl. fr. 5. p. 327. Red. lil. t. 428. N. odorus bot. mag. t. 78. non L. (v. s.)

IV. LEUCOIUM L. gen. 402.

Perigonium tubo brevi, limbo campanulato sexpartito æquali, lacinii apice crassiusculis; stigma simplex.

§. 1. *Stylo apice clavato.*

1. L. VERNUM (L. sp. 414.) spathâ uniflorâ.  $\mathcal{Y}$  in pratis humidis et umbrosis montium. — Fl. fr. 3. p. 233. Lam. ill. t. 230. f. 1. (v. v.)

2. L. ÆSTIVUM (L. sp. 414.) spathâ multiflorâ.  $\mathcal{Y}$  in pratis umbrosis Gallie australis. — Fl. fr. 3. p. 233. Lam. ill. t. 230. f. 2. eng. bot. t. 621. (v. v. DC.)

§. 2. *Stylo filiformi.*

3. L. AUTUMNALE (L. sp. 414.) spathâ monophyllâ biflorâ pedicello supremo breviorè, perigonii lobis ovalibus obtusis, internis apice 3-dentatis stamina duplò triplòve superantibus.  $\mathcal{Y}$  in saxosis Corsicæ, Nicæ, Mompelii ex Gouan. — Fl. fr. 3. p. 233. lil. t. 150. f. 2. — Ren. spec. t. 100. (v. s.)

4. L. HYEMALE (DC. fl. fr. 5. p. 327.) spathâ diphyllâ 1-2-florâ pedicellis longiorè, perigonii lobis lanceolatis acutis integris stamina paulò superantibus.  $\mathcal{Y}$  in saxosis.

a. *alba*, flore albo. — Nicæ. (v. s.)

β. *rosea*, flore roseo. — In Corsicâ. — L. roseum Lois. gall. t. 8. L. tri-chophyllum DC. syn. 1986\*. Red. lil. t. 150. f. 1. non Schousb. (v. s.)

V. GALANTHUS L. gen. 401.

Perigonium sexpartitum, laciniiis tribus interioribus externis duplò brevioribus emarginatis; stigma simplex.

1. G. NIVALIS (L. sp. 413.)  $\mathcal{Y}$  in pratis umbrosis montanis. — Fl. fr. 3. p. 234. Lam. ill. t. 230. — Moris. s. 4. t. 9. f. 11. n. 23. Scapus 1-florus. Perigonii segmenta 3 externa oblònga alba, interna viridantia. (v. v.)

## 458 MONOCOTYL. seu ENDOGENÆ.

Ordo CI. ASPARAGÆE *Juss. gen. 40. DC. fl. fr. 3. p. 172. — Asparagineæ* *Ach. Rich. in dict. class. 2. p. 20.*

Flores hermaphroditi monoici aut dioici. Perigonium petaloideum, liberum aut ovario adhærens sæpius 6-partitum interdum 6-fidum aut 4-8-partitum. Stamina tot quot perigonii laciniae, iisdem basi adhærentia; filamenta libera, rarissime (in Rusco) in urceolum coalita. Ovarium unicum 3-loculare 1-aut- $\infty$ -ovulatum, ovulis angulo interno adfixis. Stylus 1-4-5; stigmata 3-4. Bacca aut capsula sphaerica 3-4-locularis abortu aliquando unilocularis, loculis 1-3-spermis. Embryo in albumine carneo aut carnoso, sæpius minimus. — Herbæ aut suffrutices habitu vario; folia non vaginantia interdum verticillata, rarius ad axillam stipulae nascentia. Flores ad axillam bracteolæ siti, variè dispositi.

### † FLORES HERMAPHRODITI; OVARIUM LIBERUM.

#### I. ASPARAGUS *Tourn. inst. 154. L. gen. 424.*

Perigonium liberum 6-partitum, stamina 6; bacca trilocularis loculis dispermis rarius abortu monospermis.

1. *A. OFFICINALIS* (L. sp. 448. excl. var.  $\beta$ .) caule herbaceo tereti erecto ramossimo, foliis setaceis fasciculatis, stipulis minimis ovatis acuminatis, floribus sæpius abortu diclinis, pedunculis medio articulatis.  $\mathcal{U}$  in arenosis maritimis Galliae australis et austro-occidentalis, in hortisque cultus. — Fl. fr. 3. p. 173. — Black. 1. 33a. Fasciculi foliis 4-6 constant. (v. v.)
2. *A. AMARUS* (DC. cat. monsp. 81.) caule herbaceo tereti erecto subscabro, foliis rigidis setaceis fasciculatis, stipulis spinescentibus patulis subuncinatis, floribus abortu diclinis, pedunculis medio articulatis.  $\mathcal{U}$  in arenosis maritimis Monspeli, Aquarum - mortuarum. — Fl. fr. 5. p. 309. Red. lil. t. 446. — Clus. hist. 2. p. 179. ic. Fasciculi foliis 4-6 constant. (v. v. DC.)
3. *A. TENUIFOLIUS* (Lam. dict. 1. p. 294.) caule herbaceo tereti erecto ramossimo, foliis tenuissimis fasciculatis, floribus hermaphroditis, pedunculis longis sub flore articulatis.  $\mathcal{U}$  in pratis umbrosis et montosis Cebennarum, Avenionis, Gratianopolis ex cl. Mirbel. — Fl. fr. 3. p. 173 et 5. p. 309. Fasciculi foliis 10-20 constant. (v. v. DC.)
4. *A. ACUTIFOLIUS* (L. sp. 449.) caule fruticoso inermi angulato, foliis aciformibus brevibus rigidulis perennantibus mucronatis aequalibus.  $\mathcal{U}$  in sterilibus et lapideis Galliae mediterraneæ. — Fl. fr. 3. p. 173. — Clus. hist. 2. p. 178. f. 1. (v. v.)
5. *A. ALBUS* (L. sp. 449.) caule fruticoso spinoso tereti flexuoso, aculeis solitariis rectis, foliis fasciculatis triquetris obtusis deciduis, floribus pedunculatis aggregatis, baccis 1-2-spermis.  $\mathfrak{f}$  in Corsicâ propè Ajaccio detexit cl. Soleirol. — Clus. hist. 179. ic. — Moris. s. 1. t. 1. f. 3. (v. s.)

#### II. STREPTOPUS *Mich. fl. bor. amer. 1. p. 200. DC. fl. fr. 3. p. 174. — Uvulariæ sp. L.*

Perigonium 6-partitum, laciniis foveâ basi nectariferâ instructis; stamina 6; antheræ filamentis longiores; bacca lævis, cortice tenui.

1. *S. AMPLEXICAULIS* (DC. l. c.) foliis amplexicaulibus cauleque glabris.  $\frac{1}{2}$  in Vogesi, Jurasco, Alpibus, montibus Arvernæ, Pyrenæis. — *Uvularia amplexifolia* L. *Convallaria dichotoma* Thib. in Pers. ench. t. p. 373. Fl. fr. 5. p. 309. — Barr. t. 719 et 720. (v. v.)

III. *PARIS* Lin. gen. 500.

Perigonium expansum 8-partitum, laciniis 4 externis latioribus calycem, 4 internis angustioribus corollam mentientibus; stam. 8; antheræ mediis filamentis adnatæ; stigmata 4; bacca 4-locularis, loculis 6-8-spermis.

1. *P. QUADRIFOLIA* (L. sp. 527.)  $\frac{1}{2}$  in nemorosis. — Fl. fr. 3. p. 175. — Tournef. t. 117. — Moris. s. 13. t. 3. f. 15. n. 6. (v. v.)

- IV. *CONVALLARIA* Roth. germ. 1. p. 147. DC. fl. fr. 3. p. 175. — *Convallariæ* sp. L.

Perigonium globosum vel cylindricum 6-dentatum; stamina 6; bacca globosa ante maturitatem maculata 3-locularis loculis 1-spermis.

Sect. 1. *POLYGONATUM* Tourn. inst. 14.

Flores cylindricæ.

1. *C. VERTICILLATA* (L. sp. 451.) caule recto, foliis verticillatis ellipticis.  $\frac{1}{2}$  in umbrosis. — Fl. fr. 3. p. 175. — Clus. hist. 1. p. 277. f. 1. (v. v.)  
2. *C. POLYGONATUM* (L. sp. 451.) glabra, caule ancipiti, foliis ovato-lanceolatis alternis amplexicaulibus, pedunculis axillaribus subunisporis.  $\frac{1}{2}$  in nemorosis et dumetis. — Fl. fr. 3. p. 176. Fl. dan. t. 377. (v. v.)  
3. *C. MULTIFLORA* (L. sp. 450.) glabra, caule tereti, foliis alternis amplexicaulibus ovato-ellipticis, pedunculis axillaribus multifloris.  $\frac{1}{2}$  in nemorosis et dumetis. — Fl. fr. 3. p. 176. *C. latifolia* Hoffm. germ. 3. p. 162. Fl. fr. 3. p. 176. non Jacq. — Clus. hist. 1. p. 275. f. 2. (v. v.)  
*β. bracteata*, floribus bracteis suffultis. — In Vogesi propè Thann ex el. Ræper. (v. s. ex Helv.)

Sect. II. *COILOCRINON*. — *Lilium Convallium* Tourn. inst. 14.

Flores campanulati.

4. *C. MAJALIS* (L. sp. 451.) scapo nudo, foliis ovato-lanceolatis, floribus racemoso-spicatis, pedicellis nutantibus.  $\frac{1}{2}$  ad sepes, in nemorosis. — Fl. fr. 3. p. 177. Lam. ill. t. 248. — Cam. sp. 618. ic. (v. v.)  
V. *MAYANTHEMUM* Roth. germ. 1. p. 70. DC. fl. fr. 3. p. 177. — *Smilacinae* sp. Desf. — *Convallariæ* sp. L.

Perigonium 4-partitum laciniis patentibus; stamina 4; bacca sæpius ante maturitatem maculata 2-locularis, loculis 1-spermis.

1. *M. BIFOLIUM* (DC. fl. fr. 3. p. 177.) foliis alternis cordatis subpetiolatis.  $\frac{1}{2}$  in sylvis montanis. — *M. Convallaria* Roth. l. c. 2. p. 196. *Convallaria bifolia* L. sp. 452. Fl. dan. t. 291. — Moris. s. 13. t. 4. f. 10. n. 7. (v. v.)

†† FLORES DIOICI; OVARIVM LIBERUM.

VI. *SMILAX* L. gen. 1120.

Perigonium campanulato-patens, 6-partitum. Masc. Stamina 6, distincta. Fem. Ovarium 1; styli 3; stigmata 3; bacca globosa 3-locularis.

460 MONOCOTYL. SEU ENDOGENÆ.

1. *S. ASPERA* (L. sp. 1458.) caule subdumoso spinoso hinc indè flexo, foliis cordiformibus dentato-aculeatis. ♂ in provinciis mediterraneis et propè Bayonam. — Fl. fr. 3. p. 178. — Clus. hist. 1. p. 112. f. 2. p. 113. f. 1. (v. v.)
2. *S. MAURITANICA* (Poir. itin. 2. p. 363.) caule scandente subspinoso, foliis cordatis subinermibus. ♂ in Corsicâ et insulis Stæchadum. — Fl. fr. 3. p. 179. *S. excelsa* Duh. arb. ed. nov. 1. t. 54. excl. syn. (v. s.)

VII. RUSCUS *Tourn. inst.* 15. *L. gen.* 1139.

Perigonium 6-partitum sæpius expansum; filamenta coalita in tubum apud femineos flores nudum, apud mares antheras 6 gerentem; stylus 1; stigma 1; bacca globosa 3-locularis, loculis 2-spermis.

1. *R. ACULEATUS* (L. sp. 1474.) caule rigido ramoso, foliis suprà floriferis nudis. ♂ in umbrosis. — Fl. fr. 3. p. 180. *Lam. ill.* t. 835. — *Lob. ic.* t. 637. f. 2. (v. v.)
2. *R. HYPOGLOSSUM* (L. sp. 1474.) caule flexuoso subramoso, foliis nitidis medio floriferis, glomerulis florum basi foliolo acuto instructis. ♂ in lapidosis et sterilibus circà Nicæam ex All. — Fl. fr. 3. p. 180. — *Lob. ic.* t. 638. f. 1. (v. s. c.)

+++ FLORES DIOICI; OVARIUM ADHÆRENS.

VIII. TAMUS *Lin. gen.* 1119. — *Tamnus Tourn. inst.* 28.

Perigonium campanulatum 6-partitum, in maribus hexandris patens, in femineis ovario adhærens et suprà constrictum; stylus 1; stigmata 3; bacca 3-locularis.

1. *T. COMMUNIS* (L. sp. 1458.) foliis cordatis indivisis. ♀ ad sepes et in nemorosis. — Fl. fr. 3. p. 181. *Lam. ill.* t. 817. (v. v.)



Ordo CII. LILIACEÆ DC. *theor. elem. ed.* 1. p. 249. — *Lilia et Asphodeli Juss. gen.* — *Liliacearum tribus I et II. DC. fl. fr.* 3. p. 198.

Flores hermaphroditi. Perigonium petaloideum liberum sæpè tubulosum 6-fidum aut 6-partitum, segmentis duplici serie dispositis. Stamina 6 perigonii laciniis opposita et cum iisdem sæpius adnata. Ovarium 1 liberum sessile trigonum multiovulatum, ovulis longitudinaliter 2-seriatis. Stylus unicus longitudinaliter trisulcatus, rariùs nullus. Stigmata 3 aut 1 triquetrum. Capsula 3-ocularis 3-valvis, valvulis medio septiferis. Semina plurima angulo loculi interno adfixa, tegumento crustaceo membranaceo aut spongioso instructa. Embryo intrà albumen carnosum aut cartilagineum. Radicula ad hylum versa. — Herbæ nunc bulbosæ, foliis radicalibus, nunc caulescentes foliis caulinis. Folia vaginantia aut sessilia, nervis sæpissimè parallelis simplicibus. Flores spathacei aut bracteati, racemosi aut umbellati.

Trib. I. *TULIPACEÆ*. — *Liliaceæ Juss. gen.* 48.

Semina plana, stigmata 3.

I. *TULIPA* Tourn. *inst.* 199. *L. gen.* 415.

Perigonium campanulatum 6-petalum aut 6-partitum laciniis basi non nectariferis; stigmata crassa sessilia; capsula oblonga trigona.

\* *Bulbis lanuginosis.*

1. *T. SUAVEVOLENS* (Roth. cat. 1. p. 45.) foliis lato-lanceolatis, caule unifloro foliorumque paginâ superiore pubescentibus, flore erecto, perigonii segmentis ovatis acutis glabris.  $\mathcal{L}$  in hortis culta. — *Fl. fr.* 3. p. 199. *Red. lil.* 1. 111. — *Tulipa pumilio* Lob. *ic. t.* 127. (v. v. c.)
2. *T. GESSNERIANA* (L. sp. 438.) foliis lanceolatis undulatis, caule unifloro glabro, flore erecto, staminum filamentis glabris, perigonii segmentis obovatis obtusis glabris.  $\mathcal{L}$  in hortis culta, spontanea in montibus Nicææ (Stire). — *Fl. fr.* 3. p. 200. *Lam. ill. t.* 244. — *Moris. s.* 4. t. 17. f. 1. 2. Flores mirum in modum colore variant. (v. s. ex mont. Nicæensibus.)
3. *T. OCLUS-SOLIS* (St. Am. rec. soc. agric. agen. 1. p. 75. et fl. agen. p. 145. t. 3.) caule unifloro glabro, foliis oblongo-lanceolatis, inferioribus lato-lanceolatis undulatis, segmentis ovali-lanceolatis tribus exterioribus acuminatis apice glabris, interioribus apice obtusiusculis.  $\mathcal{L}$  in arvis cultis circà Agiunum, Mompelium, in Galloprovinciâ et circà Nicæam (Stire). — *Fl. fr.* 3. p. 200. *Lil. 1. t.* 60. Segmenta rubra basi maculâ atro-cæruleâ margine flavâ notata. (v. s.)
4. *T. CLUSIANA* (DC. in *Red. lil.* 1. t. 37.) caule unifloro glabro, foliis linearibus acuminatis glabris, infimo vaginante, flore erecto, perigonii segmentis lanceolatis exterioribus acutis.  $\mathcal{L}$  in vineis circà Burdigalam ex cl. Des Moulins, Mompelium ex cl. Benth., Telonem, Grasse, Nicæam (Stire). — *Fl. fr.* 5. p. 314. — *Clus. cur. post. p.* 9. *ic.* Segmenta externa dorso purpurea, margine albida, interna alba ungue purpureo. (v. s.)

\*\* *Bulbis lanugine carentibus.*

5. *T. CALYSTAWA* (DC. in Red. lil. 1. t. 38.) caule unifloro glabro, foliis lanceolato-linearibus acuminatis canaliculatis, flore erecto, perigonii segmentis oblongo-lanceolatis acutis apice glabris, staminibus basi hirsutis. ♀ in pratis provinciarum mediterranearum. — Fl. fr. 5. p. 313. Fl. flavi. An sequentis varietas. (v. s.)
6. *T. SYLVESTRIS* (L. sp. 438.) caule unifloro glabro, foliis lato-linearibus acuminatis, flore subnutante, segmentis lato-lanceolatis acutis apice barbatis, staminibus basi hirsutis. ♀ in pratis montanis paschum. — Fl. fr. 3. p. 199. fl. dan. t. 375. Fl. flavi. (v. v.)

## II. FRITILLARIA L. gen. 411.

Perigonium campanulatum 6-partitum, laciniis basi foveâ nectariferâ ovatâ excavatis.

Sect. I. MELEAGRIS. — *Fritillaria Lin. hort. cli. 119. Juss. gen. 49.*

Fovea nectarifera oblonga. — Radix 2-tuberculata, flores terminales.

1. *F. MELEAGRIS* (L. sp. 436.) foliis omnibus alternis linearibus canaliculatis, caule unifloro. ♀ in pascuis humidis provinciarum occidentaliûm, Alpium, Jurassi raribus. — Fl. fr. 3. p. 201. Lam. ill. t. 245. f. 1. — Moris. s. 4. t. 18. f. 1. 2 et 3. Fl. tesserati. (v. v.)
2. *F. PYRENAICA* (L. sp. 436.) foliis linearibus canaliculatis inferioribus oppositis, caule 2-3-floro rarius unifloro. ♀ in Pyrenæis, montibus Delphin. Gallop. — Fl. fr. 3. p. 201. Fl. violacei, minus tesserati. An satis à priori distincta? (v. v.)

Sect. II. PETILIUM *Lin. hort. cli. 119. — Imperialis Juss. gen. 49.*

Fovea nectarifera rotunda. — Radix tuberosa tunicata, caulis apice foliosis, flores axillares cernui.

3. *F. IMPERIALIS* (L. sp. 435.) racemo comoso infernè nudo, floribus reclinatis. ♀ ex Oriente orta, in hortis culta. — Fl. fr. 3. p. 201. Lam. ill. t. 245. f. 2. — Moris. s. 4. t. 19. f. 2 et 4. (v. v. c.)

## III. LILIUM *Tourn. inst. 195. L. gen. 410.*

Perigonium campanulatum, laciniis 6 profundis rectis aut extus revolutis, suprâ sulco longitudinali fimbriato vel nudo ad basin excavatis.

### Sect. I. LILIASTRUM.

Flores campanulati, segmentis sessilibus.

1. *L. CANDIDUM* (L. sp. 433.) foliis lanceolatis sparsis undulatis, floribus pedunculatis terminalibus intus glabris. ♀ in hortis cultum, in Jurasso forte spontaneum. — Fl. fr. 3. p. 202. Black. herb. t. 11. — Moris. s. 4. t. 21. f. 1. (v. v. c.)
2. *L. SULCIFERUM* (L. sp. 433.) foliis sparsis lineari-lanceolatis planis, floribus campanulatis erectis terminalibus intus scabris. ♀ in Alsatiæ, Delphin., Galloprov. montosis humidis, in Alpibus et Pyrenæis rariis, propè Noyon (cl. Graves). — Fl. fr. 3. p. 202. Jacq. aust. t. 226. Red. lil. t. 210. — Lob. ic. t. 164 et 165. Fl. crocei nigro maculati. (v. v.)

### Sect. II. MARTAGON.

Perigonia revoluta.

3. *L. POMPONIUM* (L. sp. 434.) foliis sparsis lineari-subulatis, infimis lanceo-

- latis obtusis, floribus paucis intus papilloso-dentatis, segmentis lineari-lanceolatis acutis.  $\mathcal{L}$  in montibus Delph. ex Villars, Gallopr. ex Gér. — Fl. fr. 3. p. 202. lil. t. 7. Fl. coccinei. (v. s. ex mont. Nicænsis.)
4. *L. PYRÆNAICUM* (Gou. obs. 25.) foliis sparsis lineari-lanceolatis, floribus paucis subumbellatis, segmentis lineari-lanceolatis obtusis.  $\mathcal{L}$  in Pyrenæis rariùs. — Fl. fr. 3. p. 203. — Clus. hist. app. p. 256. f. 1 et 2. Fl. flavi. (v. v. DC.)
5. *L. MARTAGON* (L. sp. 435.) foliis verticillatis ovato-lanceolatis, floribus cernuis.  $\mathcal{L}$  in sylvis montanis. — Fl. fr. 3. p. 203. Jacq. aust. t. 351. Red. lil. t. 146. — Moris. s. 4. t. 20. f. 2. n. 7. Fl. rubelli aut albicantes maculati. (v. v.)

Trib. II. *ASPHODELEÆ*. Juss. gen. 51.

Semina rotundata vel angulosa, testâ crustaceâ.

## † STIGMATA TRIA.

IV. *ERYTHRONIUM* L. gen. 414. DC. fl. fr. 5. p. 313.

Perigonium campanulatum patentissimum 6-partitum, laciniis petaloideis, tribus interioribus basi intus bicallosis; stylus 1 elongatus; stigmata 3; capsula globosa basi angustior; semina rotundata.

1. *E. DENS-CANIS* (L. sp. 437.) foliis lato-lanceolatis maculatis, perigonii segmentis oblongo-ellipticis integerrimis.  $\mathcal{L}$  in umbrosis montanis Bennesiæ, Crisæ-Desæ Vocontiorum, Nicæ (Stire), Monselpii, Pyrenæorum prope Bayonam ex Lois. — Fl. fr. 3. p. 197. Lam. ill. t. 244. f. 1. — Moris. s. 4. t. 5. f. 1 et 2. (v. v.)

## †† STIGMA UNICUM.

V. *ASPHODELUS* Tourn. inst. 178. L. gen. 421.

Perigonium patens 6-partitum; staminum filamenta basi dilatata fornicato-recurvata ovarium obtegentia; capsula globosa, loculis monospermis.

1. *A. LUTEUS* (L. sp. 443.) caule folioso, foliis triquetris striatis lineari-subulatis, perigonii segmentis sublinearibus. ② ex Sicilia ortus, in hortis cultus. — Fl. fr. 3. p. 204. Jacq. vind. t. 7. (v. v. o.)
2. *A. RISTULOSUS* (L. sp. 444.) caule nudo supernè subramoso, foliis strictis subulatis striatis subfistulosis, perigonii segmentis lineari-ellipticis.  $\mathcal{L}$  in Galliâ australi. — Fl. fr. 3. p. 204. Cav. ic. 3. t. 201. (v. v.)
3. *A. RAMOSUS* (Willd. sp. 2. p. 133.) caule nudo ramoso ramis strictis erectis, foliis ensiformibus carinatis lævibus, floribus subsessilibus racemoso-spicatis confertis, bracteis membranaceis lanceolatis acuminatis pedicellos superantibus.  $\mathcal{L}$  in rupestribus provinc. Mediterranearum. — Fl. fr. 3. p. 204. *A. ramosus* & L. sp. 444. — Clus. hist. 1. p. 196. f. 2. (v. v.)
4. *A. ALBUS* (Willd. sp. 2. p. 133.) caule nudo simplici, foliis lineari-ensiformibus carinatis lævibus, floribus pedunculatis racemosis, pedicellis confertis longitudine bractearum, capsulis rugosis.  $\mathcal{L}$  in Alpibus Galloprovinciæ, Pyrenæis et in prov. occident. — Fl. fr. 3. p. 205. *A. ramosus*  $\beta$  L. — Clus. hist. 1. p. 197. f. 2. (v. v.)

VI. *CZACKIA Andreiosky ex. mon.*

Perigonium campanulatum 6-partitum, segmentis unguiculatis; filamenta declinata; stigma 3-lobum; capsula subhexagona; semina angulata.

## 464 MONOCOTYL. sive ENDOGENÆ.

1. *C. LILIASTRUM* (Andr. l. c.).  $\mathcal{U}$  in pascuis Alpium et Pyrenæorum. — *Hemerocallis Liliastrium* L. ed. 1. p. 324. Fl. fr. 3. p. 206. *Anthericum Liliastrium* L. sp. 445. — *Moris. s. 4. t. 1. f. 7. n. 8.* Flores subsecundi albi, folia plana scapum simplicem subsequantia. (v. v.)

### VII. PHALANGIUM *Tourn. inst. 193. Juss. gen. 52.*

Perigonium sexpartitum plus minusve patens; staminum filamenta recta filiformia sæpissimè glabra, laciniarum basi inserta; stigma simplex; capsula dorso 3-sulcata sub 3-gona ovato-globosa; semina 3-quetra 2-seriata. — Radices fasciculatæ.

1. *P. LILIAGO* (Schreb. spic. 36.) caule simplicissimo monophyllo, foliis radicalibus planis canaliculatisve angustis acuminatis basi scariosis vaginantibus, floribus racemosis, pedicellis bracteis lanceolatis subulatas superantibus subæquantibusve, pistillo declinato, perigonii patuli segmentis lanceolatis.  $\mathcal{U}$  in sylvis montosis herbosis. — Fl. fr. 3. p. 210. Lam. ill. t. 240. f. 2. *Anthericum Liliago* L. sp. 445. — *Moris. s. 4. t. 1. f. 9. n. 10.* (v. v.)
  2. *P. RAMOSUM* (Lam. dict. 5. p. 250.) caule ramoso, foliis linearibus angustis acuminatis, floribus laxè paniculatis, bracteis subulatis minimis, perigonii patentis segmentis lanceolatis, pistillo erecto, capsulæ loculis 2-spermis.  $\mathcal{U}$  in montana sterilibus. — Fl. fr. 3. p. 210. *Anthericum ramosum* L. — Lob. ic. t. 47. f. 2. (v. v.)
  3. *P. BICOLOR* (DC. fl. fr. 3. p. 209.) caule nudo spice ramoso, foliis planis linearibus acutis scapum superantibus, floribus laxè paniculatis, bracteis pedicellis brevioribus, perigonii segmentis lineari-ellipticis, filamentis pubescentibus, capsulæ loculis 2-spermis.  $\mathcal{U}$  ad sepes, in arenosis Galliæ occident., Galloprovinc. circà Olbias ex Ziz. Cannas (Stire). — *Anthericum bicolor* Desf. atl. 1. t. 90. *A. planifolium* L. mant. 442. Fl. intus albi, extus purpurei. (v. v.)
- β. canaliculatum*, foliis canaliculatis tortilibus scapo brevioribus. — In agro Syrtico. (v. v.)
4. *P. SEROTINUM* (Lam. dic. 5. p. 241.) caule 1-floro 3-4-folio, foliis crassiusculis subulato-linearibus, radicalibus caulem superantibus, perigonii patentis segmentis ovato-lanceolatis, capsulâ oblongâ, loculis polyspermis.  $\mathcal{U}$  in Alpibus Delphinatûs. — Fl. fr. 3. p. 210. *Anthericum serotinum* L. — Ray. angl. t. 17. f. 1. (v. s.)

### VIII. SCILLA *Sm. brit. n. 174. DC. fl. fr. 3. p. 211.*

Perigonium 6-partitum, sæpius patens et caducum; staminum filamenta glabra filiformia basi perigonii inserta; semina rotundata. — Radix bulbosa.

#### Sect. I. SCILLOIS.

Flores patentés.

1. *S. AUTUMNALIS* (L. sp. 443.) foliis filiformibus linearibus scapo brevioribus, floribus racemosis, racemo brevi, bracteis nullis, perigonii segmentis elliptico-lanceolatis obtusis stamina subæquantibus.  $\mathcal{U}$  in siccis arenosis aut glareosis Gallie mediz, occidentalis australisque. — Fl. fr. 3. p. 212. Cav. ic. t. 274. f. 2. — Lob. ic. t. 102. f. 1. (v. v.)
2. *S. UMBELLATA* (Ram. bull. phil. n. 41. p. 130. t. 8. f. 6.) foliis linearibus crassiusculis subcarinatis erectis scapo cylindrico subbrevioribus, floribus paucis racemoso-umbellatis, pedicellis bracteam membranaceam linearem acuminatam æquantibus superantibusve, perigonii segmentis lanceolatis obtusiusculis.  $\mathcal{U}$  in Pyrenæis et subulatis circà Aquas-Tarbellicas, Brivatem (Brest). — Fl. fr. 3. p. 213. lil. t. 165. *S. verna* Huds. angl. 142. — Radb. elys. p. 36. f. 16. (v. s.)

3. *S. BARTOLOMÆI* (Duby mas.) foliis linearibus angustis obtusis scapum subsequantibus, racemo brevi pauci-(4-10)-floro, pedicellis bracteas binas membranaceas subulatas sequantibus rariùs superantibus, perigonii segmentis lanceolato-linearibus obtusis.  $\frac{1}{2}$  circà Niczam. — *S. Italica* All. Bertol. amæn. ital. 144. non L. nec Fl. fr. nec Red. lil. t. 304 nec Curt. bot. mag. t. 663. (v. s.)
4. *S. BYPOLIA* (L. sp. 443.) foliis lanceolato-linearibus subbinis scapo vix longioribus basi vaginantibus, racemo paucifloro, floribus ebracteatis, pedicellis inferioribus longioribus, perigonii segmentis lineari-lanceolatis obtusis stamina superantibus.  $\frac{1}{2}$  in umbrosis et pascuis. — Fl. fr. 3. p. 212. eng. bot. t. 24. — Moris. s. 4. t. 12. f. 1. n. 15. (v. v.)
5. *S. HYACINTHOIDES* (L. f. supp. 201.) foliis lato-linearibus acutis, racemo longissimo multifloro, bractea minimà obtusà, pedicellis ante florescentiam erectis confertis, florentibus patulis laxis, perigonii segmentis lanceolatis obtusis stamina subsuperantibus.  $\frac{1}{2}$  in siccis circà Grasse detexit cl. A. Richard. — bot. mag. t. 1440. malè. — Clns. hist. 1. p. 172. ic. Fl. intense cærulei. (v. s.)
6. *S. UNDULATA* (Desf. atl. 1. p. 300. t. 88.) foliis lineari-lanceolatis undulatis scapo brevioribus, racemo elongato laxo, bractea lineari-subulatà minimà pedicello multò breviorè, perigonii segmentis lineari-spathulatis obtusis.  $\frac{1}{2}$  in Corsicà. — Fl. purpurei. (v. s.)
7. *S. LILIO-HYACINTHUS* (L. sp. 442.) bulbo squamato, foliis lato-lanceolatis acutis humi adpressis scapo brevioribus, floribus paucis racemosis, pedicello bracteam membranaceam linearem superante subsequanteve, perigonii segmentis linearibus obtusis.  $\frac{1}{2}$  in Gallià australi, Pyrenæis, sabaletis circà Aquas-Tarbellicas, Aureliam ex Dub. — Fl. fr. 3. p. 213. — Moris. s. 4. t. 12. f. 9. n. 21. Fl. cærulei. (v. v. DC.)
8. *S. AMÆNA* (L. sp. 443.) foliis lanceolatis obtusis, scapo angulato paucifloro, floribus laxis subcernuis, bracteis obtusis brevissimis, perigonii segmentis lanceolatis obtusis staminibus subduplò longioribus.  $\frac{1}{2}$  in sterilibus et arenosis propè *Marenzin* in agro Syrtico ex Thore. — Fl. fr. 3. p. 212. lil. t. 130. bot. mag. t. 341. non Jacq. (v. s. c.)
9. *S. MARITIMA* (L. sp. 442.) foliis lato-lanceolatis, racemo conico elongato multifloro, bracteis linearibus refractis subtus calcaratis pedicello multò brevioribus.  $\frac{1}{2}$  in littoribus arenosis utriusque maris ex Lois. — Fl. fr. 3. p. 214. Black. t. 591. — Moris. s. 4. t. 16. f. 11. n. 1. (v. v. c.)

## Sect. II. LYCORIS. DC. herb.

Floris lacinie in tubum basi conniventes.

10. *S. PATULA* (DC. fl. fr. 5. p. 209.) foliis elongato-lanceolatis acutis basi attenuatis scapum superantibus patulis, racemo conico brevi recto, floribus 6-9 confertis, bracteis membranaceis binis inæqualibus lineari-lanceolatis linearibusve acuminatis, floribus campanulatis, lacinii lanceolatis obtusis basi coalitis.  $\frac{1}{2}$  in Gallià australi? colitur nunc in hortis. — Fl. fr. 3. p. 211. lil. t. 225. *Hyacinthus patulus* Desf. *Scilla hyacinthoides* Jacq. ic. rar. t. 65. non L. f. (v. s. ex agro Florentino.)
11. *S. RUTANS* (Sm. brit. 1. p. 367.) foliis lato-linearibus basi attenuatis subacutis scapum subsequantibus, racemo immaturo cernuo, florente erecto laxo 3-10-floro, bracteis binis membranaceis inæqualibus lineari-lanceolatis subulatis, floribus campanulatis, lacinii obtusis lanceolato-linearibus apice revolutis basi coalitis.  $\frac{1}{2}$  in pratis et nemoribus Galliæ mediæ et occidentalis passim. — Fl. fr. 3. p. 211. eng. bot. t. 377. *S. non-scripta*. Red. lil. t. 224. *Hyacinthus non-scriptus* L. sp. 453. (v. s.)
12. *S. PARTIATA* (Viv. cors. app. 1.) foliis linearibus angustis scapum filiformem superantibus, racemo brevi denso 3-8-floro, bracteis amplexantibus ovato-acuminatis, perigonii tubo urocolato pellucido, lacinii lanceolatis obtusis stamina superantibus.  $\frac{1}{2}$  in montibus Corsicæ. (v. s.)

IX. HYACINTHUS *Tourn. inst. 180. Desf. atl. 1. p. 307.*

Perigonium tubulosum 6-fidum, laciniis patulis; stamina circa mediam perigonii longitudinem inserta; capsula obtusè trigona, loculis polyspermis.

1. *H. ORIENTALIS* (L. sp. 454.) foliis linearibus obtusis scapo brevioribus, racemo laxo erecto 4-10-floro, bracteis membranaceis geminatis lanceolatis pedicello brevioribus, floribus infundibuliformibus basi ventricosis vix 6-fidis, laciniis oblongis obtusis.  $\mathcal{U}$  in Galloprovincia orientali et circa Aquas-Tarbellicas (cl. Grateloup) spontaneus, in hortis frequenter colitur. — Fl. fr. 3. p. 207. — Moris. s. 4. t. 11. f. 9 et 10. n. 10 et 11. (v. s. s.)
2. *H. ROMANUS* (L. mant. 224.) foliis linearibus canaliculatis scapum longè superantibus, bracteis refractis obtusis, racemo ovato-cylindrico multifloro, floribus ovato-cylindricis ultra medium 6-fidis, lobis ovato-lanceolatis apice callosis.  $\mathcal{U}$  in pratis circa Tolosam, Castres, Agatham, Telonem. — Fl. fr. 5. p. 314. Red. lil. 6. t. 334. Bellevalia operculata Lapeyr. Journ. phys. dec. 1808. p. 425. t. 1. — Clus. hist. 1. p. 180. f. 2. — Lob. ic. t. 107. f. 1. (v. s.)
3. *H. AMETHYSTINUS* (L. sp. 454.) foliis linearibus angustis, floribus laxè racemosis, bracteis solitariis linearibus membranaceis pedicello sublongioribus, perigonii campanulatis basi cylindricis 6-lobis, lobis ovatis obtusis.  $\mathcal{U}$  in Pyrenæis centralibus. — Fl. fr. 3. p. 206. lilac. 1. t. 14. (v. s.)
4. *H. SEROTINUS* (L. sp. 453.) foliis linearibus canaliculatis scapo brevioribus, floribus racemosis secundis, bracteis ovatis acuminatis pedicello longioribus, perigonii campanulatis ultra medium 6-partitis, laciniis oblongis exterioribus longioribus patulis, interioribus connatis.  $\mathcal{U}$  in lapidosis provinc. mediterr. passim, Pyren. propè Valletriam. — Fl. fr. 3. p. 207. lil. t. 202. Cav. ic. t. 30. *H. lividus* Pers., *Lachenalia serotina* Willd. (v. v.)

X. MUSCARI *Tourn. inst. 180. Desf. atl. 1. p. 308.*

Perigonium ovatum medio inflatum, sexdentatum; capsula trigona, angulis prominulis, loculis 2-spermis.

1. *M. AMBROSIAEUM* (Moench. meth. 633.) foliis sublinearibus caule sublongioribus, floribus confertis ovatis omnibus æqualibus subsessilibus apice callosis hexagonis.  $\mathcal{U}$  Monspelii, Nemausi cultum. — Fl. fr. 3. p. 200. lil. t. 132. *M. moschatum* bot. mag. t. 734. *Hyacinthus muscari* L. sp. 454. — Lob. ic. t. 109. f. 2. (v. s. c.)
2. *M. RACEMOSUM* (Mill. dict. n. 3.) foliis laxis dependentibus carinatis linearibus scapum superantibus, floribus confertis ovatis, summis sessilibus.  $\mathcal{U}$  in cultis. — Fl. fr. 3. p. 208. lil. t. 232. *Hyacinthus racemosus* L. — Clus. hist. 1. p. 181. f. 1. — Lob. ic. t. 107. f. 2. (v. v.)
3. *M. BOTRYOIDES* (Mill. dict. n. 1.) foliis strictis lato-linearibus basi angustatis scapum superantibus, floribus globosis inferioribus remotioribus.  $\mathcal{U}$  in Gallia australi. — Fl. fr. 3. p. 208. *Hyac. botryoides* L. — Clus. hist. 1. p. 181. f. 2. — Lob. ic. t. 108. f. 1. (v. v. DC.)
4. *M. COMOSUM* (Mill. dict. n. 2.) foliis lato-linearibus scapum superantibus, racemo elongato, floribus angulato-cylindricis, summis sterilibus longius pedicellatis.  $\mathcal{U}$  in arvis et cultis. — Fl. fr. 3. p. 208. *Hyac. comosus* L. Jacq. aust. t. 126. — Lob. ic. t. 106. f. 2. (v. v.)

XI. GAGEA *Salisb. in ann. of bot. 2. p. 555.*

Perigonium calyciforme 6-partitum persistens basi connivens supernè patens; stamina 6, filamentis basi non dilatatis; stigma biulcum; capsula trigona. — Flores corymbosi lutei, bractæ foliaceæ.

1. *G. LUTEA* (Gawl. in bot. mag. p. 1200. excl. ic.) bulbo aggregato, foliis radicalibus 1-2 linearibus scapum angulatum aphyllum longè superantibus, floribus paucis corymbosis, pedicellis solitariis, bracteis lanceolatis lanceolato-linearibusve acuminatis pedicellos superantibus, perigonii segmentis lanceolatis obtusis.  $\mathcal{N}$  in cultis, pratis et hortis. — *G. bracteolaris* Salisb. l. c. *Ornithogalum luteum* L. sp. 440. Fl. fr. 3. p. 214. *O. pratense* Pers. in Ust. ann. 5. p. 3. t. 2. f. 1. (v. v.)  
 $\beta$ . *sylvatica*, bulbo solitario, folio radicali solitario lineari-lanceolato. — In sylvis provinc. orientalium. — *G. lutea* Gawler in bot. mag. t. 1200. *G. fascicularis* Salisb. l. c. excl. syn. *Ornithogalum sylvaticum* Fl. fr. 5. p. 315. (v. s.)
2. *G. VILLOSA* (Dub. mss.) foliis radicalibus 2 linearibus angustissimis scapum longè superantibus, floribus numerosis corymbosis, pedicellis subramosis villosis, bracteis ciliatis lineari-lanceolatis linearibusve pedicellos superantibus, perigonii segmentis elliptico-lanceolatis subacutis dorso pubescentibus.  $\mathcal{N}$  in locis cultis et arvis. — *G. stellaris* Salisb. l. c. *O. villosum* Willd. enum. 368. *O. minimum* Fl. fr. 3. p. 215. an *L?* sp. 440. *O. arvense* Pers. in Ust. ann. 5. p. 8. t. 1. f. 2. (v. v.)
3. *G. VERTULOSA* (Dub. mss.) foliis radicalibus filiformibus fistulosis scapum superantibus, corymbo 1-4-floro, bracteis lato-lanceolatis lanceolatisve pedicellis pubescentibus brevioribus, perigonii segmentis lanceolatis obtusiusculis glabris.  $\mathcal{N}$  in Alpium et Pyrenæorum udia. — *G. pygmæa* Salisb. l. c. *Ornithogalum fistulosum* Ram. in DC. fl. fr. 3. p. 215. *O. bohemicum* Zechm. act. boh. 2. p. 121. ic. (v. v.)

XII. ORNITHOGALUM. — *Ornithogali* sp. *L. DC.*

Perigonium petaloideum sexpartitum marcescens basi connivens superne patens; stamina 6, 3 externorum filamenta basi dilatata; stigma minimum capitatum; ovarium obtusè trigonum. — Flores racemosi albi aut viridescentes; bracteæ membranaceæ.

1. *O. UMBELLATUM* (L. sp. 441.) foliis linearibus angustissimis, racemo subcorymboso paucifloro, bracteis membranaceis lanceolatis linearibus acuminatis pedicellis duplò triplòve brevioribus, perigonii segmentis elliptico-lanceolatis obtusis, filamentis subulatis.  $\mathcal{N}$  in arvis et pratis. — Fl. fr. p. 217. Jacq. aust. t. 343. Flores sequentis, sed minores. (v. v.)
2. *O. MUTANS* (L. sp. 441.) foliis linearibus caulem subsuperantibus, floribus racemosis secundis pendulis, racemo paucifloro, bracteis lanceolato-linearibus acuminatis pedicellis duplò triplòve longioribus, segmentis lanceolatis obtusis, filamentis staminum alternis majoribus bicornibus.  $\mathcal{N}$  in pratis Alsatiæ, Lugduni, Gratianopolis, Aureliæ, Abbavillæ, Nemausis. — Fl. fr. 3. p. 218. Jacq. aust. t. 301. — Clus. app. 2. p. 9. t. 9. — Moris. s. 4. t. 13. f. 6 et 7. n. 8 et 9. Flores magni albi, segmentis dorso viridescentibus. (v. v.)
3. *O. NARBONNENSIS* (L. sp. 440.) foliis lato-linearibus caule longioribus, racemo longissimo, bracteis membranaceis lanceolatis acuminatis pedicellis patentibus multò brevioribus, perigonii segmentis patentibus lineari-lanceolatis obtusis staminibus duplò ovario triplò longioribus.  $\mathcal{N}$  in provinciis australibus usque ad Tolosam. — Fl. fr. 3. p. 218. — Moris. s. 4. t. 13. f. 5. n. 6. Fl. albi. (v. v.)
4. *O. PYRÆNAICUM* (L. sp. 440.) foliis linearibus, racemo longissimo, bracteis membranaceis basi dilatatis acutissimis, perigonii segmentis linearibus obtusis stamina superantibus ovario duplò longioribus.  $\mathcal{N}$  — Fl. fr. 3. p. 216. Flores ochroleuci.  
 $\alpha$ . *flavescens*, floribus remotis, bracteis pedicello brevioribus, staminibus æqualibus. — In provinc. orientalibus, mediis occidentibusque passim. — Jacq. aust. 2. t. 103. *O. flavescens* Lam. ill. t. 242. f. 2. (v. v.)  
 $\beta$ . *stachyoides*, floribus subconfertis, bracteis pedicellum æquantibus, sta-

## 468 MONOCOTYL. sive ENDOGENÆ.

minibus inæqualibus. — In prov. australibus et Pyrenæis. — *O. stachyoides* Ait. *kew.* p. 441. — *Ren. spec. t. go.* (v. v.)

5. *O. ARABICUM* (L. sp. 441.) foliis lanceolato-linearibus caule brevioribus, racemo subcorymboso multifloro, bracteis membranaceis basi dilatatis acuminatis pedicello duplò brevioribus, perigonii segmentis obovato-lanceolatis obtusis staminibus duplò ovario triplò longioribus, filamentis subulatis.  $\frac{1}{2}$  in agro Corsico, Nicæensi, Ruscinonensi propè Prades (Corder). — Fl. fr. 3. p. 217. lll. 2. t. 63. — *Clus. hist.* 166. ic (v. s.)

### XIII. ALLIUM L. gen. 409.

Perigonium 6-partitum sæpius patens; stigma simplex; capsula triquetra, loculis profundè bipartitis, axi filiformi post valvularum dehiscientiam superstitè. — Flores umbellati, terminales, è spathâ bivalvi orti.

#### §. 1. Foliis planis, staminibus alternis tricuspidatis.

1. *A. PORRUM* (L. sp. 423.) bulbo tunicato, caule tereti, foliis crassiusculis lanceolatis acutis subcarinatis, umbellâ capsuliferâ globosâ. (1) (2) ex Helvetiæ vineis ortum, in hortis cultum. — Fl. fr. 3. p. 219. — *Black. t.* 421. — *Lob. ic. t.* 154. f. 2. Flores albi aut purpurei. (v. v. c.)
2. *A. AMPHLOPRAEUM* (L. sp. 423.) bulbo sobolifero, caule tereti, foliis crassiusculis linearibus carinatis acutis, umbellâ capsulifera globosâ.  $\frac{1}{2}$  in provinciis australibus ex Lam. — Fl. fr. 3. p. 219. Præcedenti nimiam affine. (v. v. c.)
3. *A. ROTUNDUM* (L. sp. 423.) bulbo ovato tunicato, foliis planis linearibus angustis, umbellâ capsuliferâ globosâ, spathâ brevi, perigonii segmentis lanceolatis subacutis stamina pistillumque subæquantibus, externis dorso crenulatis.  $\frac{1}{2}$  in aridis Gallop., Delph., Nicæ ex All. — Fl. fr. 5. p. 315. Fl. purpurei. (v. v. DC.)
4. *A. ACUTIFLORUM* (Lois. not. 557) foliis planis linearibus angustis, umbellâ capsuliferâ subglobosâ, spathâ ovato-lanceolatâ acutâ, perigonii segmentis lanceolatis acuminatis stamina ciliata pistillumque superantibus.  $\frac{1}{2}$  in aridis Nicæ, Galloprov. orientalis et insulæ Ratoneau propè Massiliam (DC.) — Fl. fr. 5. p. 316. Fl. rosei. (v. v. DC.)
5. *A. SATIVUM* (L. sp. 425.) bulbo sobolifero, foliis linearibus integerrimis, umbellâ bulbiferâ globosâ, spathâ ovato-rotundâ, perigonii segmentis ovatis obtusis, pistillo staminibusque exsertis.  $\frac{1}{2}$  in hortis cultum, forsè in Galloprovinciæ maritimis spontaneum. — Fl. fr. 3. p. 219. — *Lob. ic. t.* 158. f. 1. Flores albi-purpureascentes. (v. v. c.)
6. *A. SCORODORRIZUM* (L. sp. 425.) foliis linearibus undulato-subcrenatis, caule antè maturitatem contorto, umbellâ bulbiferâ globosâ spatham subæquante.  $\frac{1}{2}$  in provinciis australibus spontaneum, in hortis cultum. — Fl. fr. 3. p. 220. *A. arenarium* L. ex Trevir. all. 5. — *Moris. s.* 4. t. 15. f. 2. n. 10. (v. v. c.)
7. *A. MULTIFLORUM* (DC. fl. fr. 5. p. 316.) foliis planis lato-linearibus acutis, umbellâ capsuliferâ subglobosâ 30-60-florâ, floribus longè pedicellatis, perigonii segmentis elliptico-lanceolatis obtusiusculis stamina non ciliata pistillumque superantibus.  $\frac{1}{2}$  in arenosis et cultis Nicæ, Telonis, Narbonæ, agri Ruscinonensis, Tolosæ. — Fl. purpurei. (v. v.)

#### §. 2. Foliis teretibus, staminibus alternis tricuspidatis.

8. *A. ASCALONICUM* (L. sp. 429.) scapo tereti, foliis subulatis, umbellâ capsuliferâ globosâ, spathâ bivalvi, perigonii segmentis linearibus acutis pistillum staminaque superantibus.  $\frac{1}{2}$  ex Oriente ortum, in hortis cultum. — Fl. fr. 3. p. 228. — *Moris. s.* 4. t. 14. f. 3. (v. v. c.)
9. *A. SPHEROCEPHALUM* (L. sp. 426.) caule tereti in parte inferiori folifero, foliis semi-teretibus, spathâ 2-valvi, valvis ovatis acutis umbellâ capsuliferâ glo-



bosâ brevioribus, perigonii segmentis elliptico-lanceolatis obtusis, staminibus pistilloque exsertis. ② in montanis et arvis sterilibus passim. — Fl. fr. 3. p. 228. — Clus. hist. r. p. 195. f. 1. — Moris. s. 4. t. 14. f. 4. Fl. purpurei. (v. v.)

10. A. VINCALE (L. sp. 428.) caule tereti, foliis teretibus fistulosis, umbellâ bulbiferâ globosâ spathâ acuminatâ breviori, perigonii segmentis lanceolato-linearibus stamina subæquantibus pistilloque brevioribus. ② in vineis et sepi-  
bus passim. — Fl. fr. 3. p. 228. — Lob. ic. t. 155. f. 2. et t. 156. f. 2. Fl. ru-  
belli. (v. v.)

11. A. CEPA (L. sp. 431.) caule fistuloso infernè ventricosso foliis teretibus fistu-  
losis longiore, umbellâ capsuliferâ globosâ, perigonii segmentis lineari-ellip-  
ticis obtusis pistillo staminibusque brevioribus. ② in hortis cultum. — Fl. fr.  
3. p. 225. — Lob. ic. t. 150. f. 1 et 2. — Moris. s. 4. t. 14. f. 1 et 2. (v. v. c.)

§. 3. *Foliis teretibus, staminibus omnibus simplicibus.*

12. A. OLERACEUM (L. sp. 429.) caule foliifero tereti, foliis semiteretibus sul-  
catis, umbellâ bulbiferâ laxâ, spathâ 2-valvi, valvis ovatis fistuloso-acumi-  
natis umbellam superantibus, perigonii segmentis lanceolatis obtusis stamina  
superantibus stylo brevioribus. ② in vineis et cultis. — Fl. fr. 3. p. 226. eng-  
bot. t. 480. A. virens Lam. — Hall. all. n. 27. t. 2. f. 2. — Moris. s. 4. t. 14.  
f. 6. n. 4. Fl. viridescentes. (v. v.)

13. A. SCHÆNOPRASUM (L. sp. 432.) foliis teretibus subulatis scapos subnudos  
æquantibus, spathâ 2-valvi, valvis ovato-rotundis acutis umbellam capsu-  
liferam confertam subæquantibus, perigonii segmentis lanceolatis acutis sta-  
mina pistillumque subduplò superantibus. ② in Alpibus et Pyrenæis, in hor-  
tis frequenter cultum. — Fl. fr. 3. p. 227 et 5. p. 319. A. foliosum Clar. in fl.  
fr. 3. p. 725. — Lob. ic. t. 154. f. 1. Fl. rosei. (v. v.)

14. A. INTERMEDIUM (DC. fl. fr. 5. p. 318.) caule foliifero, umbellâ laxâ in-  
terdum bulbiferâ, pedicellis capillaribus effusis, spathâ bivalvi, valvis linea-  
ribus acuminato-fistulosis umbellam duplò triplòve superantibus, perigonii  
segmentis lanceolatis obtusis stamina subæquantibus, stylo brevissimo. ② in  
agris et aridis Delph., Corsicæ, Narbonæ et præsertim provinc. occidentali-  
um. — A. paniculatum Vill. Lapeyr. non L. Fl. rubelli. (v. s.)

15. A. PALLENS (L. sp. 427.) caule foliifero, foliis semiteretibus, spathâ 2-  
valvi, valvis lanceolatis fistuloso-acuminatis umbellâ capsuliferâ laxâ duplò-  
quadruplòve longioribus, floribus pendulis, segmentis lanceolato-linearibus  
subtruncatis mucronatis stamina subsuperantibus, stylo brevissimo. ② in  
collibus et cultis passim præsertim in provinc. austr. et occidentali-  
bus. — Fl. fr. 3. p. 227. — Col. ecphr. 2. p. 6. t. 7. f. 2. Flores pallide flavescentes.  
(v. v.)

16. A. FLAVUM (L. sp. 428.) caule foliifero, foliis semiteretibus, spathâ 2-  
valvi, valvis lanceolatis fistuloso-acuminatis umbellam capsuliferam laxam  
superantibus, floribus pendulis, segmentis lanceolatis obtusis, pistillo sta-  
minibusque exsertis. ② in arvis sepius et dumosis Gallicæ australis, mediæ  
occidentalisque passim. — Fl. fr. 3. p. 226. Jacq. austr. t. 181. Red. lil. t.  
119. Fl. flavi. (v. s.)

17. A. MOSCHATUM (L. sp. 427.) caule tereti, foliis subulato-setaceis, spathâ 2-  
valvi, valvis lanceolatis ovatisve acuminatis umbellâ capsuliferâ planâ pau-  
ciflorâ brevioribus, perigonii segmentis oblongo-linearibus acutis stamina  
pistillumque superantibus. ② in siccis et editioribus prov. Mediterran. —  
Fl. fr. 3. p. 226. — C. Bauh. prod. p. 28. ic. Fl. albo-rosei. (v. v.)

18. A. PARVIFLORUM (Viv. cors. app. 2.) caule foliifero, foliis setaceo-filiformi-  
bus, spathâ 2-valvi, valvis ovato-lanceolatis acutis umbellâ capsuliferâ  
laxâ pauci-(3-7)-florâ brevioribus, pedunculis capillaribus inæqualibus erec-  
tis, perigonii segmentis sublinearibus acutis stamina pistillumque superan-  
tibus. ② in Corsicâ. — Fl. carnei dorso lineâ sanguineâ exarati. (v. s.)

§. 4. *Foliis planis; staminibus omnibus simplicibus.*

19. *A. CARINATUM* (L. sp. 426.) caule foliifero, foliis linearibus carinatis, spathâ 2-valvi, valvis ovato-lanceolatis acuminatis umbellam bulbiferam paucifloram longè superantibus, perigonii segmentis lineari-lanceolatis obtusis stamina subæquantibus pistillo subduplò brevioribus. 2/ in agris et vineis Gallie australis, Andegavie, propè Falaise ex cl. de Bréb., Parisius ex cl. Lois. — Fl. fr. 3. p. 220. — Hall. all. n. 27. t. 2. f. 2. — Lob. ic. t. 156. f. 1. Flores roseo-viridescens. (v. s.)
20. *A. SUAVIOLENS* (Jacq. ic. rar. 2. t. 364.) caule basi foliifero, foliis linearibus obtusis caule brevioribus, spathâ 2-valvi, valvis lanceolatis acuminatis umbellâ capsuliferâ globosâ brevioribus, perigonii segmentis lineari-lanceolatis obtusis pistillo staminibusque brevioribus. 2/ in Pyrenæis, agro Syrtico propè Dax et Burdigalam (cl. Des Moulins). — *A. ambiguum* Fl. fr. 3. p. 220. *A. ericetorum* Thore land. 123. Lapeyr. abr. suppl. 47. *A. serotinum* Lap. abr. 179. suppl. 48. Fl. albescentes aut rosei, odori. (v. v.)
21. *A. SERRACENS* (L. sp. 430.) scapo nudo aucipiti folia obtusa linearia subtus convexa superante, spathâ ovatâ umbellâ capsuliferâ hemisphæricâ brevior, perigonii segmentis ovatis obtusis stamina pistillumque subæquantibus. 2/ in humidis Alpium Delph., Gallop., montium Ligeris (Prost), Pyren. — bot. mag. t. 1150. *A. angulosum* Fl. fr. 3. p. 222. an L? — Gmel. sib. t. 11. f. 2. Fl. rosei. (v. v.)
22. *A. PEDEMONTANUM* (Willd. sp. 2. p. 77.) scapo subtetragono folia linearia obtusa radicalia superante, spathâ 2-4-valvi, valvis ovatis acutis umbellâ capsuliferâ convexâ pauciflorâ brevioribus, perigonii segmentis ovatis obtusis stamina pistillumque superantibus. 2/ in montibus Arvernae ex Lam. — Fl. fr. 3. p. 223. *A. nigrum* All. ped. t. 25. f. 1. non L. Fl. rosei. (v. s. ex Pedem.)
23. *A. ALBUM* (Santi montan. x. t. 7.) scapo subtriquetro, foliis radicalibus glabris lineari-lanceolatis carinatis, spathâ 1-2-valvi, valvis ovatis acuminatis umbellâ capsuliferâ brevioribus, perigonii segmentis obovatis obtusis subenerviis stamina pistillumque superantibus. 2/ circâ Niczam, Telonem ex cl. Robert. — Fl. fr. 5. p. 317. Red. lil. t. 300. Fl. carnel. (v. s. ex Italiâ.)
24. *A. NOSEUM* (L. sp. 432.) bulbo sæpè tunicâ punctulatâ vestito, scape tereti, foliis subradicalibus linearibus lato-linearibusve acuminatis, spathâ vaginante 4-lobâ, lobis ovato-lanceolatis acutis umbellâ capsuliferâ (rariùs bulbiferâ) planiusculâ subduplò brevioribus, perigonii segmentis lanceolatis ovato-lanceolatisve obtusissimis stamina pistillumque superantibus. 2/ in agris et vineis provinc. Mediterr., Burdigalæ (Des Moulins). — Fl. fr. 3. p. 221. — Magn. bot. monsp. p. 10. ic. *A. carneum* Bertol. Fl. rosei. (v. v.)
25. *A. GRANDIFLORUM* (Lam. dict. x. p. 68.) foliis lineari-subulatis scapo tereti brevioribus, umbellâ capsuliferâ planâ, perigonii segmentis mucronatis erectis. 2/ in saxosis Alpium Delph. ex Vill., Gallopr. ex Lam. — Fl. fr. 3. p. 222. *A. narcissifolium* Vill. dauph. 2. t. 6. Fl. albo-rosei. †
26. *A. SUMMISTUTUM* (L. sp. 424.) scapo tereti, foliis linearibus acuminatis subhirsutis glabrisve subradicalibus, spathâ 2-valvi, valvis ovatis acutis umbellâ capsuliferâ planâ laxâ brevioribus, perigonii segmentis lineari-lanceolatis obtusis stamina pistillumque superantibus. 2/ in maritimis Corsicæ, Monspeli et Narbonæ? — Fl. fr. 3. p. 221. *A. graminifolium* Lois. gall. 724. — Lob. ic. t. 160. f. 1. Fl. lactei. (v. s.)
27. *A. TRIQUETRAUM* (L. sp. 431.) scapo triquetro, foliis carinatis basi vaginantibus acutis scapum superantibus aut subæquantibus radicalibus, spathâ bi-valvi, valvis lanceolatis linearibusve acuminatis umbellam capsuliferam planam pauci-(3-5)-floram subæquantibus, perigonii segmentis lineari-lanceolatis oblongis obtusis stamina pistillumque duplò superantibus. 2/ in Corsicâ, circâ Telonem, Narbonam, Collioure ex cl. Benth. — Fl. fr. 3. p. 222. — Rudb. elys. 2. p. 159. f. 16. — Moris. s. 4. t. 16. f. 8. n. 9. Flores albi. (v. s.)

28. *A. MOLY* (L. sp. 432.) scapo tereti, foliis radicalibus oblongo-lanceolatis acutis basi vaginantibus, spathâ bivalvi, valvis ovatis acutis umbellâ capsuliferâ planâ brevioribus, perigonii segmentis ovatis acutis stamina pistillumque superantibus. 2½ in Monspelii et in Pyrenæis ex L., in Andegaviâ ex Baat, circâ Parisios ex Mer., Abbavillam ex Bouch. — Fl. fr. 3. p. 224. lil. 2. t. 97. — Moris. s. 4. t. 16. f. 2 et 3. n. 4 et 5. Fl. lutei. (v. v. c.)
29. *A. CHAMÆMOXY* (L. sp. 433.) foliis lato-linearibus ciliatis scapum basi vaginantibus multoque superantibus, spathâ 2-3-valvi, valvis ovatis acuminatis umbellam capsuliferam subradicalem paucifloram subæquantibus, perigonii segmentis linearibus obtusis stamina pistillumque superantibus. 2½ in Corsicâ, circâ Arelatem (Suffren), Massiliam (Solier). — Fl. fr. 3. p. 224. Cav. ic. 3. t. 207. f. 1. — Col. cephr. t. 326. — Moris. s. 4. t. 16. f. 7. n. 8. Fl. albi. (v. s.)
30. *A. XIPHIUM* (L. sp. 430.) bulbo sobolifero, scapo tereti, foliis lato-lanceolatis oblongis radicalibus, spathâ bivalvi, valvis lato-ovatis acutis umbellam capsuliferam convexam multifloram subæquantibus, perigonii patentis segmentis lineari-lanceolatis obtusis stamina pistillumque superantibus, staminibus basi dilatatis in urceolum monadelphum coalitis. 2½ in arvis Monspelii ex Gou. et Benth., Gallopr. ex L. — Fl. fr. 3. p. 223. lil. 2. t. 102. *A. Monspesulanum* Gou. ill. 24. t. 16. — Besl. eyst. æst. 4. t. 11. f. 1. Fl. pallidi. (v. s. c.)
31. *A. MAGICUM* (L. sp. 424.) foliis lato-lanceolatis scapo multo longioribus, interiori filiformi basi vaginante apice intus bulbifero, umbellâ omnino bulbis aggregatis constante. 2½ circâ Aginnum ex S.-Am. — Fl. fr. 5. p. 317. S.-Am. mem. agr. 1. p. 79. bonq. t. 10. — J. Bauh. hist. 2. p. 569. fig. infer. An deformatio præcedentis aut species propria? †
32. *A. URANUM* (L. sp. 431.) scapo subtriquetro, foliis radicalibus ovato-lanceolatis petiolatis basi vaginantibus, spathâ 1-valvi, valvâ ovato-lanceolatâ acutâ umbellam capsuliferam planam subsuperante, perigonii segmentis linearibus acutis stamina pistillumque superantibus. 2½ in sepiibus et pratis umbrosis. — Fl. fr. 3. p. 214. fl. dan. t. 757. — Lob. ic. t. 159. f. 1. Fl. albi. (v. v.)
33. *A. VICTORIALE* (L. sp. 424.) radice oblongâ, caule tereti basi foliifero, foliis ovali-oblongis obtusis sessilibus, spathâ umbellam capsuliferam globosam subæquante, perigonii segmentis lanceolatis obtusis staminibus pistilloque brevioribus. 2½ in Vogesiâ, Jurasco, montibus Foresii, Delph., Arvernâ, Occitanîæ, Pyr. orient. — Fl. fr. 3. p. 224. Jacq. austr. t. 216. — Moris. s. 4. t. 15. f. 7. n. 16. Fl. ochrolenci. (v. v.)

### Trib. III. *HEMEROCALLIDÆ* R. Brown. prod. 295.

Perigonium tubulosum marcescens. Semina testâ nec atrâ nec crustaceâ sæpè spongiosâ et dilatâtâ vel alâtâ. Albumen carnosum.

#### XIV. *HEMEROCALLIS* L. gen. 433.

Perigonium magnum marcescens monophyllum segmentis basi in tubum coalitis; stigma simplex; stamina declinata, capsula trigona; semina rotunda.

1. *H. FULVA* (L. sp. 462.) foliis linearibus carinatis, floris laciniis tribus interioribus obtusis undulatis, nervis exterioribus ramosis. 2½ in Galloprovinciâ, propè Tarbas, Bayonam, Burdigalam, Cænomanum. — Fl. fr. 3. p. 205. Red. lil. 1. t. 16. — Lob. ic. t. 93. f. 1. (v. s.)
2. *H. FLAVA* (L. sp. 462.) foliis linearibus carinatis, floris laciniis planis acutis, nervis indivisis. 2½ in hortis culta. — Fl. fr. 3. p. 205. Jacq. vind. t. 139. Red. lil. 1. t. 15. (v. v. c.)

## 472 MONOCOTYL. SEU ENDOGENÆ.

### XV. POLYANTHES *L. gen.* 426.

Perigonium infundibuliforme, limbo sexfido patulo; stamina 6 perigonii fauci inserta; ovarium liberum perigonio obtectum; stigma 3-fidum.

1. *P. TUBEROSA* (*L. sp.* 453.) foliis linearibus scapo brevioribus.  $\frac{1}{2}$  ex America meridionali orta, in hortis culta. — *Fl. fr.* 3. p. 234. *Lam. ill. t.* 243. *Red. ill. t.* 147. (v. v. c.)

---

### Ordo CIII. BROMELIACEÆ *Juss. gen.* 49. *dict. sc. nat.* 5. p. 347.

Perigonium tubulosum mox cum ovario adhærens, mox liberum, 6-fidum aut 6-partitum laciniis duplici serie dispositis, 3 externis brevioribus persistentibus calyciformibus, 3 internis majoribus sæpè caducis petaloideis. Stam. 6 imo perigonio inserta. Ovarium 3-loculare  $\infty$ -ovulatum. Stylus simplex. Stigma 3-lobum. Fructus baccatus 3-locularis, aut capsularis. Albumen farinosum basi embryonem elongatum recurvum fovens. — Folia alterna sæpissimè ad basin scapi fasciculata, margine dentato-spinosa.

#### I. AGAVE *L. gen.* 431.

Perigonium tubulosum infundibuliforme ovario adhærens sexpartitum; stamina 6 exserta; antheræ versatiles; capsula ovata utrinque attenuata obtusè trigona 3-locularis polysperma.

1. *A. AMERICANA* (*L. sp.* 461.) acaulis, foliis dentato-spinosis, scapo ramoso, tubo perigonii medio angustato, stylo staminibus longiore.  $\frac{1}{2}$  ex America meridionali orta, nunc spontanea in provinciis Mediterraneis. — *Fl. fr.* 3. p. 235. *Lam. ill. t.* 235. f. 1. (v. v.)

---

Ordo CIV. COLCHICACEÆ DC. *fl. fr.* 3. p. 192. — Melanthaceæ *Brown prod.* 272. — Juncorum gen. *Juss. gen.*

Flores sæpiùs hermaphroditi. Perigonium coloratum, 6-fidum aut 6-partitum, laciniis æstivatione sæpè involutis. Stamina 6 perigonii laciniis adnata; antheræ extrorsæ. Ovaria 3 mox subdistincta, mox plus minùs coalita et ovarium unicum 3-loculare oculis semina plurima angulo interno adfixa continentibus constituentia, stylo mox longissimo mox subnullo terminata; stigma glandulosum. Fructus mox folliculi 3 distincti uniloculares internè longitudinaliter dehiscentes; mox folliculis coadunatis capsula 3-valvis 3-locularis, valvularum marginibus introrsum inflexis dissepimenta constituentibus; folliculis maturitate disjunctis. Semina plurima margini valvularum interiori adfixa; testa membranacea. Embryo intrà albumen carnosum. — Herbæ habitu vario.

I. BULBOCODIUM *L. gen.* 407.

Perigonium 6-partitum, lacinarum unguibus angustis conniventibus apice staminiferis; ovarium 1; stylus simplex elongatus; stigmata 3.

1. *B. verum* (*L. sp.* 422.)  $\mathcal{U}$  in Alpibus Delph., Gallop., circà Nicæam, in editioribus Pyrenæis ex *Lapeyr.* — *Fl. fr.* 3. p. 197. Vill. dauph. 2. t. 2. Folia lanceolata cum floribus prodeuntia. (v. s.)

II. MERENDERA *Ramond bull. phil. n.* 47. *DC. fl. fr.* 3. p. 196.

Perigonium 6-partitum, laciniis unguiculatis staminiferis; antheræ erectæ hastatæ; ovarium 1; styli 3 elongati apice erecti; capsula 3-follicularis, folliculis erectis minimè inflatis.

1. *M. Bulboconium* (*Ram. l. c. t.* 12. f. 2.)  $\mathcal{U}$  in graminosis editiorum Pyrenæorum. — *Fl. fr.* l. c. Red. lil. 1. t. 25. *Bulboconium autumnale* *Lapeyr. abr. pyr.* 202. — *Clus. hist.* 1. p. 201. f. 2. Folia linearia patula cum flore subprodeuntia. (v. v.)

III. COLCHICUM *Tourn. inst.* 181. *L. gen.* 457.

Perigonium tubulosum longum bulbo innascens, limbo campanulato 6-partito; stamina apici tubi inserta; antheræ oblongæ versatiles; ovarium 1; styli 3 longissimi; stigmata totidem adunca; capsula 3-follicularis, folliculis inflatis erectis basi coalitis polyspermis.

1. *C. montanum* (*L. sp.* 485.) bulbo multifloro, foliis cum flore prodeuntibus sublinearibus, perigonii laciniis lineari-ellipticis.  $\mathcal{U}$  in montibus Corsicæ. — *Fl. fr.* 3. p. 195. excl. syn. *Hall. et Clus.* Autumno floret. (v. s.)

2. *C. alpinum* (*DC. fl. fr.* 3. p. 195.) bulbo unifloro, foliis lineari-lanceolatis obtusis strictis post flores prodeuntibus, perigonii laciniis lanceolato-ellipticis.  $\mathcal{U}$  in pratis humidis Alpium Delphinatæ? — *C. montanum* *All. ped.* t. 74. f. 2. excl. syn. (v. s. ex Alp. Pedem.)

3. *C. autumnale* (*L. sp.* 485.) bulbo submultifloro, foliis planis lato-lanceolatis post flores longè prodeuntibus, perigonii laciniis ovato-ellipticis.  $\mathcal{U}$  in pratis humidis frequens. — *Fl. fr.* 3. p. 195. *Bull. herb. t.* 19. — *Moris. s.* 4. t. 3. f. 1. (v. v.)

IV. VERATRUM *Tourn. inst.* 145. *L. gen.* 1144.

Perigonium 6-partitum; ovaria 3 basi juncta in plurimis floribus

## 474 MONOCOTYL. seu ENDOGENÆ.

abortiva; styli 3 breves; capsula oblonga 2-valves polysperma; semina margine membranacea. — Caules foliosi.

1. *V. ALBUM* (L. sp. 1479.) racemo supra-decomposito, ramulis patentibus, perigonii subpatentibus (albo-virescentibus).  $\mathcal{U}$  in pascuis montanis. — Fl. fr. 3. p. 194. Lam. ill. t. 843. Bull. herb. t. 155. (v. v.)
2. *V. NIGRUM* (L. sp. 1479.) racemo composito, ramulis strictis, perigonii patentissimis (nigricantibus).  $\mathcal{U}$  in pascuis montium Niceæ, Arvernæ ex Lois., Burgundiæ ex Dur., Alsatiæ ex Mapp. — Fl. fr. 3. p. 194. Bull. herb. t. 149. — Moris. s. 12. t. 4. f. g. n. 2. (v. s.)

*V. TOFIELDIA* Huds. ang. 157. DC. fl. fr. 3. p. 193. — *Antherici* sp. L.

Perigonium 6-partitum, involucre minimo 3-partito basi cinctum; stamina glabra; capsula 3-6-locularis, loculis polyspermis basi junctis. — Caules inferne foliosi, foliis gramineis.

1. *T. PALUSTRIS* (Huds. l. c.) foliis lineari-ensiformibus nervosis.  $\mathcal{U}$  in humidis Jurassi, Alpiam, montium Corsicæ, Pyrenæorum. — Fl. fr. 3. p. 193. eng. bot. t. 536. — *Narthecium calyculatum* Lam. ill. t. 268. *Helonias borealis* Willd. — Fl. lapp. t. 10. f. 3. (v. v.)

Ordo CV. JUNCÆ DC. fl. fr. 3. p. 155. *Brown prod.* p. 25. *De La Harpe in mem. soc. hist. nat. paris. t. 3. p. 87.* — *Juncorum* gen. *Juss.*

Flores hermaphroditi. Perigonium liberum sæpius glumaceum sexpartitum, laciniis duplici serie dispositis. Stamina 3 seu 6, perigonii laciniis opposita; filamenta subulata; antheræ 2-loculares. Ovarium 1; stylus 1; stigmata 3 filiformia aut stigma unicum 3-lobum. Capsula 3-valvis, 1-aut 3-locularis, valvulis nunc medio septiferis et polyspermis, nunc septo destitutis et semen unicum basi gerentibus. Embryo subcylindricus, albumen carnosum. — Herbæ cyperaceis habitu similes. Folia vaginantia. Flores paniculati aut corymbosi rarius spicati, bracteis scariosis glumaceis instructi.

- I. *ABAMA* Adams. fam. 2. p. 47. DC. fl. fr. 3. p. 177. — *Narthecium* Lam. non *Juss.* — *Antherici* sp. L.

Perigonium 6-partitum; stamina 6; filamenta lanuginosa persistentia; ovarium pyramidale; stylus brevis; capsula 3-locularis, 3-valvis; semina numerosa, ovato-oblonga, utrinque appendiculata.

1. *A. OSSIFRAGA* (DC. l. c.) foliis ensiformibus, floribus laxè spicatis subsessilibus.  $\mathcal{U}$  in locis humidis Pyrenæorum, Galliæ occidentalis septentrionalisque. — *Narthecium ossifragum* Lam. — Lob. ic. t. 92. f. 1. (v. v.)

### II? APHYLLANTHES Tourn. inst. 430. L. gen. 408.

Perigonium 6-partitum, laciniis basi in tubum coalitis, limbis patentibus; stamina 6; capsula turbinata, 3-locularis, 3-valvis, polysperma. — Genus ex cl. de La Harpe è Juncæis expellendum et ad Liliacearum trib. II. (Asphodeleas) amandandum.

1. *A. MONSPÉLIENSIS* (L. sp. 422.)  $\mathcal{U}$  in lapidosis Galliæ mediterraneæ. — Fl. fr. 3. p. 170. Lam. ill. t. 252. — Moris. s. 5. t. 25. f. g. n. 12. Sæpi nudi. Flores 1-2 sessiles basi squamis scariosis cincti. (v. v.)

III. *JUNCUS* Michel. gen. p. 37. t. 31. DC. fl. fr. 3. p. 162. De La Harpe l. c. p. 104.

Perigonium 6-partitum, glumaceum; stamina 3 vel 6; capsula 3-locularis 3-valvis, valvulis medio septiferis; semina numerosa septo adfixa. — Folia teretia glabra.

### §. 1. Foliis nullis.

1. *J. filiformis* (L. sp. 465.) culmo nudo filiformi debili, paniculâ laterali bracteâ subsimplici, perigonii segmentis lanceolatis acutis capsulam subrotundam obtusam subæquantibus.  $\frac{1}{2}$  in paludibus turfosis Alpium, montium Liguria (cl. Prost), Pyrenæorum. — Fl. fr. 3. p. 164. eng. bot. t. 175. *J. arcticus* Lapeyr. non L. — Pluck. t. 40. f. 8. (v. s.)
2. *J. communis* (E. Meyer junc. mon. spec. p. 20.) scapo tereti sublævi rigidiusculo, paniculâ laterali, staminibus 3, perigonii segmentis sublinearibus acuminatis capsulam obovatam retusam subæquantibus.  $\frac{1}{2}$  in paludosis.
  - a. *conglomeratus* (Mey.) paniculâ conglobatâ. — *J. conglomeratus* L. sp. 464. Fl. fr. 3. p. 163. — Cam. epit. 780. ic. — Moris. s. 8. t. 10. f. 4. n. 7. (v. v.)
  - b. *effusus* (Mey.) paniculâ effusâ suprâ-decompositâ. — *J. effusus* L. sp. 464. Fl. fr. 3. p. 163. — Lob. ic. t. 84. f. 2. t. 85. f. 1. (v. v.)
3. *J. glaucus* (Sm. brit. 575.) culmo nudo tereti striato glauco apice inflexo, paniculâ laterali effusâ erectâ, staminibus 6, perigonii segmentis angustè lanceolatis acuminatis capsulâ ellipticâ obtusâ mucronatâ obsolete trigonâ subbrevioribus.  $\frac{1}{2}$  in humidis passim. — Fl. fr. 5. p. 307. eng. bot. t. 665. *J. inflexus* Lam. dict. 3. p. 265. Fl. fr. 3. p. 164. non L. *J. tenax* Poir. *J. longicornis* Bast. — Barr. ic. 208. — Moris. s. 8. t. 10. f. 5. n. 13. (v. v.)
4. *J. rigidus* (Desf. atl. 1. p. 312.) culmo tereti lævi nudo? ultrâ paniculam producto pungente, floribus solitariis sessilibus pedicellatis in paniculam elongatam erectam suprâ-decompositam digestis, pedunculis inæqualibus compressis nodosis multifariam decompositis fasciculatis, perigonii segmentis lanceolatis acutis.  $\frac{1}{2}$  in arenosis maritimis Monspelii propè Maguelonne detexit cl. Dunal. (v. s. sine fructu nec radice.)

### §. 2. Foliis radicalibus.

5. *J. acutus* (Lam. dict. 3. p. 264.) culmo lævi nudo tereti, foliis teretibus pungentibus, floribus paucis capitatis in paniculam densam terminalem proliferam digestis, spathâ diphyllâ spinosâ erectâ, bracteis capitulum subæquantibus, perigonii segmentis lanceolato-ovatis, exterioribus acutis interioribus obtusis, capsulâ oblongâ ovoideâ mucronatâ duplò brevioribus.  $\frac{1}{2}$  in paludosis maritimis maris Mediterranei et Oceani propè Brest et Cherbourg. — Fl. fr. 3. p. 163. — C. Bauh. prod. p. 21. ic. — Moris. s. 8. t. 10. f. 7. n. 15. (v. v.)
6. *J. maritimus* (Lam. dict. 3. p. 264.) culmo nudo tereti sublævi, foliis teretibus pungentibus, floribus capitatis, capitulis paucifloris in paniculam terminalem proliferam digestis, spathâ diphyllâ spinosâ erectâ, bracteis capitulo brevioribus, staminibus 6, perigonii segmentis lanceolatis acutis capsulas ovato-rotundas obtusè trigonas mucronatas æquantibus.  $\frac{1}{2}$  in arenosis maritimis. — Fl. fr. 3. p. 162. *J. acutus* a L. — Moris. s. 8. t. 10. f. 6. n. 14. (v. v.)
7. *J. squarrosus* (L. sp. 465.) culmo subnudo rigido, foliis setaceis subcarinatis culmo triplò brevioribus, floribus solitariis in paniculas 2-3 laterales terminalesque digestis bracteis scariosis stipatis, perigonii segmentis lanceolatis subobtusis margine squarrosis capsulas ovatas obtusas æquantibus.  $\frac{1}{2}$  in humidis et paludosis passim. — Fl. fr. 3. p. 165. *J. Sprengelii* Willd. prod. t. 4. f. 8. — Lob. ic. t. 18. f. 1. (v. v.)
8. *J. ericetorum* (Poll. pal. n. 350.) culmo nudo filiformi striato, foliis filifor-

mibus canaliculatis, capitalis 1-3 terminalibus subrotundis sessilibus involu-  
cratis, bracteis inaequalibus unâ capitulum superante, staminibus 3, peri-  
gonii segmentis lanceolatis acuminatis capsulam ovatam obsolete trigonam  
mucronatam superantibus. ① in glareosis subudis Galliae australis occiden-  
talisque, circâ Parisios. — Fl. fr. 3. p. 164. J. triandrus Gou. J. gracilis  
Roth. J. capitatus Willd. J. mutabilis Cav. ic. t. 296. f. 2. excl. syn. (v. v.)

### §. 3. *Caulibus foliosis.*

9. J. *summus* (Roth. germ. 1. p. 156.) culmo striato erectiusculo ramoso, foliis  
linearibus canaliculatis, capitulis tribracteatis, bracteis capitulum superan-  
tibus, staminibus 3, perigonii segmentis lanceolatis acutis capsulâ triangu-  
lari obtusâ brevioribus. ② in paludibus semixsiccatis passim, præsertim in  
provinciis occidentilibus. — Fl. fr. 3. p. 168. Fl. dan. t. 1099 excl. syn. J.  
subverticillatus & Willd. — Moris. s. 8. t. 9. f. 10 et 11. n. 3 et 4. (v. v.)
10. J. *plurimus* (Lam. dict. 3. p. 270.) culmo bulboso tenui radicante subangu-  
lato, foliis setaceis subnodoso-articulatis, floribus capitatis, capitalis paucis  
subproliferis 3-6-floris in paniculam simplicem laxissimam dispositis, stami-  
nibus tribus, perigonii segmentis anguste lanceolatis subacutis capsulâ  
ovato-oblongâ breviter mucronatâ brevioribus. ② in fossis et paludibus aqua-  
ticiis passim. — Fl. fr. 3. p. 169. eng. bot. 301. J. uliginosus Roth. J. sub-  
verticillatus Hoffm. — Scheuchz. gram. 330. t. 7. f. 10. (v. v.)
11. J. *pyramus* (Thuil. fl. par. 11. 1. p. 178.) culmo erecto tenui ramoso lævi  
paucifolio, foliis linearibus canaliculatis, capitalis terminalibus bracteis  
minimis scariosis stipatis, infimo folio suffulto, staminibus 3, perigonii seg-  
mentis sublinearibus acuminatis capsulam elongatam trigonam acutam su-  
perantibus. ① in pratis subterfosis Galliae australis, occidentalisque et propè  
Parisios. — Fl. fr. 3. p. 168. (v. v.)
12. J. *bicorporatus* (Viv. cors. 5.) culmo ascendente tereti lævi, foliis lineari-  
filiformibus subcanaliculatis, floribus capitatis, capitalis 2 uno terminali  
altero laterali bracteis scariosis minimis stipatis, perigonii segmentis subli-  
nearibus acuminatis capsulam oblongo-prismaticam acutam superantibus. (1)  
in Corsicâ circâ Bonifacio (cl. Viviani). (v. s.)
13. J. *submontus* (L. sp. 466.) culmo angulato striato racemoso-paniculato versûs  
apicem foliis 1-2 setaceis instructo, floribus sessilibus solitariis in paniculam  
laxissimam racemosam dichotomam digestis, perigonii segmentis lanceolatis  
acuminatis capsulam ovato-oblongam obtusam superantibus. ① in pratis  
paludosis, ad vias et in humilibus frequens. — Fl. fr. 3. p. 167. eng. bot. t. 802.  
— Lob. ic. t. 18. f. 2. (v. v.)
14. J. *tenaxia* (L. f. supp. 208.) culmo tereti ramoso paniculato foliis 2-3  
erectis setaceis subcanaliculatis instructo, floribus solitariis sessilibus in pa-  
niculam laxissimam dichotomo-divaricatam dispositis, perigonii segmentis  
lanceolato-ovatis acutis capsulam ovato-rotundam retusam subæquantibus.  
① in paludibus exsiccatis passim. — Fl. fr. 3. p. 167. Fl. dan. t. 1160. —  
Vaill. bot. t. 20. f. 1. — Barr. ic. t. 747. f. 2. (v. v.)
15. J. *trifidus* (L. sp. 465.) culmo tereti lævi superuè foliis 1-2 setaceis subca-  
naliculatis instructo, floribus 1-3 subcapitatis sessilibus lateralibus, peri-  
gonii segmentis lanceolatis acuminatis, staminibus 6, capsulâ ovatâ obtusâ  
trigonâ acuminatâ. ② in glareosis Alpium et Pyrenæorum. — Fl. fr. 3. p. 165.  
Fl. dan. t. 107. J. monanthos Jacq. vind. obs. t. 4. f. 1. — Lightf. scot. t. 9.  
f. 1. (v. v.)
16. J. *gerardi* (Lois. not. 60.) culmo tereti sublævi, foliis linearibus canalicu-  
latis, summo paniculam superante, floribus subsolitariis in paniculam ter-  
minalem laxam erectam paucifloram digestis, perigonii segmentis lanceolatis  
subobtusis apice callosis capsulam ovatam mucronatam superantibus. ② in  
pratis secûs rivulos Burgundiæ (Babey), Galloprov., Corsicæ, Mimatis (Prost),  
Burdigalæ (Des Moul.). — Fl. fr. 5. p. 308 excl. syn. Barr. J. attenuatus Viv.  
cors. 5. An satis à sequente distinctus. (v. s.)



17. *J. vulbosus* (L. sp. 466.) culmo compresso sublevi, foliis linearibus canaliculatis, floribus subsolitariis in paniculas decompositas cymosas multifloras digestis, perigonii segmentis lanceolatis obtusis capsulâ ovato-rotundâ trigonâ mucronatâ duplò brevioribus.  $\frac{1}{2}$  in paludibus et humidis. — Fl. fr. 3. p. 167. eng. bot. t. 934. *J. nitidiflorus* L. Duf. in ann. sc. nat. 5. p. 86. — Moris. s. 8. t. 9. f. 14. n. 11. (v. v.)
18. *J. lamprocarpus* (Ehr. gram. p. 126.) culmo nodoso tereti lævi, foliis compressis nodoso-articulatis, floribus capitatis, capitulis sessilibus pedunculatisque in paniculam simplicem compositamve terminalem erectam digestis, perigonii segmentis lanceolatis, tribus interioribus obtusiusculis exterioribus acuminatis, capsulâ acutâ triquetrà nitidissimâ subduplò brevioribus.  $\frac{1}{2}$  in humidis. — eng. bot. t. 2143. *J. articulatus* L. sp. 465. *J. sylvaticus* var. Fl. fr. 3. p. 169. — Moris. s. 8. t. 9. f. 9. n. 2. Capsula atra. (v. v.)  
*β. heterophyllus*, foliis immersis elongatis filiformi-setaceis tenuissimè articulatis. — In fossis aquosis agri Syrtici, Corsicæ. — *J. heterophyllus* L. Duf. in ann. sc. nat. 5. p. 88. (v. s.)
19. *J. ALPINUS* (Vill. dauph. 2. p. 233.) culmo erecto tereti lævi, foliis compressis subarticulatis, floribus capitatis, capitulis paucis paucifloris in paniculam simplicem digestis, perigonii segmentis lanceolatis acutis capsulâ trigonâ ovatâ mucronatâ brevioribus.  $\frac{1}{2}$  in Alpibus et Pyrenæis. — Fl. fr. 3. p. 170. *J. ustulatus* Hoppe? Sequentis varietatem esse vult. cl. Gaudin agrost. 2. p. 223. an meritò? (v. v. DC.)
20. *J. ACUTIFLORUS* (Ehr. gram. 66.) culmo nodoso tereti lævi, foliis compressis nodoso-articulatis, floribus capitatis, capitulis sessilibus pedunculatisve in paniculam terminalem erectam suprâ-decompositam digestis, perigonii segmentis lanceolatis acuminatis capsulam ovato-oblongam acuminatam subsequantibus.  $\frac{1}{2}$  in fossis et humidis. — *J. articulatus* eng. bot. t. 238 non L. *J. sylvaticus* var. Fl. fr. 3. p. 169. — Moris. s. 8. t. 9. f. 8. n. 1. Capsula fusca. (v. v.)
21. *J. ASPÆUS* (Req. vaubl. 2. p. 253.) culmo repente, ramis extra axillas ortis, foliis teretibus nodoso-articulatis ad ramorum axillas aitis, floribus capitatis in paniculam simplicem digestis, perigonii segmentis lanceolatis acuminatis.  $\frac{1}{2}$  in humidis propè Avenionem secùs Drunetiam (cl. Requier), propè Cherbourg (cl. Le Prév.). — Fl. fr. 5. p. 308. An præcedentis var? (v. s. sine fructu.)
22. *J. OBTUSIFLORUS* (Ehr. gram. 76.) culmo tereti lævi, foliis teretibus nodoso-articulatis, floribus capitatis, capitulis sessilibus pedunculatisve in paniculam suprâ-decompositam divaricatam digestis, perigonii segmentis ellipticis obtusis capsulam ovatam triquetram acuminatam subsequantibus.  $\frac{1}{2}$  in fossis et aquosis. — *J. articulatus* Fl. fr. 3. p. 169 an L? eng. bot. 30. t. 2144. (v. v.)
23. *J. MULTIFLORUS* (Desf. atl. 1. p. 313. t. 91.) culmo tereti lævi, foliis teretibus fistulosis acuminatis culmum subsequantibus, floribus solitariis sessilibus pedicellatisve in paniculam elongatam suprâ-decompositam erectam digestis, pedunculis inæqualibus multifariam decompositis fasciculatis, perigonii segmentis lanceolatis acuminatis capsulam ovatam trigonam breviter mucronatam subsequantibus.  $\frac{1}{2}$  in maritimis circâ Telonem detexit cl. Requier, in Corsicâ ex Viv. Plantâ 3-pedalis et ultra. (v. s.)
24. *J. JACQUINI* (L. mant. 63.) culmo lævisculo foliis 1-2 instructo basi squamoso, foliis subulatis cylindricis fistulosis, capitulo terminali spathâ subulatâ breviori squamis ovatis imbricatis (nigris) nitidis cincto, perigonii segmentis lanceolatis acuminatis capsulam ovoideam obtusam mucronatam subsuperantibus.  $\frac{1}{2}$  in Alpibus Delphinatâs. — Fl. fr. 3. p. 166. *J. biglumis* Jacq. vind. t. 4. f. 2. austr. 3. t. 221. non L. — Scheuchz. gram. 323. t. 7. f. 9. (v. s.)
25. *J. TRIGLUMIS* (L. sp. 467.) culmo tereti striato basi folioso, foliis flexuosis subcanaliculatis culmo duplò brevioribus, capitulo trifloro terminali erecto

# 478 MONOCOTYL. SEU ENDOGENÆ.

aphyllo bracteato, perigonii segmentis lanceolatis subobtusis capsulâ tumidiusculâ ovato-oblongâ obtusâ mucronatâ brevioribus.  $\frac{1}{2}$  in Alpibus Delph., Galloprovinciæ. — Fl. fr. 3. p. 166. L. lapp. t. 10. f. 5. Fl. dan. t. 132. (v. s.)

IV. LUZULA DC. fl. fr. 3. p. 158. — Junci sp. L. — Juncoides Michel. gen. t. 51.

Perigonium 6-partitum glumaceum; stamina 6; capsula 1-localaris, 3-sperma, 3-valvis, valvulis septo destitutis. — Folia plana hinc inde pilosa.

1. L. NIVEA (DC. l. c.) radice repente, culmo lævi, foliis linearibus pilosis, corymbo composito, pedunculis sub-5-floris, perigonii segmentis angustè lanceolatis acutis, interioribus duplò longioribus.  $\frac{1}{2}$  in Alpibus et montanis. — Juncus niveus L. — Scheuchz. gram. 3ao. t. 7. f. 7. — Moris. s. 8. t. 9. f. 7. n. 39. Fl. albi. (v. v.)
2. L. ALBIDA (DC. fl. fr. 3. p. 159.) radice fibrosâ, culmo striato, foliis linearibus pilosis, corymbo decomposito, pedunculis subquadrifloris, perigonii segmentis angustè lanceolatis acutis, interioribus paulò longioribus, capsulâ ovatâ trigonâ acuminatâ perigonio paulo brevior.  $\frac{1}{2}$  in nemorosis passim. — Junc. albidus Hoffm. J. niveus Leers herb. t. 13. f. 6. non L. J. pilosus s. L. Fl. albid. (v. v.)
3. L. LUTEA (DC. l. c.) foliis glabris lato-linearibus acuminatis, corymbo composito conferto, pedunculis multifloris, perigonii segmentis lanceolatis mucronatis subæqualibus splendentibus capsulam ovato-rotundam trigonam obtusam mucronatam æquantibus.  $\frac{1}{2}$  in summis Alpium et Pyrenæorum pratis. — J. luteus All. Vill. dauph. 2. t. 6. Fl. lutei. (v. v.)
4. L. VERNALIS (DC. l. c.) foliis lineari-lanceolatis linearibusve acutis pilosis, corymbo subsimplici laxissimo, pedunculis unifloris nutantibus, perigonii segmentis lanceolatis acutis æqualibus capsulâ globosâ obtusâ brevioribus.  $\frac{1}{2}$  in sylvis vere frequens. — J. vernalis Ehr. J. pilosus Leers herb. t. 13. f. 10. Fl. spadicei. (v. v.)
5. L. FORSTERI (DC. ic. gall. rar. 1. t. 2.) foliis linearibus angustis acuminatis pilosis, corymbo subsimplici laxissimo, pedunculis unifloris erectis, perigonii segmentis lanceolatis acuminatis æqualibus capsulam ovatam acutè trigonam mucronatam æquantibus.  $\frac{1}{2}$  in montosis passim. — Fl. fr. 5. p. 304. J. Forsteri eng. bot. t. 1293. Fl. spadicei. (v. v.)
6. L. FLAVESCENS (Gaud. agrost. 2. p. 239.) foliis angustissimis parcè pilosis, corymbo umbellato erecto paucifloro, pedunculis subunifloris subæqualibus erectis, perigonii segmentis lanceolatis acuminatis æqualibus capsulam ovatam acutè trigonam acuminatam æquantibus.  $\frac{1}{2}$  in sylvis montanis Jurassi et Pyren. orientalium. — Fl. fr. 5. p. 304. L. Hostii Desv. Journ. 1. t. 6. f. 2. J. flavescens Host. gram. 3. t. 94. An satis à priore distincta? Fl. flavescens. (v. s.)
7. L. GLABRATA (Desv. Journ. 1. t. 5. f. 3.) foliis lanceolato-linearibus acuminatis glabris, corymbo decomposito divaricato multifloro erecto laxo, pedicellis 2-4-floris, floribus pedicellatis, perigonii segmentis lanceolatis acuminatis capsulam ovato-rotundam nitidam mucronatam subsuperantibus.  $\frac{1}{2}$  in parvis humidis Montis-Aurei, Pyren. orient. (Ph. Thomas). — Fl. fr. 5. p. 304. J. intermedius Host. gram. 3. t. 99. Fl. spadicei. (v. v. DC.)
8. L. PARVIFLORA (Desv. Journ. 1. p. 144.) foliis lanceolato-linearibus ad vaginæ faucem parcè pilosis cæterum glabris, corymbo composito divaricato erecto multifloro, pedicellis omnibus unifloris, perigonii segmentis lanceolatis acuminatis capsulam subsuperantibus.  $\frac{1}{2}$  in montosis Pyrenæorum et Alpium rarè. — Fl. fr. 5. p. 305. J. parviflorus Rostk. junc. t. 1. f. 1. Præcedenti nimium affinis. (v. s. s. fructu.)

9. *L. SPADICEA* (DC. fl. fr. 3. p. 159.) foliis glabris linearibus acuminatis, vaginis fauce pilosis, corymbo decomposito divaricato laxo, pedunculis subquadrifloris, perigonii segmentis lanceolatis acuminatis capsulam ovato-rotundam trigonam breviter mucronatam æquantibus. 2/ in pratia humidis Alpium, montium Arvernæ et Pyrenæorum. — *J. spadiceus* All. Vill. dauph. 2. t. 6. f. 3. Fl. spadicei. (v. v.)
10. *L. MAXIMA* (DC. fl. fr. 3. p. 160.) foliis lanceolato-linearibus acutis pilosis, corymbo decomposito multifloro erecto laxo, pedunculis elongatis divaricatis subtrifloris, perigonii segmentis lanceolatis acuminatis capsulâ ovatâ trigonâ acuminatâ subbrevioribus. 2/ in montosis sylvaticis. — *J. pilosus* f. L. *J. maximus* Retz. Willd. Fl. nitidi fusco et albo variegati. (v. v.)
11. *L. MULTIFLORA* (Lej. spa 169.) radice fibrosâ, foliis angustis pilosis, floribus capitatis, capitulis ovatis multifloris in corymbum umbellatum erectum dispositis, spiculis longè pedicellatis, perigonii segmentis lanceolatis acuminatis capsulâ ovatâ trigonâ mucronatâ subbrevioribus. 2/ in sylvis et paludosis passim. — Fl. fr. 5. p. 306. *J. multiflorus* Hoffm. An satis à *L. campestris* distincta? (v. v.)
12. *L. CAMPESTRIS* (DC. fl. fr. 3. p. 161.) radice repente, foliis angustis plus minus pilosis, floribus capitatis, capitulis multifloris in corymbum simplicem brevem coarctatum dispositis, spiculis pedunculatis, perigonii segmentis lanceolatis acuminatis capsulâ ovato-globosâ trigonâ acutâ longioribus. 2/ in pascuis, nemorosis, montanis. — *J. campestris* L. Leers. herb. t. 13. f. 5. Capsula flava. Flores fusco et albo variegati. (v. v.)  
*β. congesta*, floribus capitatis, capitulis in corymbum simplicem capitatum congestis. — *L. congesta* Lej. spa 163. Fl. fr. 5. p. 305. *L. campestris β.* Fl. fr. 3. p. 161. (v. v.)
13. *L. SUDETICA* (DC. fl. fr. 5. p. 306.) radice repente, foliis linearibus angustis ad faucem pilosis cæterum glabris, floribus capitatis, capitulis ovato-rotundis in corymbum simplicem erectum coarctatum folio florali brevioribus congestis, perigonii segmentis lanceolatis mucronatis capsulâ ovato-rotundâ trigonâ obtusâ nitidâ subbrevioribus. 2/ in humidis torfaceis montanis Alastiæ, Alpium, Arvernæ, Cebennarum, Pyrenæorum. — *L. nigricans* Desv. journ. 1. p. 158. *J. spicatus* Krock. sil. t. 62. Nimiùm præcedenti affinis. (v. v.)
14. *L. SPICATA* (DC. fl. fr. 3. p. 161.) foliis linearibus angustissimis glabris fauce pilosis, spicâ racemosâ nutante basi compositâ, perigonii segmentis lanceolatis acuminatis capsulam ovato-globosam obsolete trigonam obtusam brevissimè mucronatam subæquantibus. 2/ in summis pratibus Jurassi supra Gex (Babey), Alpium, Montis-Aurei (DC.), Pyrenæorum ex Benth. — *Juncus spicatus* L. lapp. t. 10. f. 4. fl. dau. t. 270. (v. v.)
15. *L. PEDIFORMIS* (DC. fl. fr. 3. p. 162.) foliis linearibus pilosis, spicâ racemosâ nutante basi compositâ et lobatâ, perigonii segmentis lanceolatis acuminatis capsulam ovato-globosam obsolete trigonam acutam subsuperantibus. 2/ in montibus Delph., Gallop., Pyren. orient. — *Juncus pediformis* Vill. dauph. 2. t. 6. (v. v.)

## 480 MONOCOTYL. seu ENDOGENÆ.

### Ordo CVI. PALMÆ *Lin. ord. nat. n. 1. Juss. gen. 37.*

*Perigonium sexpartitum persistens. Stamina sæpius 6, imis perigonii laciniis inserta. Ovarium 1 superum. Styli 1 aut 3. Fructus baccatus aut drupaceus, 1-3-locularis, 1-3-spermus. Semina ossea. Embryo minimus in cavitate albuminis magni demum cornei. — Caulis fruticosus. Folia petiolata, basi vaginantia, pinnato-aut-palmato-lobata. Spadix inter folia nascens, spathâ maximâ involutus. Flores numerosi sæpius abortu diclini.*

#### I. CHAMÆROPS *L. gen. 1219.*

*Spatha univalvis; flores nunc masculi nunc hermaphroditi; perigonii laciniæ 3 externæ minores bracteiformes; stamina 6, filamentis crassis coalitis; styli 3; baccæ 3 monospermæ.*

1. *C. nummulus* (*L. sp. 1657.*) frondibus palmatis plicatis, caudicibus spinosis.  
 3 circa Nicæam loco dicto St.-Hospice. — *Fl. fr. 3. p. 724. Lam. ill. t. 900.*  
 (v. v.)

### Ordo CVII. AROIDEÆ *Juss. gen. p. 23.*

*Flores monoici sessiles circa spadicem simplicem sæpius spathâ monophyllâ involutum, mox nudi mox squamis stipati. Perigonium nullum. Masc. Stamina definita aut indefinita. Antheræ 1-aut-2-loculares. Fem. Ovaria nunc staminibus commixta, nunc ab iisdem segregata, 1-locularia rariùs 3-locularia polysperma; styli stigmataque totidem. Fructus baccatus rotundatus aut rariùs capsula aliquando abortu monosperma. Embryo rectus in centro albuminis carnosus aut farinaceus; radícula infera. — Herbæ acaules aut caulescentes. Folia alterna aut radicalia, basi subvaginantia, interdum pedata aut basi cordata.*

#### Trib. I. AROIDEÆ VERÆ *Brown prod. p. 234.*

*Flores squamis destituti. Fructus baccatus.*

#### I. ARUM *L. gen. 1028.*

*Spadix apice nudus; antheræ multiseriales ad dimidiam spadicis longitudinem; ovaria basi spadicis inserta; bacca unilocularis monosperma.*

#### Sect. 1. DRACUNCULUS *Tourn. inst. t. 70.*

*Spatha basi non tubulosa, glandulæ acutæ (stamina abortiva) cirriformes 2-3-seriales supra antheras spadici insertæ.*

1. *A. MUSCIVORUM* (Linn. f. suppl. 410.) foliis radicalibus pedatis integerrimis, spadice cylindraceo apice crinito spathâ ovatâ planâ internè pilosâ breviorè.  $\frac{1}{2}$  in Corsicâ propè Calvi et insulis circumjacentibus (cl. Soleirol). — *A. crinitum* Willd. sp. 4. p. 477. (v. s.)
2. *A. DRACUNCULUS* (L. sp. 1367.) foliis radicalibus pedatis, lobis integris, spadice lanceolato spathâ ovatâ planâ glabrâ longiorè.  $\frac{1}{2}$  in umbrosis incultis Galliæ meridionalis, Aginni ex St.-Am., Andegaviæ propè Sanctum-Macarium ex Bast. — Fl. fr. 3. p. 151. Lam. ill. t. 740. f. 2. — Lob. ic. t. 600. f. 1. (v. v.)
3. *A. VULGARE* (Lam. fl. fr. 3. p. 537.) foliis radicalibus hastato-sagittatis, lobis deflexis, spadice clavato spathâ breviorè.  $\frac{1}{2}$  ad sepes et sylvas. — Fl. fr. 3. p. 152. *A. maculatum* L. *A. Pyrenæum* Lapeyr. ex cl. Benth. — Tab. ic. 746. — Lob. ic. 597. Variat foliis maculatis immaculatiave. (v. v.)
4. *A. ITALICUM* (Mil. dict. n. 2.) foliis radicalibus albo-venosis hastato-sagittatis, lobis auriculatis divaricatis, spadice clavato spathâ breviorè.  $\frac{1}{2}$  in sylvaticis et submontanis Galliæ australis et austro-occidentalis. — Fl. fr. 3. p. 152. — Sabb. hort. rom. 2. t. 75. (v. v.)
5. *A. PICTUM* (L. f. suppl. 410.) foliis radicalibus cordatis crassiusculis venis canaliculatis et interdum albidis insculptis, spadice clavato spathâ breviorè.  $\frac{1}{2}$  in Corsicâ propè Ajaccio. — Fl. fr. 5. p. 303. (v. s.)

Sect. II. *ARISARUM* Tourn. inst. t. 70.

Spatha basi tubulosa, glandulæ cirrhiformes nullæ; ovarium stylo coronatum.

6. *A. ARISARUM* (L. sp. 1370.) foliis radicalibus hastato-sagittatis mucronatis, lobis deflexis oblongis obtusis, spadice cylindraceo incurvato spathâ cucullatâ breviorè.  $\frac{1}{2}$  in umbrosis et petrosis Corsicæ, Niceæ, Galloprovinciæ. — Fl. fr. 3. p. 152. *Arisarum vulgare* Rich. — Barr. ic. t. 573. 1130. (v. s.)

II. *CALLA* L. gen. 1030.

Spadix undique tectus staminibus et ovariis intermixtis; bacca multilocularis polysperma.

1. *C. PALUSTRIS* (L. sp. 1373.) foliis cordatis, spathâ planâ, spadice undique antheris ovariisque tecto.  $\frac{1}{2}$  in fossis aquosis Alsatiæ, Lotharingiæ, Vogesorum. — Fl. fr. 3. p. 153. Lam. ill. t. 739. f. 1. — Lob. ic. t. 600. f. 2. — Barr. ic. t. 574. (v. s.)

Trib. II. *ORONTIACEÆ* Brown prod. 337.

Flores squamis perigoniiiformibus stipati. Fructus capsularis.

III. *ACORUS* L. gen. 434.

Spatha nulla; perigonium persistens globosum 6-partitum; stam. 6 perigonii laciniis opposita; ovarium globosum 3-loculare polyspermum; stigma sessile; capsula perigonio stipata.

1. *A. CALAMUS* (L. sp. 461. excl. var. *a.*) scapi mucrone longissimo foliaceo.  $\frac{1}{2}$  in fossis aquaticis Alsatiæ, Delphinatûs, propè Beauvais (cl. Graves), Rennes ex cl. de Bréb. — Fl. fr. 3. p. 157. Lam. ill. t. 252. — Moris. s. 8. t. 13. f. 11. n. 4. (v. v.)

Ordo CVIII. TYPHACEÆ *Juss. gen.* 25.

Flores monoici in amenta unisexualia aggregati. Perigonium triphyllum aut obsoletum. *Masc.* Stamina 3. *Fem.* Ovarium unicum liberum monospermum, ovulo pendulo; stylus 1; stigmata 1-2. Fructus monospermus. Embryo rectus in centro albuminis carnosi aut farinacei. Radicula infera. — *Herbæ* enodes aquaticæ; folia alterna ensiformia subvaginantia.

I. TYPHA *Tourn. inst.* 301. *L. gen.* 1040.

Amenta cylindrica. *Masc.* Perigonium triphyllum obsoletum; stamina 3 infernè in filamentum unicum coalita. *Fem.* Perigonium nulum, fructus pedicellatus, pedicello basi pilis longis pappi iustar cincto.

1. *T. LATIFOLIA* (L. sp. 1377.) foliis linearibus planis, spicâ masculâ femineâque approximatis, utrâque cylindricâ.  $\mathcal{U}$  in paludibus. — *Fl. fr.* 3. p. 148. *Lam. ill. t.* 748. f. 1. — *Lob. ic. t.* 81. f. 1. (v. v.)
2. *T. ANGUSTIFOLIA* (L. sp. 1377. var.  $\alpha$ .) foliis linearibus canaliculatis, spicâ masculâ femineâque remotis, utrâque cylindricâ.  $\mathcal{U}$  in paludibus. — *Fl. fr.* 3. p. 148. *Lam. ill. t.* 748. f. 2. *Fl. dan. t.* 815. (v. v.)
3. *T. MEDIA* (Schl. cat. 59.) foliis linearibus angustis planis, spicâ masculâ femineâque remotis utrâque cylindricâ.  $\mathcal{U}$  in paludosis passim. — *Fl. fr.* 5. p. 302. — *Moris. s.* 8. t. 13. f. 9. n. 2. (v. v.)
4. *T. MINIMA* (Hoppe exs. Hoffm. germ. 4. p. 251.) foliis setaceo-linearibus canaliculatis, spicâ masculâ femineâque remotis, femineâ ellipticâ.  $\mathcal{U}$  in arenosis udibus Alsatiæ, Delphin., Lugd., Arelatis, Avenionis, Nicæ. — *Fl. fr.* 3. p. 148. excl. syn. Sm. Lin. — *Lob. ic.* 81. t. 2. — *Moris. s.* 8. t. 13. f. 10. n. 3. (v. v.)

II. SPARGANIUM *Tourn. inst.* 302. *L. gen.* 1041.

Amenta globosa; perigonium triphyllum; fructus sessilis turbinatus setulis basi destitutus.

1. *S. NATANS* (L. sp. 1378.) foliis decumbentibus planis, pedunculo communi simplici, stigmate ovato brevissimo, capitulo masculino subsolitario.  $\mathcal{U}$  in rivulis stagnis et lacubus passim. — *Fl. fr.* 3. p. 150. *eng. bot. t.* 273. (v. v.)
2. *S. SIMPLEX* (Roth. germ. 3. p. 469.) foliis basi triquetris lateribus planis, pedunculo communi simplici, stigmate lineari.  $\mathcal{U}$  in stagnis. — *Fl. fr.* 3. p. 149. *eng. bot. t.* 745. *S. erectum*  $\beta$ . L. — *Lob. ic. t.* 80. f. 2. (v. v.)
5. *S. RAMOSUM* (C. Bauh. pin. 153.) foliis basi triquetris lateribus concavis, pedunculo communi ramoso, stigmate lineari.  $\mathcal{U}$  in fossis et ad ripas flaviorum. — *Fl. fr.* 3. p. 149. *eng. bot. t.* 744. *S. erectum*  $\alpha$ . L. — *Lob. ic. t.* 80. f. 1. (v. v.)

Ordo CIX. CYPERACEÆ *Juss. gen.* 26.

Flores glumacei spicati, hermaphroditi aut diclini. Gluma aut squama univalvis. Perigonium proprium nullum. Stamina 3, filamentum capillare; antheræ apice acuminatæ basi cordatæ. Ovarium liberum simplex; stylus unicus; stigmata 2-3. Fructus (akenium) triangularis aut compressus monospermus non dehiscens. Albumen farinaceum; embryo minimus ad basin albuminis. — Herbæ perennes gramineis similes sed plerumque enodes; folia vaginantia, vaginâ integrâ.

I. CYPERUS *Tourn. inst.* 299. *L. gen.* 66.

Flores hermaphroditi; glumæ paleaceæ distiche imbricatæ carinatæ; spiculæ compressæ; stylus deciduus; semen basi nudum.

1. *C. MONTEI* (L. supp. 102.) culmo triquetro basi folioso, involuero umbellam decompositam paniculatam superante, foliolis 3-6 carinato-plicatis inæqualibus, pedunculis longis nudis, spiculis alternis lanceolato-linearibus laxis, stigmatibus 2, glumis demum distinctis obtusis, seminibus ovato-rotundis subcompressis breviter mucronatis.  $\mathcal{L}$  secus Varum (Stire), Rhodanum inferiorem, ad ripas Aturis inferioris. — Fl. fr. 3. p. 146. et 5. p. 302. — Monti gram. 12. t. 1. f. 2. (v. v.)
2. *C. LONGUS* (L. sp. 67.) culmo triquetro folioso, involuero foliolis 3-4 linearibus planis plicatis umbellam foliosam supra-decompositam longe superantibus, pedunculis nudis longissimis, spiculis alternis linearibus angustis laxiusculis, stigmatibus 3, glumis dense imbricatis retusis, seminibus oblongis trigonis submucronatis.  $\mathcal{L}$  in paludosis Galliæ occidentalis australisque rariis, Lotharingæ ex Vill., circa Parisios ex Mérat. — Fl. fr. 3. p. 145. Jacq. ic. rar. 2. t. 297. — Lob. ic. t. 75. f. 2. — Moris. s. 8. t. 11. f. 7. n. 13. (v. v.)
3. *C. ACUTUS* (L. sp. 67.) radicem fibrillis hinc inde in tuberculos ovatos inflatis, culmo triquetro nudo, involuero foliolis 3-4 linearibus planis plicatis umbellam decompositam paniculatam superantibus, spiculis alternis elongato-linearibus laxiusculis, stigmatibus 3, glumis acutiusculis, seminibus ovoideis 3-gonis obtusis.  $\mathcal{L}$  in omni regione mediterraneâ. — Fl. fr. 3. p. 146. et 5. p. 301. *C. olivaris* Targ. *C. esculentus* Geu. — Scheuchz. gram. t. 9. f. 3. — Camer. epit. p. 10. ic. (v. v.)
4. *C. ESCULENTUS* (L. sp. 67.) radicem tuberibus subrotundis terminalibus, culmo triquetro nudo, umbellâ foliosâ.  $\mathcal{L}$  in Galloprovinciâ cultus. — Fl. fr. 3. p. 146. — Lob. ic. t. 78. f. 1. 2. — Moris. s. 8. t. 11. f. 6. n. 10. (v. s. c. sine flore.) †.
5. *C. FUSCUS* (L. sp. 69.) culmo triquetro, involuero 3-phylo, foliolis planis umbellam subcompositam longè superantibus, spiculis linearibus confertis, stigmatibus 3, glumis acutiusculis, seminibus acutè triquetris. (I) in paludosis. — Fl. fr. 3. p. 145. — Moris. s. 8. t. 11. f. 14. n. 38. — J. Bauh. hist. 2. p. 471. ic. (v. v.)
6. *C. FLAVASCENS* (L. sp. 68.) culmo triquetro, involuero triphylo, foliolis planis plicatisve umbellam subcompositam superantibus, spiculis lanceolatis confertis, stigmatibus 2, glumis obtusis, seminibus rotundatis subcompressis mucronatis. (I) in paludosis. — Fl. fr. 3. p. 145. Lam. ill. t. 38. f. 1. — Moris. s. 8. t. 11. f. 13. n. 37. (v. v.)
7. *C. FASCICULARIS* (Rothb. cyp. 39. t. 11. f. 1.) culmo triquetro basi foliis 2-3 carinato-plicatis instructo, involuero polyphylo umbellam longè superante,

## 484 MONOCOTYL. SEU ENDOGENÆ.

spiculis lineari-lanceolatis corymboso-capitatis confertissimis, glumis obtusis, seminibus lanceolato-ovatis breviter mucronatis.  $\mathcal{U}$  circa Nicæam secus Varum. — Fl. fr. 3. p. 722. Lam. ill. t. 38. f. 2. — Pluck. t. 416. f. 6. (v. s.)

8. *C. JUNCIFORMIS* (Cav. ic. t. 204. f. 1.) culmo subtereti, spiculis lineari-lanceolatis sessilibus aggregatis, spathâ diphyllâ foliolo altero spiculis breviora, glumis mucronatis.  $\mathcal{U}$  in humidis circa Nicæam. — Fl. fr. 3. p. 144. Desf. atl. 1. t. 7. f. 1. — *C. distachyos* All. auct. t. 2. f. 5. (v. v. DC.)

### II. SCHÆNUS L. gen. 65.

Flores hermaphroditi; glumæ paleaceæ undique imbricatæ, inferiores steriles; semen nudum aut setulis glumâ brevioribus basi cinctum.

Sect. I. SCHÆNIS. — Schænus Vahl. enum. 2. p. 207.

Stylus deciduus.

\* *Semine setulis basi cincto.*

1. *S. VERBUGINUS* (L. sp. 64.) culmo tereti nudo, capitulo lineari-elliptico, spiculis 2-3, involucri diphylli valvulis subæqualibus spicas non superantibus.  $\mathcal{U}$  in montibus Jurassi et Galloprovinciæ. — Fl. fr. 3. p. 142. excl. syn. Moris. Schrad. germ. 1. t. 1. f. 4. (v. v.)
2. *S. NIGRICANS* (L. sp. 64.) culmo tereti nudo, capitulo ovato, spiculis numerosis, involucri diphylli valvulâ alterâ subulatâ longiore capitulum superante.  $\mathcal{U}$  in paludibus et inundatis. — Fl. fr. 3. p. 142. eng. bot. t. 1121. — Schenckz. agr. t. 7. f. 13. 14. 15. — Magn. monsp. p. 144. ic. (v. v.)
3. *S. COMPRESSUS* (L. sp. 65.) culmo teretiusculo basi folioso, foliis planis, spiculis numerosis sessilibus in spicam oblongam disticho-compressam approximatis.  $\mathcal{U}$  in pratis humidis passim. — eng. bot. t. 791. Poll. pal. t. 1. f. 3. Scirpus caricis Retz. Fl. fr. 3. p. 137. Host. gram. 3. t. 57. Gluma infima plerumque sterilis. (v. v.)

\*\* *Semine basi nudo.*

4. *S. MUCRONATUS* (L. sp. 63.) culmo tereti nudo, foliis canaliculatis, spiculis ovatis sessilibus densè fasciculatis, involucri polyphylo patente umbellam longè superante.  $\mathcal{U}$  in maritimis meridionalibus. — Fl. fr. 3. p. 144. — Lob. ic. t. 87. f. 1. (v. v.)
5. *S. MARISCUS* (L. sp. 62.) culmo tereti folioso, foliis margine dorsoque aculeato-serratis, spiculis bifloris ovato-acutis sessilibus capitatis, capitulis pedicellatis sessilibusve in corymbis numerosos simplices decompositosve digestis.  $\mathcal{U}$  in paludibus passim. — Fl. fr. 3. p. 143. eng. bot. t. 950. Cladium germanicum Schrad. germ. 1. t. 5. f. 7. — Lob. ic. t. 76. f. 1. — Moris. s. 8. t. 11. f. 12. n. 24. (v. v.)

Sect. II. RHYNCHOSPORA Vahl. enum. 2. p. 229.

Semen coronatum stylo persistente indurato, basi latitudine seminis, setulis basi cinctum.

6. *S. ALBUS* (L. sp. 65.) radice fibrosâ, culmo trigono folioso diviso, foliis carinatis angustissimis, floribus fasciculatis (albis), fasciculis terminalibus subcorymbosis involucri subæquantibus.  $\mathcal{U}$  in tarfosis et paludosis passim. — Fl. fr. 3. p. 143. eng. bot. t. 985. Rhynchospora alba Vahl. l. c. — Schenckz. agr. t. 11. f. 11. (v. v.)
7. *S. FUSCUS* (L. sp. 1664.) radice repente, culmo teretiusculo folioso diviso, foliis canaliculato-setaceis, floribus fasciculatis (fuscis), fasciculis terminalibus subgemminatis involucri longè minoribus.  $\mathcal{U}$  in pratis humidis Gallie occidentalis et boreali occidentalis. — Fl. fr. 3. p. 143. eng. bot. t. 1575. Rhynch. alba  $\beta$  fusca Vahl. R. fusca Rœm. et Schult. 2. p. 88. — Moris. s. 8. t. 11. f. ult. n. 40. (v. s.)



III. SCIRPUS L. *gen.* 67.

Flores hermaphroditi; glumæ paleaceæ undique imbricatæ, omnes fertiles; semen nudum aut setulis glumâ brevioribus basi cinctum.

Sect. I. FIMBRISTYLIS Vahl. *enum.* 2. p. 285.

Stylus bifidus basi bulbosus compressus margine ciliatus, setæ hypogynæ nullæ.

1. *S. annuus* (All. ped. t. 88. f. 5.) culmo triquetro nudo basi folioso, foliis setaceis pubescentibus, paniculâ corymbosâ terminali foliaceâ, spiculis oblongis acutis pedunculatis, intermediâ sessili, glumis ovatis mucronatis. (I) *Nicea secus Varum* (Stire). — Fl. fr. 3. p. 141. *Fimbristylis dichotoma* Vahl. F. *annua* Rœm. et Sch. 2. p. 95. — Pluck. t. 119. f. 3. (v. s.)

Sect. II. ELEOCHARIS R. Brown *prod.* 1. p. 224.

Setæ hypogynæ (4-12) denticulatæ rarò nullæ; fructus basi styli dilatata incurvata coronatus.

2. *S. palustris* (L. sp. 70.) radice repente, culmo tereti basi vaginâ aphyllâ instructo, spicâ ovato-oblongâ nudâ terminali, glumis imbricatis obtusis, seminibus ovato-globosis subcompressis. ♀ in paludibus et fossis. — Fl. fr. 3. p. 133. eng. bot. t. 131. — Lob. ic. t. 86. f. 1. *Eleocharis palustris* Rœm. et Sch. 2. p. 151. — Scheuchz. gram. t. 7. f. 17. (v. v.)
3. *S. multicaulis* (Sm. brit. 1. p. 48.) radice repente, culmo tereti striato basi vaginâ aphyllâ instructo, glumis obtusis externis æqualibus spicâ terminali sæpè viviparâ brevioribus, floribus triandris, seminibus ovato-globosis trigonis. ♀ in aquosis propè Lutetiam, Rothomagum, in agro Soloniensi, Monte-Aureo, Lozerâ, Corsicâ. — Fl. fr. 5. p. 299. St.-Hil. in *journal. bot.* 1804. 3. t. 21. An satis à sequente distinctus? (v. v.)
4. *S. Bzothryxon* (L. supp. 103.) radice fibrosâ, culmo tereti striato basi vaginâ aphyllâ instructo, glumis obtusis, externis spicâ terminali ovatâ brevioribus inæqualibus, seminibus ovatis 3-gonis. ♀ in paludosis. — Fl. fr. 3. p. 135. *S. campestris* Roth. Fl. fr. 3. p. 136. *S. Halleri* Vill. *S. pauciflorus* eng. bot. t. 1122. — Scheuchz. gram. t. 7. f. 21. (v. v.)
5. *S. ovatus* (Roth. cat. 1. p. 5.) radice fibrosâ, culmo tereti filiformi basi vaginâ aphyllâ instructo, spicâ ovatâ nudâ, floribus diandris, glumis arcetè imbricatis obtusissimis, seminibus obovato-globosis subcompressis. (I) in humidis passim. — Fl. fr. 3. p. 134. et 5. p. 299. *Eleocharis ovata* Rœm. et Sch. 2. p. 152. *S. turgidus* Pers. — Moris. s. 8. t. 10. f. 19. n. 34. (v. s.)

Sect. III. ISOLEPIS R. Brown *prod.* 1. p. 221.

Setæ hypogynæ nullæ; stylus deciduus.

\* *Spiculis terminalibus solitariis.*

6. *S. fluitans* (L. sp. 71.) culmo tereti folioso flaccido, foliis setaceis planis, pedunculis alternis nudis, spicis ovato-globosis, glumis ovatis obtusis. ♀ in fossis et stagnis Paris., Andeg., Monsp. ex Gouan, Galloprovinciæ ex Ger. — Fl. fr. 3. p. 139. *Isolepis fluitans* R. Br. l. c. — Moris. s. 8. t. 10. f. 13. n. 31. — Pluck. t. 35. f. 1. (v. v. DC.)
7. *S. acicularis* (L. sp. 71.) culmo simplici quadrangulo gracili basi vaginâ aphyllâ instructo, spicâ ovatâ acutâ terminali, glumis acutis exterioribus acuminatis majoribus. ♀ in paludibus et inundatis passim. — Fl. fr. 3. p. 138. eng. bot. t. 749. *Eleocharis acicularis* Rœm. et Sch. 2. p. 154. — Moris. s. 8. t. 10. f. 21. n. 37. — Pluck. t. 40. f. 7. (v. v.)

\* \* *Spiculis lateralibus aggregatis raro solitariis.*

8. *S. setaceus* (L. sp. 73.) culmo setaceo gracili simplici nudiusculo, spicis ad apicem calmi lateralibus subgeminis sessilibus ebracteatis, glumis ovatis obtusis rariùs mucronulatis, seminibus obovato-globosis longitudinaliter

- striatis. (X) in inundatis passim. — Fl. fr. 3. p. 139. *Isolepis setacea* Br. 1. p. 221. — Moris. s. 8. t. 10. f. 10. n. 23. (v. v.)
9. *S. SUPINUS* (L. sp. 73.) culmo tereti simplici nudo, spicis 3-6 ovatis sessilibus in capitulum laterale in medio culmo sessile glomeratis, glumis ovatis mucronatis, seminibus obovato-globosis 3-gonis breviter mucronatis transversè rugulosis. (X) in inundatis circà Argentoratnm (cl. Nestler), Parisios. — Fl. fr. 3. p. 140. *Isol. supina* Br. 1. c. (v. v. D. ad lacum Genev.)
- \* \* \* *Spiculis terminalibus aggregatis.*
10. *S. HOLOSCHÆNUS* (L. sp. 72.) culmo elongato tereti nudo basi vaginis 2-3 acuminatis instructo, capitulis subglobosis pluribus pedunculatis sessilibusque in paniculam umbellatam simplicissimam digestis, involnro diphylo inæquali mucronato, seminibus minutissimis. ♀ in pratis humidis provinc. austr. occidentaliūque. — Fl. fr. 3. p. 140. *Isolepis holoschænus* Rœm. et Sch. 2. p. 115. — Moris. s. 8. t. 10. f. 8. n. 17. — Pluck. t. 40. f. 4. (v. v.)
- ℓ. *Romanus*, capitulo laterali sessili. — *S. Romanus* L. sp. 73. Fl. fr. 3. p. 140. — Barr. t. 255. f. 3. (v. v.)
11. *S. MICHELIANUS* (L. sp. 76.) culmo triquetro basi folioso, foliis planis, spiculis in capitulum globosum aggregatis, involnro polyphylo capitulum longè superante, seminibus sublinearibus compressis acutis. (X) in pratis et arenis humidis Gallix occidentalis, agri Ruscinonensis ex Lapeyr., Mouspeliensis ex Gon. — Fl. fr. 3. p. 141. *Isol. Micheliana* Rœm. et Sch. 2. p. 114. — Till. pis. t. 21. f. 5. (v. v.)
12. *S. PUBESCENS* (Desf. atl. 1. t. 10.) culmo triquetro apice pedunculis apicisque pubescentibus, spicis confertis ovatis, glumis latè ovatis aristato-mucronatis, seminibus globosis trigonis retusis. ♀ in humidis Corsicæ propè Ajaccio. — Fl. fr. 5. p. 301. *Isol. pubescens* Rœm. et Sch. 2. p. 118. (v. s.)

Sect. IV. SCIRPOIS. — *Scirpus R. Br. l. c.*

Setæ hypogynæ squamis breviores; stylus deciduus.

\* *Spicis solitariis.*

13. *S. CESPITOSUS* (L. sp. 71.) radice fibrosâ, culmo tereti sulcato rigido basi vaginis mucronatis instructo, glumis obtusiusculis externis maximis spicam ovali-oblongam terminalem subæquantibus, seminibus ovato-globosis trigonis brevissimè mucronatis. ♀ in paludosis montanis et ericetis turfosis. — Fl. fr. 3. p. 135. eng. bot. t. 1029. — Scheuchz. gram. t. 7. f. 18. — Pluck. t. 40. f. 6. (v. v.)

\* \* *Spicis pluribus.*

14. *S. TRIQUETER* (L. mant. 29.) radice repente, culmo triquetro nudo ultra cymam producto erecto, cymâ laterali confertâ, spiculis ovato-globosis in capitula sessilia pedunculatave digestis, glumis ovatis breviter mucronatis, seminibus obovatis breviter mucronatis. ♀ in palustribus rariis. — Fl. fr. 3. p. 136. *S. mucronatus* Poll. Roth. non L. — Pluck. t. 40. f. 2. — Moris. s. 8. t. 10. f. 9. n. 20. (v. s.)
15. *S. LITTORALIS* (Schrad. germ. 1. t. 5. f. 7.) culmo triquetro apice rectiusculo basi vaginis 1-2 foliiferis instructo, spiculis oblongo-ovatis pedicellatis in cymam lateralem decompositam congestis, stigmatibus 2, glumis latè ovatis breviter mucronatis, seminibus oblongo-obovatis breviter mucronatis. ♀ in paludosis arenosis subsalsis secùs Mediterraneum passim. — Fl. fr. 5. p. 300. (v. v. DC.)
16. *S. MUCRONATUS* (L. sp. 73.) culmo triangulo nudo apice ultra capitulum producto subpatulo, spiculis ovatis in capitulum sessile laterale conglomeratis, glumis lato-ovatis obtusis vix mucronatis, stigmatibus 3, seminibus obovato-globosis compresso-trigonis breviter mucronatis. ♀ in pratis humidis rariis. — Fl. fr. 3. p. 137. — Scheuchz. gram. t. 9. f. 14. (v. s.)
17. *S. LACUSTRIS* (L. sp. 72.) culmo tereti elongato nudo ad basin vaginis 2-3

- scariosis instructo, cymâ terminali decompositâ, spiculis ovatis sessilibus pedunculatisque in capitula pedicellata agglomeratis, glumis lato-ovatis breviter mucronatis, seminibus ovato-globosis subcompressis breviter mucronatis.  $\mathcal{L}$  in paludibus et lacubus. — Fl. fr. 3. p. 136. eng. bot. t. 666. — Lob. ic. t. 85. f. 2. — Moris. s. 8. t. 10. f. 1. (v. v.)
18. *S. tenuifolius* (DC. fl. fr. 5. p. 300.) radice repente, culmo triquetro lævi basi foliis 1-2 planis apice triquetris instructo ultra capitulum producto erecto, spiculis 1-3 sessilibus in capitulum laterale sessile conglobatis, glumis lato-ovatis aristato-mucronatis externis apice trifidis, stylo bifido, seminibus ovato-globosis compressis breviter mucronatis.  $\mathcal{L}$  in paludibus maritimis pagi dicti Teste de Buch propè Burdigalam. (v. v. DC.)
19. *S. maritimus* (L. sp. 74.) culmo triquetro folioso, foliis linearibus planis, spiculis ovato-oblongis sessilibus in capitula sessilia pedunculata aggregatis, capitulis in paniculam cymosam simplicem terminalem conglobatis, glumis mucronatis externis trifidis, dente intermedio anbulato, seminibus obovatis trigonis breviter mucronatis.  $\mathcal{L}$  in lacubus et fossis. — Fl. fr. 3. p. 137. et 5. p. 301. eng. bot. t. 342. *S. tuberosus* Desf. — Schenckz. gram. t. 9. f. 7-10. — Lob. ic. t. 20. f. 1. (v. v.)
20. *S. sylvaticus* (L. sp. 75.) culmo triquetro folioso, foliis lato-linearibus planis, paniculâ corymbosâ terminali suprâ-decompositâ foliaceâ, spiculis ovatis sessilibus pedunculatisve confertis, glumis ovatis obtusis, seminibus minimis triquetris.  $\mathcal{L}$  in nemorosis humidis. — Fl. fr. 3. p. 138. eng. bot. t. 319. Lam. ill. t. 38. f. 2. — Moris. s. 8. t. 11. f. 8. n. 15. (v. v.)

IV. *ERIOPHORUM* L. gen. 68.

Flores hermaphroditi; glumæ paleaceæ undique imbricatæ; semen basi setis longissimis cinctum.

§. 1. *Spicis pluribus pedunculatis.*

1. *E. polystachyum* (L. sp. 76.) culmo tereti, foliis lato-linearibus planis apice triquetris, pedunculis maturis nutantibus sæpe ramosis, glumis obtusis, seminibus trigonis oblancoatis basi attenuatis obtusis.  $\mathcal{L}$  in pratis palustribus frequens. — Fl. fr. 3. p. 131. eng. bot. t. 563. *E. latifolium* Hoppe, Gand. fl. helv. 1. p. 129. *E. vulgare* Pers. ench. 1. p. 70. (v. v.)
- $\beta$ . *Vaillantii*, pedunculis abbreviatis spicâ brevioribus. — *E. Vaillantii* Poit. et Turp. t. 52. Fl. fr. 5. p. 298. *E. angustifolium*  $\beta$  *Vaillantii* Gaud. l. c. (v. s.)
2. *E. gracile* (Roth. cat. 2. p. 259.) culmo triquetro, foliis canaliculato-plicatis triquetris, pedicellis simplicibus erectis, glumis obtusis, seminibus oblongo-linearibus acutè triquetris retusis.  $\mathcal{L}$  in pratis paludosis rariùs. — Fl. fr. 3. p. 132. eng. bot. t. 2402. *E. triquetrum* Hoppe. (v. s.)
3. *E. angustifolium* (Reich. mænogr. n. 34.) culmo tereti, foliis linearibus canaliculato-plicatis superne triquetris, pedunculis maturis nutantibus omnibus simplicibus, glumis oblongo-lanceolatis acuminatis, seminibus subacatis lanceolatis acutè triquetris.  $\mathcal{L}$  in pratis paludosis rariùs. — Fl. fr. 3. p. 131. eng. bot. t. 564. *E. intermedium* East. in Journ. bot. 1814. 3. p. 19. Fl. fr. 5. p. 298. (v. v.)

§. 2. *Spicis solitariis terminalibus.*

4. *E. vaginatum* (L. sp. 76.) radice fibrosâ cespitosâ, culmo vaginato trigono, spicâ ovatâ scariosâ, antheris linearibus elongatis.  $\mathcal{L}$  in paludosis turfosis montosis. — Fl. fr. 3. p. 132. eng. bot. t. 873. *E. cespitosum* Host gram. 1. t. 39. — Scheuchz. ag. t. 7. f. 1. 2. 3. (v. v.)
5. *E. capitatum* (Hoffm. germ. 3. p. 26.) radice repente, culmo basi vaginato tereti, spicâ subglobosâ, antheris cordato-ovatis brevibus.  $\mathcal{L}$  in paludibus Alpium et Pyrenæorum. — Fl. fr. 3. p. 133. eng. bot. t. 2387. *E. Scheuchzeri* Hoppe. Pers. — Scheuchz. prod. t. 7. (v. v.)
6. *E. alpinum* (L. sp. 77.) radice subrepente, culmo angulato gracili nudius-

## 488 MONOCOTYL. SEU ENDOGENÆ.

culo, spicâ ovato-cylindricâ, setis ad quodque semen subæmis crispis.  $\mathcal{U}$  in paludosis Vogesorum, Jurassi, Alpium, montium Ligeris. — Fl. fr. 3. p. 133. eng. bot. t. 311. — Scheuchz. prod. t. 8. f. 1. agr. t. 7. f. 4. (v. v.)

V. KOBRESIA Willd. sp. 4. p. 205. DC. fl. fr. 5. p. 298.

Flores monoici, masculi et feminei in iisdem spicis permixti et sæpius sub iisdem squamis geminati et squamulâ distincti; spiculæ bifloræ; semina triquetra in urceolo minime inclusa.

1. K. SCIRPINA (Willd. sp. 4. p. 205.) spicâ solitariâ simplici tereti.  $\mathcal{U}$  in summis Alpibus Delph. et Pyrenæis. — Fl. fr. 5. p. 298. Carex Bellardi All. ped. t. 92. f. 2. Fl. fr. 3. p. 103. C. myosuroides Vill. dauph. 2. t. 6. C. Dufourii Lapeyr. abr. supp. 140. Elyna spicata Schrad. (v. v.)

VI. CAREX Gærtn. carp. 1. p. 12. t. 2. Good. trans. lin. 2.

Flores monoici aut rarius dioici, spiculis nunc androgynis nunc unisexualibus; stigmata 2-3; urceolus persistens ovarium et postea semen includens apice perforatus, capsulæformis.

### §. 1. Spicâ unicâ simplici; stigmatibus 2.

1. C. DIOICA (L. sp. 1379.) radice repente, foliis setaceis carinato-triquetris culmisque lævibus, spicis dioicis cylindricis angustis, fructibus erectiusculis confertis ovalis striatis mucronatis margine supernè serrulatis.  $\mathcal{U}$  in pratis spongiosis montanis. — Fl. fr. 3. p. 100. eng. bot. t. 643. Schk. trad. n. 1. A. t. A. n. 4. C. linneana Host. 3. t. 77. — Mich. gen. t. 32. f. 2. (v. v.)
2. C. DAVALLIANA (Smith. trans. lin. 5. p. 266.) radice fibrosâ, foliis setaceis carinato-triquetris culmisque retrorsum scabris, spicis masculis androgynisve linearibus, fructibus recurvatis confertis oblongo-subulatis triquetris nervosis margine supernè scabris patulis demum reflexis.  $\mathcal{U}$  in pratis spongiosis montanis. — Fl. fr. 3. p. 101. Schk. car. trad. n. 1. B. t. A. 2. W. n. 2. — Scheuchz. gram. t. 11. f. 10. (v. v.)
3. C. FULICARIS (L. sp. 1380.) foliis setaceis convoluto-canaliculatis, spicâ androgynâ supernè masculâ, fructibus oblongis laxis teretiusculis glabris utrinque attenuatis demum retroflexis squamâ persistenti longioribus.  $\mathcal{U}$  in pascuis limosis. — Fl. fr. 3. p. 101. Schk. car. trad. n. 3. t. A. n. 3. eng. bot. t. 105. — Mich. gen. t. 33. f. 1. (v. s.)
4. C. MACROSTYLA (Lapeyr. abr. 562.) foliis setaceis convoluto-canaliculatis, spicâ lineari androgynâ supernè masculâ, fructibus oblongis teretiusculis glabris utrinque attenuatis erectis squamâ duplò longioribus, stylo longè exserto.  $\mathcal{U}$  in Pyrenæis editioribus. — Fl. fr. 5. p. 287. (v. s.)

### §. 2. Spicâ unicâ simplici; stigmatibus 3.

5. C. PYRÆNICA (Wahl. act. holm. 1803. p. 139.) foliis linearibus canaliculatis, spicâ ovatâ androgynâ supernè masculâ, fructibus confertis oblongis acuminatis primùm erectis demum patenti-reflexis squamâ caducâ vix longioribus.  $\mathcal{U}$  in Pyrenæis. — Fl. fr. 5. p. 287. C. Fontanesiana et C. Ramondiana Fl. fr. 3. p. 102. C. acutissima Degl. in Lois. gall. 628. C. Marchandiana Lapeyr. (v. v.)
6. C. RUPESTRIS (All. ped. t. 92. f. 1.) culmo acutangulo, foliis rigidis linearibus striatis planiusculis, spicâ androgynâ supernè masculâ cylindricâ, fructibus laxiusculis obovatis squamis obtusissimis brevioribus.  $\mathcal{U}$  in rupestribus editorum Alpium Delphinatûs (DC.). — Fl. fr. 5. p. 288. C. petræa Gaud. agr. 2. p. 78. (v. v. DC.)
7. C. PAUCIFLORA (Light. scot. 2. t. 6. f. 2.) foliis setaceis carinato-plicatis, spicâ subovatâ androgynâ, floribus 5-8 uno alterove superiori masculo, fructibus laxis utrinque subulatis divericatis patulis squamas femineas caducas

vix superantibus.  $\frac{1}{2}$  in pratis paludosis Vogesorum, Jurassi et Alpium. — Fl. fr. 3. p. 102. Schk. trad. t. A. n. 4. C. leucoglochis L. f. supp. 413. (v. v.)

§. 3. *Spicis pluribus androgynis supernè masculis; stigmatibus 2.*

8. C. ARENARIA (L. sp. 1381.) foliis linearibus planiusculis striatis, culmo incurvo triquetro, spiculis plurimis ovatis alternis sessilibus approximatis, superioribus masculis, inferioribus femineis, intermediis androgynis, fructibus confertis ovato-oblongis marginatis acuminatis bifidis margine scabris squamas ovatas acuminatas æquantibus.  $\frac{1}{2}$  in arenosis maritimis Oceani et Corsicæ. — Fl. fr. 5. p. 288 et 3. p. 103. excl. syn. Bell. eng. bot. t. 928. Schk. trad. t. B et Dd. n. 6. (v. v.)
9. C. DISTICHA (Huds. ang. 403.) foliis linearibus planis, culmo erecto acutè triquetro angulis scabro, spiculis plurimis ovatis alternis sessilibus in spicam ovato-cylindricam imbricato-congestis, inferioribus remotiusculis, superioribus et inferioribus femineis, intermediis androgynis, fructibus ovato-oblongis acuminatis marginatis bifidis ciliato-serratis squamas ovato-oblongas acuminatas æquantibus.  $\frac{1}{2}$  in paludosis. — Fl. fr. 3. p. 104. C. intermedia Good. trans. lin. 2. p. 154. Schk. trad. t. B. n. 7. Gand. agr. 2. p. 80. C. arenaria Leers. herb. t. 14. f. 2. non L. (v. v.)
10. C. VULPINA (L. sp. 1382.) foliis lato-linearibus planis acuminatis, culmo erecto acutissimè triquetro, bracteis serrulatis, spiculis ovatis numerosis sessilibus in spicam oblongam interruptam subapproximatis, fructibus ovatis acuminatis bidentatis compresso-triquetris divergentibus margine scabris squamas ovatas acuminatas subsuperantibus.  $\frac{1}{2}$  in paludosis. — Fl. fr. 3. p. 105. eng. bot. t. 307. Schk. trad. t. 6. n. 10. — Mich. t. 33. f. 13. 14. (v. v.)
11. C. DIVISA (Huds. ang. 405.) radice repente, foliis linearibus carinato-plicatis scabris, culmo acutè trigono, bractea foliaceâ erectâ strictâ, spiculis 3-6 sessilibus oblongo-ovatis in spicam ovatam approximatis, fructibus ovatis subulatis acuminatis bidentatis adpressis squamas ovatas aristato-acuminatas subæquantibus.  $\frac{1}{2}$  ad vias et in arenosis Gallicæ mediterranæ, occidentalis et boreali-occidentalis. — Fl. fr. 3. p. 105 et 5. p. 289. Good. tr. lin. 2. t. 19. f. 2. eng. bot. t. 1096. C. schænoides Host. 1. t. 45. C. splendens Pers. C. Bertolonii Schk. app. (v. v.)
12. C. DIVULSA (Good. tr. lin. 2. p. 160.) foliis linearibus planis, culmo erecto acutè triquetro striato scabriusculo, spiculis ovatis superioribus approximatis, inferioribus distantibus, fructibus erectiusculis ovatis plano-convexis marginatis scabris acutis bidentatis glumas ovatas acutas superantibus.  $\frac{1}{2}$  in nemorosis humidis. — Fl. fr. 3. p. 105. Schk. car. trad. t. Dd. n. 89. eng. bot. t. 629. — Mich. t. 33. f. 10. An satis à sequenti distincta? (v. v.)
13. C. MURICATA (L. sp. 1382.) foliis linearibus planis, culmo erecto acutè triquetro striato scabro, spiculis ovatis in spicam cylindricam interruptam subapproximatis, fructibus scabris divergentibus ovatis plano-convexis bidentatis basi marginatis glumas ovatas acuminatas superantibus.  $\frac{1}{2}$  in nemorosis humidis. — Fl. fr. 3. p. 106. Schk. trad. t. E. n. 22. eng. bot. t. 1097. C. virens Lam. dict. 3. p. 384. Fl. fr. 3. p. 106. C. nemorosa Host. (v. v.)
14. C. FETIDA (All. ped. n. 2297.) foliis linearibus acuminatis planis ant carinato-plicatis, spiculis numerosis in capitulum ovato-globosum densè aggregatis, fructibus ellipticis teretiusculis acuminatis bifidis, bracteis squamisque ovatis mucronatis.  $\frac{1}{2}$  in humidis summorum Alpium et Pyrenæorum. — Fl. fr. 3. p. 106. Schk. car. trad. t. Hh. n. 96. — Scheuchz. prod. t. 4. it. alp. f. 18. (v. v.)
15. C. CHORODORIZA (Ehr. phyt. n. 77.) radice repente, culmo basi subramoso, foliis linearibus planiusculis acuminatis culmeis erectis, spiculis adpressis in capitulum ovatum densè congestis, fructibus ovatis acuminatis, bracteis lanceolato-ovatis acutis.  $\frac{1}{2}$  in turfosis Jurassi. — Fl. fr. 3. p. 107. Schk. trad. t. G et Ji. n. 31. (v. s.)

16. *C. JUNCTIFOLIA* (All. ped. t. 92. f. 4. non Schk.) radice repente, culmo teretiusculo, foliis setaceis canaliculatis, spiculis in capitulum subrotundum dense aggregatis, fructibus ovatis semi-globosis lævibus rostratis ore subintegerrimis squamas ovatas obtusas aut submucronatas superantibus. ♀ in editioribus Pyrenæis ex Lapeyr. — Fl. fr. 3. p. 107. eng. bot. t. 927. *C. incurva* Lightf. scot. 2. t. 24. f. 1. Schk. car. trad. t. Hh. f. 95. (v. s. ex monte Cenisio.)
17. *C. LOBATA* (Schk. car. trad. t. D. f. 18. non Vill.) foliis linearibus planiusculis canaliculatisve convolutis, culmo trigono erecto, spiculis 3 oblongis alternis in capitulum parvum trilobum dense congestis, fructibus ellipticis acuminatis nervosis. ♀ in Alpibus Galloprov. — *C. tripartita* Fl. fr. 3. p. 188. non All. nec Sut. (v. v.)
18. *C. PARADOXA* (Willd. mem. p. 32. t. 1. f. 1.) foliis linearibus carinato-plicatis, culmo erecto trigono-compresso striato, spiculis sessilibus parvis oblongis in spicas oblongas angustas congestis, spicis in paniculam angustam elongatam digestis, inferioribus remotiusculis, fructibus subrotundo-ovatis rostellatis bidentatis margine supernè ciliato-serratis, squamis ovatis acutis. ♀ in limosis secùs rivulos rariùs. — Fl. fr. 3. p. 108. Schk. car. trad. t. E. f. 21. *C. canescens* Host. gram. t. 57. (v. s. ex Austria.)
19. *C. TERETIUSCULA* (Good. tr. lin. 2. t. 19. f. 3.) radice repente, foliis linearibus carinato-plicatis, culmo erecto obtusè trigono subtereti, spiculis ovatis simplicibus rarè compositis in spicam tereti-oblongam approximatis, fructibus trigonis plano-convexis apice attenuatis subbidentatis squamas ovatas acutas superantibus. ♀ in turfosis Jurassi, circà Tolosam secùs canalè Occitanicum ex Lapeyr. — Fl. fr. 5. p. 289. Schk. car. t. D. f. 19. (v. v.)
20. *C. PANICULATA* (L. sp. 1383.) foliis linearibus duris planis aut carinato-plicatis, culmo erecto triquetro scaberrimo, spiculis oblongo-ovatis 3-gonis adpressis in spicas oblongas congestis, spicis in paniculam plus minùsve coarctatam digestis, fructibus ovatis supernè marginatis acuminatis bidentatis margine ciliato-serratis squamas ovatas acutas æquantibus. ♀ in paludibus. — Fl. fr. 3. p. 108. Schk. trad. t. D. f. 20. eng. bot. t. 1064. f. 1. — Mich. gen. t. 33. f. 7. — Scheuchz. prod. t. 8. f. 2. (v. v.)

§. 4. *Spicis pluribus androgynis apice femineis; stigmatibus 2.*

21. *C. CYPEROIDES* (L. supp. 413.) foliis linearibus, culmo triquetro striato, spiculis ovato-oblongis aggregatis globoso-capitatis involucreatis, fructibus lanceolatis bicuspidatis, squamis lanceolatis aristatis ciliato-scabris. ♀ in humidis arenosis propè *Sézanne en Brie* ex cl. Desfont. et Mér. — Fl. fr. 3. p. 109. Schk. car. trad. t. A. f. 5. *Bohemica* Schreb. gram. t. 28. f. 3. *Schelhameria capitata* Mœnch supp. 119. — Mich. gen. t. 33. f. 19. (v. s. ex Bohemià.)
22. *C. OVALIS* (Good. tr. lin. 2. p. 148.) foliis linearibus planis, culmo erecto obsoletè trigono fistuloso striato, spiculis subsenis sessilibus ovalibus alternis in spicam ovoideam subapproximatis, fructibus ovatis acuminatis marginatis bidentatis ciliato-serratis glumas ovato-lanceolatas subacutas subsequantibus. ♀ in pratis ndis. — Fl. fr. 3. p. 110. Schk. trad. t. B. f. 8. eng. bot. t. 306. *C. leporina* Huds. Leers. herb. t. 14. f. 6. an L? — Scheuchz. gram. t. 10. f. 15. (v. v.)
23. *C. APPROXIMATA* (Hoppe exs. Hoffm. germ. 4. p. 201.) foliis linearibus planis, culmo triquetro incurvo, spiculis tribus sessilibus subrotundo-ellipticis alternis approximatis, fructibus ellipticis compressis acuminatis ore integris margine glabris glumas ovatas obtusiusculas æquantibus. ♀ in editioribus Alpibus Delphinatùs, Pyrenæorum ex Lapeyr. — Fl. fr. 6. p. 290. *C. leporina* L. lapp. 322. Schk. car. t. F. ff. f. 129. DC. syn. n. 1718. (v. s. ex monte Cenisio.)
24. *C. SCHREBERI* (Willd. mém. p. 22.) radice repente, foliis angustissimis planiusculis, culmo trigono angulis scabriusculo, spiculis 5-7 sessilibus ovato-cylindricis alternis in spicam ovatam confertis, fructibus ovatis bidentatis

- striatis acutis squamas ovato-lanceolatas acutissimas æquantibus.  $\frac{1}{2}$  in arenosis graminosis siccis. — Fl. fr. 3. p. 110. Schk. car. trad. t. B. f. g. C. tenella Thuil. — Seg. ver. t. t. r. f. 2. (v. v.)
25. *C. BRIZOIDES* (L. sp. 1381.) radice fibrosâ, foliis linearibus planis, culmo nudo triquetro gracillimo scabro, spiculis 5-7 sessilibus alternis subdistichis curvatis oblongo-lanceolatis approximato-contignis, fructibus elongato-acuminatis divergentibus marginatis bifidis squamis ovato-oblongis obtusiusculis longioribus.  $\frac{1}{2}$  secus sylvas et sepes demissiorum Pyrenæorum, agri Syrtici et Jurassi circâ Salins (Babey). — Fl. fr. 3. p. 111. Schk. car. t. C. U. n. 12. (v. s.)
26. *C. CURTA* (Good. tr. lin. 2. p. 145.) foliis linearibus planis carinatis plicatisve, culmo triquetro striato scabriusculo, spiculis subsenis sessilibus alternis remotiusculis demùm ovato-cylindraceis, superioribus subapproximatis, fructibus ovatis plano-convexis acutiusculis lævibus ore integris erectis squamas ovatas acutas superantibus.  $\frac{1}{2}$  in graminosis subhumidis passim. — Fl. fr. 3. p. 111. Schk. car. trad. t. C. f. 13. — C. tenella Hoffm. germ. 1. p. 318. — Lœs. pruss. t. 32. (v. v.)
27. *C. STELLULATA* (Good. tr. lin. 2. p. 144.) foliis linearibus canaliculatis, culmo erecto obsolete trigono læviusculo, spiculis sessilibus ovato-globosis subquaternis remotiusculis, fructibus ovato-acuminatis bidentatis margine scabris stellatim divergentibus squamas ovatas acutas superantibus.  $\frac{1}{2}$  in pratis humidis passim. — Fl. fr. 3. p. 112. eng. bot. t. 806. C. stellata Schk. trad. t. C. f. 14. — Schenchz. gram. t. 11. f. 3. (v. v.)
28. *C. REMOTA* (L. sp. 1383.) foliis linearibus planiusculis, culmis gracilibus obsolete trigonis læviusculis, spiculis sessilibus ovato-oblongis alternis remotis solitariis, bracteis foliaceis longissimis, fructibus ovatis acuminatis integris bidentatisve compressiusculis margine scabris squamas lanceolatas acutas superantibus.  $\frac{1}{2}$  in umbrosis subhumidis. — Fl. fr. 3. p. 112. eng. bot. t. 832. Schk. trad. t. E. f. 23. — Mich. gen. t. 33. f. 15. 16. (v. v.)
29. *C. ELONGATA* (L. sp. 1383.) foliis linearibus planis carinato-plicatisve, culmo triquetro asperimo, spiculis sessilibus subnodosis alternis cylindricis remotiusculis, fructibus ovato-acuminatis emarginatis integrisve nervosis patulis margine subscabris squamas ovatas acutas obtusave superantibus.  $\frac{1}{2}$  in nemorosis humidis. — Fl. fr. 3. p. 112. Schk. car. trad. t. 5. f. 25. — Schenchz. gram. t. 11. f. 4. (v. v.)

§. 5. *Spiculis pluribus androgynis, stigmatibus 3.*

30. *C. CURVOLA* (All. ped. t. 92. f. 3.) foliis subsetaceis canaliculatis duris curvulis, spiculis adpressis sessilibus in capitulum oblongum densissimè aggregatis, fructibus ovatis compressis acuminatis, squamis ovatis mucronatis.  $\frac{1}{2}$  in summis Alpibus et Pyrenæis. — Fl. fr. 3. p. 109. Schk. car. trad. t. D. Hb. f. 17. (v. v. DC.)
31. *C. GYNOMANE* (Bert. dec. 2. p. 43.) foliis setaceis planiusculis debilibus, bracteis foliaceis infimâ scapum superante, spiculis 2-3 sessilibus linearibus gracilibus distantibus, fructibus 3-5 laxiusculis erectis triquetris ovato-acuminatis squamis lanceolatis acuminatis brevioribus.  $\frac{1}{2}$  secus vias et in montosis agris Ruscinonensis, Monspelii, Telonis et Corsicæ. — Fl. fr. 5. p. 289. C. tuberosa Degl. in Lois. gall. 2. p. 629. C. Linckii Schk? (v. v.)
32. *C. ATRATA* (L. sp. 1386.) foliis lato-linearibus acuminatis planis striatis, culmo acutè triquetro, spiculis tribus quiniseve ovatis approximatis, fructiferis nutantibus, fructibus subrotundo-ovatis breviter rostratis bidentatis squamis ovatis acutis brevioribus.  $\frac{1}{2}$  in Alpium et Pyrenæorum pascuis. — Gaud. agrost. 2. p. 114. — Schenchz. gram. t. 11. f. 1 et 2.
- a. pedunculata*, spicis androgynis sublonge pedunculatis demùm pendulis, culmo elatiori. — C. atrata Fl. fr. 3. p. 113. Schk. trad. t. X. f. 77. (v. v.)

- β. *dubia* (Gaud.) spicâ supremâ masculâ (paucioribus intermixtis femineis floribus) reliquis merè femineis, fructibus atro-purpureis. (v. v.)  
 γ. *nigra* (Gaud.) spicis confertis subsessilibus erectis, terminali masculâ solitariâ, inferioribus femineis, culmo humiliori. — *C. nigra* All. ped. 2310. Fl. fr. 3. p. 113. (v. v.)

§. 6. *Spicis pluribus sexu distinctis; stigmatibus 2.*

33. *C. mucronata* (All. ped. n. 2318.) foliis setaceis canaliculatis, culmo flexuoso vix trigono læviusculo, spicâ masculâ solitariâ sessili, femineâ subsolitariâ approximâtâ elliptico-subrotundâ sessili, fructibus ovato-oblongis plano-convexis incurvato-rostratis bidentatis margine ciliatis glumas ovato-oblongas subacutas subæquantibus. ½ in pascuis summis Alpinum Delphin. ex Vill., Jurassi ex Hall. — Fl. fr. 3. p. 114. Schk. trad. t. K. f. 44. (v. v. D. in Italiâ.)  
 34. *C. cespitosa* (L. sp. 1388.) foliis linearibus planis glaucescentibus patulis, culmo erecto acutè trigono læviusculo, spicâ masculâ subsolitariâ, spicis femineis 3-4 sessilibus cylindraceis attenuatis distantibus, fructibus densè imbricatis glabris compressis ellipticis obtusis ore pertusis squamas oblongas obtusas aut brevissimè mucronatas superantibus. ½ in limosis. — Fl. fr. 3. p. 114. Good. tr. lin. 2. t. 21. f. 8. Schk. trad. t. Aa Bb. n. 85. (v. v.)  
 35. *C. trinervis* (Degl. in Lois. gall. 731.) radice repente, foliis rigidis carinato-triquetris canaliculatis culmum erectum incurvum 3-gonum lævem superantibus, spicis masculis 1-2, femineis 3-4 ovato-cylindricis sessilibus subdistantibus, mediis interdum apice masculis, fructibus densissimè imbricatis compressis ovatis extus trinerviis glabris nitidis pellucido-punctatis ore integris squamas oblongas subacutas superantibus. ½ in arenis uliginosis maritimis Vasconiz, Picardiz. — Fl. fr. 5. p. 291. (v. v. DC.)  
 36. *C. stricta* (Good. tr. lin. 2. t. 21. f. 9.) foliis linearibus glaucis planis aut carinato-plicatis strictis reticulo basi connexis, culmo erecto acutè triquetro supernè scabriusculo, spicâ masculâ subsolitariâ elongatâ, spicis femineis 2-3 cylindraceis sessilibus attenuatis distantibus, fructibus densè imbricatis glabris glaucis ovatis compressis acutiusculis ore pertusis squamas lanceolatas acutiusculas superantibus. ½ in paludibus. — Fl. fr. 3. p. 114. eng. bot. t. 914. Schk. trad. t. V. f. 73. (v. v.)  
 37. *C. acuta* (L. sp. 1388.) foliis linearibus planis margine scabris, culmo erecto acutè trigono, spicis masculis binis terniave elongatis, femineis subquaternis subpedunculatis subnutantibus cylindraceis filiformi-elongatis remotis, fructibus supernè densè imbricatis infernè laxioribus oblongo-ovatis glabris nitidis nervosis brevissimè rostellatis ore integro pertuso squamas oblongas acutas subæquantibus. ½ in pratis paludosis passim. — eng. bot. t. 580. Schk. car. trad. t. Ee Ff. f. 92. *C. gracilis* Curt. Fl. fr. 3. p. 115. Spicæ femineæ superiores sæpè androgynæ. (v. a.)

§. 7. *Spicis pluribus sexu distinctis; stigmatibus 3; fructu pubescente aut villosa.*

\* *Spicâ masculâ solitariâ.*

38. *C. tomentosa* (L. mant. 123.) foliis linearibus angustis planis aut margine revolutis, culmo erecto trigono, spicis femineis subgeminatis ovato-cylindraceis subapproximatis sessilibus, fructibus subrotundis tomentosis squamam ovatam acuminatam subæquantibus. ½ in pratis umbrosis et sylvis. — Fl. fr. 3. p. 116. Schk. trad. t. F. f. 28. (v. v.)  
 39. *C. montana* (L. succ. 845.) foliis linearibus planis, culmo obsolete trigono læviusculo, spicis femineis subgeminis approximatis ellipticis sessilibus, fructibus ovato-oblongis brevissimè rostratis pubescentibus squamas ovato-rotundas obtusas superantibus. ½ in nemorosis montanis passim. — Fl. fr. 3. p. 116. Schk. trad. t. F. f. 29. *C. conglobata* All. non Willd. *C. globularis*



- Willd. act. her. 1794. p. 45. t. 2. f. 1. — Scheuchz. gram. t. 10. f. 8 et 9. (v. v.)
40. *C. FILULIFERA* (L. sp. 1385.) foliis linearibus planiusculis, culmo triquetro læviusculo, spicis femineis subternis approximatis elliptico-subrotundis sessilibus, fructibus subrotundis breviter rostratis pubescentibus squamâ oblongâ acutâ brevioribus.  $\mathcal{L}$  in nemorosis. — Fl. fr. 3. p. 117. eng. bot. t. 885. Schk. trad. t. 1. f. 39. (v. s.)
41. *C. ERICETORUM* (Poll. pal. n. 886.) foliis duris linearibus planis, culmo incurvo lævi obtusè trigono, spicis femineis subgeminis approximatis oblongo-sessilibus, fructibus subrotundo-obovatis pubescentibus brevissimè mucronatis squamas obovatas obtusissimas margine scarioras superantibus.  $\mathcal{L}$  in collibus siccis et ericetis Palatinatus, Alpium, Pyren. orientalium (DC.) rariis. — Fl. fr. 3. p. 117. *C. ciliata* Schk. trad. t. J. f. 42. Gaud. agr. 2. p. 133. — Scheuchz. gram. t. 10. f. 10. (v. s.)
42. *C. PRÆCOX* (Jacq. aust. t. 446.) foliis duris linearibus planis, culmo erecto læviusculo obsolete trigono, spicis femineis subternis oblongo-ovatis subsessilibus, fructibus subgloboso-triangularibus pubescentibus breviter obtusè mucronatis squamam ovatam acutam æquantibus.  $\mathcal{L}$  in siccis sterilibus montosis. — Fl. fr. 3. p. 115. Schk. trad. t. F. f. 27. *C. filiformis* Leers. herb. t. 16. f. 5. (v. v.)
43. *C. UMBROSA* (Host. gram. t. 69.) caespitosa, foliis linearibus planis erectis scapum subsuperantibus, culmo erecto lævi compresso-trigono, spicis femineis subgeminis oblongis subsessilibus remotiusculis, fructibus ovato-triangularibus pubescentibus rostratis ore bidentatis squamas ovatas mucronatas dorso ciliato-asperas æquantibus.  $\mathcal{L}$  Circâ Burdigalam (cl. Des Monl.). — Fl. fr. 3. p. 723. *C. præcox*  $\beta$ . Gaud. agr. 2. t. 131. (v. s.)
44. *C. CYNOBASIS* (Vill. dauph. 2. p. 206. excl. syn.) foliis linearibus firmis carinatis margine scabris subrevolutis, culmo gracili trigono, spicis femineis 3-5-floris rotundis, superioribus breviter pedunculatis approximatis, unâ radicali longissimè pedunculatâ, fructibus obovatis vix pubescentibus nervosis breviter mucronatis mucrone apice truncato squamas ovato-lanceolatas acutiusculas subæquantibus.  $\mathcal{L}$  in collibus et humilioribus montibus Gallie meridionalis et Juraasi. — Fl. fr. 3. p. 118. *C. diversiflora* Host. gram. t. 70. *C. Halleriana* Ass. arr. t. 9. f. 2. (v. v.)
45. *C. HUMILIS* (Leyss. hall. n. 952.) foliis setaceo-canaliculatis scabriusculis culmum longè superantibus, spicis femineis binis subtrifloris remotis subsessilibus subinclusis bracteis margine membranaceis obliquis vaginatis, fructibus obovatis obtusis breviter pubescentibus squamam scarioras obovatam mucronatam subæquantibus.  $\mathcal{L}$  in collibus siccis et rupibus. — Fl. fr. 3. p. 117. *C. clandestina* Good. Schk. trad. t. k. f. 43. Gaud. agr. 2. p. 140. — Scheuchz. gram. t. 10. f. 1. (v. v.)
46. *C. DIGITATA* (L. sp. 1384.) foliis lato-linearibus planis acutis striatis, culmo erecto teretiusculo læviusculo, spicis femineis subternis linearibus subdigitatis subapproximatis, inferioribus pedicellatis, bracteis vaginatis pedicellis dimidiò brevioribus, fructibus obovatis breviter rostratis triquetris pubescentibus squamam obovatam brevissimè mucronatam æquantibus.  $\mathcal{L}$  in nemorosis umbrosis. — Fl. fr. 5. p. 291. et 3. p. 119. — Mich. t. 32. f. 9. (v. v.)
- $\beta$ . *ornithopoda*, spicis femineis digitatis subsessilibus, fructibus squamas superantibus. — *C. ornithopoda* Willd. *C. pedata* Fl. fr. 3. p. 119. non L. — Mich. t. 32. f. 14. (v. v.)
- \* \* *Spicis masculis duabus aut plurimis.*
47. *C. FILIFORMIS* (L. sp. 1385.) foliis prælongis erectis linearibus planis canaliculatis margine scabris, culmo erecto teretiusculo basi lævi, spicula masculâ subsolitariâ longissimâ, femineis geminis ovato-oblongis sessilibus distantibus, fructibus confertis ellipticis bifurcatis laevibus squamam oblongo-

lanceolatam mucronatam superantibus aut subæquantibus.  $\mathcal{L}$  in paludibus circa Parisios et in Alpibus. — Fl. fr. 3. p. 119. eng. bot. t. 904. Good. tr. lin. 2. t. 20. f. 5. C. lasiocarpa Ehr. Gaud. agr. 2. p. 127. (v. s.)

48. C. OLAUCA (Scop. caru. n. 1157.) radice repente, foliis linearibus planis glaucescentibus margine scabris, culmo erecto obtusè trigono, spicis masculis geminis, femineis 2-3 distantibus cylindricis pedunculatis pendulis rariùs sessilibus, fructibus confertis ovatis obovatisve ciliato-scabriusculis brevissimè rostratis rostro retuso squamam ovatam obtusam enaspidatamve subæquantibus.  $\mathcal{L}$  in humidis et paludosis vulgatissima. — Fl. fr. 3. p. 120. C. flacca Schk. trad. t. O. P. f. 57. a. b. C. recurva Huds. — Mich. t. 32. f. 5. Planta polymorpha. (v. v.)
49. C. HIRTA (L. sp. 1389.) foliis linearibus carinato-plicatis subhirsutis, culmo obtusè trigono læviusculo, spicis masculis 2-3 hirsutis inequalibus, femineis 3 remotis cylindricis breviter pedunculatis, fructibus oblongis acuminatis hirsutis bicuspidatis squamā oblongā aristatā majoribus.  $\mathcal{L}$  in sabulosis. — Fl. fr. 3. p. 121. Leers. herb. t. 16. f. 3. Schkr. trad. t. Un. f. 108. — Moris. s. 8. t. 12. f. 6. n. 10. (v. v.)

§. 8. *Spicis pluribus sexū distinctis, stigmatibus tribus, fructibus glabris aut ad angulos tantum ciliato-scabris.*

\* *Spicā masculā solitariā.*

50. C. FLAVA (L. sp. 1384.) foliis lato-linearibus planis aut carinato-plicatis, culmo erecto teretiusculo lævi, spicis femineis subternis ovato-ellipticis inclusè pedunculatis, superioribus subapproximatis, fructibus ovatis reflexis longè rostratis rostro curvato bidentato squamā ovato-lanceolatā acuminatā longioribus.  $\mathcal{L}$  in pratis humidis. — Fl. fr. 3. p. 121. Schk. trad. t. H. f. 36. (v. v.)
- ℓ. *Oederi* (Gand.) spicis femineis subquaternis, omnibus approximatis. — C. Oederi Ehr. Schk. trad. t. F. f. 26. non Retz. (v. v.)
51. C. NITIDA (Host. gram. 1. t. 71.) foliis linearibus planis striatis, culmo erecto triquetro scabro, spicis femineis 2-3 ovatis subdistantibus, imā exsertè pedunculatā, fructibus subrotundo-ovatis nitidis brevissimè rostratis ore obliquo squamam ovatam obtusam æquantibus.  $\mathcal{L}$  in siccioribus montanis. — Fl. fr. 5. p. 294. C. alpestris Fl. fr. 3. p. 122. excl. syn. Hall. All. Sut. C. verna Schk. trad. t. L. f. 46. (v. v.)
52. C. EXTENSA (Good. tr. lin. 2. t. 21. f. 7.) foliis linearibus carinato-plicatis striatis subconvolutis rigidis, culmo erecto stricto subtrigono lævi, spicis femineis subternis elliptico-cylindricis subsessilibus, superioribus subapproximatis, fructibus ovatis nervosis rostro brevi recto bidentato squamas obovatas mucronatas superantibus.  $\mathcal{L}$  in udis maritimis Oceani et Mediterranei. — Fl. fr. 5. p. 292. Spica mascula aliquando gemina. (v. v.)
53. C. BASTARDIANA (DC. fl. fr. 5. p. 293.) foliis lato-linearibus planis striatis, culmo erecto acutè trigono, spicis femineis binis ovatis parvis approximatis, infimæ bracteæ foliaceæ spicam masculam æquante, fructibus ovato-triquetris margine hispidis squamā lanceolatā sublongioribus.  $\mathcal{L}$  in paludosis Andegavizæ septentrionalis (cl. Bastard). (v. s. specimen imperf.) †.
54. C. FIRMA (Host. aust. 509 et gram. 1. t. 75.) foliis brevibus firmis patulis linearibus planis striatis, culmo lævi teretiusculo, spicis femineis subtribus ovato-cylindricis distantibus, infimā exsertè pedunculatā, fructibus oblongis acutè trigonis margine hispidis rostratis ore obliquo squamā lanceolatā mucronatā longioribus.  $\mathcal{L}$  in editioribus Pyrenæis (DC.). — Fl. fr. 3. p. 122. Schk. trad. t. O. Y. f. 54. (v. v. DC.)
55. C. SEMPERVIRENS (Vill. dauph. 2. p. 214.) foliis linearibus firmis planis striatis margine scabriusculis, culmo obsoletè trigono lævi, spicis femineis subternis oblongis erectis distantibus, binis inferioribus pedunculatis, fructibus oblongis compresso-triquetris angulis ciliatis longè rostratis rostro apice

- scarioso bilobo, squamâ ovatâ longioribus, spicâ masculâ acutâ.  $\mathcal{L}$  in Jurasso, Alpihus et Pyrenæis. — *C. ferruginea* Schk. trad. t. M. f. 48. Fl. fr. 3. p. 123. excl. syn. Vill. *C. erecta* Fl. fr. 3. p. 120. *C. varia* Host. gram. t. 80. Fructus facies supernè pilis nonnullis adpressis conspersæ. (v. v.)
56. *C. SCOROLIANA* (Willd. sp. 4. p. 292.) foliis longissimis linearibus carinatis planisve scabriusculis, culmo erecto obsolete trigono lævisculo apice nutante, spicis femineis tribus distantibus subteretibus pedicellatis subnutantibus, fructibus ovatis obsolete trigonis angulis ciliatis mucronatis ore colorato bilobo squamâ ovali-subacutâ longioribus.  $\mathcal{L}$  in Jurassi monte Dole et Alpihus umbrosia. — Gaud. agr. 2. p. 168. *C. spadicea* Fl. fr. 3. p. 120. non Schk. *C. ferruginea* Host. gram. t. t. 81. non Willd. *C. Mielichhoferi* Willd. Schk. — Schenck. gram. t. 10. f. 6. (v. v.)
57. *C. BRACHYSTACHYS* (Schranck bay. 294.) foliis linearibus angustissimis convolato-setaceis canaliculatisve, culmo erecto filiformi lævi obsolete trigono, spicis femineis 2-4 oblongo-cylindricis distantibus pedunculatis, fructibus lanceolatis glabris rostratis bidentatis squamâ oblongâ mucronatâ longioribus.  $\mathcal{L}$  in Alpihus, montibus Ligeris (cl. Prost). Pyrenæis orientalibus ex Lapeyr. — Fl. fr. 3. p. 123. Schk. trad. t. P. f. 58. — Schenck. gram. t. 10. f. 7. (v. v.)
58. *C. STRIGOSA* (Good. tr. lin. 2. t. 20. f. 4.) foliis linearibus planis, culmo triquetro lævi, spicis femineis 3-4 pendulis filiformibus distantibus, inferioribus exserte pedunculatis, fructibus oblongo-lanceolatis utrinque attenuatis nervosis glabris ore oblique truncatis squamam ovato-lanceolatam acutam subsequantibus.  $\mathcal{L}$  in sylvaticis udibus Alsatiz secus Rhenum. — Fl. fr. 5. p. 295. Schk. trad. t. N. f. 53. non All. *C. leptostachys* L. f. supp. 414. (v. s.)
59. *C. FULVA* (Good. trans. lin. 2. p. 177.) foliis linearibus planis striatis scabris, culmo trigono erecto scabriusculo, spicis femineis subternis ovato-oblongis distantibus, supremâ sessili, reliquis pedunculatis, fructibus densè imbricatis glaberrimis subglobosis nitidis longè rostratis bidentatis squamâ ovatâ longioribus.  $\mathcal{L}$  in pratis humidis rariis. — Fl. fr. 3. p. 126. Schk. trad. t. T. f. 67. eng. bot. 1295. *C. trigona* All. ped. t. 89. f. 4. (v. v.)
60. *C. PUNCTATA* (Gaud. agr. 2. p. 152.) foliis linearibus planis lævibus, culmo trigono erecto lævi, spicis femineis 2-3 oblongo-cylindricis distantibus, inferioribus breviter exserto-pedunculatis erectis, fructibus ovato-globosis nitidis lævissimis pellucido-punctatis binerviis breviter rostratis rostro lævisimo bidentatis squamâ ovatâ mucronatâ longioribus.  $\mathcal{L}$  in agro Syrtico circâ S.-Sever perierit cl. Dufour, circâ la Teste de Buch cl. Des Moulins, propè Nicæam D. Stire. (v. s.)
61. *C. BILIGULARIS* (DC. cat. monsp. 88.) foliis lato-linearibus planis striatis, ligulis 2 oblongis aliâ liberâ oppositifoliâ, alterâ disco folii adnatâ, culmo trigono lævi, spicis femineis tribus oblongo-cylindricis distantibus, superioribus inclusè inferioribus exserte pedunculatis, infimâ nutante, fructibus ovatis acuminatis nervosis glabris rostratis rostro bicuspidato margine subciliato-scabro squamâ ovatâ acuminatâ vix longioribus.  $\mathcal{L}$  in nemoribus humidis circâ Falaise ex cl. de Bréb., Cænomanum, Andegavum. — Fl. fr. 5. p. 296. (v. s.)
62. *C. MINORIS* (Smith. brit. 993.) foliis linearibus planis, culmo erecto acutè triquetro lævi, spicis femineis tribus ovato-cylindricis valdè distantibus, supremâ inclusè reliquis exserte pedunculatis, stigmatibus tribus, fructibus ovato-ellipticis glabris nervosis nervis 2 proeminentibus rostratis rostro bicuspidato margine ciliato-scabro squamâ obovatâ obtusâ mucronatâ longioribus.  $\mathcal{L}$  in pratis humidis Parisiorum et provinc. occidentaliū passim. — Fl. fr. 5. p. 296. An satis à sequente distincta (v. s.)
63. *C. DISTANS* (L. sp. 1387.) foliis linearibus planis, culmo erecto acutè triquetro lævi, spicis femineis tribus erectis oblongo-ovatis valdè distantibus, supremâ sessili, reliquis breviter inclusè pedunculatis, fructibus ovatis angulosis glabris rostratis rostro bicuspidato margine ciliato-scabro squamam

- ovatam acuminatam mucronatamve superantibus.  $\frac{1}{2}$  in sepiibus et pratis. — Fl. fr. 3. p. 126. Schk. trad. t. Yy. f. 78. eng. bot. 1234. — Moris. s. 8. f. 10. n. 18. (v. v.)
64. *C. BREVICOLLIS* (DC. fl. fr. 5. p. 295.) foliis linearibus planis margine scabris, culmo erecto acutè triquetro lævi, spicâ masculâ erectâ, femineis 1-3 cylindricis 6-10-floris pedicellatis bracteis vaginatis, fructibus obovatis ventricosis glabris sublævibus rostro brevissimo truncato aut 2-dentato squamis oblongis acuminatis subbrevioribus.  $\frac{1}{2}$  ad rupes Rhodano proximas in monte Parve propè Belley. (cl. V. Auger). (v. s.)
65. *C. MICHELII* (Schk. tab. P et Vv. f. 59.) foliis firmis linearibus planis striatis, culmo triquetro læviusculo, spicâ masculâ clavatâ, femineis subbinis 3-5-floris remotis, infimâ exsertè pedunculatâ, fructibus grossis laxis sublævibus glabris globosis ventricosis longè rostratis rostro bicuspidato apice ciliato-scabro squamam scariosam oblongam acuminatam subsuperantibus.  $\frac{1}{2}$  in pratis montosis circâ Parisios ex cl. Merat. — Fl. fr. 5. p. 295. Host. gram. 1. t. 73. — Mich. gen. t. 32. f. 5. (v. s. ex Austriâ.)
66. *C. DEPAUPERATA* (Good. tr. linn. 2. p. 181.) foliis linearibus planis, culmo erecto 3-gono lævi, spicâ masculâ gracili, spicis femineis 2-3, 2-5-floris remotis, hinis inferioribus exsertè pedunculatis, fructibus grossis laxis ovato-globosis nervosis inflatis glabris longè rostratis ore obliquo squamâ oblongâ scariosâ acuminatâ longioribus.  $\frac{1}{2}$  in sylvis circâ Parisios, Andegavum et in Pictaviâ superiore. — Fl. fr. 5. p. 294. *C. triflora* Willd. phyt. 2. t. 1. f. 2. (v. s.)
67. *C. FRIGIDA* (All. ped. n. 2331.) foliis linearibus planis lævibus margine scabris, culmo erecto trigono læviusculo, spicâ masculâ obtusâ, spicis femineis ternis ovato-cylindricis distantibus, 2 inferioribus pedunculatis nantantibus, fructibus (nigris) lanceolatis triquetris angulis ciliatis acuminatis bicuspidatis squamâ oblongâ longioribus.  $\frac{1}{2}$  in humidis summorum Alpium et Pyrenæorum. — Fl. fr. 3. p. 124. (v. v.)
68. *C. LIMOSA* (L. sp. 1386.) radice repente, foliis linearibus angustis carinatis, culmo erecto acutè triquetro læviusculo, spicis femineis binis ovatis pedunculatis pendulis, bracteis amplexicaulibus, fructibus confertis ellipticis compresso-triquetris nitidos striatos brevissimè rostellatos ore integro squamis subrotundo-ovatis cuspidatis subæquantibus superantibusve.  $\frac{1}{2}$  in paludosis turfosis rariis. — Fl. fr. 3. p. 127. Schk. trad. t. X. f. 78. — Scheuchz. gram. t. 10. f. 13. (v. v.)
69. *C. ALBA* (Scop. carn. 1148.) foliis setaceo-canaliculatis scabriusculis, culmo erecto obsolete trigono scabriusculo, bracteis membranaceis vaginatis hyalinis, spicis femineis 1-2 pedunculatis sub-5-floris distantibus scariosis masculam subæquantibus, fructibus laxis obovatis glabris sulcatis breviter rostratis oblique truncatis squamas membranaceas obtusas superantibus.  $\frac{1}{2}$  in montosis sylvaticis Alpium, Jurassi. — Fl. fr. 3. p. 124. Schk. trad. t. O. f. 55. — Scheuchz. gram. t. 10. f. 3. 4. 5. — Pluck. t. 9x. f. 2. (v. v.)
70. *C. PANICKA* (L. sp. 1367.) foliis firmis planis glaucis scabris, culmo obtusè trigono læviusculo, spicis femineis subbinis cylindricis distantibus, supremâ subincluse infimâ breviter exsertè pedunculatâ, fructibus laxis ovatis glabris lævibus vix mucronatis mucrone retuso squamam ovatam acutam superantibus.  $\frac{1}{2}$  in pratis udis. — Fl. fr. 3. p. 127. Schk. trad. t. Ll. f. 100. — Mich. gen. t. 34. f. 11. Variat spicâ infimâ radicali longissimè pedunculatâ. (v. v.)
71. *C. PALLESCENS* (L. sp. 1386.) foliis linearibus planis subpubescentibus, culmo erecto acutè triquetro scabriusculo, spicis femineis subternis ovato-oblongis subdistantibus, fructiferis sæpè cernuis, inferioribus pedunculatis, fructibus confertis obovato-oblongis subventricosis obtusis glabris nitidis squamam ovato-oblongam cuspidatam subæquantibus.  $\frac{1}{2}$  in pratis humidis. — Fl. fr. 3. p. 127. Schk. trad. t. Kk. f. 99. — Mich. gen. t. 33. f. 13. (v. v.)
72. *C. PILOSA* (All. ped. n. 2323.) foliis lato-linearibus striatis margine pilosis

- scabrisque, culmo erecto obtusè trigono striato læviusculo, spicis femineis 2-3 gracilibus distantibus binis inferioribus remotis, fructibus laxis glabris ovatis rostratis, ore membranaceo obliquo emarginato squamâ ovatâ acuminatâ longioribus.  $\frac{1}{2}$  in sylvis circâ Parisios ex Mérat, Arbois (Babey), in Pyrenæis ex Lapeyr. — Fl. fr. 3. p. 123. (v. s.)
73. *C. FATULA* (Scop. carn. n. 1160. t. 59.) foliis linearibus planis margine scabris, culmo erecto triquetro lævi, spicis femineis quaternis remotis filiformibus pedunculatis cernuis, fructibus laxiusculis oblongo-ovatis rostratis bifidis squamâ ovato-lanceolatam membranaceam acuminatam subsequantibus.  $\frac{1}{2}$  in sylvis vulgè. — Fl. fr. 3. p. 128. *C. sylvatica* Huds. Schk. trad. t. Ll. f. 101. eng. bot. t. 995. *C. drymeia* L. f. supp. 1414. — Moris. s. 8. t. 12. f. 13. n. 9. (v. v.)
74. *C. CAPILLARIS* (L. sp. 1386.) foliis linearibus angustis planis margine læviusculis, culmo erecto subtrigono lævissimo, spicis femineis subternis longè pedunculatis cernuis subsexfloris superioribus approximatis masculam superantibus, fructibus laxis ovatis glabris lævibus rostratis ore obliquo, squamâ membranaceâ ovatâ obtusâ longioribus.  $\frac{1}{2}$  in pratis udis summaram Alpium ex Vill. et Pyrenæorum ex Ram. — Fl. fr. 3. p. 125. Scop. carn. n. 1152. t. 59. Schk. trad. t. O. f. 56. — Seg. véron. 3. t. 3. f. 1. (v. s. ex Alpiibus Helveticis.)
75. *C. PSEUDO-CYPERUS* (L. sp. 1387.) foliis lato-linearibus planis margine scabris, culmo erecto trigono scabro, spicis femineis quaternis geminatis pedunculatis pendulis cylindricis, fructibus dense imbricatis ovato-lanceolatis nervosis divergentibus bicuspidatis squamam lanceolatam aristatam subsequantibus.  $\frac{1}{2}$  in udiset fossis passim. — Fl. fr. 3. p. 128. Schk. trad. t. Mm. f. 102. eng. bot. t. 242. — Lob. ic. t. 76. f. 2. (v. v.)
76. *C. LÆVIGATA* (Sm. brit. 3. p. 1005.) foliis glaucescentibus firmis longissimis lato-linearibus carinato-caniculatis infernè subrevolutis, culmo erecto acutè triquetro lævi, spicâ masculâ subsolitariâ, femineis 2-4 distantibus elongato-cylindricis erectis, inferioribus breviter pedunculatis, fructibus ovato-tri-quetris glabris rostro brevissimo 2-dentato squamam ovatam mucronatam subsequantibus.  $\frac{1}{2}$  in Corsicæ paludosis. — Schk. car. t. Ses. f. 162. *C. patula* Schk. car. n. 97. t. Bbb. f. 116. *C. æthiopica* Schk. car. n. 91. t. Z. f. 83. Variat spiculis masculis binis. (v. s.)
77. *C. MAXIMA* (Scop. carn. 2. p. 1166.) foliis lineari-lanceolatis planis margine scabris, culmo erecto triquetro lævi, spicis femineis subquinis elongato-cylindricis curvato-pendulis sessilibus, infimâ exsertâ pedunculatâ, fructibus minimis dense imbricatis lævibus ellipticis brevissimè rostratis ore bidentatis squamam ovatam submucronatam æquantibus.  $\frac{1}{2}$  in sylvaticis udis. — Fl. fr. 3. p. 125. *C. pendula* Huds. Schk. trad. t. 2. f. 60. *C. agastachys* L. f. supp. 1414. — Barr. ic. t. 45. (v. v.)

\*\* *Spicis masculis duabus aut pluribus.*

78. *C. HORDEISTICHOS* (Vill. dauph. 2. t. 6.) foliis linearibus planis culmum erectum teretiusculum superantibus, spicis masculis 2-3 oblongo-ovatis, femineis tribus oblongo-ovatis remotiusculis, infimâ subradicali, fructibus imbricatis ovatis compressis acuminatis bidentatis angulis ciliato-serratis squamam ovatam scariosam superantibus.  $\frac{1}{2}$  in paludosis Delphinatûs et agri Parisiensis. — Fl. fr. 3. p. 129. *C. hordeiformis* Thuill. Host. gram. 1. t. 76. (v. s.)
79. *C. VESICARIA* (L. sp. 1388. a.) foliis longissimis linearibus planis margine scabris, culmo erecto acutè triquetro, spicis masculis 2-3 gracilibus, femineis 2-3 pedunculatis erectis cylindraceis, fructibus imbricatis nervosis oblongo-conicis inflatis rostratis bicuspidatis squamâ lanceolatâ majoribus.  $\frac{1}{2}$  in paludosis sylvaticis. — Fl. fr. 3. p. 129. Schk. trad. t. Sa. f. 106. eng. bot. t. 779. (v. v.)
80. *C. AMPULLACKA* (Good. tr. lin. 2. p. 207.) foliis longissimis linearibus an-

gustis subcanaliculatis, culmo erecto obtusè triquetro lævi, spicis masculis 2-3 gracillimis, femineis binis cylindraceis breviter pedunculatis erectis, fructibus imbricatis subglobosis inflatis rostratis, rostro filiformi bifurco, squamam lanceolatam acutam superantibus.  $\mathcal{N}$  in paludibus montosis. — Fl. fr. 3. p. 130. Schk. trad. t. Ft. f. 107. eng. bot. 780. (v. v.)

81. *C. PALUDOSA* (Good. tr. lin. 2. p. 202.) foliis longissimis glaucescentibus planis lato-linearibus, culmo stricto acutè triquetro, spicis masculis 3-4 inæqualibus, femineis 3-4 erectis cylindraceis attenuatis inferioribus pedunculatis, fructibus glabris densissimè imbricatis ovatis nervosis breviter mucronatis emarginatis squamâ lanceolatâ acuminatâ latioribus sublongioribusque.  $\mathcal{N}$  in paludibus. — Fl. fr. 3. p. 130. Schk. trad. t. Oo Vv. f. 103. eng. bot. t. 807. *C. acuta* Cart. lond. 4. t. 61. (v. v.)

82. *C. KOCHIANA* (DC. cat. monsp. 89.) foliis longissimis glaucescentibus planis lato-linearibus scaberrimis, culmo stricto acutè triquetro angulis scabro, spicis masculis 2 inæqualibus, femineis 3-5 erectis cylindraceis, inferioribus pedunculatis, fructibus glabris densissimè imbricatis nervosis oblongo-acuminatis apice bidentatis squamam lanceolatam acuminato-aristatam subæquantibus.  $\mathcal{N}$  ad fossas et rivulos Gallie boreali-orientalis et propè Mimatem (cl. Frost.). — Fl. fr. 5. p. 297. (v. s.)

83. *C. RIPARIA* (Cart. lond. 4. t. 60.) foliis longissimis planis lato-linearibus asperissimis, culmo stricto acutè triquetro scaberrimo, spicis masculis 4-5 subæqualibus, femineis 3-4 erectis cylindraceis attenuatis inferioribus pedunculatis, fructibus glabris imbricatis oblongo-ovatis rostratis multinerviis bifurcatis, latitudine et longitudine squamæ lanceolatæ aristatæ.  $\mathcal{N}$  ad fluviorum ripas. — Fl. fr. 3. p. 130. Schk. trad. t. 29. R. f. 105. eng. bot. t. 579. — Mich. t. 32. f. 6. 7. (v. v.)

84. *C. SOLEIROLIS* (Duby mss.) foliis glaucescentibus firmis longis planis lato-linearibus lævibus margine scabris, culmo stricto acutè triquetro lævissimo, spicis masculis 3-4, femineis 3-4 cylindricis oblongis sessilibus erectis, fructibus ovato-rotundis compressis margine superne ciliatis brevissimè rostellatis ore bidentato, squamâ lineari-lanceolatâ obtusiusculâ brevioribus.  $\mathcal{N}$  in Corsicâ circa Bonifacio reperiit cl. Soleirol. — Affinis *C. longearistatæ* Riv. Bern. sic. manip. IV. p. 8. t. 2. sed squamæ non aristatæ. (v. s.)

Ordo CX. GRAMINEÆ *Juss. gen.* 28. — Gramina *L.*

Flores glumacei, hermaphroditi aut rariùs diclini. Gluma exterior sæpiùs bivalvis, uniflora aut flores plurimos distiche spicatos includens. Singulo flori perigonium (glumella) glumæ analogum, plerumque bivalve. Stamina sæpiùs 3 hypogyna. Ovarium 1 liberum; stigmata 2. Fructus siccus monospermus semen nudum simulans. Embryo parvus monocotyledoneus, ad basin albuminis farinacei. — Culmi cylindrici, nodis hinc indè distincti, herbacei sæpiùs simplices; folia alterna vaginantia, vaginis longitudinaliter fissis, plerumque ad basin internam membranâ (ligulâ) instructa; flores paniculati aut spicati.

† FLORES SEXU DISTINCTI NEMPÈ MASCULI ET FEMINEI IN DIVERSIS SPICULIS.

I. MAYS *Tourn. inst.* 353. *Gærtn. carp.* 1. p. 6.

Spiculæ masculæ bifloræ paniculatæ terminales; femineæ unifloræ spicatæ laterales; stigmata longissima; semina subrotunda lævia seriatim disposita.

1. *M. Zea* (*Gærtn. fruct.* 1. p. 6. t. 1.). ① volitur passim. — *Fl. fr.* 3. p. 98. *Zea* *Mays*. *L. sp.* 1133. — *Moris. s.* 8. t. 13. f. 1. 2. — *Fuchs. hist.* 473. ic. (v. v. c.)

II. HOLCUS *Schreb. gram.* 129. t. 18. *DC. fl. fr.* 3. p. 97. — *Holci sp. L.*

Spiculæ 1-2-floræ, aliæ masculæ membranaceæ muticæ, aliæ hermaphroditæ coriaceæ aristatæ. — Flores paniculati.

1. *H. Aleppensis* (*L. sp.* 1485.) paniculâ laxâ, spiculis subglabris, aristis è receptaculo ortis. ② ex Oriente ortus, in Galliâ meridionali nunc indigenus. — *Fl. fr.* 3. p. 98. *Host. gram.* 1. t. 1. *Sorghum Aleppense* *Pers.* — *Pluck. t.* 32. f. 1. (v. v. c.)
2. *H. Sorghum* (*L. sp.* 1484.) paniculâ sublaxâ ovatâ, spiculis villosis, aristis ex apice glumarum ortis. ① in Galliâ meridionali passim cultus. — *Fl. fr.* 5. p. 286. *Sorghum vulgare* *Pers. ench.* 1. p. 101. *Host. gram.* t. 4. t. 2. — *Fuchs. hist.* 771. ic. (v. v. c.)

III. ANDROPOGON *L. gen.* 1145.

Spiculæ unifloræ, aliæ masculæ pedicellatæ muticæ, alteræ hermaphroditæ sessiles ex apice perigonii aristatæ; gluma basi extus pilosa. — Flores spicato-digitati aut subpaniculati.

1. *A. GAVILLUS* (*L. sp.* 1480.) paniculâ laxâ, spiculis ternis pilis flavescentibus basi circumdati, intermediâ hermaphroditâ biaristatâ. ② in siccis provinciis australium, Arvernæ ex *Delarb.* — *Fl. fr.* 3. p. 95. *Aploda Gryllus* *P. de Beauv.* *Pollinia Gryllus* *Spreng.* — *Lam. ill.* t. 840. f. 1. — *Scheuchz. gram.* t. 6. f. 1. (v. v.)
2. *A. ISCHÆMUM* (*L. sp.* 1483.) foliis pilosis, spicis 6-10 erectis subdigitatis, pedicellis villosis, spiculis geminis (purpurascens), alterâ hermaphroditâ uniaristatâ, glumis obtusiusculis glabriusculis basi et apice pilorum fasci-

## 502 MONOCOTYL. seu ENDOGENÆ.

X. CALAMAGROSTIS Roth. germ. 1. p. 33. DC. fl. fr. 3. p. 24. — Arundinis sp. Sm. Schrad. Gaud.

Gluma uniflora bivalvis; perigonium bivalve, extus basi aut dorso villis longis sericeis onustum. — Flores paniculati.

1. C. ERIGENOS (Roth. germ. 2. p. 1. p. 91.) foliis scabris, paniculâ angustâ strictâ subpatente, spiculis laxè imbricatis, glumis acuminatis ciliato-scabris perigonio subduplò longioribus, aristâ dorsali rectâ pilis glumam subæquantibus breviori.  $\frac{1}{2}$  in siccis humidique. — Fl. fr. 5. p. 257. Arundo epigeios L. sp. 120. Schrad. germ. 1. t. 4. f. 1. — Schenckz. prod. 21. t. 5. (v. v.)
2. C. LANCEOLATA (Roth. germ. 1. p. 34.) culmo subramoso, foliis angustis leviusculis, paniculâ laxâ diffusâ, spiculis subsparsis, glumis acuminatis perigonio subduplò longioribus, perigonii basi pilosi valvulâ exteriori apice aristatâ, aristâ rectâ exiguâ, pilis glumellâ longioribus glumâve subbrevioribus.  $\frac{1}{2}$  in sylvis et pratis humidis rarius. — Fl. fr. 5. p. 259. 3. p. 26. var.  $\alpha$ . Arundo Calamagrostis L. sp. 121? Schrad. germ. 1. t. 4. f. 4. (v. s. ex Ardenis.)
3. C. LITTORÆA (DC. fl. fr. 5. p. 255.) foliis linearibus margine scabris, paniculâ laxâ diffusâ, glumis acuminatis perigonio longioribus, aristâ terminali brevi pilisque glumam subæquantibus.  $\frac{1}{2}$  in arenosis humidis secus flumina passim. — Arundo littoræa Schrad. germ. 1. t. 4. f. 2. A. pseudo-phragmites Gaud. non Schrad. (v. v.)
4. C. MONTANA (DC. fl. fr. 5. p. 254.) foliis planis lato-linearibus margine scabris, paniculâ erectâ laxâ diffusâ, glumis acutis perigonium subæquantibus, aristâ subbasilari geniculatâ glumam excedente, glumellâ basi pilosâ, pilorum penicillo ad valvulâ interioris basin erecto glumellâ subæquali.  $\frac{1}{2}$  in montosis. — C. arundinacea Fl. fr. 3. p. 25. C. acutiflora Fl. fr. 5. p. 255. Arundo montana Gaud. A varia et acutiflora Schrad. germ. p. 216. t. 4. f. 6. et p. 217. (v. v.)
5. C. SYLVATICA (DC. fl. fr. 5. p. 253.) foliis angustis scabrinusculis, paniculâ erectâ plus minusve contractâ, spiculis laxè imbricatis, glumis acutis perigonium æquantibus, aristâ dorsali geniculatâ glumam excedente, pilorum penicillo ad valvulâ interioris basin erecto glumellâ subduplò breviori.  $\frac{1}{2}$  in sylvatilis montosis Jurassi, agri Mimatensis (cl. Prost), Pyrenæorum. — Arundo sylvatica Schrad. germ. 1. t. 4. f. 7. Agrostis arundinacea L. sp. 91? Gaud. agr. helv. 1. p. 73. (v. s.)
6. C. ARGENTEA (DC. fl. fr. 3. p. 25.) culmo sæpius ramoso, foliis linearibus levibus, paniculâ laxâ diffusâ, spiculis sparsis, glumis acuminatis perigonio longioribus perigonii extus lanati valvulâ exteriori apice aristatâ, aristâ geniculatâ glumâ duplò triplòve longiore, pilis glumam subæquantibus.  $\frac{1}{2}$  in siccis et calidis Alpium inferiorum, Monapeli, Pyrenæorum. — Fl. fr. 5. p. 253. Agrostis calamagrostis L. Gaud. agr. Stipa Calamagrostis Wahlb. Gaud. helv. 1. p. 172. Arundo speciosa Schrad. germ. t. 5. f. 8. (v. v.)
7. C. ARENARIA (Roth. germ. 1. p. 34.) radice repente, foliis involutis mucronato-pungentibus, paniculâ spiciformi elongatâ, glumis nitidis acutis perigonium muticum basi villosum æquantibus, pilis glumellâ triplò brevioribus.  $\frac{1}{2}$  in arenosis maritimis. — Fl. fr. 3. p. 24. Arundo arenaria L. Ammophila arundinacea Host. gram. 4. t. 4r. Psamma arenaria P. de Beauv. — Schenckz. gram. t. 3. f. 1. (v. v.)

### XI. AGROSTIS L. gen. n. 80. — Agrostidis sp. Lam. DC.

Gluma uniflora bivalvis; perigonium 2-valve aut (abortu?) univalve, caducum, glabrum aut puberulum, valvulâ alterâ interdum dorso aut basi aristatâ; semen liberum. — Flores paniculati.



Sect. I. *VILFA* Adans. *Pal. de Beauv.* 16. t. 5. f. 8. — Decandollia  
Bast. *ess.* p. 28.

Perigonium 2-valve valvulis moticis.

1. *A. PUNGENS* (Schreb. *gram.* 2. t. 27. f. 3.) culmo ramoso repente, foliis involutis rigidis distichis, ligulis pubescentibus, paniculâ contractâ, pedunculis ramosis, glumæ valvulis inæqualibus lævibus perigonio subbrevioribus.  $\frac{1}{2}$  in arenosis maritimis à Nicmâ ad Hispaniam. — Fl. fr. 3. p. 23. Cav. ic. t. III. (v. v.)
2. *A. MARITIMA* (Lam. *dict.* 1. p. 61.) foliis involutis rigidis, ligulis glabris, paniculâ contractâ, glumæ valvulis æqualibus carinâ hispidiusculis perigonium paulò superantibus.  $\frac{1}{2}$  in arenosis maritimis Narbonæ, Vasconie. — Fl. fr. 3. p. 23. et 5. p. 253. (v. v.)
3. *A. STOLONIFERA* (L. sp. 93. non DC.) culmo ramoso repente, foliis planis margine scabris, ligulâ brevi truncatâ fimbriato-lacerâ, paniculâ coarctatâ, spiculis congestis, glumis æqualibus obtusiusculis breviter ciliatis pubescentibusque perigonium superantibus.  $\frac{1}{2}$  in humidis arenosis provine australium et Burdigalæ (Des Moul.). — Eng. bot. t. 153a. *A. verticillata* Vill. (v. v.)
4. *A. DECUMBENS* (Gaud. *agr.* 1. p. 78.) culmo repente ramoso, foliis planis atriatis scabris, ligulâ oblongâ truncatâ, paniculâ (dessiccatione) contractiori, glumis acuminatis ciliato-scabris æqualibus perigonium paulò superantibus, perigonii valvulâ internâ duplò minore emarginatâ.  $\frac{1}{2}$  in agris humidis et ad fossas. — *A. stolonifera* Fl. fr. 3. p. 23 non L. *A. alba decumbens* Gaud. *helv.* 1. p. 187. Nimium a subsequentibus affinis. (v. v.)
5. *A. ALBA* (L. sp. 93.) culmo basi repente, foliis margine scabris, ligulâ oblongâ truncatâ, paniculâ laxâ, glumæ valvulis acutis æqualibus carinâ subhispidis perigonium superantibus, perigonii valvulâ interiore subduplò minore retusâ.  $\frac{1}{2}$  in humidis. — Fl. fr. 3. p. 22. An reverâ à sequente distincta? (v. v.)
6. *A. VULGARIS* (Hoffm. *germ.* 3. p. 36.) culmo erectiusculo, foliis planis angustis, ligulâ brevi truncatâ bifidâ, paniculâ laxâ subtrichotomâ, glumæ valvulis acuminatis dorso hispidiusculis æqualibus, perigonii valvulâ interiore duplò brevior retusâ.  $\frac{1}{2}$  in pratis, arvis et sylvis. — Gaud. *agr.* 1. p. 82. Schenckz. *gram.* t. 3. f. 3. B.
  - a. elongata*, culmo elongato ascendente, paniculâ capillari erecto-patente. — *A. vulgaris* Fl. fr. 3. p. 22. et 5. p. 252. *eng. bot.* t. 1671. *A. capillaris* Vill. Fl. dan. t. 163 non L. (v. v.)
  - β. pumila* (Gaud.) culmis fasciculatis erectis humilibus. — *A. pumila* L. *mant.* 31. Fl. fr. 3. p. 22. (v. v.)
  - γ. dubia*, perigonio interdum aristâ dorsali brevi instructo. — *A. dubia* Lætz herb. t. 4. f. 4. Fl. fr. 3. p. 21. *A. compressa* Willd. (v. v.)

Sect. II. *TRICHOIDIUM* Mich. *fl. bor. amer.* 1. p. 41. Schrad. *germ.* 1. p. 198.

Perigonium 1-valve.

\* *Perigonii valvulâ naticâ.*

2. *A. ELEGANS* (Thore in Lois. *not.* t. 1. f. 1.) foliis involuto-subulatis, culmo erecto simplici, paniculâ laxâ capillari, pedicellis lævibus tenuissimis, glumis lævibus acutis, perigonio obtuso 1-valvi. ① in ericetis circa Aquas-Tarbellicas. — Fl. fr. 5. p. 252. *Trichodium elegans* Thore in Rœm. et Sch. 2. p. 283. (v. s.)

\*\* *Perigonii valvulâ aristatâ.*

8. *A. RUBRA* (L. sp. 92. excl. syn. Pluck.) culmo erecto, foliis planis angustissimis margine scabriusculis, ligulâ oblongâ obtusâ, paniculâ per anthesin patente antè et post eam contractâ, pedunculis scabris, glumæ valvulis acuminatis, externâ dorso ciliato-scabrâ perigonium superante, perigonii valvulâ

## 504 MONOCOTYL. SEU ENDOGENÆ. AGROSTIN.

- apice retusâ bimucronulatâ, aristâ dorsali geniculatâ spiculam paulô superante. (1) in pratis et ad vias. — Fl. fr. 3. p. 19. A. vinealis Bast. ess. — Schenchn. gram. t. 3. f. 11. C. Non diversam à sequente nec veram A. rubram Linnæi volent cl. Wahlberg Willd. etc. (v. s.)
9. A. CANINA (L. sp. 92.) culmo prostrato, foliis inferioribus fasciculatis setaceis, superioribus planis, ligulâ obtusâ fimbriato-lacerâ, paniculâ per anthesin patente, antè et post eam contractâ, glumæ valvulis acuminatis inæqualibus externâ dorso hispido-scabrâ, perigonii valvulâ breviter 2-aristatâ, aristâ dorsali spiculam superante. 2/ in pratis subhumidis. — Fl. fr. 3. p. 20. Hoffm. germ. 3. t. 6. eng. bot. t. 1856. Trichodium caninum Schrad. (v. v.)
10. A. GLAUCINA (Bast. suppl. 25.) foliis angustissimis planis glaucis, paniculâ contractâ, pedunculis scabris, glumæ valvulis acuminatis subæqualibus perigonium paulô superantibus, glumellâ apice 2-mucronulatâ, aristâ basilari geniculatâ spiculam paulô superante. 2/ in ericetis Andegaviæ. — Fl. fr. 5. p. 251. Trichodium glaucinum Rœm. et Sch. 2. p. 281. (v. s.)
11. A. SETACEA (Curt. lond. 6. t. 12.) foliis setaceis, ligulâ oblongâ acuminatâ, culmo erecto stricto, paniculâ contractâ, pedicellis scabris, glumæ valvulis acuminatis inæqualibus perigonium paulô superantibus, glumellâ apice breviter 2-aristatâ, aristâ basilari geniculatâ spiculam paulô superante. 2/ in ericetis Vasconiz, Armoraciz. — Fl. fr. 5. p. 251. eng. bot. t. 1188. Trichodium setaceum Rœm. et Sch. 2. p. 280. (v. v. D.)
12. A. ALPINA (Leyss. hal. n. 67.) culmo erecto, foliis setaceis, paniculâ primo coarctatâ demum patulâ, pedunculis lævibus, glumæ valvulis acuminatis subæqualibus perigonium superantibus, glumellâ apice 2-mucronulatâ dorso aristatâ, aristâ spiculâ longioribus. (1) in pratis Alpium, montium Coricæ, Arverniz, Pyrenæorum. — Fl. fr. 5. p. 20. Host. gram. 3. t. 49. Trichodium alpinum Schrad. germ. t. 3. f. 4. (v. v.)
13. A. NUPASTRA (All. ped. n. 2161.) culmo erecto, foliis setaceis, paniculâ patulâ, pedunculis scabris, glumæ valvulis acuminatis subinæqualibus perigonium superantibus, glumellâ apice breviter 2-aristatâ basi externè aristatâ, aristâ spiculâ duplò longioribus. (1) in pratis siccioribus Alpium, montium Mimatis (Prost), Pyrenæorum. — Fl. fr. 3. p. 21. A. filiformis Vill. Fl. fr. 3. p. 20. Trichodium rapestre Schrad. germ. t. 3. f. 5. Schenchn. prod. t. 4. f. 1. (v. v.)
14. A. PALLIDA (DC. fl. fr. 5. p. 251. non Schk. Hoffm.) culmo erecto geniculato, foliis planis angustissimis, ligulâ oblongâ 2-dentatâ, paniculâ ramossissimâ, glumæ valvulis subæqualibus acuminatis, externâ dorso subciliato-scabrâ, perigonio duplò longioribus, glumellâ obtusâ brevissimè 2-mucronulatâ, aristâ dorsali rectâ glumâ duplò longiore. (1) in Corsicâ et Galloprovinciâ circâ Forum-Julii. (v. s.)
15. A. INTERRUPTA (L. sp. 92.) culmo erecto basi geniculato, foliis planis angustis scabris, paniculâ elongatâ coarctatâ strictâ subinterruptâ, glumæ valvulis acuminatis inæqualibus dorso ciliato-scabris perigonium vix superantibus, glumellâ subciliato-scabrâ infrâ apicem aristatâ, aristis rectis longissimis. (1) in agris arenosis passim. — Fl. fr. 3. p. 19. Host. gram. 3. t. 48. — Vaill. par. t. 17. f. 4. (v. v.)

### Sect. III. ANEMOCHLOA DC. herb.

Perigonium 2-valve, valvulâ externâ aristatâ.

16. A. SPICA-VENTI (L. sp. 91.) foliis planis scabris, paniculâ elongatâ patulâ multiflorâ, glumæ valvulis æqualibus acuminatis glabris perigonium superantibus, perigonii valvulâ externâ acuminatâ infrâ apicem aristatâ, aristis rectis strictis longissimis, internâ paulô breviorè breviter 2-mucronulatâ. (1) inter segetes. — Fl. fr. 3. p. 18. Leers herb. t. 4. f. 1. — Lob. ic. t. 3. f. 1. (v. v.)

## XII. MILIUM Linn. gen. 79. Schrad. germ. 1. p. 194. — Agrostidis sp. DC. fl. fr.

Gluma 2-valvis uniflora ventricosa; perigonium 2-valve glumâ minus persistens glabrum aut puberulum, valvulâ externâ sub apice aristatâ rarius muticâ internam involvente; semen perigonio corticatum. — Fl. paniculati.

## \* Perigonio aristato.

1. M. ? LENDIGERUM (L. sp. 91.) foliis planis scabris linearibus, paniculâ coarctatâ spicæformi ovato-cylindricâ, glumæ valvulis nitidis basi ventricosis aristato-acuminatis margine scabris perigonium longè superantibus, perigonii valvulâ externâ pilosâ infrâ apicem aristatâ aristâ glumam subsuperante. ① in arvis Galliæ australis, occidentalis, mediæ usque ad Parisios passim. — eng. bot. t. 1107. Schreb. gram. t. 23. f. 3. Agrostis lendigera Fl. fr. 3. p. 18. Gastridium australe P. de Beauv. — Pluck. t. 33. f. 6. An reverâ Millii species aut potiùs genus distinctum? (v. v.)
2. M. MULTIFLORUM (Cav. dem. bot. p. 36.) foliis planis margine scabris, ligulis brevioribus truncatis, paniculâ elongatâ laxâ diffusâ multiflorâ, pedicellis semiverticillatis, glumæ valvulis glabris acuminatis subæqualibus, aristâ terminali rectâ strictâ spiculâ vix duplò longiore. ① in arenosis Galliæ mediterraneæ. — Agrostis miliacea L. sp. 91. Fl. fr. 3. p. 19. Host. gram. 3. t. 45. Piptatherum multiflorum P. de Beauv. (v. v.)
3. M. THOMASII (Duh. mss.) culmo stricto basi geniculato glauco, foliis convolutis lævibus linearibus angustis, ligulâ brevi truncatâ, paniculâ elongatâ laxâ subcoarctatâ, pedicellis semiverticillatis, glumæ valvulis glabris acuminatis subæqualibus, aristâ terminali rectâ spiculam paulò superante. 2½ in rupestribus Corsicæ ad promontorium Capo-Corso dictum reperiit Ph. Thomas. — Semiverticillorum inferiorum flosculi plerùmque abortiunt et panicula involucri setiformi cincta videtur. (v. s.)
4. M. PARADOXUM (Schreb. gram. t. 28. f. 2.) foliis planis lato-linearibus scabris, ligulâ brevissimâ, paniculâ laxissimâ, pedicellis geminis, glumæ valvulis glabris subæqualibus acuminatis nitidis, perigonii valvulâ externâ breviter pubescente, aristâ terminali rectâ spiculâ subtriplo longiore. 2½ in agro Nicænsi, in Galloprovincia et Delphin. australiori, circâ Monspelium. — Agrostis paradoxa L. sp. 62. Fl. fr. 3. p. 17. Piptatherum paradoxum P. de Beauv. — Pluck. t. 32. f. 2. (v. v.)
5. M. CÆRULESCENS (Desf. atl. 1. t. 12.) foliis convolutis setaceis angustissimis glaucis, ligulâ oblongâ obtusâ, paniculâ laxissimâ, pedicellis solitariis geminisve ramosis, glumis acuminatis æqualibus glabris nitidis, perigonii valvulâ externâ nitidâ, aristâ terminali caducâ glumam subæquantē. 2½ in lapideis apricis Nicæ, Telonis, Arelatis, agri Ruscinonensis. — Agrostis cærulescens Fl. fr. 5. p. 250. Piptatherum cærulescens P. de Beauv. (v. v. DC.)

## \*\* Perigonio mutico.

6. M. EFFUSUM (L. sp. 90.) foliis lato-linearibus planis lævibus, ligulâ oblongâ fimbriato-lacertâ, paniculâ laxissimâ, pedicellis semiverticillatis patulis, spiculis dispersis lævibus, glumæ valvulis ovatis acutis perigonio glabro paulò longioribus. 2½ in sylvis. — eng. bot. t. 1106. Agrostis effusa Lam. DC. fl. fr. 3. p. 21. — Moris. s. 8. t. 5. f. et n. 10. (v. v.)

## XIII. STIPA L. gen. 90.

Gluma uniflora bivalvis acuta; perigonium 2-valve; glumâ minus, valvulâ exteriore convolutâ apicē aristam longam basi articulatam deciduam gerente. — Panicula subsimplex.

1. S? ARISTELLA (All. auct. p. 39. t. 2. f. 4. an L? non Gouan.) foliis superioribus planis, aristis glabris flosculo vix duplò longioribus. 2½ in locis siccis

## 506 MONOCOTYL. SEU ENDOGENÆ.

- petrosia Galliae mediterraneæ passim. — Fl. fr. 3. p. 28. et 5. p. 258. Agrostis bromoides L. Gouan. ill. t. 1. f. 3. — An vera Stipa? (v. v.)
2. *S. JUNCEA* (L. sp. 116.) foliis convolutis filiformibus intus pubescentibus, paniculâ laxâ pallidâ, aristis longissimis basi subortilibus pubescentibus.  $\frac{1}{2}$  in collibus aridis provinc. australium. — Fl. fr. 3. p. 27. (v. v.)
  3. *S. CAPILLATA* (L. sp. 116.) foliis convolutis subfiliformibus intus villosis, paniculâ laxâ demum rufescente, aristis longissimis glabris basi rectis apice tortilibus.  $\frac{1}{2}$  in collibus aridis passim. — Fl. fr. 3. p. 27. et 5. p. 257. Host gram. 3. t. 5. — Scheuchz. gram. t. 3. f. 13. A. (v. v.)
  4. *S. TORTILIS* (Desf. atl. 1. t. 31. f. 1.) foliis convolutis filiformibus, paniculâ spiciformi folio superiore spathæformi involutâ, aristis longissimis basi tortilibus pubescentibus.  $\frac{1}{2}$  in apricis lapideis Corsicæ, agri Nicœnensis et Ruscinonensis. — Fl. fr. 5. p. 257. Desf. atl. t. 31. f. 2. — Spartium, etc. Bocc. mus. p. 138. t. 97. (v. v.)
  5. *S. PENNATA* (L. sp. 115.) foliis filiformibus, aristis longissimis plumosis basi glabris.  $\frac{1}{2}$  in siccis montanis lapideis passim. — Fl. fr. 3. p. 27. Lam. ill. t. 41. f. 1. — Moris. s. 8. t. 7. f. 21. n. 9. (v. v.)

### XIV. PANICUM Juss. gen. 29. — Panici sp. L.

Gluma uniflora bivalvis, valvulâ tertiâ extus ad basin floris additâ; perigonium bivalve persistens; semen perigonio corticatum. — Panicula laxa aut spiciformis.

1. *P. CAPILLARE* (L. syst. veg. 106.) vaginis hirsutissimis, paniculâ capillari ramosissimâ patente, glumis acuminatis lævibus. ① circa Nicæm (Stiræ). — Fl. fr. 3. p. 15. (v. s.)
2. *P. REPENS* (L. sp. 87.) vaginis glabris, foliis in paginâ superiori et ad oras pubescentibus, paniculâ erectâ coarctatâ laxiflorâ.  $\frac{1}{2}$  ad littora maris prope Olbias (cl. Robert). — Fl. fr. 5. p. 250. Cav. ic. 2. t. 110. (v. s.)
3. *P. MILLIACUM* (L. sp. 36.) foliorum vaginis hirtis, paniculâ laxâ nutante, glumis unneronatis nervosis, seminibus lævissimis. ① ex Indiâ ortum, in Galliâ frequenter cultum. — Fl. fr. 3. p. 15. — Lob. ic. t. 39. f. 1. (v. v. c.)
4. *P. CAUS-GALLI* (L. sp. 83.) paniculâ spiciformi decompositâ, spiculis alternis conjugatisque subdivisis, glumis aristatis hispidis, rachis subquingulari. ① in arvis et cultis. — Fl. fr. 3. p. 14. eng. bot. t. 876. Echinochloa crus-galli Röm. et Sch. 2. p. 478. — Moris. s. 3. t. 4. f. 5. n. 16. (v. v.)
5. *P. ITALICUM* (L. sp. 83.) paniculâ spiciformi densâ basi interruptâ subramosâ, spiculis glomeratis, rachis hirsutâ, involuclis subbisetosis. ① ex Indiâ ortum in Galliâ meridionali cultum. — Fl. fr. 3. p. 14. P. germanicum Willd. Setaria Italica Röm. et Sch. a. p. 493. — Lob. ic. t. 42. f. 1. (v. v. c.)
6. *P. GLAUCUM* (L. sp. 83.) paniculâ spiciformi elongato-cylindricâ, spiculis confertis, involuclis bifloris multisetosis scabriusculis, semine transversim rugoso. ① ad margines agrorum et viarum. — Fl. fr. 3. p. 13. Host. gram. 2. t. 16. Setaria glauca Röm. et Sch. 3. p. 490. — Lob. ic. t. 13. f. 2. (v. v.)
7. *P. VIRIDE* (L. sp. 83) culmis suberectis, paniculâ spiciformi cylindricâ, racemulis laxiusculis, involuclis bifloris multisetosis aurorum scabriusculis, seminibus minutissimè muricatis. ① in arvis. — Fl. fr. 3. p. 13. eng. bot. t. 875. Pennisetum viride R. Brown. Setaria viridis Röm. et Sch. — Moris. s. 8. t. 4. f. 1. n. 10. (v. v.)
8. *P. VERTICILLATUM* (L. sp. 82.) culmis subdiffusis, paniculâ spiciformi cylindricâ, spiculis verticillatis quaternis, involuclis unifloris bisetosis retrorsum scabris. ① in arvis. — Fl. fr. 3. p. 13. Lam. ill. t. 43. eng. bot. t. 874. Pennisetum verticillatum R. Brown. Setaria verticillata Röm. et Sch. — Moris. s. 8. t. 4. f. 2. n. 11. (v. v.)

XV. PHALARIS Willd. sp. 1. p. 326. DC. fl. fr. 3. p. 8. — Phlei et Phalaris sp. L.

Gluma uniflora 2-valvis, valvulis æqualibus carinatis; perigonium bivalve, valvulis inæqualibus concavis acutis glumâ minoribus. — Panicula spiciformis.

§. 1. *Glumis carinatis dorso alato-appendiculatis nec ciliatis.*

1. P. ARUNDINACEA (L. sp. 80.) culmo striato, foliis lato-linearibus striatis margine scabris, paniculâ elongatâ plus minùsve laxâ, pedunculis ramosis; glumæ valvulis trinerviis glabriusculis acutis, perigonio hirsuto.  $\frac{1}{2}$  in pratis humidis. — Host. gram. 2. t. 33. Schrad. germ. t. 6. f. 5. Calamagrostis colorata Sibth. Fl. fr. 3. p. 26. — Scheuchz. gram. t. 3. f. 4. A-C. (v. v.)
2. P. AQUATICA (L. amœn. 4. p. 264.) culmo lævi, paniculâ spiciformi oblongâ, glumâ glabrâ carinatâ acutâ dorso denticulatâ, perigonio adpressæ sericeo villosa.  $\textcircled{1}$  in aquaticis Nicææ, Telonis, Narbonæ. — Fl. fr. 5. p. 249. Host. gram. t. 39. (v. s.)
3. P. CANARIENSIS (L. sp. 79.) culmo lævisculo, paniculâ spiciformi ovatâ vaginâ superiore non involutâ, glumis navicularibus integris lato-ovatis acutis, perigonio inermibus.  $\textcircled{1}$  in maritimis Galloprovinciæ et Occitanæ. agri Ruscinonensis ex Lapeyr., circâ Aginnum ex St.-Am. — Fl. fr. 3. p. 19. eng. bot. t. 1310. — Moris. s. 8. t. 3. f. 15. n. 1. (v. v. c.)
4. P. VULBOSA (L. sp. 79.) culmo basi bulboso lævi, paniculâ laxâ spicatâ ovato-cylindraceâ sæpè basi vaginâ superiore involutâ, glumæ valvulis glabriusculis apice mucronulatis, perigonio glaberrimo.  $\frac{1}{2}$  circâ Caunæ et Antipolium reperiit cl. Balbis, in Corsicâ cl. Requien. — Fl. fr. 5. p. 249. Cav. ic. 1. t. 64. — Barr. ic. t. 700. f. 1. (v. s.)
5. P. PARADOXA (L. dec. t. 18.) paniculâ arctè spicatâ oblongo-cylindricâ è superiore vaginâ dilatâ vix exsertâ, glumæ appendicibus acutis, flosculis in singulo ramo 5-7, intermedii hermaphroditi glumæ valvulis acuminatis, reliquorum imperfectorum valvulis acutis.  $\textcircled{1}$  in Galloprovinciâ, circâ Narbonam, in agro Ruscinonensi ex Lapeyr. — Fl. fr. 3. p. 10. Schreb. gram. t. 12. — Pluck. t. 33. f. 5. (v. s.)
6. P. CYLINDRICA (DC. fl. fr. 3. p. 11.) culmo ascendente lævi, paniculâ arctè spicatâ cylindricâ elongatâ, glumis glabris acutis nervosis integris margine membranaceis.  $\textcircled{1}$  in maritimis Galloprovinciæ, agri Ruscinonensis ex Lapeyr. — Fl. fr. 5. p. 249. P. tennis Host. gram. 2. t. 36. Achnodonton Bellardi P. de Beauv. — Barr. ic. t. 14. f. 1. (v. v. DC.)

§. 2. *Glumis dorso ciliatis nec appendiculatis.*

7. P. TRIGYNA (Host. gram. aust. 4. t. 20.) culmo erecto lævi, vaginis folisque scabriusculis, paniculâ densè spicatâ cylindricâ elongatâ, glumis carinatis acuminatis dorso longè ciliatis.  $\frac{1}{2}$  circâ Nicæam reperiit Stire. — Phleum cuspidatum Willd. supp. enum. p. 5. Chilochloa cuspidata Rœm. et Sch. 2. p. 409. (v. s.)
8. P. ALPINA (Hæneke in Jacq. coll. 2. p. 91.) culmo lævissimo, paniculâ spicatâ basi laxiusculâ cylindricâ elongatâ, glumis carinatis acuminatis dorso longè ciliatis.  $\frac{1}{2}$  in summo Jurasso ex Hall., alpibus Delphin. ex Vill. — Fl. fr. 3. p. 9. Phleum Michellii Allion. Schrad. germ. t. 1. f. 2. eng. bot. t. 2265. Phleum hirsutum Gaud. (v. v. D. in Alpibus Italicis.)
9. P. PHELOMES (L. sp. 80.) culmo simplici, vaginis subventricosis, paniculâ arctè spicatâ elongato-cylindricâ, glumis glabriusculis dorso breviter ciliatis apice subtruncatis mucronatis.  $\frac{1}{2}$  in pratis et sylvarum marginibus. — Fl. fr. 3. p. 9. excl. syn. Moris. Host. gram. 2. t. 34. Phleum phalaroides Gaud. Chilochloa Boehmeri Rœm. et Sch. 2. p. 408. — Barr. ic. t. 21. f. 1. (v. v.)
10. P. ARRHARIA (Kœl. gram. 42.) culmo lævissimo basi ramoso, vaginis ventri-

## 508 MONOCOTYL. seu ENDOGENÆ.

cosis, paniculâ densè spicatâ oblongo-obovatâ, glumis ciliatis acuminatis. ① in arenosis præsertim maritimis. — Fl. fr. 3. p. 8. eng. bot. t. 222. Phleum arenarium L. Chilochloa arenaria Röm. et Sch. l. c. — Pluck. t. 33. f. 8. (v. v.)

XVI. PHLEUM Willd. sp. 1. p. 354. DC. fl. fr. 3. p. 6. — Phlei sp. L.

Gluma uniflora bivalvis, valvulis apice retusis, carinis in aristam prominentibus bicuspidata; perigonium 2-valve, glumâ minus inermis. — Panicula spiciformis.

1. P. PRATENSE (L. sp. 87.) paniculâ densè spicatâ cylindricâ, glumæ valvulis breviter pubescentibus carinâ ciliatis aristâ multò longioribus. ② in pratis ad margines viarum, etc. — Schreb. gram. 1. t. 14. Host. gram. 3. t. 9.  
 $\alpha$ . elongatum, radice minus nodosâ, paniculâ elongatâ. — P. pratense Fl. fr. 3. p. 6. — Moris. s. 8. t. 4. f. 18. n. 1. (v. v.)  
 $\beta$ . nodosum (Gaud.) radice nodosâ, paniculâ brevi. — P. nodosum L. Fl. fr. 3. p. 6. — Moris. s. 8. t. 4. f. 20. n. 3. (v. v.)
2. P. ASPERUM (Vill. dauph. 2. t. 2. f. 4.) culmo erecto basi ramoso, paniculâ spicatâ cylindricâ elongatâ basi subvaginatâ, glumæ valvulis glabris subcartilagineis apice incusatis, aristâ brevissimâ. ① in agris à Moguntia ad Galloprovinciam. — Fl. fr. 3. p. 7. Phleum paniculatum eng. bot. t. 1077. Chilochloa aspera Röm. et Sch. (v. v.)
3. P. ALPIMUM (L. sp. 88.) ligulâ supremâ oblongâ acutâ, paniculâ densè spicatâ ovato-cylindricâ, glumæ valvulis pubescentibus carinâ ciliatis ferè aristæ retrorsum ciliatæ longitudine. ② in pascuis Jurassi, Alpium, Montis-Aurei, Pyrenæorum. — Fl. fr. 3. p. 7. Vill. dauph. 2. t. 2. f. 5. eng. bot. t. 519. (v. v.)
4. P. COMMUTATUM (Gaud. agr. 1. p. 40.) ligulâ supremâ brevissimâ, paniculâ densè spicatâ ovatâ, glumis pubescentibus subasperis carinâ ciliatis in aristas glabras glumæ longitudine desinentibus. ② in summis Pyrenæis frequens, in Alpibus rarum. — Fl. fr. 5. p. 248. (v. v. DC.)

XVII. POLYPOGON Desf. atl. 1. p. 67. — Alopecuri sp. L.

Gluma uniflora bivalvis, valvulâ utrâque aristatâ; perigonium 2-valve glumâ minus, valvulâ alterâ aristatâ.

1. P. LAGASCE (Röm. et Sch. 2. p. 326.) paniculâ coarctatâ elongatâ, glumæ valvulis acutis indivisis glabris dorso ciliatis aristâ terminali vix duplò brevioribus. ① ad oras stagnorum circâ Monspelium rarissimè (Saltzmann). — P. elongatum Lag. non Humb. et Bonpl. (v. s.)
2. P. MONSPELIENSE (Desf. atl. 1. p. 67.) paniculâ subcoarctatâ, glumæ valvulis pubescentibus margine ciliatis apice emarginatis aristâ triplò brevioribus. ① in maritimis incultis et humidis. — Fl. fr. 3. p. 5. Alopecurus Monspelien-sis et panicus L. sp. 90. — Barr. ic. t. 115. f. 1. et 2. (v. v.)
3. P. MARITIMUM (Willd. nov. act. nat. cur. t. 3.) paniculâ coarctatâ oblongâ, glumæ valvulis apice bifidis valdè ciliatis aristâ è fundo fissaræ ortâ triplò brevioribus. ① in maritimis Corsicæ, Occitanie, Oceani rariùs. — Fl. fr. 5. p. 248. An satis à priore distinctum? (v. v.)  
 $\beta$ . subspathaceum, paniculis basi vaginâ folii supremi dilatâ inclusis. — In insulâ Laveiro propè Bonifacio. — P. subspathaceum Req. in ann. sc. nat. 5. p. 386. (v. s.)

XVIII. ALOPECURUS Desf. atl. 1. p. 63. DC. fl. fr. 3. p. 4. — Alopecuri sp. L.

Gluma uniflora bivalvis inermis; perigonium glumam subæquans, univalve, valvulâ basi dorso aristatâ. — Panicula spiciformis.

1. *A. PRATENSIS* (L. sp. 88.) culmo lævi, paniculâ spicatâ crassâ cylindricâ, glumæ valvulis villosis acutis basi connatis perigonii aristâ subduplò brevioribus.  $\frac{1}{2}$  in pratibus. — Fl. fr. 3. p. 4. Lam. ill. t. 42. — Lob. ic. t. 8. f. 2. (v. v.)
2. *A. AGRESTIS* (L. sp. 89.) culmo erecto scabriusculo, paniculâ densè spicatâ elongatâ cylindricâ, glumæ valvulis glabris acutis basi connatis.  $\frac{1}{2}$  in pratibus, agris, vineis. — Fl. fr. 3. p. 4. Schreb. gram. t. 19. f. 2. — Lob. ic. t. 9. f. 2. (v. v.)
3. *A. UTRICULATUS* (Pers. ench. 1. p. 80.) culmo erecto lævi, foliis supremi vaginâ ventricosâ, paniculâ densè spicatâ ovatâ, glumæ valvulis glabris acutis basi marginato-dilatatis perigonii aristâ multò brevioribus. ① in pratibus humidis Burgundicæ, Lugduni, Occitanicæ, agri Ruscinonensis ex Lapeyr. — *Phalaris utriculata* L. Fl. fr. 3. p. 10. Host. gram. 3. t. 7. — Moris. s. 8. t. 4. f. 11. n. 19. (v. v.)
4. *A. BULBOSUS* (L. sp. 1665.) culmo erecto lævi basi bulboso, paniculâ densè spicatâ cylindricâ, glumis villosis obtusis.  $\frac{1}{2}$  in pratibus paludosis aut subsalsis. — Fl. fr. 3. p. 5. eng. bot. t. 1249. — Barr. ic. t. 699. f. 1. et 630. f. 1. et 2. (v. v.)
5. *A. GERICULATUS* (L. sp. 89.) culmo geniculato lævi, paniculâ densè spicatâ cylindricâ elongatâ, glumæ valvulis liberis obtusis carinâ pubescentibus perigonii aristâ paulò brevioribus.  $\frac{1}{2}$  in paludosis et fossis, interdum natans. — Fl. fr. 3. p. 4. eng. bot. t. 1250. — Lob. ic. t. 13. f. 1. (v. v.)
6. *A. GERARDI* (Will. dauph. 2. t. 2. f. 6.) foliorum vaginis ventricosis, paniculâ densè spicatâ subrotundâ, glumis lanceolatis villosis aristam superantibus.  $\frac{1}{2}$  in editioribus Alpibus et Pyrenæis. — *Phleum Gerardi* All. Jacq. ic. rar. 2. t. 301. Fl. fr. 3. p. 7. (v. v.)

XIX. *CRYPISIS* *Ait. Lam. ill. t. 42.* — *Phlei, Anthoxanthi et Schæni*  
*sp. L.*

Gluma uniflora bivalvis compressa; perigonium bivalve, valvulis inæqualibus lanceolatis inermibus glumâ longioribus; stamina 2 seu 3. — Panicula spiciformis.

1. *C. ALOPECUROIDES* (Schrad. germ. 1. p. 167.) culmis teretibus subsimpli-  
cibus, paniculâ oblongâ nudâ, floribus triandris. ① in provinciis occiden-  
talibus. — Fl. fr. 5. p. 248. *Heleochloa alopecaroides* Host. gram. 1. t. 29.  
(v. s.)
2. *C. SCHÆNOIDES* (Lam. ill. t. 42. f. 1.) culmis ramosis subcanaliculatis, pani-  
culâ ovatâ vaginâ foliaceâ cinctâ.  $\frac{1}{2}$  in locis petrosis humidis Occitanicæ,  
Andegavie ex Bast. — Fl. fr. 3. p. 3. *Phleum schænoides* L. Cav. ic. t. 52.  
*Heleochloa schænoides* Host. gram. 1. t. 30. (v. v.)
3. *C. ACULEATA* (Lam. ill. t. 42. f. 2.) culmis ramosis canaliculatis, paniculâ  
subrotundâ foliis vaginantibus brevibus pungentibus cinctâ.  $\frac{1}{2}$  in arenosis  
maritimis provinc. mediterranearum, secus Ligerim ex Bast. — Fl. fr. 3.  
p. 3. Host. gram. 1. t. 31. *Schœnus aculeatus* L. Cav. ic. t. 52. (v. v.)

XX. *ANTHOXANTHUM* *L. gen. n. 42.*

Gluma uniflora bivalvis; perigonium bivalve, valvis acutis dorso aristulatis; stamina 2. — Panicula spiciformis.

1. *A. ODORATUM* (L. sp. 40.) paniculâ ovato-oblongâ spiciformi, flosculis aristâ  
longioribus.  $\frac{1}{2}$  in pratibus. — Fl. fr. 3. p. 2. et 5. p. 247. Lam. ill. t. 23.  
(v. v.)

## 510 MONOCOTYL. SEU ENDOGENÆ.

### +++ FLORES PANICULATI HERMAPHRODITI, RARIUS POLYGAMI; SPICULÆ MULTIFLORE.

XXI. LAMARCKIA *Mench. meth.* 201. DC. fl. fr. 3. p. 30. — *Cynosuri* sp. L.

Spiculæ aliæ steriles ternæ pendulæ muticæ ad basin fertilium involucrum mentientes, aliæ fertiles geminatæ; gluma 2-valvis, 2-3-flora; perigonium 2-valve, valvulâ exteriorē longè aristatâ.

1. L. AUREA (Mench. l. c.). ① ad rupes Nicæ, Galloprovinciæ australioris, agri Ruscimonensis circâ Collioure (Ph. Thomas), Corsicæ. — Fl. fr. 3. p. 30. *Cynosurus aureus* L. sp. 107. *Chrysurus cynosuroides* Pers. (v. v.)

XXII. MELICA *Mench. meth.* 172. DC. fl. fr. 3. p. 30. — *Melicæ* sp. L.

Gluma bivalvis scariosa 2-3-flora, floribus 1-2 hermaphroditis, 1 terminali abortivâ; perigonium bivalve subventricosum.

#### §. 1. Perigonio glabro.

1. M. UNIFLORA (Retz. obs. 1. p. 10.) foliis planis, vaginis oppositifoliis, paniculâ laxâ paniculâ ramosâ, spiculis erectis floribus hermaphroditis solitariis. ½ in sylvis. — Fl. fr. 3. p. 30. eng. bot. t. 1058. M. nutans Lam. ill. t. 44. non L. (v. v.)
2. M. MONTANA (L. sp. 98 exel. pl. syn.) foliis planis, paniculâ nutante secundâ simplici, spiculis pendulis, floribus hermaphroditis binis. ½ in sylvis montanis. — eng. bot. t. 1059. M. montana Huds. angl. 37. Fl. fr. 3. p. 31. (v. v.)
3. M. RAMOSA (Vill. dauph. 2. p. 91.) foliis convolutis rigidis, paniculâ laxâ pyramidatâ, floribus hermaphroditis binis. ½ ad rupes apricas Delphin., Galloprov., Narbonæ, agri Ruscimonensis. — Fl. fr. 3. p. 31. M. aspera Desf. — Barr. t. 95. f. 1. (v. v.)

#### §. 2. Perigonio extus villosa.

4. M. CILIATA (L. sp. 97.) paniculâ spiciformi elongatâ, perigonis ciliato-villosissimis. ½ in collibus sterilibus saxosis. — Fl. fr. 3. p. 32. — Schenckz. gram. t. 3. f. 16. (v. v.)
5. M. BAUMERI (All. auct. p. 43.) paniculâ laxâ subsecundâ ramulis inferis divaricatis, perigonis ciliato-villosis. ½ ad rupes apricas Galliæ mediterraneæ passim. — Fl. fr. 3. p. 32. Host. gram. 4. t. 23. (v. v.)

XXIII. AIROPSIS *Desv. journ. bot.* 1. p. 200. DC. fl. fr. 5. p. 262.

Glumæ bifloræ bivalves nitentes concavæ obtusæ; flores ante maturitatem inclusi mutici.

1. A. GLOBOSA (Desv. l. c.) culmis erectis filiformibus cespitosis, foliis setiformibus, vaginâ superiore subtriculoâ, paniculâ coarctatâ, perigonis ciliatis. ① in ericetis circâ Aquas Tarbellicas, propè Forum-Julii (cl. Solier). — Fl. fr. 5. p. 262. *Aira globosa* Thore *journ. bot.* 1. t. 7. f. 3. 4. *Milium tenellum* Cav. ic. 3. t. 274. f. 1. (v. v. D.)
2. A. AGROSTIDEA (DC. fl. fr. 5. p. 262.) culmis ramosis basi geniculatis radican- tibus, foliis planis, paniculâ expansâ laxâ, perigonis membranaceis gla- bris. ½ in humidis herbosis secus rivos et stagna Armoracæ, Andegaviæ, Soloniæ. — *Poa agrostide* aDC. ic. rar. t. 1. *Airopsis Candollii* Desv. (v. s.)



XXIV. AIRA *Mench. meth.* 182. *DC. fl. fr.* 3. p. 42. — *Airæ sp. L.*

Gluma biflora bivalvis; perigonium bivalve, valvulâ exteriori a basi aristam geniculatam emittente. — Flores paniculati nitentes.

1. *A. CRISPATA* (L. sp. 96.) foliis planis nervosis, ligulâ bifidâ, paniculâ elongatâ ramosissimâ patente, glumæ valvulis acutis flosculos basi pilosos subæquantibus, flosculo altero pedicellato, aristis rectis glumâ vix longioribus.  $\mathcal{H}$  in pratis et sylvis. — *Fl. fr.* 3. p. 43. *eng. bot.* t. 1453. *Deschampsia cespitosa* P. de Beauv. — *Scheuchz. gram.* t. 2. f. 2 et 3. (v. v.)
2. *A. FLEXUOSA* (L. p. 96.) foliis setaceis, ligulâ bifidâ, paniculâ laxâ pauciflorâ flexuosâ, glumæ valvulis subacutis flosculos basi villosos subæquantibus, flosculo altero pedicellato, aristis glumâ duplò longioribus.  $\mathcal{H}$  in sylvis et montanis. — *Fl. fr.* 3. p. 43. *eng. bot.* t. 1519. *Aira montana* L. — *Scheuchz. prod.* t. 4. f. 4. et t. 6. f. 1. (v. v.)
3. *A. CAPILLARIS* (Host. *gram.* 4. t. 35.) foliis setaceis, ligulâ oblongâ acuminatâ, paniculâ subcorymbosâ capillari patentissimâ, glumâ truncatâ mucronulatâ flosculos sessiles superante, flosculo superiore aristato aristâ glumam paululum excedente. (1) in Corsicâ et insulis Stæchadam (cl. Requier). — *A. elegans* Willd. *ined.* in Röm. et Sch. 2. p. 682. *Variat mutica.* (v. s.)
4. *A. CARIOPHYLLA* (L. sp. 97.) foliis setaceis, ligulis acuminatis integris, vaginis scabriusculis, paniculâ divaricatâ trichotomâ, glumæ valvulis acutis flosculos sessiles glabros superantibus, aristis glumâ paulò longioribus.  $\mathcal{H}$  in siccis et arenosis. — *Fl. fr.* 3. p. 44. *Lam. ill.* t. 44. *eng. bot.* t. 812. — *Moris. s.* 8. t. 5. f. 15. n. 11. (v. v.)
5. *A. ARTICULATA* (Desf. *atl.* 1. t. 13.) foliis planis siccitate convolutis, ligulâ oblongâ subacutâ, paniculâ primo coerctâ demùm patulâ, glumis acuminatis, aristis medio articulatis apice clavatis glumam subæquantibus. (1) in arenosis maritimis Corsicæ, Nicæ, Telonis, Narbonæ (cl. Requier), agri Ruscitonensis propè Prades (Ph. Thomas). — *Fl. fr.* 5. p. 261. *Corynephorus articulatus* P. de Beauv. (v. s.)
6. *A. CANESCENS* (L. sp. 97.) foliis setaceis involutis, ligulâ oblongâ truncatâ, paniculâ coerctâ, glumis acuminatis, aristis medio articulatis apice clavatis glumam subæquantibus. (1) in arenosis passim. — *Fl. fr.* 3. p. 44. *eng. bot.* t. 1190. *Corynephorus canescens* P. de Beauv. — *Moris. s.* 8. t. 3. f. 2. n. 10. (v. v.)
7. *A? MEDIA* (Gou. *ill.* 3.) culmo erecto 2-3-folio, foliis glaucis convulso-setaceis, radicalibus fasciculatis, ligulâ oblongâ acuminatâ, paniculâ patente, glumis acutis perigonium truncatum basi pilosiusculum subæquantibus, aristâ dorsali recta glumâ subbreuiore. (1) in aridis Monspelii, Galloprovinciæ propè Sisteron et Delphin. propè Vapincum ex Vill. — *Fl. fr.* 5. p. 261. *Deschampsia media* et *D. juncea* Röm. et Sch. 2. p. 687. — *An vera Aira?* (v. s.)
8. *A. PRÆCOX* (L. sp. 97.) culmo erecto pusillo, foliis setaceis involutis, paniculâ coerctâ subspiciformi, glumæ valvulis acutis glumellam acuminatam subæquantibus, aristis medio articulatis apice filiformibus glumâ subduplò longioribus. (1) in arenosis Galliæ occidentalis, mediæ et australis. — *Fl. fr.* 3. p. 44. *eng. bot.* t. 1296. *Avena præcox* P. de Beauv. — *Ray. ang.* t. 22. f. 2. (v. v.)

XXV. AVENA *Kœl. gram.* 276. *DC. fl. fr.* 3. p. 34. — *Avenæ et Holci sp. L.*

Gluma 2-valvis 2-aut multiflora; flores omnes hermaphroditi aut interdum abortu masculi; perigonium 2-valve, valvulâ exteriori dorso aristam geniculatam gerente.

## 512 MONOCOTYL. SEU ENDOGENÆ AVENA.

Sect. I. PSUD'HOLCUS. — *Holcus Gaud. agrost. 1. p. 135. Holci sp. L.*

Glumæ 2-3-floræ sæpius polygamæ, flosculis dissimilibus altero mutico altero aristato.

1. *A. ODORATA* (Kœl. gram. 299.) foliis planis glabris, paniculâ laxâ, spiculis trifloris, flosculis glumæ valvulas ovatas subæquales membranaceas mucronatas æquantibus, lateralibus masculis intermedio hermaphrodito.  $\frac{1}{2}$  in pratis humidis et fossis Arvernæ, Mimatis, Monspelii. — Fl. fr. 3. p. 42. *Holcus odoratus* L. *Hierochloa odorata* et *H. repens* P. de Beauv. — Schenck. gram. t. 4. f. 26. Variat flosculis 1-2-muticis aristatisve, staminibus 2-3. (v. a.)
2. *A. MOLLIS* (Kœl. gram. 301.) foliis planis, vaginis glabriusculis, nodis villosis, paniculâ contractâ cylindricâ, spiculis 2-floris, glumæ valvulis æqualibus carinâ ciliatis flosculos superantibus, glumellæ valvulâ externâ glabrâ basi pilosâ.  $\frac{1}{2}$  in pratis. — Fl. fr. 3. p. 41. *Holcus mollis* L. sp. 1485. eng. bot. t. 1170. — Schenck. gram. t. 4. f. 25. (v. v.)
3. *A. LANATA* (Kœl. gram. 303.) foliis planis vaginisque lanuginosis, paniculâ contractâ cylindricâ, spiculis 2-floris, glumæ valvulis obtusis subæqualibus apice ciliatis glumam superantibus, glumellæ valvulâ externâ glabrâ.  $\frac{1}{2}$  in pratis. — Fl. fr. 3. p. 41. *Holcus lanatus* L. eng. bot. t. 1169. — Moris. s. 8. t. 7. f. 5. n. 34. (v. v.)
4. *A. ELATOR* (L. sp. 117.) radice repente, foliis planis nodisque glabris, paniculâ subcontractâ nutante, spiculis 2-floris, glumæ valvulis glabris lævibus acutis, internâ flosculos subæquante, externâ duplò minore, glumellæ valvulâ externâ 2-dentatâ basi pilosâ.  $\frac{1}{2}$  in pratis. — Fl. fr. 3. p. 41. *Holcus avenaceus* Smith. eng. bot. t. 813. *Gaud. Arrhenatherum avenaceum* P. de Beauv. — Moris. s. 8. t. 7. f. 11. n. 37. (v. v.)  
 $\beta$ . *bulbosa*, radice bulbiferâ, culmi nodis inferioribus plerumque pubescentibus. — *A. bulbosa* Willd. DC. fl. fr. 5. p. 261. *A. precatoria* Thail. — Moris. s. 8. t. 7. f. 12. n. 38. (v. v.)
5. *A. THOREI* (Duby mss.) foliis angustis involutis rigidis longissimis inferioribus vaginis nodisque pubescentibus, paniculâ erectâ subcontractâ, spiculis 2-floris, flosculis glumam subæquantibus inferno aristato, superiore mutico, axi piloso, glumæ valvulis inæqualibus acuminatis, glumellæ valvulâ externâ pubescenti acuminatâ integrâ.  $\frac{1}{2}$  in ericetis siccis et collibus circâ Aquas Tarbellicas. — *A. longifolia* Thore. (v. v. D.)

### Sect. II. AVENASTRUM. — *Avena* L.

Glumæ 3-7 floræ, flosculis plerumque omnibus aristatis, perigonii valvulâ externâ integrâ, dentatâ aut bifidâ.

6. *A. FLAVESCENS* (L. sp. 118.) foliis planis mollibus, inferioribus pubescentibus, paniculâ subcontractâ (flavescente), spiculis 2-5-floris, flosculis glumam superantibus, axi pubescente, glumæ valvulis acuminatis, externâ subduplò minore, glumellæ valvulâ externâ glaberrimâ apice 2-fidâ.  $\frac{1}{2}$  in collibus et pratis siccis. — Fl. fr. 3. p. 40. *Trisetum pratense* Pers. ench. 97. — Moris. s. 8. t. 7. f. 19. n. 42. (v. v.)  
 $\beta$ . *alpestris*, radice fibrosâ, paniculâ contractiore purpurascente. — In Alpibus. — *A. Alpestris* Host. DC. fl. fr. 5. p. 260. *A. sesquiteria* Fl. fr. 3. p. 40. non L. (v. v.)
7. *A. TENUIS* (Mœnch. meth. 195.) foliis glabris angustis brevibus, paniculâ tenui laxâ, spiculis 2-4-floris, flosculis glumam subsuperantibus, axi glabro, glumæ valvulis acuminatis externâ subduplò minore, flosculi inferioris glumellæ valvulâ externâ apice in aristam rectam desinente, cæterorum valvulâ externâ apice aristato-bifidâ basi breviter pilosâ. ① in campis aridis Delphin., Mimatis (cl. Prost.). — Fl. fr. 3. p. 39. *A. triaristata* Vill. dauph. 2. t. 4. *Trisetum striatum* Pers. (v. s.)

8. *A. SETACEA* (Vill. dauph. 2. t. 5.) foliis convolutis setaceis longissimis, vaginis tomentosis, paniculâ erectâ contractâ, spiculis 2-3-floris, glumæ valvulis subinæqualibus acuminatis flosculos subæquantibus, axi piloso, glumellæ valvulâ externâ 2-fidâ basi pilosâ.  $\mathcal{L}$  in Alpibus Delphinatûs. — Fl. fr. 3. p. 37. (v. s.)
9. *A. PANICKA* (Lam. ill. 117.) foliis planis mollibus vaginisque pilosis, paniculâ ovato-elongatâ densâ, spiculis 4-6-floris, glumæ valvulis inæqualibus acuminatis flosculos subæquantibus, axi glabro, glumellæ valvulâ externâ glaberrimâ apice aristato-bifidâ. ① in arenosis maritimis Corsicæ. — *A. neglecta* Savi pis. 1. t. 1. f. 4. Spiculæ pro in genere minimæ. (v. s.)
10. *A. AIROIDES* (Koel. gram. 208.) culmo apice tomentoso, foliis planis angustis glabris, paniculâ erectâ cylindricâ spicæformi, spiculis 2-3-floris, glumæ valvulis subæqualibus acutis flosculos subæquantibus, axi parçè piloso, glumellæ valvulâ externâ glabrâ bidentatâ.  $\mathcal{L}$  in summis Alpibus et Pyrenæis. — Fl. fr. 3. p. 37. *A. subspicata* Clairv. Gaud. fl. helv. *Aira subspicata* L. Jacq. coll. t. 19. f. 3. *Trisetum airoides* P. de Beauv. — Scheuchz. prod. t. 6. f. 2. (v. s. ex Helv.)
11. *A. FRAGILIS* (L. sp. 119.) foliis planis villosis, spicâ fragili articulatâ, spiculis 4-6-floris sessilibus adpressis alternis, glumæ valvulis valdè inæqualibus obtusiusculis, glumellæ valvulâ externâ glabrâ brevissimè 2-dentatâ. ① in provinciis orientalibus, australibus usque ad Tolosam, propè Nannetem (cl. Hectot). — Fl. fr. 3. p. 38. Schreb. gram. t. 24. f. 3. *Gaudinia fragilis* P. de Beauv. — Scheuchz. gram. t. 1. f. 7. G. (v. v.)
12. *A. PRATENSIS* (L. sp. 119.) foliis subinvolutis glabris margine asperis, caulibus brevissimis, paniculâ erectâ subspicatâ, pedicellis solitariis aut geminis, spiculis 5-8-floris, glumæ valvulis inæqualibus acutis flosculis brevioribus, axi parçè piloso, glumellæ valvulâ glabrâ ant. basi pilosulâ apice 2-dentatâ.  $\mathcal{L}$  in pratis ad sylvarum margines. — Fl. fr. 3. p. 38. eng. bot. t. 1204. *A. bromoides* L. — Scheuchz. gram. t. 4. f. 21. 22. (v. v.)
13. *A. DISTICHOPHYLLA* (Vill. dauph. 2. t. 4.) culmo ascendente basi ramoso, foliis angustis planis plicatisve, ramorum juniorum distichis, ligulâ brevissimâ truncatâ, paniculâ erectâ angustâ, spiculis 2-3-floris, glumæ valvulis subæqualibus acutis, flosculis glumam subæquantibus, axi villosis, glumellæ valvulâ externâ 2-dentatâ basi longè pilosâ.  $\mathcal{L}$  ad rupes in Alpibus. — Fl. fr. 3. p. 36. *Trisetum distichophyllum* P. de Beauv. (v. s.)
14. *A. PUBESCENS* (L. sp. 1665.) foliis planis inferioribus pubescentibus, ligulis superioribus oblongis obtusis, paniculâ erectâ subsimplici, spiculis 3-floris, flosculis glumam subæquantibus, glumæ valvulis acuminatis, externâ duplò minore, glumellæ valvulâ externâ glabrâ apice eroso-dentatâ.  $\mathcal{L}$  in pratis montanis et collinis. — Fl. fr. 3. p. 36. *A. amethystina* Fl. fr. 3. p. 37 et 5. p. 260. — Scheuchz. gram. t. 4. f. 20. (v. v.)
15. *A. VASICOLOR* (Vill. dauph. 2. t. 4.) foliis planis angustis, ligulâ oblongâ subcutâ, paniculâ laxâ erectâ angustâ, spiculis 5-6-floris, glumæ valvulis inæqualibus acuminatis flosculis brevioribus, axi piloso, glumellæ valvulâ externâ glaberrimâ acutâ.  $\mathcal{L}$  in Alpibus, montibus Lugdunensibus, Arverniciâ, Pyrenæis. — Fl. fr. 3. p. 36. *A. Scheuchzeri* Host. gram. 2. t. 52. — Scheuchz. prod. t. 3. (v. v.)
16. *A. SEMPERVIRENS* (Vill. dauph. 2. t. 5.) foliis longissimis angustis rigidis subinvolutis intus striatis, ligulâ brevissimâ vaginæque oris pilosis, paniculâ subpatulâ nutante, spiculis 3-floris, flosculis glumam subæquantibus, 2 inferioribus tantum aristatis, axi piloso, glumellæ valvulâ externâ bifidâ basi lanatâ.  $\mathcal{L}$  in montosis apricis Delphin., Pyrenæorum. — Fl. fr. 3. p. 35. (v. s.)
17. *A. SEDUKENSI* (DC. fl. fr. 3. p. 719 et 5. p. 260.) foliis angustis involutis culmo multò brevioribus, ligulâ brevi truncatâ, paniculâ erectâ contractâ, pedicellis geminis, spiculis 2-3-floris, glumæ valvulis inæqualibus acuminatis flosculos subæquantibus, axi piloso, glumellæ valvulâ externâ 2-fidâ

# 514 MONOCOTYL. SEU ENDOGENÆ.

basi pilosâ. ♀ in Alpibus Galloprov., Delphin., monte Celtico, Pyrenæis. (v. v.)

18. *A. LONGIFOLIA* (Req. et DC. ined.) foliis angustis planis involutivæ in paginâ superiore asperis, ligulâ oblongâ acuminatâ, paniculâ erectâ angustâ contractâ, pedicellis solitariis geminisve, spiculis 5-8-floris, glumæ valvulis inæqualibus acuminatis flosculis brevioribus, axi parçè piloso, glumellæ valvulâ externâ breviter bidentatâ basi pilosâ. ♀ inter pineta propè Narbonam reperiit cl. Requier. (v. s.)
19. *A. ALBA* (Vahl. symb. 2. p. 24.) foliis linearibus planis, ligulâ brevi obtusâ, paniculâ laxâ erectâ angustâ, spiculis brevibus bifloris, glumæ valvulis lævibus inæqualibus externâ acutâ duplò minore, internâ obtusâ flosculos subæquante, flosculo altero mutico, glumellæ valvulâ externâ 7-nerviâ acuminatâ basi breviter pilosâ. ♀ in agro Ruscinonensi propè Prades (cl. Coder). — Fl. fr. 5. p. 259. (v. s.)
20. *A. BREVIS* (Willd. 1. p. 445.) radice fibrosâ, foliis linearibus planis, ligulâ brevi obtusâ, paniculâ secundâ, spiculis brevibus subbifloris, flosculis glumam æquantibus, glumellæ valvulâ externâ basi glabrâ apice pilosulâ bidentatâ. ① in agris circâ Parisios ex Merat, Neustris propè Vire ex cl. Le Prév. et Lenormand. — Fl. fr. 5. p. 258. Host. gram. 3. t. 52. (v. s. c.)
21. *A. ORIENTALIS* (Schreb. spic. 52.) radice fibrosâ, paniculâ secundâ contractâ, spiculis bifloris, flosculis glumâ minoribus, altero mutico, axi glabro, glumellæ valvulâ externâ glabrâ breviter 2-dentatâ. ① colitur frequenter. — Fl. fr. 5. p. 258. Host. gram. 3. t. 44. (v. v. c.)
22. *A. RUDA* (L. sp. 118.) paniculâ laxâ, spiculis 3-floris, flosculo ultimo mutico, axi glabro, glumellæ valvulâ externâ glabrâ 2-fidâ, seminibus lævibus denudatis. ① in agris culta, interdum abortu mutica. — Fl. fr. 3. p. 34. Moris. s. 8. t. 7. f. 2. n. 4. — Lob. ic. t. 32. f. 1. (v. v. c.)
23. *A. SATIVA* (L. sp. 118.) paniculâ laxâ, spiculis dispersis, axi glabro, glumellæ valvulâ externâ glabrâ aristato-2-fidâ, seminibus lævibus perigonio involutis. ① in agris culta, interdum abortu mutica. — Fl. fr. 3. p. 34. — Lob. ic. t. 31. f. 2. (v. v. c.)
24. *A. STERILIS* (L. sp. 118.) radice fibrosâ, foliis lato-linearibus vaginisque inferioribus glabris, ligulâ ovatâ acutâ, paniculâ secundâ, spiculis sub-3-floris, axi glabro, flosculis glumâ minoribus, inferioribus aristatis et pilosis, superioribus muticis glabrisque, glumellæ valvulâ externâ aristato-bifidâ. ① inter segetes Occitanis ex Lam. et Benth. — Fl. fr. 5. p. 259. Jacq. ic. rar. 1. t. 23. Host. gram. 2. t. 57. (v. v. c.)
25. *A. FATUA* (L. sp. 118.) foliis linearibus planis vaginisque inferioribus pubescentibus, ligulâ ovatâ truncatâ, paniculâ laxâ, spiculis 3-5-floris, flosculis glumam æquantibus, omnibus aristatis basi hirsutissimis, axi hirsuto, glumellæ valvulâ externâ apice fissâ biaristatâ. ① in arvis. — Fl. fr. 3. p. 35. Schreb. gram. t. 45. Host. gram. 2. t. 58. (v. v.)
26. *A. STRIGOSA* (Schreb. spic. 52.) foliis glabris planis scabris, ligulâ ovatâ obtusâ, paniculâ tenui laxâ, spiculis 2-floris, flosculis glumam subæquantibus omnibus aristatis, axi glabriusculo, glumæ valvulis subæqualibus aristato-acuminatis, glumellæ valvulâ externâ apice striatâ et scabrâ, internâ duplò minore. ① in provinciis australibus. — Fl. fr. 3. p. 40. eng. bot. t. 1266. Danthonia strigosa P. de Beauvois. (v. s.)

XXVI. DANTHONIA DC. fl. fr. 3. p. 32. — Danthonia et Triodia R. Brown.

Gluma multiflora bivalvis, valvulis maximis concavis; perigonium bivalve, valvulâ exteriore apice fissâ, et ex imâ fissurâ aristam nunc longam tortam; nunc ferè abortivam emittente.

i. *D. PROVINCIALIS* (DC. l. c.) foliis glabris ore pilosis, inferioribus convolutis setaceis, spiculis 5-6-floris, aristâ longâ tortili. ♀ in Galloprovinciâ propè

*Segetesteronem*, in Delphinatâ propè Vapincum. — *Avena calycina* Vill. daph. 2. t. 2. f. 9. non Lam. — Monti prod. t. 2. f. 1. (v. s.)

2. *D. decumbens* (DC. l. c.) foliis pubescentibus planis, spiculis 3-4-floris, aristâ subabortivâ. 2 in collibus et sylvis siccioribus. — *Festuca decumbens* L. *Poa decumbens* Eng. bot. t. 792. *Triodia decumbens* P. de Beauv. (v. v.)

## XXVII. BROMUS L. gen. 89.

Gluma multiflora bivalvis; perigonium bivalve; valvulâ exteriore majore concavâ, aristam sub apice gerente; interiore minore duplici ciliorum serie extus onustâ.

1. *B. caesus* (Desf. ined. in DC. fl. fr. 3. p. 68.) foliis firmis pubescentibus vaginis sulcatis, paniculâ laxâ subnutante, spiculis ovatis compressis pubescentibus 8-12-floris, flosculis demum laxis, glumæ valvulis ovatis externâ acutâ internâ mucronatâ, glumellæ valvulâ externâ ovatâ obtusâ 2-fidâ, aristâ rectâ glumellam subæquante. ① in sterilibus secus vias. — Fl. fr. 5. p. 275. *B. velutinus* Schrad. (v. v.)
2. *B. secalinus* (L. sp. 112.) foliis mollibus subpubescentibus, vaginis sulcatis inferioribus hirsutis, paniculâ laxissimâ nutante, spiculis ovatis glabris compressis 6-12-floris, flosculis demum laxissimis, glumæ valvulis ovatis externâ acuminatâ internâ mucronatâ, glumellæ valvulâ externâ subobtusâ bifidâ, aristâ rectâ glumellam subæquante. ① in arvis. — Fl. fr. 3. p. 67. Lam. ill. t. 46. f. 2. Eng. bot. t. 1171. — Moris. s. 8. t. 7. f. 4. n. 16. (v. v.)
3. *B. multiflorus* (Weig. obs. t. 1. f. 1.) foliis mollibus pubescentibus, vaginis præsertim inferioribus tomentosis, paniculâ laxâ suberectâ, spiculis oblongo-lanceolatis adpressè pubescentibus 12-14-floris, flosculis imbricatis, glumæ valvulis ovatis subacutis, glumellæ valvulâ externâ obtusâ bifidâ, aristâ rectâ glumam subæquante. ① in collibus et arvis ex Fl. fr. 3. p. 69. — *B. secalinus* Leers herb. t. 11. f. 2. — Scheuchz. gram. t. 5. f. 9. (v. s. ex Austria.)
4. *B. mollis* (L. sp. 112.) foliis lævibus vaginisque mollissimè villosis, paniculâ contractâ erectâ, pedunculis ramosis, spiculis ovato-oblongis 8-10-floris pubescentibus, flosculis imbricatis, glumæ valvulis ovatis acutis, glumellæ valvulâ externâ ovatâ subobtusâ 2-dentatâ, aristâ rectâ glumellæ longitudine. ① in pratis siccis secus vias. — Fl. fr. 3. p. 68. Lam. ill. t. 46. f. 1. Eng. bot. t. 1078. — Scheuchz. gram. t. 5. f. 12. — Moris. s. 8. t. 7. f. 5. n. 18. (v. v.)
5. *B. squarrosus* (L. sp. 112.) foliis mollibus vaginisque pubescentibus, paniculâ laxâ nutante simplicissimâ, spiculis ovatis compressis 9-10-floris, flosculis densè imbricatis, glumæ valvulis ovatis obtusis, glumellæ valvulâ externâ obtusâ integrâ, aristis demum divaricatis glumellâ sub brevioribus. ① ad agrorum margines imprimis in provinciis australibus. — Fl. fr. 3. p. 69. Eng. bot. t. 1885. — Scheuchz. gram. t. 5. f. 11. Variat spiculis glabris velutinisve. (v. v.)
6. *B. divaricatus* (Rohde in Lois. not. 22.) foliis mollibus glabriusculis, vaginis pubescentibus, paniculâ erectâ subcontractâ simplicissimâ, spiculis lanceolatis 10-15-floris pubescentibus, flosculis densè imbricatis, glumæ valvulis ovato-lanceolatis subacutis, glumellæ valvulâ exteriore bifidâ acutâ, aristis basi contortis demum divaricatis glumellâ sub brevioribus. ① in aridis arenosis à Nicæâ ad Monspelium. — Fl. fr. 5. p. 276. *B. lanuginosus* Poir. (v. v.)
7. *B. erectus* (Huds. ang. 49.) foliis radicalibus angustissimis ciliato-pilosis, vaginis pilosis, superioribus latioribus glabris, paniculâ erectâ laxâ, pedicellis subsimplicibus, spiculis lanceolato-oblongis 5-9-floris scabris, glumæ valvulis subæqualibus acuminatis, glumellæ valvulâ externâ acuminatâ bifidâ, aristâ rectâ glumellâ multò breviori. 2 in pratis. — Fl. fr. 3. p. 69. et 5. p. 276. Eng. bot. t. 471. — Vaill. bot. t. 18. f. 2. (v. v.)

## 516 MONOCOTYL. SEU ENDOGENÆ. BROMUS.

8. *B. ARVENSI* (L. sp. 113.) foliis firmis in paginā superiore pilosis, vaginis sulcatis, paniculā erectā laxā, pedunculis elongatis ramosis, spiculis lanceolatis glabrescentibus 6-8-floris, flosculis imbricatis, glumæ valvulis ovatis obtusis mucronatis, glumellæ valvulā externā ovato-rotundā obtusā, aristā rectā glumellā subbreviore. (1) in arvis. — Fl. fr. 3. p. 70. eng. bot. t. 1924. non Lam. — Scheuchz. gram. t. 5. f. 15. (v. v.)
9. *B. PRATENSIS* (Ehr. gram. 116.) foliis mollibus vaginisque molliter pubescentibus, paniculā erecto-patente, spiculis ovato-lanceolatis compressis 7-9-floris glabris, flosculis imbricatis, glumæ valvulis lanceolatis subobtusis, glumellæ valvulā externā ovatā brevissimē 2-dentatā, aristā rectā flosculi longitudine. 2/ in pratis et arvis. — Fl. fr. 3. p. 70. (v. v.)
10. *B. RACEMOSUS* (L. sp. 114.) foliis mollibus vaginisque pubescentibus, paniculā erectiusculā diffusā, spiculis ovatis 6-8-floris glabris, flosculis densē imbricatis, glumæ valvulis ovatis externā acutā internā obtusā aut mucronulatā, glumellæ valvulā externā ovatā obtusā, aristā rectā glumellam subsequente. (1) in pratis et arvis passim. — Fl. fr. 5. p. 275. eng. bot. t. 1079. *B. simplex* Gand. agr. 1. p. 296. (v. v.)  
*β. elongatus*, vaginis (exceptā infimā) glabris. — *B. elongatus* Gaud. agr. 1. p. 305. Fl. fr. 5. p. 275. (v. v.)
11. *B. ASPER* (L. snpp. 111.) foliis duris asperis lato-linearibus in paginā inferiori pubescentibus, vaginis hispidis pilis deflexis, paniculā laxā ramosā nutante, spiculis linear-oblongis 8-10-floris subpubescentibus, flosculis laxē imbricatis, glumæ valvulis inæqualibus externā acuminatā internā mucronatā, glumellæ valvulā externā acuminatā bidentatā, aristā subterminali rectā glumellā breviore. 2/ in sepibus et umbrosis. — Fl. fr. 3. p. 71. eng. bot. t. 1172. (v. v.)
12. *B. GIGANTEUS* (L. sp. 114) foliis lato-linearibus vaginisque glaberrimis, paniculā nutante subsecundā laxā, spiculis linearilanceolatis 4-5-floris glaberrimis, flosculis imbricatis, glumæ valvulis subæqualibus acuminatis, glumellæ valvulā externā acuminatā bidentatā, aristā subterminali rectā undulatā glumellā subduplō longiore. 2/ in nemorosis et pratis umbrosis passim. — Fl. fr. 3. p. 71. Host. gram. 1. t. 66. Vaill. bot. t. 18. f. 3. (v. v.)
13. *B. STERILIS* (L. sp. 113.) foliis mollibus vaginisque inferioribus pubescentibus, paniculā laxā, spiculis oblongis nutantibus 5-9-floris glabris, flosculis laxis nervosis sulcatis, glumæ valvulis inæqualibus acuminatis, glumellæ valvulā externā acuminatā bifidā, aristā subterminali glumellā longiore. (1) secus sepes et muros, in locis sterilibus. — Fl. fr. 3. p. 71. eng. bot. t. 1030. — Moris. s. 8. t. 7. f. 8. n. 11. — Scheuchz. gram. t. 5. f. 14. (v. v.)
14. *B. TECTORUM* (L. sp. 114.) foliis mollibus vaginisque pubescentibus, paniculā laxā subsecundā semiverticillatā, spiculis linearibus pendulis pubescentibus 5-7-floris, flosculis laxē imbricatis, glumæ valvulis inæqualibus acuminatis, glumellæ valvulā externā acuminatā bifidā aristam subsequente. (1) ad tecta et in locis sterilibus. — Fl. fr. 3. p. 72. Leers. herb. t. 10. f. 2. — Pluck. t. 299. f. 2. (v. v.)
15. *B. POLYSTACHYUS* (DC. fl. fr. 5. p. 276.) foliis mollibus ciliatis vaginisque inferioribus breviter pubescentibus, paniculā patente, pedicellis fasciculatis simplicibus ramosisve, spiculis linearibus 8-12-floris, flosculis laxis, glumæ valvulis inæqualibus acuminatis, glumellæ valvulā externā acuminatā bifidā, aristis rectis flosculi longitudine. (1) in versuris agri Monspelienis, Ruscinonensis, Burdigalensis, Nannetensis. — *B. multispicatus* Rœm. et Sch. Variat spiculis pubescentibus et glabris. (v. s.)
16. *B. MADRITENSIS* (L. sp. 114.) foliis mollibus vaginisque pubescentibus, paniculā erecto-patente coarctatā subramosā, pedicellis ciliato-scabris apice incrassatis, spiculis linearibus retrorsum scabris 5-7-floris, flosculis plerumque diandris, glumæ valvulis acuminatis inæqualibus, glumellæ valvulā externā acuminatā bifidā aristā rectā duplō breviore. (1) in arvis Galloprovinciæ, circā Manoscā, Avenionem, Massiliam, propē Monspelium,

Nannetum. — Fl. fr. 3. p. 72. et 5. p. 277. B. rigidus Roth. cat. 1. p. 17. Fl. fr. 5. p. 278. — Barr. ic. t. 76. f. 1. (v. v.)

β. *maximus*, pedicellis velutinis, aristis glumellâ subtriplo longioribus. — In arenosis circa Aquas-Mortuas, secus canalē Occitanicam ex cl. Benth. — B. *maximus* Desf. atl. 1. t. 26. Fl. fr. 5. p. 277. (v. v. DC.)

17. B. *RUPENS* (L. sp. 114.) foliis substrictis vaginisque glabriusculis aut brevissimè pubescentibus, paniculâ fasciculatâ simplici, spiculis breviter pedicellatis 5-7-floris villosis, glumæ valvulis acuminatis inæqualibus, glumellæ valvulâ externâ acuminatâ 2-fidâ aristæ rectæ subæquali. ① in arenosis Galliæ mediterraneæ. — Fl. fr. 3. p. 73. et 5. p. 278. Cav. ic. 1. t. 45. f. 2. Festuca rubens Pers. — Variat spiculis glabris, ciliato-scabris aut villosis. (v. v.)

### XXVIII. FESTUCA L. gen. 88.

Gluma bivalvis multiflora; perigonium bivalve, valvulis acutissimis, aristâ terminali sæpè donatis.

#### §. 1. *Glumæ valvulis inæqualibus, perigonii valvulis aristam iisdem longioribus gerentibus.*

1. F. *UNIGLUMIS* (Ait. kew. 1. p. 108.) foliis angustis involutis, paniculâ spiciformi erectâ, pedicellis compressis dilatatis, spiculis oblongis 4-6-floris glabris, glumæ valvulâ externâ brevissimâ, internâ aristato-acuminatâ, glumellæ valvulis subæqualibus acuminatis, externâ aristatâ, aristâ glumellâ duplo longiore. ① in arenosis Galliæ australis, occidentalis rariùs, propè Parisios. — Fl. fr. 3. p. 55. eng. bot. t. 1430. — Ray. syn. t. 16. f. 2. Stamina 3. (v. v.)
2. F. *CILIATA* (Danth. in DC. fl. fr. 3. p. 55.) foliis angustis involutis, paniculâ elongatâ coarctatâ subnutante, spiculis 4-6-floris ciliato-hirsutis, flosculis monandris, glumæ valvulis brevissimis, externâ minimâ, glumellæ valvulis æqualibus acuminatis, externâ aristatâ aristâ subduplo breviorē. ① in rupestribus Galliæ mediterraneæ, Aginni ex St.-Am., Burdigalæ (Des Moulins). — Fl. fr. 3. p. 55. et 5. p. 268. Host. gram. 4. t. 65. — Scheuchz. gram. t. 6. f. 12. (v. v.)
3. F. *MYRUS* (L. sp. 109.) foliis angustis involutis, paniculâ elongatâ coarctatâ subnutante, spiculis 4-8-floris glabriusculis, flosculis monandris, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis æqualibus acuminatis, externâ aristatâ aristâ duplo breviorē. ① ad muros, in locis glareosis et arenosis. — Fl. fr. 3. p. 54. eng. bot. t. 1412. Vulpia Myrus Gmel. — Barr. ic. t. 99. f. 1. 2. (v. v.)
4. F. *BROMODES* (L. sp. 110.) foliis setaceis involutis, ligulis obsoletis, paniculâ simplicissimâ secundâ spiciformi erectâ, spiculis 3-4-floris scabris, flosculis monandris, glumæ valvulis inæqualibus acuminatis, externâ aristatâ aristâ subduplo breviorē. ① in campis et pratis arenosis passim. — Fl. fr. 3. p. 55. Lam. ill. t. 46. f. 4. eng. bot. t. 1411. Vulpia sciuroides Gm. — Pluck. t. 33. f. 10. An satis à priori distincta? (v. s.)

#### §. 2. *Glumæ valvulis subæqualibus, perigonii valvulis aristam iisdem breviorē gerentibus.*

5. F. *STIPOIDES* (Desf. atl. 1. p. 90.) foliis linearibus convolutis, paniculâ subcontractâ secundâ, pedicellis apice dilatatis compressis, spiculis 3-5-floris lanceolatis, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis subæqualibus internâ bidentatâ, externâ acuminatâ aristatâ aristæ longitudine. ① in arenosis Corsicæ, inter Niceam et Telonem, propè Monspelium (Salzmänn). — Fl. fr. 5. p. 267. Bromus stipoides L. mant. 567. — Barr. ic. t. 76. f. 2. (v. s.)
6. F. *SABULICOLA* (L. Dufour in ann. sc. nat. 5. p. 85.) glauca, radice repente, foliis convolutis angustis erectis, paniculâ elongatâ subsecundâ laxiusculâ,

## 518 MONOCOTYL. SEU ENDOGENÆ. *FESTUCA.*

- spiculis compressis lanceolatis tomentosis 4-9-floris, glumæ glumellæque valvulis inæqualibus aristato-acuminatis.  $\mathcal{L}$  in sabulosis agri Syrtici. — *F. arenaria* Askel. diss. supp. scand. p. 4? non Lam.? *Schœnolurus phœnicoides* Rœm. et Sch. 2. p. 699. (v. s.)
7. *F. GLAUCA* (Lam. dict. 2. p. 459.) foliis glaucis filiformibus convolutis rigidis extus lævibus intus vaginisque inferioribus pubescentibus, paniculâ glaucâ patente strictâ pyramidatâ, pedicellis scabris, spiculis 2-5-floris, glumæ valvulis inæqualibus, externâ acuminatâ, internâ macronatâ apice ciliatâ, glumellæ valvulis æqualibus glabris aut apice pubescentibus, internâ 2-fidâ, externâ aristatâ, aristâ subduplò longiore.  $\mathcal{L}$  in arenosis siccis passim. — Fl. fr. 3. p. 31. (v. v.)
8. *F. DURIUSCULA* (L. sp. 108.) radice fibrosâ, foliis strictis setaceis, inferioribus angulato-convolutis, superioribus laxius convolutis suprâ glabris, paniculâ strictâ subsecundâ, axi pedunculisque lævisculis, spiculis 5-6-floris, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis æqualibus acuminatis, externâ aristatâ aristâ triplò duplòve longiore.  $\mathcal{L}$  in pratis siccis.
- a. glabra*, spiculis glabris. — *F. duriuscula* DC. fl. fr. 3. p. 50. *F. stricta* a Gaud. agr. 1. 237. *F. curvula* Gaud. l. c. p. 239. (v. v.)
- β. cinerea*, spiculis pubescentibus. — *F. cinerea* Vill. DC. fl. fr. 3. p. 51. *F. dumetorum* L. F. dura Host. gram. 2. t. 87. Fl. fr. 5. p. 266. *F. stricta* β Gaud. agr. 1. t. 237. (v. v.)
9. *F. RUBRA* (L. sp. 109.) radice repente, foliis inferioribus convolato-setaceis, superioribus angustissimis planiusculis suprâ pubescentibus, paniculâ erectâ angustâ sublaxâ, axi pedunculisque angulato-scabris, spiculis 5-7-floris glaberrimis, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis æqualibus acuminatis externâ aristatâ aristâ triplò longiore.  $\mathcal{L}$  in siccis et sterilibus. — Fl. fr. 3. p. 50. *Nimium* var. *α* præcedentis affinis. (v. v.)
10. *F. HETEROPHYLLA* (Lam. fl. fr. 3. p. 600.) foliis longissimis laxis capillaribus mollioribus, superioribus planis interdum pubescentibus, paniculâ laxâ subpatulâ nante subsecundâ, axi pedunculisque ciliato-scabris, spiculis 5-6-floris glabriusculis, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis æqualibus internâ acutâ aut 2-fidâ, externâ aristato-acuminatâ aristâ longiore.  $\mathcal{L}$  in sylvaticis umbrosis.
- a. Lamarchii*, aristâ glumellâ vix duplò brevior. — *F. heterophylla* Hencke in Jacq. coll. Fl. fr. 3. p. 52. (v. v.)
- β. Hostii*, aristâ glumellâ multò brevior. — *F. heterophylla* Host. gram. 3. t. 18. (v. s.)
11. *F. OVINA* (L. sp. 108.) culmo subtetragono, foliis capillaribus scabris, paniculâ secundâ coarctatâ strictâ, pedunculis ciliato-scabris, spiculis 3-8-floris glaberrimis pubescentibusve, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis æqualibus internâ breviter bifidâ, externâ acuminatâ brevissimè aristatâ.  $\mathcal{L}$  in pascuis siccioribus et apricis. — Fl. fr. 3. p. 50. et 5. p. 264. *Leers herb.* t. 8. f. 3. (v. v.)
- β. tenuifolia*, foliis tenuissimis longioribus, spiculis muticis. — *F. tenuifolia* Smith ox. 138. Fl. fr. 5. p. 264. *Poa capillata* Merat paris. ed. 2. t. 2. p. 38. *P. setacea* Kœl. — *Moris. s.* 8. t. 3. f. 3. p. 13. (v. v.)
12. *F. FLAVESCENS* (Bell. act. tur. 5. p. 217.) culmo tereti, foliis setaceis elongatis subpuigentibus, paniculâ erectâ coarctatâ, spiculis 4-5-floris glaberrimis, glumæ valvulis inæqualibus externâ acuminatâ internâ subobtusâ, perigonii valvulis subæqualibus acuminatis margine membranaceis, externâ enerviâ acuminato-aristatâ.  $\mathcal{L}$  in editioribus Pyren. orientalium. — Fl. fr. 3. p. 719. et 5. p. 265. (v. v.)
13. *F. PUMILA* (Vill. dauph. 2. p. 102.) foliis setaceis duriusculis, paniculâ collectâ strictâ, pedicellis ciliato-scabris, spiculis 4-floris subteretibus oblongis scabris, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis æqualibus,



internâ 2-fidâ ciliatâ externâ acuminatâ brevissimè aristatâ.  $\mathcal{L}$  in Alpibus et Jurassi summo jugo Thoiry ex cl. Gaudin. — Fl. fr. 3. p. 52. *Schœnoderus pumilus* Rœm. et Sch. (v. v.)

14. *F. HALLERI* (All. ped. n. 2245.) culmo subnudo, foliis capillari-setaceis, paniculâ strictâ angustâ, axi pedunculisque angulato-scabris, spiculis 4-5-floris glabris teretiusculis oblongis, rachi glabrâ, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis æqualibus acuminatis externâ cristatâ aristâ glumellam subsequente.  $\mathcal{L}$  in Alpibus, monte Pilato, Pyrenæis. — Fl. fr. 3. p. 53. (v. v.)

15. *F. monescaze* (Lam. dict. 2. p. 460.) radice fibrosâ, culmo stricto, foliis convolutis setaceis, radicalibus longiusculis, culmeis brevibus extus glabris intus tenuissimè pubescentibus, paniculâ erectâ ramosâ laxiusculâ, spiculis 5-8-floris, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis acuminatis apice tenuissimè ciliato-scabris, internâ bifidâ, externâ aristatâ aristam subsequente.  $\mathcal{L}$  in pratis montanis Alpium, montis Aurei ex Lam., Jurassi ex Gaud. — Fl. fr. 5. p. 266. Spiculæ atro violaceo et albo variegatæ. (v. s.)

16. *F. RHETICA* (Sat. helv. 1. p. 52.) foliis involutis setaceis, paniculâ erectâ flexuosâ, pedicellis semi-verticillatis, spiculis 4-5-floris oblongis glabris, rachi infra fasciculos pilosâ, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis subæqualibus, internâ ciliatâ breviter 2-fidâ, externâ acuminatâ sub apice brevissimè aristatâ.  $\mathcal{L}$  in Pyren. orient. monte Canigon, montibus Ligeris (cl. Prost), Corsicæ (cl. Requier). — Fl. fr. 3. p. 53. et 5. p. 267. *F. pilosa* Gaud. *F. poæformis* Host. gram. 2. t. 81. *Poa violacea* Bell. act. tur. 5. t. 3. *Schœnoderus poæformis* P. de Beauv. Sequenti affinis sed debilior. (v. s.)

17. *F. EAKIA* (Ram. in DC. fl. fr. 3. p. 52. et 5. p. 267.) foliis involuto-complacatis nitidis pangentibus, paniculâ erectâ collectâ, spiculis 6-10-floris compressis oblongis glabris, rachi puberulâ, glumæ valvulis inæqualibus acutis glumellæ valvulis subæqualibus internâ ciliatâ breviter 2-fidâ, externâ acuminatâ aut breviter aristatâ.  $\mathcal{L}$  in Pyrenæis. — *F. lubrica* Lapeyr. *F. varia* Lapeyr. (v. v.)

§. 3. *Perigonii valvulis acutis aristâ destitutis.*

18. *F. SEROTINA* (L. sp. 111.) foliis convolutis rigidis patentibus pangentibus vaginisque glaberrimis, paniculâ primo erectâ demum divaricatâ, spiculis 3-5-floris, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis subæqualibus, externâ aristato-acuminatâ.  $\mathcal{L}$  in siccis apricis Galloprovinciæ, propè Monspelium loco dicto *Capouladoux* ex cl. Benth. — Fl. fr. 3. p. 46. Host. gram. 2. t. 42. *Schœnoderus serotinus* P. de Beauv. *Molinia serotina* Mert. et Koch. Gaud. fl. helv. 1. p. 217. Glumæ violaceæ. (v. v.)

19. *F. MARITIMA* (DC. fl. fr. 3. p. 47.) foliis convolutis erectis vaginisque glaberrimis, paniculâ subramosâ erectâ strictâ, pedicellis angulatis primo erectis demum divaricatis incrassatis, spiculis 6-8-floris compressis lanceolatis glaucis, glumæ valvulis subæqualibus acutis, glumellæ valvulis subæqualibus acuminatis. ①? in arenosis secus mare Mediterraneum. — *Triticum maritimum* L. sp. 128. *Brachypodium maritimum* Rœm. et Sch. — Bocc. mus. 2. t. 95. (v. v.)

20. *F. SPADICEA* (L. mant. 732.) foliis angustis lævibus subpangentibus, radicalibus involutis superioribus planiusculis, paniculâ erectâ laxiusculâ, pedunculis lævibus, spiculis 3-5-floris compressis ovatis, glumæ valvulis subæqualibus acuminatis, glumellæ valvulis subæqualibus internâ breviter 2-fidâ, externâ acuminatâ.  $\mathcal{L}$  in pratis montanis Arvernæ et Galliæ meridionalis. — Fl. fr. 3. p. 47. Host. gram. 3. t. 20. *Schœnoderus spadiceus* P. de Beauv. — Ger. gallop. t. 2. f. 1. Spiculæ nitido-aureæ. (v. v. DC.)

21. *F. LOLIACRA* (Curt. lond. 6. n. 66.) foliis planis linearibus vaginisque glabris, paniculâ simplici erectâ, spiculis 7-11-floris remotis distichis alternis subsessilibus, glumæ valvulis inæqualibus subacutis, glumellæ valvulis æqualibus

## 520 MONOCOTYL. sive ENDOGENÆ.

- acutis.  $\mathcal{L}$  in pratis humidis passim. — Fl. fr. 3. p. 48. eng. bot. t. 1821. *Schœnodoros loliaceus* P. de Beauv. (v. s.)
22. *F. INERMIS* (DC. fl. fr. 3. p. 49.) foliis planis lato-linearibus vaginisque glaberrimis, paniculâ erectâ laxâ demum diffusâ, spiculis 10-15-floris linearibus compressis, glumæ valvulis inæqualibus acutis, glumellæ valvulis subæqualibus internâ retusâ, externâ acuminatâ, aristis brevissimis subterminalibus sæpius abortivis.  $\mathcal{L}$  in pratis secus rivulos rariùs. — *Bromus inermis* L. Gaud. Schreb. gram. t. t. 13. *Schœnodoros inermis* Roem. et Sch. (v. s.)
23. *F. SYLVATICA* (Vill. dauph. 2. t. 2. f. 8.) culmis collo squamatis, foliis planis lato-linearibus, vaginisque inferioribus margine scabris, paniculâ erectâ laxâ patente ramosissimâ, spiculis 3-5-floris oblongis, glumæ valvulis subæqualibus acuminatis, glumellæ valvulis subæqualibus internâ apice breviter bifidâ, externâ acuminatâ.  $\mathcal{L}$  in sylvis Delphinatûs, Jurassî, Vogesorum. — Fl. fr. 3. p. 48. et 5. p. 263. *Poa trinervata* Fl. fr. 3. p. 58. *F. calamaria* eng. bot. t. 1005. *Schœnodoros calamarius* Roem. et Sch. (v. v.)
24. *F. ARUNDINACEA* (Schreb. spic. 57.) foliis lato-linearibus planis vaginisque glaberrimis, paniculâ ramosâ erectâ subdiffusâ, spiculis 4-6-floris ovatis compressis glaberrimis, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis subæqualibus acuminatis, externâ sub apice brevissimè aristatâ.  $\mathcal{L}$  secus rivulos passim. — Fl. fr. 3. p. 49. *F. elatior* Schrader. L. sp. 111. non fl. succ. *Bromus littoreus* Host. gram. t. t. 8. — Scheuch. gram. t. 5. f. 18. (v. v.)
25. *F. ELATIOR* (L. succ. 3a. non sp.) foliis planis linearibus vaginisque glaberrimis, paniculâ erectâ secundâ subcontractâ, spiculis 7-9-floris subteretibus, glumæ valvulis inæqualibus acuminatis, glumellæ valvulis subæqualibus internâ acutâ, externâ acuminatâ muticâ aut sub apice breviter aristatâ.  $\mathcal{L}$  in pratis. — Fl. fr. 3. p. 48. Host. gram. 2. t. 79. *F. pratensis* eng. bot. t. 1592. *Schœnodoros pratensis* P. de Beauv. — Moris. s. 8. t. 2. f. 19. n. 15. (v. v.)
26. *F. CÆRULEA* (DC. fl. fr. 3. p. 46.) culmo subuninodi, foliis planis vaginisque glaberrimis, paniculâ ramosâ coarctatâ, spiculis 2-3-floris cylindricis, glumæ valvulis inæqualibus ovatis obtusis mucronatisve, glumellæ valvulis æqualibus subobtusis.  $\mathcal{L}$  in sylvulis et pratis umbrosis. — Fl. fr. 3. p. 46. *Aira cærulea* L. sp. *Molinia cærulea* Moench. *Enodium cæruleum* Gaud. agr. — Moris. s. 8. t. 5. f. 20. n. 22. (v. v.)

### XXIX. ARUNDO Roth. germ. 1. DC. fl. fr. 3. p. 45. — Arundinis sp. L.

Gluma multiflora bivalvis; perigonium 2-valve extus villosissimum. — Flores paniculati.

Sect. I. PHRAGMITES Trin. fund. agrost. — Arundo P. de Beauv. t. 13. f. 2.

Glumæ 5-7-floræ; flosc. inferior masculus seu neuter; glumellæ valv. externa acuminata.

1. *A. PHRAGMITES* (L. sp. 120.) glumæ valvulis 5-7-floris, paniculâ laxâ, culmo herbaceo rigido.  $\mathcal{L}$  in lacubus, fluviis, stagnis puris, etc. — Fl. fr. 3. p. 45. *Phragmites communis* Trin. Gaud. fl. helv. t. p. 203. — Lab. ic. t. 51. f. 1. (v. v.)

Sect. II. DONAX P. de Beauv. t. 16. f. 4 et 5. — Scolochloa Mert. et Koch. deut. fl. 1. p. 530.

Glumæ 3-7-floræ, flosc. omnibus hermaphroditis; glumellæ valv. externa trisetâ.

2. *A. DONAX* (L. sp. 120.) culmo suffruticoso, paniculâ densâ, spiculis 3-6-floris, glumæ valvulis æqualibus, acuminatis flosculos æquantibus, glumellæ valvulis inæqualibus, externâ aristato-acuminatâ internâ duplò minore retusâ.  $\mathcal{L}$  in collibus Galloprovinc., Occitanie, agri Ruscinonensis. — Fl. fr. 3. p. 44. Host. gram. 4. t. 38. *Donax arundinaceus* P. de Beauv. — Moris. s. 8. t. 8. f. 1. n. 5. (v. v.)

3. *A. MAURITANICA* (Desf. atl. 1. p. 106.) culmo fruticoso, paniculâ densâ, spiculis 1-3-floris, glumæ valvulis subæqualibus acuminatis flosculos subæquantibus, glumellæ valvulis subæqualibus, externâ aristulatâ internâ acutâ.  $\frac{1}{2}$  in maritimis propè Antipolim reperiit cl. Gay, ad littus occidentale insulæ Sanctæ-Luciæ propè Narbonam indicat cl. Benth. — *Donax Mauritania* P. de Beauv. *Calamagrostis Plinii* Gmel. — Zannoni hist. t. 19. f. 2. (v. s.)

XXX. *DACTYLIS* Mich. DC. fl. fr. 3. p. 73. — *Dactylis* sp. L.

Gluma multiflora bivalvis, valvulis inæqualibus carinatis acutis; perigonium bivalve, valvulis carinatis alterâ apice aristam brevissimam gerente. — Panicula secunda glomerata.

1. *D. GLOMERATA* (L. sp. 105.) culmo erecto, foliis linearibus planis margine scabris.  $\frac{1}{2}$  in pratibus ubique frequens. — Fl. fr. 3. p. 73. Lam. ill. t. 44. — Scheuchz. t. 6. f. 15. (v. v.)

$\beta$ . *glaucescens*, foliis carinatis glaucescentibus rigidis. — In sylvaticis Galloprov. orientalis. — *D. glaucescens* Wild. en. ber. 1. p. 111. (v. s.)

$\gamma$ . *Hispanica*, culmo humiliori, foliis angustis margine lævibus. — In apricis maritimis. — *D. Hispanica* Roth. DC. fl. fr. 3. p. 278. (v. v.)

XXXI. *KOELERIA* Pers. ench. 1. p. 96. DC. fl. fr. 5. p. 268.

Gluma bivalvis compressa carinata 2-aut pauciflora; perigonium bivalve valvulâ exteriori carinatâ acuminatâ aut breviter aristatâ internam angustam acutiusculam plicatam amplexante; semen nudum. — Panicula contracta spiciformis.

1. *K. MACILENTA* (DC. fl. fr. 5. p. 270.) foliis angustis convolutis culmisque velutinis, paniculâ angustâ sublaxâ, spiculis 3-4-floris, glumæ valvulis valde inæqualibus, externâ lanceolatâ carinatâ acuminatâ, internâ quadruplò brevior subulatâ, glumellæ valvulâ externâ sub apice aristatâ, internâ duplò minore acutâ. ① in arenosis maritimis Monspelii, Corsicæ (cl. Requier). (v. s.)

2. *K. CALYCINA* (DC. fl. fr. 5. p. 271.) foliis setaceis ad vaginam et collum pilosis, paniculâ laxiusculâ oblongâ subinterruptâ, spiculis linearilanceolatis 2-5-floris, glumæ valvulis subæqualibus obtusiusculis margine membranaceis flosculos superantibus, glumellæ valvulis æqualibus obtusis, externâ nervosâ basi puberulâ. ① in arvis Galloprov. et ægri Ruscinonensis. — *Festuca calycina* Cav. ic. 1. t. 44. f. 2. *Schismus marginatus* P. de Beauv. (v. v. D.)

3. *K. VILLOSA* (Pers. ench. 1. p. 97.) foliis planis molliter villosis, paniculâ densè spiciformi ovali vel cylindricâ, spiculis linearibus 2-floris, glumæ valvulis æqualibus acuminatis dorso villosis flosculos subæquantibus, glumellæ valvulis subæqualibus externâ acutâ infrâ apicem aristatâ, internâ acutâ aut bidentatâ. ① in maritimis humidis secus Mediterraneum. — Fl. fr. 5. p. 270. *Phalaris pubescens* Lam. dict. 1. p. 92. DC. fl. fr. 3. p. 81. — Ger. gallop. 77. t. 1. (v. v.)

$\beta$ . *cinerea*, pusilla, foliis brevibus, spicâ ovatâ. — In arenosis Avenionis et Narbonæ (cl. Requier). (v. s.)

4. *K. PHELOIDES* (Pers. ench. 1. p. 97.) culmo glabro, foliis planis pilosis, paniculâ spiciformi cylindricâ, spiculis lanceolatis 2-6-floris, glumæ valvulis inæqualibus glabris acuminatis flosculis brevioribus, perigonii valvulis æqualibus internâ acuminatâ externâ extus piloso-exasperatâ sub apice aristatâ, aristâ molli flosculo longiore. ① in cultis arenosisve provinc. mediterr. et austro-occidentalium. — Fl. fr. 5. p. 271. *Festuca phleoides* Vill. dauph. t. 2. f. 7. Fl. fr. 3. p. 54. — Schenchz. gram. t. 5. f. 5. *Panicula interdum ramosa*. (v. v.)

5. *K. BRACHYTACHIA* (DC. cat. monsp. 120.) culmo glabro, foliis planis pilosis, paniculâ spiciformi ovato-cylindricâ, spiculis lanceolatis 6-9-floris, glumæ valvulis inæqualibus dorso ciliato-scabris acutis flosculis brevioribus, perigonii valvulis subæqualibus internâ acutâ externâ glabrâ sub apice aristatâ, aristâ molli flosculo brevior.  $\frac{1}{2}$  secus littora maris Mediterranei ex cl. Benth. (v. s. c.)

## 522 MONOCOTYL. SEU ENDOGENÆ.

6. *K. ALBESCENS* (DC. cat. monsp. 117. et fl. fr. 5. p. 269.) culmo pubescente, foliis angustis involutis inferioribus vaginisque pubescentibus, paniculâ elongatâ spiciformi hinc inde subinterruptâ basi folio superiore subvaginatâ, spiculis 2-3-floris lineari-lanceolatis, glumæ valvulis subæqualibus acutis dorso scabris, glumellæ valvulis inæqualibus acutis.  $\frac{1}{2}$  in sterilibus Tolosæ (DC.), Burdigalæ (Des Moulins), Andegaviæ (DC.). (v. v. DC.)
7. *K. SETACEA* (Pers. ench. 1. p. 97.) culmo longè exserto apice volutino, foliis inferioribus convolutis setaceis glaberrimis, paniculâ ovato-cylindricâ densè spicatâ, spiculis 2-3-floris lineari-lanceolatis, glumæ valvulis subæqualibus acuminatis dorso scabris, glumellæ valvulis subinæqualibus, externâ acuminatâ dorso subciliatâ internâ acutiusculâ.  $\frac{1}{2}$  in montosis apricis Galloprov., Pyrenæorum. — Fl. fr. 5. p. 269. *K. tuberosa* Pers. *Poa pectinacea* Lam. (v. v.)
8. *K. CRISTATA* (Pers. ench. 1. p. 97.) culmo glabriusculo, foliis planis inferioribus pubescentibus, paniculâ elongatâ spiciformi basi interruptâ, spiculis 3-4-floris lineari-lanceolatis, glumæ valvulis inæqualibus lanceolatis acuminatis dorso scabris, glumellæ valvulis subæqualibus acuminatis.  $\frac{1}{2}$  in collibus, pasosis, sterilibus. — Fl. fr. 5. p. 268. *Aira cristata* L. *Poa cristata* æ. Fl. fr. 3. p. 65. Host. gram. 2. t. 75. — Moris. s. 8. t. 4. f. 21. n. 7. (v. v.)

XXXII. *POA* *Karl. gram. p. 154. DC. fl. fr. 3. p. 56.* — *Poa* et *Airæ* sp. L.

Gluma bivalvis multiflora; spiculæ muticæ; perigonium bivalve, valvulis margine scariosis externâ carinatâ internam linearem plicatam amplexante. — Fl. paniculati.

### Sect. 1. SCHLEROCHLOA.

Spiculæ 2-5-floræ; glumæ obtusæ truncatæve flosculi breviores; glumellæ valvula interna integra; semen rostratum. — Panicula secundâ spicato-congesta.

\* *Glumellæ valvulâ externâ obtusâ.* — *Schlerochloa* P. de Beauv. t. 19. f. 4.

1. *P. DURA* (Scop. carn. 1. p. 70.) culmo compresso, foliis planis vix acutis, paniculâ rigidâ secundâ congestâ spiciformi, spiculis 3-5-floris glabris adpressis subsessilibus erecto-patulis, valvulis inæqualibus obtusis margine scariosis, glumellæ valvulis subæqualibus obtusis, internâ subciliatâ. ① in glareosis aridis Delphinatûs circâ Gratianopolim et Vapincum ex Vill. — Fl. fr. 3. p. 66. *Cynosurus durus* L. Poll. pal. t. 1. f. 1. Eleusine dura Lam. Gaud. agr. *Schlerochloa dura* P. de Beauv. Gaud. fl. helv. — Barr. ic. t. 50. (v. s. ex Valesiâ.)
2. *P. PROCUMBENS* (Smith. brit. 1. p. 98.) caule ascendente, foliis planis acutis, paniculâ rigidâ secundâ congestâ, pedunculis scabris erecto-patulis, spiculis 5-floris glabris teretiusculis subimbricatis, glumæ valvulis inæqualibus obtusis 5-nerviis, glumellæ valvulis æqualibus obtusis, internâ ciliatâ. ① in maritimis occidentali-boreslibus usque ad Armoraciâ meridionalem. — Fl. fr. 5. p. 275 eng. bot. t. 532. *Schlerochloa procumbens* P. de Beauv. (v. v. DC.)
- \*\* *Glumellæ valvulâ externâ acutâ.* — *Dactylis* et *Sesleria* sp. P. de Beauv.
3. *P. DISTICHA* (Jacq. ic. rar. 1. t. 19.) culmo 1-2-phylo, foliis radicalibus caespitosis involutis setaceis, paniculâ secundâ spiciformi compressâ ovatâ, spiculis imbricatis 4-5-floris patentibus subsessilibus distichis, glumæ valvulis subæqualibus ovatis truncatis, glumellæ valvulis æqualibus acutis basi pilosis, internâ ciliatâ.  $\frac{1}{2}$  in pratis editioribus Pyrenæorum. — Fl. fr. 3. p. 62. *Poa seslerioides* All. ped. t. 91. f. 1. non Lam. *Sesleria disticha* Pers. (v. v.)
4. *P. LITTORALIS* (Gou. monsp. 470.) culmo prostrato, foliis distichis apice involutis, paniculâ secundâ densè contractâ spiciformi, spiculis sessilibus confertis patentibus 4-8-floris teretiusculis, glumæ valvulis concavis æqualibus truncatis, glumellæ valvulis æqualibus externâ acutâ internâ truncatâ.  $\frac{1}{2}$  in maritimis Monpelii et Galloprovinciæ. — Fl. fr. 3. p. 63. Lam. ill. t. 45. f. 5. *Dactylis littoralis* Willd. P. de Beauv. (v. v.)

Sect. II. DIACHORISMOS. — *Sclerochloæ* sp. *P. de Beauv.*

Spiculæ lanceolatæ; glumæ valvula valdè inæquales, externa brevissima truncata interna subacuta; glumellæ valvulæ obtusæ. — Panícula divaricata trichotoma.

5. *P. DIVARICATA* (Gou. ill. 4. t. 2. f. 1.) foliis filiformibus, paniculâ erectâ demùm divaricatâ, pedicellis rigidis trichotomis apice incrassatis, spiculis 4-floris lanceolatis minimis, glumæ valvulis membranaceis valdè inæqualibus, glumellæ valvulis æqualibus obtusis. ① in arenosis circâ Monspelium, Arelatem, Telonem. — Fl. fr. 3. p. 65. *Sclerochloa divaricata* P. de Beauv. (v. v. DC.)

Sect. III. POASTRUM. — *Poa* *P. de Beauv.* t. 14. f. 10.

Glumæ 2-12-floræ glumellam plerumque subæquantes, perigonii valvula interna breviter bifido-dentata; semen sulcatum. — Panícula composita plus minus ramosa et effusa.

6. *P. MARITIMA* (Huds. ang. 42.) culmo ascendente, foliis angustis, ligulâ obtusâ brevi integrâ, paniculâ contractâ aut patulâ subsecundâ, spiculis 5-12-floris teretiusculis, flosculis demùm laxis, glumæ valvulis inæqualibus obtusis, glumellæ valvulis æqualibus, externâ obtusissimâ internâ ciliatâ. ② in arenosis maritimis. — Fl. fr. 3. p. 57. eng. bot. t. 1140. *P. arenaria* Retz. (v. v.)

β. *distans*, ligulâ 2-fidâ, paniculâ erectâ demùm divaricatâ, spiculis 3-6-floris. — In pratis humidis salis rariis. — *P. distans* L. mant. 32. Fl. fr. 3. p. 58. eng. bot. t. 986. *P. salina* Poll. (v. v. DC.)

7. *P. COMPRESSA* (L. sp. 101.) radice repente, culmo ascendente compresso, foliis carinato-plicatis vaginisque lævibus, ligulis brevibus obtusissimis, paniculâ coarctatâ secundâ, spiculis ovatis 3-9-floris, flosculis laxis, glumæ valvulis subæqualibus acutis, glumellæ valvulis æqualibus, externâ obtusâ basi subpubescente internâ subacutâ. ② in pratis siccioribus et muris. — Fl. fr. 3. p. 61. Leers. herb. t. 5. f. 4. — Vaill. bot. t. 18. f. 5. (v. v.)

8. *P. VULBOSA* (L. sp. 102.) culmo basi bulboso, foliis planis angustis, culmeis abbreviatis, ligulâ elongatâ acutâ, paniculâ brevi subsecundâ patentiusculâ subflexuosâ, spiculis ovatis 4-floris, glumæ valvulis subæqualibus acuminatis, glumellæ valvulis inæqualibus obtusis basi villo connexis. ② in pratis. — Fl. fr. 3. p. 61. eng. bot. t. 1071. — Vaill. bot. t. 17. f. 8. Panícula sæpè vivipara. (v. v.)

9. *P. TRIVIALIS* (L. sp. 99.) radice fibrosâ, foliis planis vaginisque scabris, ligulâ oblongâ lanceolatâ acutâ, paniculâ pyramidalî diffusâ, pedunculis semiverticillatis, spiculis 3-4-floris ovatis, glumæ valvulis acutis subæqualibus, glumellæ valvulis æqualibus internâ acutâ, externâ obtusâ basi pubescente. ② in pratis frequens. — eng. bot. t. 1072. *P. scabra* Ehr. Fl. fr. 3. p. 59. *P. dubia* Leers. herb. t. 6. f. 5. (v. v.)

10. *P. SAROTINA* (Ehr. gram. 82.) foliis planis scabris, vaginis culmoque lævibus, ligulâ oblongâ obtusâ, paniculâ pyramidalî laxiusculâ, pedunculis semiverticillatis, spiculis 3-4-floris ovatis, glumæ valvulis subæqualibus acuminatis, glumellæ valvulis æqualibus obtusiusculis, externâ basi pubescente. ② in pratis paludosis rariis. — *P. palustris* Fl. fr. 3. p. 60 et 5. p. 272. non L. (v. s.)

11. *P. PRATENSIS* (L. sp. 99.) radice repente, foliis superioribus planis plicatis ve, vaginis culmoque lævibus, ligulâ abbreviatâ truncatâ, paniculâ diffusâ, spiculis 3-4-floris ovatis, glumæ valvulis subæqualibus acutis, glumellæ valvulis æqualibus basi villo connexis, externâ acutâ, internâ subobtusâ. ② in pratis. — Host. gram. 2. t. 61.

α. *vulgaris* (Gaud.) foliis omnibus planis — *P. pratensis*. Fl. fr. 3. p. 60. eng. bot. t. 1073. (v. s.)

- β. angustifolia* (Roem. et Sch.) foliis inferioribus convolutato-setaceis. — *P. angustifolia* L. sp. 99. Fl. fr. 3. p. 60. (v. v.)
12. *P. MEMORALIS* (L. sp. 102.) foliis angustis basi plicatis, ligulis subnullis, paniculâ elongatâ attenuatâ, spiculis 2-5-floris lineari-lanceolatis lanceolatis ovatisve, glumæ valvulis subæqualibus acuminatis, glumellæ valvulis subæqualibus.  $\frac{1}{2}$  in nemoribus.
- α. vulgaris* (Gaud.) spiculis pallidis lineari-lanceolatis, glumellæ valvulis subacutis externâ basi glabriusculâ. — *P. nemoralis* Fl. fr. 3. p. 61. eng. bot. t. 1265. — Schenckz. prod. t. 2. f. 2. (v. v.)
- β. glauca* (Gaud.) spiculis ovatis 3-4-floris coloratis, glumellæ valvulis subobtusis, externâ basi villosâ. — In montibus. — *P. glauca* Fl. dan. t. 964. Fl. fr. 5. p. 273. *P. miliacea* Fl. fr. 3. p. 64. excl. syn. (v. v.)
- γ. coarctata* (Gaud.) paniculâ coarctatâ strictâ, spiculis coloratis lanceolatis 3-5-floris, glumellæ valvulis subobtusis, externâ basi villosâ. — *P. coarctata* Fl. fr. 5. p. 273. (v. v.)
13. *P. ANNUA* (L. sp. 99.) culmo obliquo compresso, foliis planis flaccidis obtusis, ligulâ oblongâ obtusâ, paniculâ brevi subsecundâ patulâ, pedunculis geminis divaricatis, spiculis 3-6-floris ovatis, flosculis laxis, glumæ valvulis subæqualibus ovatis acutis, glumellæ valvulis inæqualibus, externâ obtusiusculâ basi pubescente, internâ acutâ ciliatâ. ① in ruderalis et cultis frequens. — Fl. fr. 3. p. 59. Lam. ill. t. 45. f. 3. eng. bot. t. 1141. — Schenckz. gram. t. 3. f. 17. (v. v.)
14. *P. LAXA* (Willd. sp. 1. p. 386. non Lam.) culmo erecto compresso, foliis plicatis acutis, ligulâ lanceolatâ elongatâ acutâ, paniculâ brevi flexuosâ laxâ pauciflorâ, spiculis 3-floris ovatis, flosculis laxis, glumæ valvulis subæqualibus acuminatis, glumellæ valvulis subæqualibus, internâ angustissimâ, externâ ovatâ obtusiusculâ.  $\frac{1}{2}$  in Alpibus Galloprov. et in Pyrenæis. — Host. gram. 3. t. 15. *P. elegans* DC. fl. fr. 3. p. 62. — Schenckz. prod. t. 4. f. 2. Spiculæ ex viridi violaceo et albo variegatæ. (v. v.)
15. *P. ALPINA* (L. sp. 99.) culmo erecto, foliis planis, ligulis infimis truncatis brevissimis, superioribus lanceolatis, paniculâ brevi contractâ demum diffusâ, spiculis 4-6-floris ovato-cordatis, flosculis laxis, glumæ valvulis subæqualibus ovatis acutis, glumellæ valvulis subæqualibus, externâ ovatâ acutâ basi et margine pubescenti, externâ acuminatâ ciliatâ.  $\frac{1}{2}$  in pratis montanis et Alpibus. — Fl. fr. 3. p. 62. eng. bot. t. 1003. — Schenckz. prod. t. 3. f. 4. (v. v.)
- β. brevifolia* (Gaud.) foliis radicalibus rigidis, culmeis abbreviatis, paniculâ angustâ. — *P. brevifolia* DC. fl. fr. 3. p. 274. *P. collina* Host. gram. 2. t. 66. (v. v.)
16. *P. CERNUA* (All. auct. p. 40) radice repente, culmo ascendente compresso, foliis planis rigidis erecto-patentibus, culmorum steriliū distinctis, ligulis latis obtusis, paniculâ brevi subdiffusâ, spiculis 6-7-floris ovatis, flosculis laxis, glumæ valvulis subæqualibus acuminatis, glumellæ valvulis subæqualibus acutis, externâ basi et dorso pubescente.  $\frac{1}{2}$  in Pyrenæorum summo jugo *Pic du Midi* ex cl. Ramond (mém. du mus. 7. p. 244.). — Fl. fr. 3. p. 720 et 5. p. 274. *P. distichophylla* Gaud. (v. s. ex monte Cenisio)
17. *P. SUNDRICA* (Willd. sp. 1. p. 389.) culmo recto compresso, vaginis compressis carinatis, foliis planis lato-linearibus glaberrimis acutis, paniculâ diffusâ patulâ, spiculis 4-5-floris lanceolatis acutis, glumæ valvulis lanceolatis inæqualibus acuminatis, glumellæ valvulis æqualibus externâ subacutâ sub 5-nerviâ, internâ subtruncatâ ad nervos ciliato-scabrâ.  $\frac{1}{2}$  in sylvaticis Vogesorum, Jurassi, Alpium, montis Aursi, Pyren. — Fl. fr. 5. p. 272. *P. rubens* Willd. Fl. fr. 3. p. 59. *P. sylvatica* Vill. dauph. 2. t. 3. *Festuca compressa* DC. fl. fr. 5. p. 263. (v. v.)
18. *P. AQUATICA* (L. sp. 98.) culmo erecto tereti, foliis planis latissimis glaberrimis acuminatis, paniculâ elongatâ ampliâ diffusâ, spiculis 6-8-floris lineari-

lanceolatis, glumæ valvulis inæqualibus ovatis obtusis, glumellæ valvulis æqualibus obtusis, externâ 5-7-nerviâ.  $\mathcal{L}$  ad oras lacuum et fossarum passim. — Fl. fr. 3. p. 58. eng. bot. t. 1315. Leers. herb. t. 5. f. 5. — Scheuchz. gram. t. 4. f. 1. (v. v.)

Sect. IV. ERAGROSTIS. — *Eragrostis* et *Megastachya* P. de Beauv. t. 14. f. 11. et t. 15. f. 5.

Spiculæ elongatæ, flosculis distiche imbricatis, glumæ 4-20-floræ flosculis breviores; semen haud sulcatum. — Panicula composita plus minus effusa.

19. P. *ARGIDA* (L. sp. 101.) foliis angustis acutis, paniculâ secundâ lanceolatâ coarctatâ rigidâ, rachi marginatâ, spiculis 8-14-floris linearibus glabris rigidis, flosculis distichis laxissimis, glumæ glumellæque valvulis subæqualibus angustis acutis. ① in arenosis aridis. — Fl. fr. 3. p. 65. eng. bot. t. 1371. *Megastachya rigida* P. de Beauv. — Scheuchz. gram. t. 6. f. 2. 3. (v. v.)

20. P. *MEGASTACHYA* (Kœl. gram. 181.) foliis planis oris pilosis vaginis glabrisculis, paniculâ elongatâ patulâ, pedunculis basi glanduloso-pubescentibus, spiculis lanceolatis compressis sub-20-floris, flosculis laxis, glumæ valvulis subæqualibus ovatis acutis, glumellæ valvulis obtusis. ① in arenosis, muris et marginibus viarum passim. — Fl. fr. 3. p. 56. P. *Eragrostis* Cav. ic. t. 92 non L. *Briza Eragrostis* L. *Megastachya Eragrostis* P. de Beauv. — Scheuchz. gram. t. 4. f. 4. (v. v.)

21. P. *ERAGROSTIS* (L. sp. 100.) foliorum oris et vaginis pilosis, paniculâ patente, pedunculis basi nudis, spiculis oblongo-linearibus sub-10-floris compressis, flosculis laxè imbricatis, glumæ valvulis subæqualibus ovatis acutis, glumellæ valvulis æqualibus subobtusis, externâ 3-nerviâ. ① in incultis Parisiorum ex Merat, Andegaviæ ex Bast., Arvernæ, Ligeris (Prost), Galliæ australis. — DC. syn. p. 130. Schreb. gram. 2. t. 38. *Eragrostis pœoides* P. de Beauv. (v. s.)

22. P. *PILOSA* (L. sp. 100.) foliorum oris pilosis vaginis glabris, paniculâ elongatâ patente tenuissimâ, pedicellis flexuosis, primariis basi pilosis, spiculis linearibus compressis 7-8-floris oblongis, glumæ valvulis valdè inæqualibus subacutis, glumellæ valvulis inæqualibus obtusis. ① in arenosis Galliæ mediæ, australis occidentalisque. — Fl. fr. 5. p. 272. *Eragrostis pilosa* P. de Beauv. — Scheuchz. gram. t. 4. f. 3. (v. v.)

Sect. V. CATABROSIA P. de Beauv. t. 19. f. 8.

Glumæ 2-5-floræ obtusæ flosculis breviores; perigonii valvula externa eroso-dentata, interna subtrifida. — Paniculâ composita.

23. P. *ATROIDES* (Kœl. gram. 194.) paniculâ laxâ, spiculis 2-floris lanceolatis, glumæ valvulis membranaceis inæqualibus obtusis.  $\mathcal{L}$  in fossis aquaticis. — Fl. fr. 3. p. 64. *Aira aquatica* L. eng. bot. t. 1557. *Catabrosia aquatica* P. de Beauv. — Vaill. par. t. 17. f. 7. (v. v.)

Sect. VI. GLYCERIA R. Brown. prod. p. 179. P. de Beauv. t. 19. f. 7.

Glumæ truncatæ 7-11-floræ flosculis breviores, perigonii valvula externa crosa navicularis, interna bifido-dentata. — Panicula subsimplex elongata.

24. P. *FLUITANS* (Kœl. gram. 204.) paniculâ erectâ, spiculis 7-12-floris elongato-linearibus tereti-compressis.  $\mathcal{L}$  in fossis aquaticis. — Fl. fr. 3. p. 57. eng. bot. t. 1520. *Festuca fluitans* L. *Glyceria fluitans* R. Brown. — Moris. s. 8. t. 3. f. 4. n. 16. (v. v.)

XXXIII. *BRIZA* Kœl. gram. p. 148. DC. fl. fr. 3. p. 66. — *Briza* sp. L.

Gluma bivalvis multiflora; perigonium bivalve, valvulis ventricosis cordatis obtusis, interiore minutâ.

## 526 MONOCOTYL. SEU ENDOGENÆ.

1. *B. MAXIMA* (L. sp. 103.) spiculis 15-17-floris cordatis. ① in provinciis mediterraneis. — Fl. fr. 3. p. 66. *B. Monspessulana* Gon. — Moris. s. 8. t. 6. f. 13. n. 48. (v. v.)
2. *B. MEDIA* (L. sp. 103.) ligulâ brevissimâ obtusâ, paniculâ basi nudâ, pedunculis lævisculis, spiculis 5-7-floris ovatis. ② in pratis et sylvis frequens. — Fl. fr. 3. p. 67. eng. bot. t. 340. — Moris. s. 8. t. 6. f. 10. n. 45. Spiculæ coloratæ. (v. v.)
3. *B. MINOR* (L. sp. 101.) ligulâ longissimâ lanceolatâ, paniculâ basi folio superiore involutâ, pedunculis scabris. ① in sylvis rarior. — eng. bot. t. 1316. *B. virens* Fl. fr. 3. p. 67. an L? — Moris. s. 8. t. 6. f. 12. n. 47. Spiculæ virides. (v. v.)

† † † FLORES HERMAPHRODITI; SPICULÆ SESSILES, SECUS RACHIN SPICATÆ.

### XXXIV. CYNOSURUS *Mench. meth. DC. fl. fr. 3. p. 75. — Cynosuri sp. L.*

Bractea foliacea lacera ad basin spiculæ; gluma bivalvis, 2-5-flora; perigonium bivalve, valvulis integris.

1. *C. CRISTATUS* (L. sp. 105.) spiculis in racemum spicatum simplicem linearem digestis, bracteis pinnatifidis muticis. ② in sterilibus secus vias. — Fl. fr. 3. p. 75. Lam. ill. t. 47. f. 1. — Moris. s. 8. t. 4. f. 22. n. 6. (v. v.)
2. *C. ECHINATUS* (L. sp. 105.) spiculis in paniculam ovatam glomeratam aggregatis, bracteis pinnatis scariosis aristatis ① in incultis Gallis australis, ad littora Oceani propè St.-Malo et Cherbourg (cl. de Bréb.). — Fl. fr. 3. p. 75. Lam. ill. t. 47. f. 2. *Chrysurus echinatus* P. de Beauv. — Moris. s. 8. t. 4. f. 5. n. 13. (v. v.)
3. *C. ELÆGANS* (Desf. atl. 1. t. 17.) spiculis in paniculam ovatam secundum laxam fasciculatis, bracteis pinnatis setiformibus, glumæ valvulâ alterâ aristatâ. ① in montibus Corsicæ. — *C. gracilis* Viv. cors. 1. *Chrysurus elegans* P. de Beauv. (v. s.)

### XXXV. ECHINARIA *Desf. atl. 2. p. 385.*

Gluma bivalvis membranacea 2-3-flora, flosculis aliis hermaphroditis, aliis masculis; perigonium bivalve, valvulâ exteriorē 4-5-fidâ, laciniis rigidis longis subulatis, interiore minore 2-3-fidâ.

1. *E. CAPITATA* (Desf. f. c.) capitulo terminali subrotundo. ① in aridis Gallis australis occidentalisque. — Fl. fr. 3. p. 74. *Cenchrus capitatus* L. *Sesleria echinata* Host. gram. 3. t. 8. — Moris. s. 8. t. 5. f. 4. n. 1. (v. v.)

### XXXVI. SESLERIA *Scoep. carn. 1. p. 63. DC. fl. fr. 3. p. 75. — Cynosuri sp. L.*

Bractea nulla; gluma 2-3-flora bivalvis; perigonium bivalve, valvulâ exteriorē apice trifidâ, interiore bifidâ. — Flores capitati vel densè spicati.

1. *S. CÆRULEA* (Ard. sp. 2. t. 6. f. 3. 4. 5.) foliis subobtusis, ligulâ oblongâ obtusâ, spicâ subdensâ oblongâ. ② in rupibus et pratis montium. — Fl. fr. 3. p. 76. et 5. p. 280. Lam. ill. t. 47. f. 1. *Cynosurus cæruleus* L. (v. v.)
2. *S. CYLINDRICA* (DC. syn. p. 134.) foliis acutissimis, ligulâ nullâ, spicâ densâ cylindricâ elongatâ. ② in calidis montosis Nicæ. — Fl. fr. 3. p. 279. *Festuca argentea* Savi in Ust. ann. 1800. ic. (v. s.)



XXXVII. TRACHYNOTIA *Michaux bor. am. 1. p. 64. DC. fl. fr. 3. p. 74.*

Gluma uniflora bivalvis, valvulis angustis carinatis alterâ apice breviter aristatâ; perigonium bivalve, valvulis glumæ conformibus.

1. *T. STRICTA* (DC. fl. fr. 3. p. 74.) culmo foliisque strictis, spicis geminis erectis, spiculis secundis pubescentibus.  $\frac{1}{2}$  ad maris littora circâ Rupellam, Carnac, Noirmoutiers, Vire. — *Dactylis stricta* Ait. *Spartina stricta* Roth. *Limnætia pungens* Pers. Host. gram. 4. t. 66. (v. v.)
2. *T. ALTERNIFLORA* (DC. fl. fr. 5. p. 279.) culmo stricto, foliis planis apice involutis, spicis paniculatis coerctatis gracilibus, rachi flexuosâ, spiculis alternis distinctis.  $\frac{1}{2}$  in limosis maritimis Bayonnæ. — *Spartina alterniflora* Lois. (v. v. D.)

XXXVIII. CHAMAGROSTIS *Wib. wett. 126. DC. fl. fr. 3. p. 77.*

Gluma uniflora bivalvis, valvulis æqualibus oblongis obtusis truncatis; perigonium membranaceum pubescens minimum urceolatum margine lacerum. — Flores spicati subsecundi.

1. *C. MINIMA* (Wib. l. c.) ① in arenosis primo vere passim. — Fl. fr. l. c. *Agrostis minima* L. *Sturmia minima* Sturm germ. fasc. 7. t. 1. *Knappia agrostidea* eng. bot. t. 1127. *Mibora verna* Adams. et P. de Beauv. — J. Bauh. hist. 2. p. 469. f. 3. (v. v.)

XXXIX. NARDUS L. *gen. 69.*

Gluma uniflora 2-valvis acutissima; perigonium nullum; stigma unicum.

1. *N. STRICTA* (L. sp. 77.) spicâ setacâ rectâ secundâ, floribus approximatis, glumis breviter aristatis.  $\frac{1}{2}$  in aridis et montanis. — Fl. fr. 3. p. 77. eng. bot. t. 290. — Moris. s. 8. t. 7. f. 22. n. 8. (v. v.)
2. *N. ANISTATA* (L. syst. 145.) spicâ setacâ secundâ flexuosâ aut nutante, floribus distantibus, glumis longè aristatis. ② in aridis Delph., Monspelii, agri Ligericini (Prost). — Fl. fr. 3. p. 78. Vill. dauph. 2. t. 2. *Rotibolla monandra* Cav. ic. 1. t. 39. f. 1. *Monerma monandra* P. de Beauv. — Barr. ic. t. 117. f. 1. (v. v.)

XL. ROTTBOLLA L. *gen. supp. 114.*

Gluma mutica nunc univalvis et unicum florem hermaphroditum gerens, nunc bivalvis et flores 2 unum hermaphroditum alterum masculum includens; perigonium bivalve glumâ minus, valvulis inæqualibus.

1. *R. INCURVATA* (L. supp. 114.) foliis planiusculis, vaginâ sulcatâ, spicâ tereti subulatâ curvulâ, glumâ bivalvi uniflorâ subadpressâ subulatâ perigonium acuminatum subæquante. ① in campis Galliæ australis et maritimis prov. occid. — Fl. fr. 3. p. 78. Cav. ic. t. 213. eng. bot. t. 760. *Ophiurus incurvatus* P. de Beauv. (v. v.)
2. *R. FILIFORMIS* (Roth. catal. 1. p. 21.) foliis involutis, vaginâ lævisculâ, spicâ filiformi subulatâ subcompressâ erectâ, glumâ bivalvi acuminatâ uniflorâ patente perigonium acuminatum subæquante. ① in maritimis secus Mediterraneum, agri Syrtici circâ la Teste (Des Moul.), propè Caen ex Lenormand. — *R. erecta* Savi giorn. pis. 4. f. 5. 6. Fl. fr. 5. p. 280. *Ophiurus filiformis* Roem. et Sch. — Barr. ic. t. 6. (v. v.)
3. *R. SUBULATA* (Savi. giorn. pis. 4. f. 4. 8.) spicâ tereti strictâ subulatâ, glumâ 1-valvi ensiformi acuminatâ per anthesin patente. ① in maritimis mediterraneis propè Cette. — Fl. fr. 5. p. 281. *Monerma subulata* P. de Beauv. — Barr. ic. t. 5. (v. s.)

## 528 MONOCOTYL. SEU ENDOGENÆ.

### XLII. ÆGILOPS L. *gen.* 1150.

Gluma triflora coriacea bivalvis, valvulâ exteriorē apice 3-5 aristis rigidas gerente, flosculo intermedio masculo; perigonium bivalve, valvulâ externâ apice 3-4 aristas gerente. — Flores in rachi excavatâ semi-immersi.

1. *Æ. OVATA* (L. sp. 1489.) spicâ ovatâ aristatâ aristis breviorē, glumarum omnium valvulâ exteriorē nervosâ adpressâ ciliato-scabrâ 3-4-aristatâ. ② secus vias in provinciis meridionalibus et propè Bourg (Gironde) (Des Moul.). — Fl. fr. 3. p. 79. Lam. ill. t. 839. f. 1. — Scheuchz. gram. t. 1. f. 2. (v. v.)
2. *Æ. TRUNCIALIS* (L. sp. 1489.) spicâ oblongâ aristatâ aristarum ferè longitudine, glumarum inferiorum valvulâ exteriorē nervosâ adpressâ pubescenti biaristatâ. ♀ in sterilibus propè Parisios ex Merat. et provinc. australiorum usque ad Aginum. — Fl. fr. 3. p. 79. Lam. ill. t. 839. f. 3. — Vaill. bot. t. 17. f. 1. (v. v.)
3. *Æ. SQUARROSA* (L. sp. 1489.) spicâ oblongâ aristis longiorē, aristis inferioribus brevissimis, superioribus longioribus, glumarum valvulâ exteriorē 1-2-aristatâ sulcatâ nervosâ glabriusculâ. ♀ circa Monspelium propè Pont-Juve-  
nal reperiit Requien. — Fl. fr. 3. p. 721. Schreb. gram. t. 72. f. 2. (v. s.)

### XLIII. TRITICUM DC. *fl. fr.* 3. p. 80. — Triticum et Bromi *sp.* L.

Rachis dentata; spiculæ in quoque racheos dente splitariæ, rachi oppositæ, gluma bivalvis multiflora; perigonium 2-valve.

Sect. I. SILICO. — Triticum *P. de Beauv.* t. 20. f. 4.

Spiculæ 3-4-floræ flosculis 2 inferioribus fertilibus, supremis sæpè abortivis; glumæ valvulæ latæ naviculares dentatæ; ovarium apice barbatum. — Spica simplex densa imbricata.

1. *T. AESTIVUM* (Lam. dict. 2. p. 554.) spicâ simplici, glumis 4-floris ventricosis, flosculis omnibus hermaphroditis. ① ubique in planitiibus colitur. — Fl. fr. 3. p. 80. *T. aestivum* et *T. hybernum* L. (v. v.)
2. *T. COMPOSITUM* (L. f. supp. 115.) spicâ compositâ, glumis 3-floris ventricosis, flosculis omnibus hermaphroditis. ① colitur rarius. — Fl. fr. 3. p. 82. — Lob. ic. t. 26. f. 2. (v. v.)
3. *T. SPELTA* (L. sp. 127.) spicâ simplici, glumis 4-floris cartilagineis truncatis, floribus 2-3 hermaphroditis, 2-1 neutris. ① colitur in provinciis Helvetiæ conterminis. — Fl. fr. 3. p. 82. — Moris. s. 8. t. 6. f. 16. n. 1. (v. v.)
4. *T. MONOCOCCUM* (L. sp. 127.) spicâ simplici compressâ distichâ, glumis apice tridentatis trifloris, flore unico fertili. ① colitur in montanis aridis. — Fl. fr. 3. p. 83. — Moris. s. 8. t. 6. f. 17. n. 2. — Lob. ic. t. 31. f. 1. (v. v.)

Sect. II. AGROPYRON *Gærtn.* — Agropyron et Brachypodium *P. de Beauv.* t. 20. f. 1. et 2. et t. 19. f. 3.

Spiculæ 3-15-floræ flosculis omnibus fertilibus; glumæ valvulæ lineares aut lanceolatæ actus integræ; ovarium glabrum. — Spica plerumque composita, spiculis remotis nec imbricatis.

5. *T. CANINUM* (L. sp. ed. 1. p. 86.) radice fibrosâ, foliis planis scabris, spicâ simplici nutante densâ, spiculis 4-5-floris sessilibus adpressis glabris, glumæ valvulis subæqualibus aristato-acuminatis, glumellæ valvulis æqualibus, internâ obtusâ brevissimè ciliatâ, externâ aristatâ aristâ breviorē. ♀ in sepibus et dumosis. — eng. bot. t. 1372. *T. sepium* Lam. DC. fl. fr. 3. p. 83. *Elymus caninus* L. sp. ed. 2. Agropyrum caninum *P. de Beauv.* — Moris. s. 8. t. 1. f. 17. n. 2. (v. v.)
6. *T. REPENS* (L. sp. 128.) radice repente, foliis planis aut apice convolutis

margine scabris, spicâ elongatâ simplici strictâ densâ, rachi scabrâ, spiculis glabris 4-5-floris sessilibus compressis, glumæ valvulis æqualibus, glumellæ valvulis æqualibus, internâ obtusâ ciliatâ.  $\frac{1}{2}$  in cultis nimium frequens. — Fl. fr. 3. p. 83. Schreb. gram. t. 26. eng. bot. t. 909. *Agropyrum repens* P. de Beauv.

- a. *aristatum* (DC.) spicâ subcylindricâ, glumâ glumellæque valvulâ externâ aristatis. — Vaill. par. t. 17. f. 2. (v. v.)
- β. *muticum* (DC.) spicâ subcylindricâ, glumæ valvulis obtusis acutisve, glumellæ valvulâ externâ obtusâ submucronulatâ. (v. v.)
- γ. *pungens*, spicâ subdistichâ, glumæ valvulis subacuminatis, glumellæ valvulâ externâ acutâ. — *T. pungens* Pers. ench. i. p. 109. Fl. fr. 5. p. 283. (v. v.)
7. *T. rigidum* (Schröd. germ. i. p. 392.) radice subrepente, foliis subconvolutis, spicâ distichâ elongatâ laxiusculâ basi interruptâ, spiculis compressis 5-7-floris sessilibus adpressis, rachi scabrâ, glumæ valvulis subæqualibus retusis 5-7-nerviis, glumellæ valvulis æqualibus obtusis exaristatis, internâ tenuissimè ciliatâ, externâ obtusâ mucronulatâ.  $\frac{1}{2}$  in arenosis propè Narbonam. — Fl. fr. 5. p. 282. *Agropyrum rigidum* P. de Beauv. (v. s.)
8. *T. acutum* (DC. cat. 153.) radice repente, foliis convolutis erectis, culmo erecto stricto, spicâ distichâ continuâ, spiculis compressis 5-7-floris sessilibus subadpressis, rachi lævi, glumæ valvulis subæqualibus subacutis 5-nerviis, glumellæ valvulis æqualibus subacutis, internâ tenuissimè ciliatâ.  $\frac{1}{2}$  in arenosis limosisve maritimis secus Oceanum et Mediterraneum. — Fl. fr. 5. p. 282. (v. v.)
9. *T. junceum* (L. sp. 128.) radice repente, culmo erecto stricto, foliis convolutis longissimis erectis, spicâ elongatâ distichâ subcontinua, spiculis compressis 7-9-floris sessilibus adpressis, rachi lævi, glumæ valvulis subæqualibus obtusissimis 9-nerviis, glumellæ valvulis inæqualibus internâ obtusâ tenuissimè ciliatâ, externâ obtusâ brevissimè mucronatâ.  $\frac{1}{2}$  in arenosis secus mare Mediterraneum. — Fl. fr. 5. p. 281. *T. farctum* Viv. fragm. i. t. 26. f. 1. *Agropyrum junceum* P. de Beauv. (v. v.)
10. *T. pinnatum* (Moench. hass. n. 102.) radice repente, foliis planis pubescentibus, spicâ simplici rectâ distichâ, spiculis 9-18-floris glabris pubescentibusve brevissimè pedicellatis teretiusculis demum distichis erecto-patulis, glumæ valvulis inæqualibus acutis, glumellæ valvulis æqualibus, internâ ciliatâ obtusâ, externâ acutâ aristatâ aristâ longiore.  $\frac{1}{2}$  in sylvis, glareosis et rudibus. — Fl. fr. 5. p. 283. 3. p. 84. *T. gracile* Fl. fr. 3. p. 84. *Bromus pinnatus* L. Gaud. *Brachypodium pinnatum* P. de Beauv. — Scheuchz. gram. t. 1. f. 7. (v. v.)
11. *T. sylvaticum* (Moench. hass. n. 103.) radice fibrosâ, foliis planis villosis, spicâ simplici rectâ distichâ, spiculis 8-9-floris villosis glabrisve brevissimè pedicellatis erectis, glumæ valvulis inæqualibus aristato-acuminatis, glumellæ valvulis æqualibus internâ ciliatâ obtusâ, externâ acuminatâ aristatâ aristis supremis glumellâ longioribus.  $\frac{1}{2}$  in sylvis. — Fl. fr. 3. p. 85. et 5. p. 283. *Bromus sylvaticus* Poll. Gaud. eng. bot. t. 729. *Brachypodium sylvaticum* P. de Beauv. (v. v.)
12. *T. ciliatum* (DC. fl. fr. 3. p. 85.) culmo lævi stricto, foliis planis subpilosis ciliatis, spicâ erectâ distichâ, spiculis paucis 8-20-floris compressis rigidis sessilibus, glumæ valvulis inæqualibus acuminatis, perigonii valvulis subæqualibus, internâ obtusâ apice longè ciliatâ, externâ dorso ciliato-scabrâ acuminatâ aristatâ aristâ breviorè. ① secus vias et agros prov. Medietr. — Fl. fr. 5. p. 284. *Bromus distachyos* L. *Festuca monostachya* Desf. atl. i. t. 24. f. 2. *Brachypodium distachyon* P. de Beauv. — Ger. gallop. t. 3. f. 1. (v. v.)
13. *T. phœnicoides* (DC. fl. fr. 3. p. 85. excl. syn., 5. p. 284.) culmo erecto lævi, foliis convolutis pungentibus erectis, spicâ simplici rectâ distichâ, spiculis 10-20-floris erectis brevissimè pedicellatis, glumæ valvulis inæqualibus acu-

## 530 MONOCOTYL. seu ENDOGENÆ.

- minatis, glumellis valvulis subæqualibus, internâ obtusâ tenuissimâ ciliatâ, externâ acuminato-aristatâ aristâ multò longiore. ① in glareosis secus mare Mediterraneum, in Delphin. propè S.-Spiritus pontem. — Ger. gallop. t. 2. f. 2. (v. v.)
14. *T. cæspitosum* (DC. cat. 153. fl. fr. 5. p. 284.) radice repente, culmo basi ramosissimo, foliis convolutis glaucis tenuissimis rigidis patulis, spicâ erectâ simplici, spiculis paucis alternis 6-12-floris brevissimâ pedicellatis, glumæ valvulis inæqualibus acutis, glumellis valvulis æqualibus internâ obtusâ tenuissimâ ciliatâ, externâ aristatâ, aristâ brevî. ① in saxosis apricis provinc. austrak. — *Festuca cæspitosa* Desf. atl. 1. t. 24. *T. Placknæii* AH. — Pluck. t. 33. f. 2. (v. v.)
15. *T. Poa* (DC. fl. fr. 3. p. 86.) culmo tenello stricto, foliis angustis involutis strictis, spicâ simplici rectâ elongatâ strictâ, spiculis 6-9-floris glabris ovatis sessilibus adpressis, glumæ valvulis lanceolato-linearibus subæqualibus acutis, glumellis valvulis æqualibus obtusiusculis muticis. ① in Vogesia, circâ Lugdunum, in Galliâ mediter. et occid. passim. — *T. Halleri* Viv. frag. t. 26. f. 1. *Brachypodium Poa* P. de Beauv. — Pluck. t. 32. f. 7. (v. v. DC.)
- β. Festuca*, spicâ basi ramosâ. — Circâ Cenomannum. — *T. Festuca* DC. fl. fr. 3. p. 87. (v. a.)
16. *T. UNILATERALE* (L. mant. 35.) culmo tenello subsolitario, foliis angustis involutis vaginisque puberulis, spicâ simplici rectâ unilateralî, spiculis 4-6-floris sessilibus adpressis lineari-lanceolatis, glumæ valvulis linearibus inæqualibus acuminatis, glumellis valvulis æqualibus acuminatis exaristatis. ① in arvis propè Monspelium, Avenionem. — Fl. fr. 5. p. 285. (v. v. DC.)
17. *T. ROTUNDOLLA* (DC. fl. fr. 3. p. 86.) culmis plurimis humillimis, foliis plicatis convolutis vaginisque glaberrimis, spicâ densâ simplici erectâ, rachi flexuosâ, spiculis 5-6-floris glabris oblongis secundis sessilibus adpressis, floeculis imbricatis, glumæ valvulis lanceolato-ovatis subæqualibus obtusis, glumellis valvulis æqualibus obtusis muticis. ① in arenosis secus mare Mediter. et provine. boreali-occidentalium. — Fl. fr. 2. p. 86. et 5. p. 285. *T. unilaterale* Viv. frag. 1. t. 23. f. 1. *T. loliaceum* Smith. (v. v.)
18. *T. Nardus* (DC. fl. fr. 3. p. 87.) foliis setaceis involuto-canaliculis, spicâ simplici rectâ, spiculis 5-7-floris subglabris secundis sessilibus adpressis, glumæ valvulis linearibus inæqualibus acuminatis, glumellis valvulis æqualibus acuminatis, externâ sæpius aristatâ. ① in sterilibus Parisiorum, prov. occident. rarior et australi. — *Brachypodium tenellum* et *tenuiflorum* Rostk. et Schmidt. (v. v.)
19. *T. TRIVICULUM* (Lols. not. 27.) culmo simplici recto apice nudo, foliis angustis involutis vaginisque glabris, spicâ simplici rectâ strictâ, spiculis sessilibus adpressis 7-9-floris ovato-oblongis glaberrimis, glumæ valvulis subæqualibus lineari-lanceolatis acutis, glumellis valvulis æqualibus, internâ obtusiusculâ, externâ acutâ aristatâ aristâ subduplò longiore. ① in agris circâ Mennotæ, Andegavum. — Fl. fr. 5. p. 285. *T. Hispanicum* Viv. fragm. t. 23. f. 2. *T. festuoides* Bert. (v. s.)

### XLIII. SECALE *L. gen. 97.*

Rachis dentata; spiculis in quoque racheos dente solitariæ, gluma 2-et-rarius-3-flora, bivalvis; perigonium bivalve, valvulâ exteriorē apice aristatâ.

1. *S. CEREALE* (L. sp. 124.) glumis setaceis, perigonis scabris. ① colitur ubique. — Fl. fr. 3. p. 88. Lam. ill. t. 49. (v. v.)
2. *S. VILLOSUM* (L. sp. 124.) glumis truncatis aristatis, perigonis villosis. ① in agri, Lugdani ex Latour, Mounpelli, Biteris, in Corsicâ. — Fl. fr. 3. p. 88. (v. s.)

XLIV. LOLIUM *L. gen.* 95.

Rachis subdentata; spiculæ solitariae rachi parallelæ; gluma multiflora 2-valvis, valvulâ externâ magnâ, internâ minimâ sæpè abortivâ; perigonium 2-valve, valvulâ internâ ciliato-scabrâ.

1. *L. PERENNIS* (*L. sp.* 122.) culmo lævi, spiculis 6-12-floris subcompressis muticis glumâ longioribus.  $\frac{1}{2}$  secus vias et in pascuis sterilibus. — *Fl. fr.* 3. p. 89. *Lam. ill.* t. 48. (v. v.)
2. *L. TENUE* (*L. sp.* 122.) culmo gracili lævi, foliis angustis, spiculis 3-4-floris subcylindricis muticis glumam subæquantibus.  $\frac{1}{2}$  secus vias et prata sicca. — *Fl. fr.* 3. p. 89. — An ut voluit cl. Schrader et Gaudin prioris var. ? (v. v.)
3. *L. TEMULENTUM* (*L. sp.* 122.) culmo scabro, spiculis 5-9-floris compressis aristatis glumam subæquantibus. ① inter segetes. — *Fl. fr.* 3. p. 89. *Lam. ill.* t. 48. f. 2. (v. v.)
4. *L. MULTIFLORUM* (*Lam. fl. fr.* 3. p. 621.) culmo sublævi, spiculis 20-25-floris compressis, superioribus aristatis, glumâ duplò longioribus. — In agris et pratis passim. — *Fl. fr.* 3. p. 90. Variat muticum. (v. v.)

XLV. ELYMUS *L. gen.* 96.

Rachis dentata; spiculæ 2-3 in quoque racheos dente; gluma 2-4-flora bivalvis, valvulis sæpè patulis et involucrum mentientibus; flosculi superiores interdum masculi; perigonium bivalve.

1. *E. ARNARIUS* (*L. sp.* 122.) spiculis geminis, spicâ erectâ arctâ, glumis pubescentibus flosculo mutico longioribus.  $\frac{1}{2}$  in arenosis maritimis. — *Fl. fr.* 3. p. 90. — *Gm. sib.* 1. t. 25. (v. s.)
2. *E. EUROPEUS* (*L. mant.* 35.) spiculis ternis, spicâ erectâ, glumis scabris setaceis, perigonio aristato.  $\frac{1}{2}$  in pratis et secus vias. — *Fl. fr.* 3. p. gr. *Cuviera Europæa* Koel. (v. v.)

XLVI. HORDEUM *L. gen.* 98.

Rachis dentata; spiculæ in quoque racheos dente ternæ, laterales sæpius masculæ pedicellatæ, media sessilis hermaphrodita; gluma uniflora bivalvis, valvulis sæpè patulis et involucrum mentientibus.

1. *H. VULGARE* (*L. sp.* 125.) flosculis omnibus hermaphroditis aristatis, ordinibus indistinctis. ① Colitur autumnò satum. — *Fl. fr.* 3. p. 90. — *J. Bauh. hist.* 2. p. 418. f. 3. (v. v.)
2. *H. HEXASTICHON* (*L. sp.* 125.) flosculis omnibus hermaphroditis aristatis, spicâ crassâ sexfariam sulcatâ. ① Colitur cum priori. — *Fl. fr.* 3. p. 92. *Host. gram.* 3. t. 35. (v. v.)
3. *H. DISTICHUM* (*L. sp.* 125.) flosculis lateralibus masculis muticis, seminibus imbricatis, spicâ elongatâ compressâ. ① Colitur verè satum. — *Fl. fr.* 3. p. 92. — *C. Bauh. theat.* 440. ic. (v. v.)
4. *H. ZEACARTON* (*L. sp.* 125.) flosculis lateralibus masculis muticis, seminibus patentibus, spicis basi crassioribus subpyramidatis. ① Colitur rariùs. — *Fl. fr.* 3. p. 93. *Schreb. gram.* t. 17. — *J. Bauh. hist.* 2. p. 430. f. 1. (v. v.)
5. *H. MURINUM* (*L. sp.* 126.) flosculis lateralibus masculis aristatis, intermediis glumis setaceis ciliatis. ① in ruderatis. — *Fl. fr.* 3. p. 93. *eng. bot.* t. 1971. — *Moris. a. 8. t. 6. f. 25. n. 4.* (v. v.)
6. *H. SECALINUM* (*Schreb. spic.* 148.) flosculis lateralibus masculis aristatis, glumis omnibus setaceis scabris. ① in pratis siccis et sterilibus. — *Fl. fr.* 3. p. 94. — *Vaill. par. t. 17. f. 6.* (v. v.)
7. *H. MARITIMUM* (*Vahl. symb.* 2. p. 25.) flosculis lateralibus masculis aristatis

## 532 MONOCOTYL. SEU ENDOGENÆ.

dorso pubescentibus, glumis scabris lateralibus lanceolatis.  $\frac{1}{2}$  in arenosis maritimis à Niceâ usque ad Abbatis-villam sed rarius. — Fl. fr. 3. p. 94. H. geniculatum All. ped. t. 91. f. 3. (v. v.)

8. *H. castrum* (Desf. atl. 1. p. 113.) flosculis geminis fertilibus aequalibus longè aristatis, flosculorum superiorum aristis involucri 4-phyllo aristato longioribus. ① in arenosis secus vias Fori-Julii. — H. jubatam Lois. DC. fl. fr. 5. p. 286. non *L. Elymus crinitus* Schreb. gram. 2. t. 24. (v. a.)

### Ordo CXI ? LEMNACEÆ *Duby mss.* — *Lemna Ach. Rich. dict. class. 9. p. 276.*

Flores monoici in spathâ sessili monophyllâ compressâ tenui membranaceâ primò inclusi. *Fl. masc.* pauci (1-2). Perigonium 0. Stamen 1; filamentum cylindricum pistillo longius; antheræ 2 juxtapositæ globulosæ 1-loculares longitudinaliter dehiscentes. *Flos femineus* unicus. Pistillum 1. Ovarium ovoideum compressum 1-loculare, ovulis 2-5-erectis. Stylus crassus teres. Stigma concavum truncatum. Capsula indehiscens 1-∞-sperma. Semina raphe notata; integumentum crassum. Embryo 1-cotyledoneus. Albumen 0. — Herbæ minimæ annuæ natantes, folio aut fronde compressâ lenticulari constantes. Flores et radiculae lateraliter nascentes. — Hujus ordinis in Monocotyledoneis phanerogamis locus incertus et idcirco ad hujus classis finem rejectus. Lemnam Pistiæ affinem vult cl. Hooker scot. 2. p. 191. et inter Aroideas inserit et revera Aroideæ videntur ordo quocum majores affinitates habent Lemnaceæ.

#### I. LEMNA *L. gen. n. 1038.*

Character isdem ac ordinis.

1. *L. trisulca* (L. sp. 1376.) foliis lanceolatis in petiolum attenuatis cruciatim coherentibus, radicibus solitariis. ① in aquis pigris puris. — Fl. fr. 2. p. 588. eng. bot. t. 926. — Mich. gen. t. 11. f. 5. (v. v.)
2. *L. minor* (L. sp. 1376.) foliis ellipticis utrinque planis basi coherentibus, radicibus solitariis. ① ad superficiem aquarum quietarum. — Fl. fr. 2. p. 389. eng. bot. t. 1093. — Vaill. bot. t. 20. f. 3. — Mich. gen. t. 11. f. 3. (v. v.)
3. *L. gibba* (L. sp. 1377.) foliis ellipticis subtus bullato-convexis basi coherentibus, radicibus solitariis. ① ad superficiem aquæ quietæ. — Fl. fr. 2. p. 589. eng. bot. t. 1233. — Mich. gen. t. 11. f. 2. (v. v.)
4. *L. polyrrhiza* (L. sp. 1377.) foliis ellipticis planis basi coherentibus, radicibus fasciculatis. ① ad superficiem aquarum quietarum. — Fl. fr. 2. p. 590. — Vaill. bot. t. 20. f. 2. — Mich. gen. t. 11. f. 1. (v. v.)
5. *L. aruzza* (L. mant. 294.) foliis elliptico-subrotundis geminis subtus convexis eradicatis. ① ad superficiem aquarum quietarum. — Fl. fr. 2. p. 590. bull. phil. n. 78. f. 2-8. — Mich. gen. t. 11. f. 4. (v. v.)

Classis Tertia. MONOCOTYLEDONEÆ seu  
ENDOGENÆ CRYPTOGRAMMÆ DC. or-  
gan. 1. p. 128.

Truncus medullâ centrali, radiis medullaribus corticeque verâ plane destitutus, fibris sparsis nec concentricè dispositis constans, apice tantum crescens, in margine magis quam in centro induratus, interdum abortivus subterraneus radiciformis. Folia sæpè vaginata integra nervis simplicibus, aut lobata nervis ramosis, nunquàm verè composita. Flores indistincti. Embryo cotyledone unico cellulis tantum nec tracheis constante. (DC. mss.)

Ordo CXII? CHARACEÆ L. C. Rich. Ad. Brongn. in dict.  
class. 3. p. 474.

Flores axillares perigonio destituti, monoici. Masc? Disci solitarii orbiculares rubri, annulo albo cincti, extus membranâ reticulatâ translucidâ formati, intus mucilaginosi filamenta articulata aut tubos cylindricos foventes. Fem. Capsula 1-locularis 1-sperma, integumento duplici, externo membranaceo tenuissimo translucido supernè 5 dentibus rosaceis terminato, interno duro sicco opaco-valvis 5 angustis spiraliter contortis. — Plantæ aquaticæ submersæ, caulibus ramosis debilibus duris fragilibus, ramis verticillatis in verticillis superioribus 3-4-5 capsulas remotas basi 2-3 bracteis (ramis abortivis?) stipatas gerentibus. — Ordo ex cl. Leman Onagrariis et Salicariis, ex cl. Agardh Bory-St.-Vincent etc. Algis affinis, hic propter ramorum dispositionem in vicinio Equisetacearum collocatus.

I. CHARA Vaill. acad. 1719. p. 23. L. gen. n. 1203.

Character isdem ac ordinis.

1. C. VULGARIS (L. sp. 1624.) caulibus striatis inermibus, ramulis basi nudis, fructibus subquaternis spiraliter striatis bracteam superantibus. ♀ in aquis pigris. — Fl. fr. 2. p. 584. et 5. p. 246. Lam. ill. t. 742. f. 1. — Vaill. acad. 1719. t. 3. f. 1. (v. v.)
2. C. TOMENTOSA (L. sp. 1624.) caulibus sulcatis apice subhispidis, ramulis basi foliosis, fructibus solitariis spiraliter striatis bractea brevioribus. ♂ in fossis. — Fl. fr. 2. p. 585. — Moris. s. 15. t. 4. f. 13. n. 9. (v. v.)

## 534 MONOCOTYL. SEU ENDOGENÆ.

3. *C. HISPIDA* (L. sp. 1624.) caulibus sulcatis hispidis, ramulis basi subfoliosis, fructibus solitariis spiraliter striatis bractea brevioribus. ① in fossis. — Fl. fr. 2. p. 585. Lam. ill. t. 376. f. 3. — Vaill. acad. 1719. t. 3. f. 3. (v. v.)
4. *C. GALLOIDES* (DC. cat. 93 et fl. fr. 5. p. 246.) caulibus tenuibus non sulcatis apice setis raris hispidis, fructibus solitariis lævibus bracteis ternis circumdatis et iisdem subbrevioribus. ① in fossis aquæ subdulcis Occitanie maritimæ, in Neustrie flumine Orne ex Lenormand. (v. v. DC.)
5. *C. CAPILLACRA* (Thunb. par. II. 1. p. 474.) caulibus lævibus semipellucidis, ramis verticillatis elongatis, fructibus solitariis ovoideis sublævibus bractea subbrevioribus. ① in aquis pigris. — Fl. fr. 2. p. 585. (v. v.)
6. *C. FLEXILIS* (L. sp. 1524.) caulibus lævibus semi-pellucidis, ramis subverticillatis elongatis, fructibus aggregatis ovoideis sublævibus bractea longioribus. ① in aquis pigris. — Fl. fr. 2. p. 586. *C. translucens* Pers. eng. bot. t. 1855. *Nitella flexilis* et *translucens* Agardh syst. alg. 124. — Vaill. acad. 1719. t. 3. f. 8. g. (v. v.)
7. *C. CANESCENS* (Lois. not. 139.) caulibus tenuissimis dichotomis albidis copiose hispidis, spinulis subulatis subverticillatis, internodiis supremis approximatis, ramulis fructiferis brevissimis verticillatis; bracteis fructu duplò longioribus. ① in aquis dulcibus stagnosis secus littora maris Mediterranei propè Telonem, Monapegium. — Fl. fr. 5. p. 246. *C. erinita* Wallr. t. 3. Ag. l. c. (v. v. DC.)
8. *C. BATRACHOSPERMA* (Thunb. par. II. 1. p. 473.) caulibus lævibus semipellucidis, ramis approximatis verticillatis, fructibus subquaternis ovoideis spiraliter striatis bractea brevioribus. ① in aquis pigris. — Fl. fr. 2. p. 586. *Nitella batrachosperma* Agardh alg. 126. (v. v.)
9. *C. HYALINA* (DC. fl. fr. 5. p. 247.) caulibus lævibus lucidis hyalinis, ramis verticillatis confertis in 7-8 ramulos divisis, ramulis trifidis in axillâ fructum ovatum breviter pedicellatum reflexum gerentibus. ① in aquis quietis circà Cænomanum, Nannetes. — *C. tenuissima* Desv? *Nitella hyalina* Ag. l. c. (v. v.)
10. *C. SYNCARPA* (Thunb. par. II. 1. p. 473.) caulibus lævibus semipellucidis, ramis verticillatis, fructibus subternis bractea destitutis. ① in aquis pigris. — Fl. fr. 2. p. 586. (v. v.)

### Ordo CXIII. EQUISETACEÆ Richard. DC. fl. fr. 2. p. 580. Vauch. monog. in mem. soc. gen. 31.

Fructificationes terminales dispositæ in amentum conicum squamis peltatis intus floriferis constans. Involucella bivalvia. Semina sphaerica (acotyledonea ex Vauch.) numerosa nuda, filamentis quatuor polliniferis hygrometricis apice dilatatis circumplicata. — Plantæ aphyllæ ramis verticillatis sulcatis articulatis, articulis vagina monophyllâ cinctis.

#### I. EQUISETUM L. gen. 1169. Vauch. l. c.

Character isdem ac ordinis.

1. *E. ARVENSE* (L. sp. 1516.) caule sterili subscabro, striis circiter duodenis, vaginæ dentibus totidem, ramis scabriusculis tetragonis, caule fructifero evanido, involucri laxi, vaginæ dentibus 12. ½ in glareosis humidis frequens. — Fl. fr. 2. p. 581. Vauch. mon. t. 1. (v. v.)



2. *E. FLUVIATILE* (L. sp. 1517.) caule sterili cylindrico glabro, dentibus et ramis circiter triginta, caule fructifero nudo evanido, vaginis amplis elongatis laxis, dentibus 30.  $\mathcal{U}$  in palustribus et ripis fluviorum. — Fl. fr. 2. p. 582. Vauch. mon. t. 2. *E. Telmateya* Ehr. crypt. 31. Fl. fr. 2. p. 581. *E. eburneum* Schreb. (v. v.)
3. *E. SYLVATICUM* (L. sp. 1516.) caule sterili fistuloso striato, dentibus et ramis circiter 12, ramis decompositis arcuato-deflexis, caule fructifero subnudo vaginis laxis, ramis paucioribus depauperatis.  $\mathcal{U}$  in montosis argillosis humidis. — Fl. fr. 2. p. 583. Vauch. mon. t. 3. (v. v.)
4. *E. UMBROSUM* (Willd. sp. 5. p. 3.) caule sterili fistuloso, dentibus et ramis circiter duodecim, ramis simplicibus arcuato-deflexis triquetris, caule fructifero nudiusculo, vaginis laxis, ramis depauperatis.  $\mathcal{U}$  in umbrosis Pyrenæorum ex Lapeyr. — Vauch. t. 4. (v. s. ex Germ.)
5. *E. PALUSTRE* (L. sp. 1516.) caule sulcato subscabro, soleis dentibusque ferè octonis, ramis totidem quadrangularibus sæpè abortivis.  $\mathcal{U}$  in pratis et humidis. — Fl. fr. 2. p. 583. *E. tuberosum* Fl. fr. 5. p. 245. Vauch. mon. t. 5. (v. v.)
  - $\beta$ . *polystachyon* (Ray ang. ed. 3. t. 5. f. 3.) ramis elongatis fructiferis. — Vauch. mon. t. 5. f. 2. (v. v.)
  - $\gamma$ . *nudum*, vaginis dilatatis. (v. v.)
6. *E. LIMOSUM* (L. sp. 1517. non herb.) caule glabro fistuloso, striis dentibusque ferè 14, ramis simplicibus sæpius abortivis.  $\mathcal{U}$  in stagnis et fossis. — Fl. fr. 2. p. 582. Ray. t. 5. f. 2. Vauch. mon. t. 7. (v. v.)
7. *E. HYEMALE* (L. sp. 1517.) caule subnudo scabro fistuloso, striis et vaginæ dentibus ferè octodecim, dentibus pilosis aut evanidis.  $\mathcal{U}$  in sylvis humidis. — Fl. fr. 2. p. 580. Vauch. monog. t. 9. Vaginæ basi apiceque nigrescentes. (v. v.)
8. *E. MULTIFORME* (Vauch. mon. t. 12.) caulibus numerosis striatis glabrisculis sulcatis apice spiciferis, ramis paucioribus irregulariter dispositis, vaginis plus minùsve sphacelatis, dentibus octonis.  $\mathcal{U}$  in siccis et arenosis.
  - $\alpha$ . *variegatum* (Vauch.) vaginis variegatis. — *E. variegatum* Schl. in DC. fl. fr. 5. p. 244. — Tabern. ic. 251. f. 1. (v. v.)
  - $\beta$ . *ramosum* (Vauch.) vaginis amplis laxis, ramis numerosioribus. — *E. ramosum* Schl. in DC. fl. fr. 5. p. 245. (v. v.)
  - $\gamma$ . *campanulatum* (Vauch.) vaginis campanulatis, caulibus cinereis profundè sulcatis. — *E. campanulatum* Poir. (v. v.)

## 536 MONOCOTYL. seu ENDOGENÆ.

Ordo CXIV. FILICES *Brown. prod. 145. DC. fl. fr. 3. p. 546.*  
— *Filicum gen. Juss.*

Fructificationes gregariæ, juniores sæpius indusio tectæ, in facie inferiore frondis sitæ, in paucis spuriis terminales spicatae. *Masc?* Antherae? minimæ sparsæ in foliis vix evolutis præsentæ, membranâ tenui tectæ. *Fem.* Capsulæ uniloculares, sæpius annulo articulado elasticè dissiliente cinctæ, rariùs bivalves, seminulis numerosis minutissimis repletæ. — Frondes (folia) alternæ, pseudo-radicales, sæpè lobato-decompositæ, juniores ab apice ad basin circinnato-convolutæ.

Trib. I. *OPHIOGLOSSÆ Hook. fl. scot. 2. p. 158.*

Capsulæ 1-loculares, basi adnatæ, subglobosæ, coriaceæ, opacæ, annulo elastico destitutæ, semibivalves.

I. *OPHIOGLOSSUM Sw. journ. Schrad. 2. p. 112. DC. fl. fr. 2. p. 570.* — *Ophioglossi sp. L.*

Capsulæ subglobosæ sessiles uniloculares transversè dehiscentes, in spicam subarticulatam disticham connatæ.

1. *O. VULGATUM* (L. sp. 1518.) fronde ovatâ.  $\mathcal{U}$  in pascuis humidis. — *Fl. fr. 2. p. 570. eng. bot. t. 108. Lam. ill. t. 364. f. 1. Barr. ic. t. 252. f. 1. (v. v.)*
2. *O. LUSTANICUM* (L. sp. 1518.) fronde lineari-lanceolatâ.  $\mathcal{U}$  in arenosis maritimis Armoraciæ circa Brivatem et Legionem, Bardigalæ (cl. Bory), Corsicæ. — *Fl. fr. 5. p. 243. Lam. ill. t. 364. f. 3. — Barr. ic. t. 252. f. 2. Præcedenti triplo quadruplè minus. (v. s.)*

II. *BOTRYCHIUM Sw. journ. Schrad. 2. p. 110. DC. fl. fr. 2. p. 569.*

Capsulæ subglobosæ distinctæ sessiles uniloculares, ab apice ad basin dehiscentes, in spicam racemosam congestæ.

1. *B. LUNARIA* (Sw. l. c.) fronde solitariâ pinnatâ, pinnis lanatis.  $\mathcal{U}$  in pascuis montanis. — *Fl. fr. 2. p. 569. Osmunda Lunaria L. eng. bot. t. 318. — Barr. ic. t. 252. f. 3. (v. v.)*
2. *B. MATRICARIOIDES* (W. sp. 4. p. 62.) frondibus geminis subradicalibus petiolatis tripartitis, partitionibus subpinnatis, segmentis obtusis subdentatis subrotundis.  $\mathcal{U}$  in sylvis Vogesorum ex Schanenb. et Pyrenæorum ex Lapeyr. — *Fl. fr. 5. p. 243. — Moris. s. 14. t. 5. f. g. n. 26. (v. s. ex Siberiâ.)*

Trib. II. *OSMUNDACÆ Hook. l. c.*

Capsulæ annulo destitutæ, vasculoso-reticulatæ pellucidæ apice radiatæ aut substriatæ, longitudinaliter dehiscentes.

III. *OSMUNDA Lam. fl. fr. 1. p. 10. DC. fl. fr. 2. p. 569.*

Capsulæ confertæ subglobosæ pedicellatæ uniloculares semibivalves in racemum seu in dorsum frondis dispositæ.

1. *O. REGALIS* (L. sp. 1521.) frondibus bipinnatis, omnibus fructiferis, racemo

terminali suprâ-decomposito.  $\mathcal{U}$  in sylvis humidis — Fl. fr. 2. p. 569. Lam. ill. t. 865. f. 2. eng. bot. t. 209. (v. v.)

Trib. III. *POLYPODIACEÆ* Hook. scot. p. 152.

Capsulæ monoculares annulo elastico longitudinali cinctæ, transversæ et irregulariter dehiscentes.

IV. CETERACH *C. Bauh. pin. 354. DC. fl. fr. 2. p. 566.*

Capsulæ sparsæ aut variè aggregatæ, squamis scarioso-fuscis membranaceis aut filiformibus obtectæ.

1. *C. OFFICINARUM* (C. Bauh. l. c.) fronde pinnatifidâ, laciniiis alternis confluentibus obtusis, subtus densè squamosis.  $\mathcal{U}$  ad rupes et muros. — Fl. fr. l. c. *Asplenium Ceterach* L. *Grammitis Ceterach* Willd. — Moris. s. 14. t. 2. f. 13. (v. v.)
2. *C. MARANTÆ* (DC. fl. fr. 2. p. 567.) fronde subbipinnatâ, pinnulis oppositocoadunatis basi subdentatis subtus densè squamosis.  $\mathcal{U}$  ad rupes volcanicas circâ Thueis (Ardèche) detexit cl. Prost, propè Tournon cl. Req. — Fl. fr. 5. p. 243. *Acrostichum Marantæ* L. — Lob. ic. t. 816. (v. s.)

V. GRAMMITIS *Sv. syn. fil. p. 3.*

Capsulæ in acervos oblongos sublineares rectis sparsæ; indusio nullo.

1. *G. LEPTOPHYLLA* (Sw. l. c. p. 3. et 218. t. 1. f. 6.) frondibus pinnatis bipinnatisque glaberrimis, pinnulis cuneiformibus rotundo-lobatis.  $\mathcal{U}$  in maritimis Corsicæ, Galloprovinciæ, Armoraciæ circâ Brivatem et Legionem. — *Acrostichum leptophyllum* DC. fl. fr. 2. p. 565. — Barr. ic. t. 431. (v. s.)

VI. POLYPODIUM *Adans. fam. 2. p. 26. DC. fl. fr. 2. p. 564. — Polypodii sp. L.*

Capsulæ in puncta rotunda sparsa aggregatæ; indusio nullo.

1. *P. VULGARE* (L. sp. 154.) fronde profundè pinnatifidâ, lobis oblongis crenulatis obtusis approximatis, caudice squammato.  $\mathcal{U}$  in muris, tectis et arborum truncis. — Fl. fr. 2. p. 564. eng. bot. t. 1149. *P. cambricum* L. — Moris. s. 14. t. 2. f. 6. n. 8. (v. v.)  
*β. serratum* (Willd.) lobis lineari-lanceolatis serratis acutis. — Fl. fr. 5. p. 242. — Barr. ic. t. 38. In agro Syrtico propè S.-Sever. (cl. Dufour), circâ Vire ex cl. Lenormand. (v. s.)
2. *P. PHÆGOTERIS* (L. sp. 1550.) fronde pinnatâ pinnis pinnatifidis basi coadunatis, infimis reflexis, segmentis lineari-lanceolatis obtusis integris ciliatis.  $\mathcal{U}$  in montosis humidis Vogesorum, Arvernæ, Cebennarum (cl. Prost), ad rupes madidas sylvarum Neustriæ ex cl. Lenormand. — Fl. fr. 2. p. 564. eng. bot. t. 2224. — Moris. s. 14. t. 4. f. 3. n. 17. (v. s.)
3. *P. RHÆTICUM* (L. sp. 1552.) fronde bipinnatâ, pinnulis oblongis obtusis pinnatifidis, lobis dentatis, soris orbiculatis.  $\mathcal{U}$  in pascuis sylvaticis Alpium Delphinatûs ex Vill., in summis præruptis herbidis Vogesorum (cl. Moug. et Nestl.) — Fl. fr. 5. p. 242. — J. Bauh. hist. 3. p. 740. f. 1. (v. s.)
4. *P. DAYOTTERIS* (L. sp. 1555.) fronde ternatâ bipinnatâ patulo-deflexâ, laciniiis obtusis subcrenatis, radice filiformi.  $\mathcal{U}$  in umbrosis sylvarum imprimis in montanis. — Fl. fr. 2. p. 565. eng. bot. t. 616. (v. v.)
5. *P. CALCARUM* (Sm. brit. 3. p. 1117.) fronde ternatâ bipinnatâ erectâ rigidulâ, laciniiis obtusis subcrenatis, maculis fructiferis confluentibus.  $\mathcal{U}$  in ericetis montosis calcaris Jurassi et Pyrenæorum. — Fl. fr. 5. p. 243. — Clus. hist. 2. p. 212. f. 1. (v. v.)

## 538 MONOCOTYL. seu ENDOGENÆ.

VII. WOODSIA R. Br. in trans. linn. t. xi. p. 170.

Capsulæ pedicellatæ in soros subrotundos aggregatæ; indusium calyciforme apertum margine crinitum, receptaculo communi elevato nullo.

1. W. HYPERBOREA (Br. l. c. t. 11.) fronde pinnatâ, pinnis tripartitis vel pinnatifido-incisæ cuneiformibus obtusis subtus hirtis, soris solitariis demum confluentibus.  $\frac{1}{2}$  in Alpinis et Pyrenæis variis. — Polypodium hyperboreum Willd. eng. bot. t. 2023. Fl. fr. 5. p. 242. Ceterach alpinum Fl. fr. 2. p. 566. — Meris. s. 14. t. 3. f. ult. n. 23<sup>p</sup> (v. s.)

VIII. POLYSTICHUM Roth. germ. 3. p. 76. DC. fl. fr. 2. p. 559. — Hypopeltis Rich. — Polypodii sp. L.

Capsulæ in puncta subrotunda sparsa dispositæ, indusio umbilicato nempe centro adfixo et margine toto deliscente tectæ.

1. P. OREOPTERIS (DC. fl. fr. 2. p. 563.) fronde pinnatâ pinnis lanceolatis glabris resinoso-glandulosis inferioribus pinnatifidis, segmentis lanceolatis obtusis integris, punctis fructiferis marginalibus subconfluentibus, stipite nudo.  $\frac{1}{2}$  in sylvis montanis et ericetis. — Polyp. oreopteris eng. bot. t. 1019. (v. v.)
2. P. THELYPTERIS (Roth. germ. 3. p. 77.) fronde pinnatâ, pinnis pinnatifidis glabris basi distinctis decussantibus, segmentis ovatis acutis integris, punctis fructiferis marginalibus confluentibus, stipite nudo.  $\frac{1}{2}$  in sylvis subpaludosis. — Fl. fr. 2. p. 563. Polyp. Thelypteris Hedw. th. retr. t. 6. eng. bot. t. 509. (v. v.)
3. P. CALLIPTERIS (DC. fl. fr. 2. p. 562.) fronde subbipinnatâ, pinnis cordato-oblongis, inferioribus pinnatis, superioribus pinnatifidis, pinnulis ovatis obtusis denticulato-serrulatis, stipite paleaceo.  $\frac{1}{2}$  in paludosis Abbeville (cl. Boucher), Parisiorum ex Loise., Jurassi. — Polypod. cristatum L. sp. 1551. Afz. act. holm. 1787. t. 9. Aspidium cristatum eng. bot. t. 2125. (v. s.)
4. P. TANAETIFOLIUM (DC. fl. fr. 2. p. 562.) fronde bipinnatâ, pinnulis oblongis pinnatifidis, laciniis linearibus apice serratis, stipite subpaleaceo.  $\frac{1}{2}$  in Vogesis, montibus Arvernæ, Corsicæ. — Polypod. tanaetifolium Hoff. (v. s.)
5. P. DILATATUM (DC. fl. fr. 5. p. 241.) fronde bipinnatâ, pinnulis oblongis distinctis inciso-pinnatifidis, segmentis mucronato-serratis, stipite paleaceo.  $\frac{1}{2}$  in sylvis montanis. — P. spinulosum Fl. fr. 2. p. 561. excl. syn. Sw. Aspidium dilatatum eng. bot. t. 1461. — Mapp. alb. 106. f. 8. (v. v.)
6. P. FILIX-MAS (DC. fl. fr. 2. p. 559.) fronde bipinnatâ, pinnulis oblongis crenatis obtusis apice serratis basi duobus punctis fructificationis utrinque donatis, rachi stipiteque paleaceis.  $\frac{1}{2}$  in sylvaticis et sterilibus. — Polyp. filix-mas L. sp. 1551. Aspidium filix-mas eng. bot. t. 1458. (v. v.)  
 $\beta$ . abbreviatum, pinnulis ovalibus obtusissimis crenatis, basi unico fructificationis puncto utrinque donatis. — In sylvaticis agri Syrtici. — P. abbreviatum DC. fl. fr. 2. p. 560. (v. s.)
7. P. RIGIDUM (DC. fl. fr. 2. p. 560.) fronde pinnatâ ovato-lanceolatâ, pinnis bipinnatifidis cordato-lanceolatis, laciniis oblongis obtusis apice dentatis, stipite paleaceo.  $\frac{1}{2}$  in sylvaticis petrosis Alpium et rariis Pyrenæorum ex Lapeyr. — Polypodium fragrans Vill. excl. syn. (v. s.)
8. P. ACULEATUM (Roth. germ. 3. p. 77.) fronde bipinnatâ, pinnulis rigidis ovatis sublunatis acuminatis basi decurrentibus margine subserratis spinulosis, rachi stipiteque paleaceis.  $\frac{1}{2}$  in sepibus sylvisque montanis. — Fl. fr. 2. p. 561. Aspidium aculeatum eng. bot. t. 1562. — Pluck. t. 180. f. 2. (v. v.)
9. P. PRUCKETHII (DC. fl. fr. 5. p. 241.) fronde pinnatâ, pinnis pinnatifidis basi subbipinnatis, lobis ovato-lunulatis ciliato-serratis, inferiore anticæ maximo,

- stipite paleaceo.  $\mathcal{L}$  in sepiibus et umbrosis. — *Polypodium Plucknetii* Lois. not. 146. — Pluckn. t. 180. f. 3. (v. s.)
10. *P. Lonchitis* (Roth. germ. 3. p. 77.) fronde pinnatâ, pinnis lanceolato-falcatis acutis ciliato-serratis basi autice acutè auriculatis, stipite paleaceo.  $\mathcal{L}$  in sylvis montanis. — Fl. fr. 2. p. 560. *Aspidium Lonchitis* eng. bot. t. 796. — Moris. s. 14. t. 2. f. 12. n. 1. An 2. præcedentes hujus tantum varietates? (v. v.)

IX. *ASPIDIUM* DC. fl. fr. 2. p. 557. — *Aspidii* sp. Sw.

Capsulæ in puncta subrotunda sparsa dispositæ, indusio ab apice ad basin utrinque dehiscente acutissimo tectæ.

1. *A. FLEBILE* (Sw. journ. Schrad. 2. p. 40.) pinnis bipinnatifidis oppositis, pinnulis ovatis obtusis incisis, lacinulis acutis serrato-dentatis.  $\mathcal{L}$  in sylvaticis montanis et fissuris rupium. — Fl. fr. 2. p. 558. excl. var.  $\beta$  et  $\delta$  et 5. p. 240. *Cyathea fragilis* eng. bot. t. 1587. — Pluckn. t. 79. f. 3 et 180. f. 5. Admodum variabile. (v. v.)
2. *A. AZEUM* (Sw. syn. 58.) fronde bipinnatâ, pinnulis obovato-oblongis lobato-pinnatifidis, lacinulis linearibus oblongis obtusis subintegerrimis.  $\mathcal{L}$  in rupibus Alpium et Pyrenæorum. — Fl. fr. 5. p. 240. *Cyathea regia* Sm. eng. bot. t. 163. — Vill. danph. 3. t. 53. f. C. An præcedentis var.? (v. s.)
3. *A. ALPINUM* (Sw. syn. 60.) frondibus triplicato-pinnatis, pinnulis pinnatifidis confluentibus, lacinulis linearibus obtusis apice bidentatis.  $\mathcal{L}$  in rupibus editorum Alpium et Pyrenæorum. — Fl. fr. 5. p. 241. *Polypodium Alpinum* Jacq. ic. rar. 3. t. 642. — Seg. veron. suppl. t. 1. f. 3. (v. v.)
4. *A. MONTANUM* (Sw. journ. Schrad. 2. p. 42.) suprâ-decompositum, foliolis ternis tripinnatifidis, lacinulis subfalcatis obtusis apice dentatis.  $\mathcal{L}$  in sylvaticis montanis. — Fl. fr. 2. p. 558. *Polypodium myrrhidifolium* Vill. danph. 4. t. 53. — Pluckn. t. 89. f. 4. (v. v.)

X. *ATHYRIUM* DC. fl. fr. 2. p. 556. — *Athyrii* sp. Roth.

1. *A. FILIX-FEMINA* (Roth. germ. 3. p. 68.) fronde bipinnatâ, pinnulis oblongo-lanceolatis inciso-serratis, serraturis 2-3-dentatisve acutis, soris oblongis angustis.  $\mathcal{L}$  in sylvaticis et montanis. — Fl. fr. 2. p. 556. *Aspidium Filix-femina* eng. bot. t. 1459. — Pluckn. t. 181. f. 2. (v. v.)

XI. *ASPLENIUM* Smith. bris. 3. p. 1126. DC. fl. fr. 2. p. 553. — *Asplenii* sp. L.

Capsulæ in lineolas rectas transversas sparsas dispositæ, indusio laterali interiùs dehiscente tectæ.

1. *A. HALLERI* (DC. fl. fr. 5. p. 240.) frondibus bipinnatis rigidis, pinnulis alternis cuneiformi-triangularibus apice mucronato-dentatis obtusis.  $\mathcal{L}$  in empibus calcareis Jurassi, Alpium, Pyrenæorum. — *Aspidium Halleri* Willd. 4. p. 274. *Athyrium fontanum* Fl. fr. 2. p. 557. — Barr. ic. t. 432. f. 1. (v. v.)
2. *A. LANCEOLATUM* (Sm. eng. bot. 240.) frondibus bipinnatis, pinnis ovato-lanceolatis, pinnulis obovatis argutè dentatis.  $\mathcal{L}$  Ad fissuras humidæ rupium Fontisbleaudi, Neustrim propè Vire et Faleise ex cl. Lenormand et de Bréb., Nigri-Monasterii, Burdigalæ (Des Moul.), Cebennarum. — Fl. fr. 5. p. 239. — Dod. pempt. 265. ic. — Vaill. bot. t. 9. f. 1. (v. s.)
3. *A. ADIANTHUM-NIGRUM* (L. sp. 1542.) frondibus subtripinnatis, pinnulis ovato-lanceolatis inciso-serratis, soris demùm confluentibus.  $\mathcal{L}$  in locis umbrosis et nemoribus humidis. — Fl. fr. 2. p. 556. eng. bot. t. 1950. (v. v.)
4. *A. RUTA-MURARIA* (L. sp. 1541.) frondibus alternatim decompositis, pinnulis cuneato-rhomboides subtrilobato-crenulatis.  $\mathcal{L}$  in muris et rupibus umbrosis frequens. — Fl. fr. 2. p. 555. eng. bot. t. 150. (v. v.)
5. *A. MARINUM* (L. sp. 1540.) fronde pinnatâ, pinnis oblongis obtusis inciso-

## 540 MONOCOTYL. SEU ENDOGENÆ.

- serratis, superioribus basi rotundis et subauriculatis.  $\mathcal{U}$  in maritimis Galliz occidentalis passim, insulis Stœckadum ex Ger. — Fl. fr. 2. p. 555. eng. bot. t. 392. — Moris. s. 14. t. 3. f. 4. n. 25. (v. s.)
6. A. *viride* (Huds. ang. 453.) frondibus pinnatis, stipitibus basi fascis, pinnis rhombeo-subrotundis obtusè serratis.  $\mathcal{U}$  ad fissuras rupium Vogesorum, Jurassi, Alpium, Pyrenæorum. — Fl. fr. 2. p. 554. eng. bot. t. 2257. (v. v.)
7. A. *fontanum* (DC. fl. fr. 5. p. 239.) frondibus pinnatis, stipitibus viridibus glabris basi fascis, pinnis inferioribus trilobis crenatis, superioribus subrotundis.  $\mathcal{U}$  ad rupes Pyrenæorum orientaliom. — Polypodium fontanum Sm. brit. 3. p. 1114. — Pluck. t. 89. f. 2. — Moris. s. 14. t. 3. f. 2. n. 11. An prioris var. (v. s.)
8. A. *trichomanes* (L. sp. 1540.) frondibus pinnatis, stipitibus nigrescentibus, pinnis rotundo-oblongis obtusè crenatis basi truncato-cuneatis.  $\mathcal{U}$  in muris et rupibus umbrosis humidis frequens. — Fl. fr. 2. p. 554. eng. bot. t. 576. — Moris. s. 14. t. 3. f. 1. n. 10. (v. v.)
- $\beta$ . *Petrarchæ*, stipitibus pubescentibus. — A. Petrarchæ Fl. fr. 5. p. 238. A. glandulosum Lois. not. 145. (v. s.)
9. A. *germanicum* (Weiss. goett. p. 299.) frondibus alternatim subdecompositis, pinnulis subcuneiformibus apice incis.  $\mathcal{U}$  in muris et rupibus submontanis Alsatiæ, Vogesorum, Jurassi ex Hall. agri Ligericini (cl. Prost). — Fl. fr. 2. p. 553. A. alternifolium Jacq. misc. 2. t. 5. f. 2. eng. bot. t. 2258. — Breyn. cent. t. 97. (v. s.)
10. A. *septentrionale* (Hoffm. germ. 2. p. 12.) stipitibus nudis glabris superne trifidis, segmentis linearibus apice laciniatis tridentatisve.  $\mathcal{U}$  in fissuris rupium. — Fl. fr. 2. p. 553. Acrostichum septentrionale L. eng. bot. t. 1017. (v. v.)
- XII. *SCOLOPENDRIUM* Smith. act. taur. 5. p. 410. DC. fl. fr. 2. p. 551. — Asplenii sp. L.

Capsulæ in lineolas sparsas lineares geminas intervenias dispositæ, indusii 2 superficialibus suturâ longitudinali dehiscentibus tectæ.

1. S. *officinale* (Sm. act. taur. 5. p. 410.) frondibus cordato-lingulatis.  $\mathcal{U}$  in humidis umbrosis. — Fl. fr. 2. p. 552. eng. bot. t. 1150. A. Scolopendrium L. — Moris. s. 14. t. 1. f. 1. (v. v.)
2. S. *agittatum* (DC. fl. fr. 5. p. 238.) frondibus hastatis basi cordatis, auriculis mediocribus obtusis.  $\mathcal{U}$  in rupibus maritimis Massiliæ, Corsicæ (cl. Soleirol). — S. Hemionitis Fl. fr. 2. p. 552. non L. — Clus. hist. 2. p. 214. f. 1. — Matth. comm. 646. f. 2. (v. v.)

XIII. *BLECHNUM* Smith. act. taurin. 5. p. 411. DC. fl. fr. 2. p. 551. — Osmundæ sp. L.

Capsulæ in lineas longitudinales solitarias continuas costæ frondis parallelas dispositæ, indusio superficiali continuo interius dehiscente tectæ.

1. B. *spicant* (Sm. l. c.) frondibus stérilibus pinnatifidis, laciniis lanceolatis obtusiusculis parallelis, fructiferis pinnatis, pinnis linearibus acuminatis.  $\mathcal{U}$  in nemoribus montanis. — Fl. fr. 2. p. 551. B. boreale Sw. eng. bot. t. 1159. Osmunda spicant L. (v. v.)

XIV. *PTERIS* Sm. act. taur. 5. p. 412. DC. fl. fr. 2. p. 549. — Pteridis et Acrostichi sp. L.

Capsulæ in lineam continuam marginalem dispositæ, indusio è margine frondis inflexo interius dehiscente tectæ.

1. P. *crispa* (All. ped. n. 2352.) frondibus stérilibus bipinnatis, pinnulis pin-

natifidis, segmentis obovato-crenatis apice inciso-dentatis, frondibus fertilibus inferne bipinnatis tripinnatisve, pinnulis linearibus oblongis subobtusis integris basi angustatis.  $\mathcal{U}$  in pratis saxosis montium. — Fl. fr. 2. p. 550. eng. bot. t. 116. *Osmunda crispa* L. — Moris. s. 14. t. 4. f. 11. n. 4. (v. v.)

2. *P. AQUILINA* (L. sp. 1533.) fronde supra-decomposita, pinnulis lineari-lanceolatis, superioribus indivisis, inferioribus pinnatifidis, segmentis oblongis obtusis.  $\mathcal{U}$  in campestribus et sylvaticis sterilibus. — Fl. fr. 2. p. 550. Hook. scot. 2. p. 156. eng. bot. t. 1679. (v. v.)
3. *P. CARTICA* (L. mant. 130.) fronde pinnata, pinnis oppositis lanceolatis elongatis basi angustatis, infimis tripartitis.  $\mathcal{U}$  in Corsica et circa Niceam. — Fl. fr. 2. p. 549. — Tourn. inst. t. 321. (v. v. DC.)

XV. ADIANTHUM L. gen. n. 1180.

Capsulæ in puncta discreta seu lineolas ad marginem frondis distributæ, indusii è margine ortis replicatis interius dehiscentibus tectæ.

1. *A. OBOVATUM* (DC. fl. fr. 5. p. 437.) frondibus bipinnatis, pinnulis ovatis obtusis sublobatis subhis nudis, stipite palaceo.  $\mathcal{U}$  ad muros et rupes humidæ Corsicæ, Galloprovinciæ orientalis, Pyrenæorum orientalium. — A. fragrans Fl. fr. 2. p. 549. Viv. frag. 9. t. 11. f. 2. *Pteris acrosticha* Balb. *Polypodium fragrans* Desf. atl. 2. t. 257. non L. *Cheilanthes odora* Sw. (v. s.)
2. *A. CAPILLUS-VENERIS* (L. sp. 1558.) frondibus decompositis, pinnulis cuneatis apice rotundatis lobatis, stipite lævi.  $\mathcal{U}$  ad fontes umbrosos Jurassi et provinc. australium. — Fl. fr. 2. p. 548. — Tourn. inst. t. 317. (v. v.)

XVI. HYMENOPHYLLUM Sm. act. taur. 5. p. 118. DC. fl. fr. 2. p. 547. — *Trichomanis* sp. L.

Capsulæ marginales sessiles receptaculo communi cylindrico insertæ, intra indusia bivalvia.

1. *H. TURNERIANUM* (Sm. l. c.) frondibus subbipinnatis ovatis decurrentibus pinnulisque serratis, fructificationibus supra-axillaribus solitariis.  $\mathcal{U}$  inter muscos, ad truncos in maritimis Armoraciæ, propè Cherbourg, Mortain (Manche) et Bèfort (cl. Lejeune). — Fl. fr. l. c. eng. bot. t. 162. — Pluck. t. 3. f. 5. 6. (v. s.)

## 542 MONOCOTYL. SEU ENDOGENÆ.

Ordo CXV. MARSILEACEÆ *Brown. prod.* 166. — Rhizospermæ *Roth. DC. fl. fr.* 3. p. 577.

Fructificationes radicales; involucrium subsphæricum non dehiscens, coriaceum aut membranaceum, uni-aut multiloculare, organa utriusque sexûs, deindè semina plura fovens. — Herbæ aquaticæ.

Trib. I. SALVINIÆ *Ad. Brongn. in dict. class. t.* 10. p. 196.

Fructificationes membranacæ bifformes; aliæ (semina) ovoideæ integumento tenui reticulato in aquâ non intumescenti; aliæ (involucra? flores masculi?) multos granulos sphæricos columnæ centrali longis pedicellis adheps, seminibus multò minoribus, reticulatis, in aquâ non disrumpes, foventes. — Folia opposita (in nostris), non convoluta.

I. SALVINIA *Mich. gen. t.* 58. *Juss. gen. p.* 16.

Involucra 4-9 aggregata ovoidea membranacea 1-locularia; semina embryone unico.

1. *S. NATANS* (Hoffm. *genu. 2. p.* 1.) foliis ovato-subcordatis obtusis subtus pilis fasciculatis scabris. — Natans in aquis quietis provinciarum australium. *Burdigalæ ex Laterr.* ad ripas Rheni (cl. Moog. et Nestl.). — *Fl. fr.* 2. p. 579. *Lam. ill. t.* 863. *Marsilea natans* L. — *Guett. acad.* 1762. t. 29. f. 1. (v. v.)

Trib. II. MARSILÆÆ. — Marsileacæ veræ *Ad. Brongn. l. c.*

Involucra coriacea crassa indehiscencia aut multivalvia intus dissepimentis membranaceis multilocularia. In quoque locali adsunt organa biformia parietibus inserta, alia pauciora (ovaria aut potius semina) integumento duplici, externo pellucido humiditate tumescenti, interno duro coriaceo, semine in substantiâ gelatinosâ nidulante et omninò libero; alia numerosiora utriculi membranacei humiditate tumescentes apice dehiscences et in muco gelatinoso globulos sphæricos numerosos seminibus multò minores foventes. — Folia juniora filicum more apice convoluta.

II. MARSILEA *Juss. gen. p.* 16. — *Marsilea sp. L.*

Involucra 2-3 pedicello communi insidentia, ovoidea, coriacea, intus dissepimentis tenuissimis multilocularia.

1. *M. QUADRIFOLIA* (L. *sp.* 1563.) foliis longè petiolatis quadrifoliolatis, foliolis margine integris. — In aquis natans passim. — *Fl. fr.* 2. p. 578. *Lam. ill. t.* 863. — *Mapp. als.* 166. ic. (v. v.)

III. PILULARIA *L. gen. n.* 1183.

Involucra solitaria subsessilia globosa coriacea quadrilocularia.

1. *P. GLOBULIFERA* (L. *sp.* 1563.) foliis filiformibus.  $\frac{1}{2}$  in inundatis ad littora paludum passim. — *Fl. fr.* 2. p. 577. *Lam. ill. t.* 862. *Vaill. bot. t.* 15. f. 6. — *Juss. acad.* 1739. t. 11. (v. v.)



Ordo CXVI. LYCOPODIACEÆ Richard. DC. fl. fr. 2. p. 571.

Fructificationes crustaceæ sessiles, ad axillas nunc foliorum et tunc axillares, nunc bractearum et tunc spicatæ dictæ. Capsulæ nunc uniformes seminula multa foventes; nunc bifformes; aliæ vulgatiores (forsan masculæ) globulis sphaericis pulveraceis repletæ; aliæ rariores (verosimiliter femineæ) includentes semina sphaerica subscabra subtus costis tribus prominulis notata.

Trib. I. ISOETIDÆ.

Capsulæ indehiscentes.

I. ISOETES L. gen. n. 1184.

Fructificationes subradicales, basi frondis immersæ; cocculi oblonga non dehiscentia polysperma. — Folia radicalia.

1. I. LACUSTRIS (L. sp. 1563.) foliis subulatis semiteretibus. ♀ in fundo lacuum Armoracæ, Aquitanicæ, Monspeliæ. — Fl. fr. 2. p. 576. Lam. ill. t. 862. eng. bot. t. 1084. — Ray ang. 1. p. 210. ic. (v. v.)

Trib. II. LYCOPODIÆ.

Capsulæ dehiscentes.

II. LYCOPODIUM L. gen. 1185.

Dioicum aut monoicum. Masc. Cocculi bivalvia pulvere plena. Fem. Cocculi quadrivalvia 1-4-sperma. — Herbarum habitu sicco vario, foliis caulinis numerosis imbricatis aut bifariam dispositis.

Sect. 1. EULYCOPIDIUM — Lycopodium Ad. Brongn. in dict. class. t. 9. p. 559.

Cocculi omnia bivalvia semina? rotunda sphaerica aut ovalia libera foventia.

1. L. ALPINUM (L. sp. 1567.) foliis quadrifariam imbricatis acutis carinatis, floralibus ovato-lanceolatis planis, ramulis fastigiatis erectis. ♀ in sylvis montanis. — Fl. fr. 2. p. 572. Fl. lapp. t. 11. f. 6. — Dill. musc. t. 58. f. 2. (v. v.)
2. L. COMPLANATUM (L. sp. 1567.) foliis quadrifariis basi connatis, duobus majoribus bifariam dispositis, duobus minimis adpressis, apicibus geminis pedunculatis. ♀ in sylvis Vogesorum et circa Parisios. — Fl. fr. 2. p. 572. — Dill. musc. t. 59. f. 3. (v. s.)
3. L. ANNOTINUM (L. sp. 1566.) foliis sparsis subserratis muticis quinquefariis, floralibus dilatatis abbreviatis, ramis floriferis annotino-articulatis. ♀ in Alpibus Delphinatibus ex Vill. — L. juniperifolium Lam. DC. fl. fr. 2. p. 572. — Dill. musc. t. 63. f. 9. Moris. s. 15. t. 5. f. 2. n. 3. (v. v. in Alp. helv.)
4. L. CLAVATUM (L. sp. 1564.) foliis sparsis apice filamentosis, floralibus dilatato-membranaceis, ramis floriferis inferne strigosis apice bitrifidisve. ♀ in ericetis et sylvis montanis. — Fl. fr. 2. p. 573. — Dill. musc. t. 58. f. 1. Moris. s. 15. t. 5. f. 1. n. 2. (v. v.)
5. L. SELAGO (L. sp. 1565.) foliis sparsis integerrimis lanceolatis muticis uniformibus octofariis, caule dichotomo erecto fastigiato. ♀ in ericetis montosis

544 MONOCOTYL. SEU ENDOGENÆ. LYCOPODIUM.

humidiusculis. — Fl. fr. 2. p. 573. — Dill. musc. t. 56. f. 1. — Moris. s. 15. t. 5. f. 7. n. 9. (v. v.)

6. *L. MUNDATUM* (L. sp. 1565.) foliis sparsis integerrimis lineari-lanceolatis muticis, floralibus basi dilatatis patentiusculis.  $\mathcal{U}$  in ericetis inundatis passim. — Fl. fr. 2. p. 574. — Dill. musc. t. 61. f. 7. — Vaill. bot. t. 16. f. 1. (v. s.)

Sect. II. *STACHYGYNANDRUM* P. de Beauv. æth. prod. p. 105. Ad. Brongn. l. c.

Coccula biformia 2-valvia (masc?) granulos tenuissimos liberos foventia; alia ovalia 4-valvia 3-4 semina 3-quetra placentæ centrali adnexa? foventia.

7. *L. SELAGINOIDES* (L. sp. 1565.) foliis sparsis ciliatis lanceolatis, floralibus majoribus patentibus.  $\mathcal{U}$  in ericetis et pascuis humidis Pyrenæorum, Alpium, Jurassi. — Fl. fr. 2. p. 574. — Dill. musc. t. 68. f. 1. — Hall. helv. t. 46. f. 1. (v. v.)

8. *L. HELVETICUM* (L. sp. 1568.) foliis ovalibus subacutis quadrifariis, duobus majoribus patulis, duobus minimis adpressis, caule radicante.  $\mathcal{U}$  in alpi- bus ad radices arborum et in rupibus umbrosis. — Fl. fr. 2. p. 575. — Dill. musc. t. 64. f. 2. Moris. s. 15. t. 6. f. 3. n. 34. (v. v.)

9. *L. DENTICULATUM* (L. sp. 1568.) foliis ovalibus acutè mucronatis irregulariter quadrifariis, duobus majoribus patulis, duobus minimis adpressis, caule radicante. (♂) ad pedes arborum Galloprovinciæ, Monspeli. — Fl. fr. 2. p. 575. — Dill. musc. t. 66. f. 1. A. (v. s.)



1

1

1

